



**HAL**  
open science

# Stannylenes and Germylenes Stabilized by Tetradentate Bis(amidine) Ligands with a Rigid Naphthalene Backbone

Alejandra Acuña, Sonia Mallet-Ladeira, Jean-Marc Sotiropoulos, Eddy Maerten, Alan R Cabrera, Antoine Baceiredo, Tsuyoshi Kato, René S Rojas, David Madec

► **To cite this version:**

Alejandra Acuña, Sonia Mallet-Ladeira, Jean-Marc Sotiropoulos, Eddy Maerten, Alan R Cabrera, et al.. Stannylenes and Germylenes Stabilized by Tetradentate Bis(amidine) Ligands with a Rigid Naphthalene Backbone. *Molecules*, 2024, 29 (2), pp.325. 10.3390/molecules29020325 . hal-04437950

**HAL Id: hal-04437950**

**<https://hal.science/hal-04437950v1>**

Submitted on 5 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Article

# Stannylenes and Germylenes Stabilized by Tetradentate Bis(amidine) Ligands with a Rigid Naphthalene Backbone †

Alejandra Acuña <sup>1,2</sup>, Sonia Mallet-Ladeira <sup>3</sup> , Jean-Marc Sotiropoulos <sup>4</sup> , Eddy Maerten <sup>1</sup> , Alan R. Cabrera <sup>2</sup> , Antoine Baceiredo <sup>1</sup> , Tsuyoshi Kato <sup>1</sup> , René S. Rojas <sup>2,\*</sup>  and David Madec <sup>1,\*</sup> 

<sup>1</sup> Laboratoire Hétérochimie Fondamentale et Appliquée (UMR 5069), Université de Toulouse, CNRS, 118 Route de Narbonne, 31062 Toulouse, Cedex 09, France; azacuna@uc.cl (A.A.); eddy.maerten@univ-tlse3.fr (E.M.); jose-antoine.baceiredo@univ-tlse3.fr (A.B.); tsuyoshi.kato@univ-tlse3.fr (T.K.)

<sup>2</sup> Departamento de Química Inorgánica, Facultad de Química y de Farmacia, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, Santiago 7820436, Chile; arcabrer@uc.cl

<sup>3</sup> Institut de Chimie de Toulouse (UAR 2599), 118 Route de Narbonne, 31062 Toulouse, Cedex 09, France; sonia.ladeira@univ-tlse3.fr

<sup>4</sup> IPREM, UMR 5254, UPPA/CNRS Technopole Helioparc-2, av. Pdt P. Angot, 64053 Pau, Cedex 09, France; jean-marc.sotiro@univ-pau.fr

\* Correspondence: rrojasg@uc.cl (R.S.R.); david.madec@univ-tlse3.fr (D.M.)

† This paper is dedicated to the memory of Dr. Jean-Pierre Férézou.

**Abstract:** An unusual series of germylenes and stannylenes stabilized by new tetradentate bis(amidine) ligands RNC(R')N-linker-NC(R')NR with a rigid naphthalene backbone has been prepared by protonolysis reaction of Lappert's metallylenes [M(HMDS)<sub>2</sub>] (M = Ge or Sn). Germylenes and stannylenes were fully characterized by NMR spectroscopy and X-ray diffraction analysis. DFT calculations have been performed to clarify the structural and electronic properties associated with tetradentate bis(amidine) ligands. Stannylene **L<sub>1</sub>Sn** shows reactivity through oxidation, oxidative addition, and transmetalation reactions, affording the corresponding gallium and aluminum derivatives.

**Keywords:** amidine; ligand; metallylene



**Citation:** Acuña, A.; Mallet-Ladeira, S.; Sotiropoulos, J.-M.; Maerten, E.; Cabrera, A.R.; Baceiredo, A.; Kato, T.; Rojas, R.S.; Madec, D. Stannylenes and Germylenes Stabilized by Tetradentate Bis(amidine) Ligands with a Rigid Naphthalene Backbone. *Molecules* **2024**, *29*, 325. <https://doi.org/10.3390/molecules29020325>

Academic Editor: Vassilis Tangoulis

Received: 18 December 2023

Revised: 3 January 2024

Accepted: 4 January 2024

Published: 9 January 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


## 1. Introduction

Amidines have been employed as ligands for numerous metallic and semi-metallic elements [1], and one of their best features is the ability to adjust both steric and electronic properties, leading to many substitution patterns of the CN<sub>2</sub> framework [2–4]. In recent years, the trend has shifted to develop bulkier systems to stabilize a broader range of metal centers or to control the complex geometry via rigid coordination, as seen with bridged bis(amidine) ligands.

Amidinate ligands have shown a remarkable ability to stabilize low-valent main group elements, such as tetrylenes, forming planar NCN-Ge<sup>(II)</sup> and NCN-Sn<sup>(II)</sup> four-membered rings [5]. The first bis(amidinato)germylene, reported by Richeson et al. in 1997, exhibited one chelating and one dangling amidinate ligand, with the three-coordinate germanium center adopting a trigonal-pyramidal geometry (Figure 1) [6]. Using the same synthetic method, Karsch et al. reported in 1998 a bis(amidinato)germylene with both amidinate ligands acting as chelates due to the less hindered substituents on the nitrogen atoms [7]. On the other hand, the first mono- and bis-(amidinato)stannylenes were reported by Tolman et al. in 2002, using a *N*-silylated benzamidinate ligand (Figure 1) [8].

More recently, the first tetrylenes with 1,4-phenylene (**I**) and 1,4-cyclohexylene (**II**) bridged bis(amidine) ligands were reported by Jones et al. in 2020 [9]. In 2021, the same group reported a bis-germylene and a bis-stannylene with a dibenzofurandiyl-linked Dipp-substituted bis(amidine) ligand (**III**) [10]. In the same year, Kretschmer et al. reported bis-germylenes and bis-stannylenes using a terphenyl-linked bis(amidine) ligand (**IV**) [11].

It is important to note that all these examples correspond to amidine fragments connected through the carbon atom, and only one bis-germylene with a bis(amidine) ligand nitrogen connected via a 1,3-phenylene bridge has been synthesized by Jones et al. (V) [10]. The main characteristic of bridged bis(amidine) ligands exhibited is that the amidine functionalities act as independent ligands, allowing the convenient preparation of stable bis-metallylenes. Moreover, there is no report of a bis(amidine) system acting as a tetradentate ligand with both amidine moieties coordinated to the same metal center. We therefore considered the hypothesis that such a bis(amidine) system could influence the stabilization and the reactivity of the corresponding tetrylenes.



**Figure 1.** Examples of germylenes and stannylenes with bis(amidine) ligands [6–11].


Herein, we report the synthesis and structural characterization of novel stannylene and germylene compounds stabilized by naphthalene-linked bis(amidine) ligands, with the amidine functionalities connected to the naphthalene bridge through the nitrogen atoms. In addition, the reactivity of synthesized stannylenes was explored in oxidation, oxidative addition, and transmetalation reactions.

## 2. Results and Discussion

### 2.1. Synthesis of Bis(amidine) Ligands with a Rigid Naphthalene Backbone


Bis(amidine) ligands  $L_1H_2$  and  $L_2H_2$ , with a rigid planar 1,8-diaminonaphthalene linker, were prepared following the reported procedures [12,13], while the synthesis of  $L_3H_2$  was carried out through the reaction between one equivalent of *N,N'*-(naphthalene-1,8-diyl)bis(2,2-dimethylpropanimidoyl chloride) and two equivalents of *p*-tolylamine in dry toluene under reflux for 4 h in the presence of Et<sub>3</sub>N (Scheme 1).  $L_3H_2$  was isolated as yellow crystals in 63% yield and characterized by NMR spectroscopy. The <sup>1</sup>H NMR

spectrum shows two NH signals at 9.43 and 8.50 ppm, similar to those previously reported bis(amidine) ligands [13,14]. In addition, CH<sub>3</sub> resonances of *p*-tolyl and *t*butyl fragments appear as singlets at 2.05 and 2.01 ppm and at 1.38 and 1.23 ppm, respectively. The <sup>13</sup>C NMR spectrum also exhibits two different C=N groups at 161.3 and 160.7 ppm, confirming the non-symmetrical character of L<sub>3</sub>H<sub>2</sub> ligand in solution due to NH···N hydrogen bonds.



**Scheme 1.** Synthesis of ligand L<sub>3</sub>H<sub>2</sub>.

The presence of an intramolecular hydrogen bond between the N-H group and the nitrogen atom of the amidine moiety [N1-(H1)···N(3) 1.95(5) Å] is confirmed in the solid-state structure of ligand L<sub>3</sub>H<sub>2</sub> (Figure 2), in line with similar observations for ligands L<sub>1</sub>H<sub>2</sub> and L<sub>2</sub>H<sub>2</sub> [12,13].




**Figure 2.** Molecular structure of L<sub>3</sub>H<sub>2</sub>. Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms (except H1 and H4A) have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: N(1)-C(11) 1.421(5); N(2)-C(11) 1.272(5); N(3)-C(23) 1.286(5); N(4)-C(23) 1.373(5); N1-(H1)···N(3) 1.95(5); N(1)-C(11)-N(2) 122.5(3); N(3)-C(23)-N(4) 126.1(3).

## 2.2. Synthesis of Stannylenes L<sub>1-3</sub>Sn

The synthesis of stannylenes L<sub>1-3</sub>Sn was carried out using protonolysis reactions due to the simplicity of the procedure [15–17]. Therefore, the corresponding bis(amidine) ligand L<sub>1-3</sub>H<sub>2</sub> reacted with one equivalent of Sn(HMDS)<sub>2</sub> in dry THF at 60 °C for 3 h (Scheme 2). Stannylene L<sub>1</sub>Sn was isolated as colorless crystals from a pentane solution at –30 °C in 77% yield. The formation of L<sub>1</sub>Sn was confirmed by <sup>1</sup>H, <sup>13</sup>C, and <sup>119</sup>Sn NMR spectroscopy and mass spectrometry. The <sup>1</sup>H NMR spectrum shows a singlet at 1.97 ppm with an integration of 6 H, corresponding to the two amidine methyl groups, and a singlet at 2.18 ppm with an integration of 12 H, corresponding to the methyl groups of 2,6-dimethylphenyl fragments. This observed NMR pattern agrees with a symmetrical structure around the tin center. The <sup>13</sup>C NMR spectrum shows a characteristic signal for the NCN fragment at 169.0 ppm, similar to those (amidinato)stannylenes previously reported [8,15,16,18]. In the <sup>119</sup>Sn NMR spectrum, a resonance at –276.7 ppm confirms the tetracoordinated nature of the tin atom, comparable to that of reported homoleptic four-coordinated bis(amidinate) tin(II) [15]. L<sub>3</sub>H<sub>2</sub>

also reacts with  $\text{Sn}(\text{HMDS})_2$  in the same manner to give the corresponding tetracoordinated stannylene  $\text{L}_3\text{Sn}$ , as shown by the NMR analysis. Indeed, the  $^1\text{H}$  NMR spectrum shows a singlet at 1.30 ppm that integrates for 18 H, corresponding to the amidine *t*-butyl groups, and a singlet at 2.29 ppm that integrates for 6 H, corresponding to the *p*-tolyl methyl groups. The  $^{13}\text{C}$  NMR spectrum shows a characteristic signal for the NCN fragment at 177.5 ppm. The symmetrical tetracoordination of the tin center is also confirmed by a resonance at  $-254.9$  ppm in the  $^{119}\text{Sn}$  NMR spectrum. In contrast, the use of the bulky bis(amidinate) ligand  $\text{L}_2\text{H}_2$  leads to the formation of a dimeric structure  $(\text{L}_2\text{Sn})_2$ .



**Scheme 2.** Synthesis of stannylene  $\text{L}_{1-3}\text{Sn}$ .


Stannylene  $\text{L}_1\text{Sn}$  was characterized in the solid state by single-crystal X-ray diffraction analysis (Figure 3). The molecular structure indicates a four-coordinate tin center with a distorted square-based pyramidal geometry. A few stannylenes with unbridged bis(amidinate) ligands display a similar geometry [16]. The acute N1–Sn–N2 and N3–Sn–N4 angles of  $57.38^\circ$  and  $57.74^\circ$  are comparable to values previously recorded for  $\text{Sn}^{\text{II}}$  amidinate complexes [15,18].

The coordinates of the experimental solid-state structure are consistent with the calculated values in DFT (see Supplementary Materials for more information). Indeed, the modeling calculates values between 2.27 and 2.42 Å for the Sn–N bonds, probably due to steric repulsion, and angles at the tin atom of  $56^\circ$ . Moreover, the energy of molecular orbitals of this species (computed using DFT) shows an electrophilic character (the first vacant orbital on the Sn atom is computed at  $-0.60$  eV) and a potential nucleophilicity, as the HOMO centered on the non-bonding pair of tin is rather accessible ( $-5.42$  eV) (Figure 4).


The modeling demonstrates that this compound can be obtained from the bis(amidinate) ligand through the addition of  $\text{Sn}(\text{HMDS})_2$ . Indeed, starting from the resulting complex, a prototropic shift can occur, leading to the facile formation of the corresponding  $\text{RSnH}$  (TS = 21.63 kcal/mol) (see Supplementary Materials for more information). Subsequently, the elimination of HMDS results in the obtaining of  $\text{L}_1\text{Sn}$ , exhibiting a higher but consistent kinetic rate with the applied experimental conditions ( $60^\circ\text{C}/3$  h).

Because of the poor solubility of stannylene dimer  $(\text{L}_2\text{Sn})_2$  in common organic solvents, the NMR characterization cannot be obtained once crystallized. Nevertheless, the X-ray diffraction analysis shows an eight-membered cyclic structure with a Sn···Sn distance of 3.7311(5) Å (Figure 5). Each Sn center has a trigonal pyramidal geometry with a sum of angles around the Sn atom of  $272^\circ$ . Interestingly, the tin atoms are coordinated to both nitrogen atoms directly linked to the naphthalene core and, simultaneously, to one of the nitrogen atoms from another amidinate fragment. Unfortunately, providing theoretical arguments for the formation of dimer  $(\text{L}_2\text{Sn})_2$  is challenging. Indeed, calculations reveal an


energetically favorable reaction pathway ( $\Delta G = -27.74$  kcal/mol) to obtain the corresponding monomer. We can postulate that the presence of the 2,6-dimethylphenyl and *t*-butyl groups in the ligand introduces significant steric hindrance, inhibiting the arrangement of a tetracoordinated tin center and thereby influencing the preferential formation of the dimer.



**Figure 3.** Molecular structure of  $L_1Sn$ . Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms have been omitted for clarity. Selected bond distances [ $\text{\AA}$ ] and bond angles [deg]: Sn(1)-N(1) 2.2435(17); Sn(1)-N(2) 2.3327(17); Sn(1)-N(3) 2.2232(17); Sn(1)-N(4) 2.3181(16); N(1)-C(11) 1.326(3); N(2)-C(11) 1.328(3); N(3)-C(21) 1.328(3); N(4)-C(21) 1.329(3); N(3)-Sn(1)-N(1) 72.57(6); N(3)-Sn(1)-N(4) 57.74(6); N(1)-Sn(1)-N(4) 107.35(6); N(3)-Sn(1)-N(2) 112.07(6); N(1)-Sn(1)-N(2) 57.38(6); N(4)-Sn(1)-N(2) 95.29(6); N(1)-C(11)-N(2) 111.85(17); N(3)-C(21)-N(4) 111.37(17).




**Figure 4.** LUMO +2 (up) and HOMO -1 (down) for  $L_1Sn$ .




**Figure 5.** Molecular structure of  $(L_2Sn)_2$ . Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms and solvent molecules have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Sn(1)-N(1) 2.243(2); Sn(1)-N(2<sup>i</sup>) 2.327(2); Sn(1)-N(3) 2.144(3); C(11)-N(1) 1.378(4); C(11)-N(2) 1.309(4); N(1)-Sn(1)-N(3) 78.80(9); N(3)-Sn(1)-N(2<sup>i</sup>) 97.73(9); N(1)-Sn(1)-N(2<sup>i</sup>) 95.21(9); C(11)-N(1)-Sn(1) 112.43(18); N(1)-C(11)-N(2) 114.2(3).

Stannylenes  $L_3Sn$  and dimer  $(L_2Sn)_2$  can also be prepared through the reaction of deprotonated bis(amidine) ligand ( $L_2K_2$  or  $L_3K_2$ ) with one equivalent of  $SnCl_2$ . The same products,  $(L_2Sn)_2$  and  $L_3Sn$ , were also obtained using two equivalents of  $SnCl_2$ , instead of the expected bis-stannylenes  $L_2Sn_2$  and  $L_3Sn_2$  (Scheme 3).



**Scheme 3.** Synthetic routes to obtain  $L_{2,3}Sn$  from deprotonated bis(amidine) ligands.

Other attempts to synthesize bis-stannylenes  $L_2Sn_2$  and  $L_3Sn_2$ , by reacting the corresponding ligand ( $L_2H_2$  or  $L_3H_2$ ) with two equivalents of  $Sn(HMDS)_2$  (via protonolysis reactions), have also failed (Scheme 4). In both cases, at room temperature, the formation of an intermediate  $LH-SnN(TMS)_2$  was first detected by  $^1H$  NMR spectroscopy. The subsequent heating at  $60^\circ C$  for 1h affords the corresponding stannylenes  $(L_2Sn)_2$  or  $L_3Sn$  (44% and 61% yield, respectively), accompanied by the formation of H-HMDS and a black solid of  $Sn^0$  as by-products.




**Scheme 4.** Synthetic routes to obtain  $L_{2,3}Sn$  by protonolysis.



### 2.3. Synthesis of Germylenes Stabilized by Bis(amidine) Ligands

Following the same synthetic strategy, we have considered the reaction of bis(amidinato) ligands  $L_{1-3}H_2$  with  $Ge(HMDS)_2$ . In the case of  $L_1H_2$ , only dimer  $(L_1Ge)_2$  was obtained as yellow crystals in 42% yield (Scheme 5). Because of poor solubility, it was only possible to characterize  $(L_1Ge)_2$  by an X-ray diffraction analysis. The eight-membered cyclic molecular structure of  $(L_1Ge)_2$  exhibits two trigonal pyramidal Ge centers with a Ge–Ge distance of 3.4984(3) Å (Figure 6).



Scheme 5. Synthesis of germylenes  $L_{1-3}Ge$ .





Figure 6. Molecular structure of  $(L_1Ge)_2$ . Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms and solvent molecules have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Ge(1)–N(1<sup>1</sup>) 2.0961(11); Ge(1)–N(2) 2.0048(10); Ge(1)–N(3) 1.9332(10); C(9)–N(1) 1.3074(16); C(9)–N(2) 1.3740(16); N(3)–Ge(1)–N(2) 87.48(4); N(3)–Ge(1)–N(1<sup>1</sup>) 95.69(4); N(2)–Ge(1)–N(1<sup>1</sup>) 96.79(4); N(1)–C(9)–N(2) 116.31(11); C(9)–N(2)–Ge(1) 112.38(8).

The reaction of  $L_2H_2$  with  $Ge(HMDS)_2$  afforded germylene  $L_2Ge$  after heating at 110 °C for 48 h to complete the reaction (Scheme 5). Germylene  $L_2Ge$  was isolated as yellow crystals from a pentane solution at  $-30$  °C in 52% yield. The  $^1H$  NMR spectrum shows a singlet at 1.60 ppm with an integration of 18 H, corresponding to the *t*butyl groups, and a singlet at 2.18 ppm with an integration of 12 H, corresponding to the methyl groups of the 2,6-dimethylphenyl substituent. In addition, the  $^{13}C$  NMR spectrum shows the



characteristic signal for the NCN fragment at 170.9 ppm. These data are in agreement with a symmetrical tetracoordinated germanium center in solution. However, in the solid state, the X-ray diffraction analysis of  $L_2Ge$  shows a tri-coordinated germanium atom with a distorted trigonal pyramidal geometry. Here, the coordination is through the two nitrogen atoms of the naphthalene bridge and one nitrogen atom of amidine moiety, leaving the remaining nitrogen atom dangling (Figure 7). The N–C distances [1.275(5)–1.417(5) Å] indicate an electron delocalization within the NCN fragments, and due to the bulky nature of the *t*butyl substituents, they are in opposite directions to each other, distorting the naphthalene ring.





**Figure 7.** Molecular structure of  $L_2Ge$ . Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Ge(1)-N(1) 1.969(3); Ge(1)-N(2) 2.073(3); Ge(1)-N(3) 1.912(3); C(11)-N(2) 1.308(5); C(11)-N(1) 1.374(5); C(24)-N(4) 1.275(5); C(24)-N(3) 1.417(5); N(1)-Ge(1)-N(2) 64.89(13); N(1)-Ge(1)-N(3) 90.95(13); N(2)-Ge(1)-N(3) 102.00(13); N(1)-C(11)-N(2) 108.0(3); N(3)-C(24)-N(4) 124.0(3).

#### 2.4. Reactivity of Stannylyene $L_1Sn$

Several tests of reactivity were carried out with stannylyene  $L_1Sn$ , which is easy to obtain in good yields. We first considered the activation of small molecules such as ethylene,  $NH_3$ , and  $CO_2$ , but without any success because of no reactions, despite the multiple conditions evaluated (temperature, time, and pressure).


However, when exposed to  $N_2O$  (five bars) in THF solution,  $L_1Sn$  slowly reacts (3 h at 70 °C) to give a dimeric amidinate-stannoxane **1a**, which was isolated as colorless crystals in 46% yield (Scheme 6). Compound **1a** is moderately soluble in THF, and the  $^1H$  NMR signals tend to be broadened. Therefore, the coupling of signals in the aromatic region cannot be clearly seen, but the integration is as expected, with two broad singlets at 2.07 and 1.85 ppm that integrate for 12 H, corresponding to the methyl groups of the 2,6-dimethylphenyl fragment. In addition, a resonance at 1.76 ppm, integrating for 12 H and corresponding to the methyl groups of the amidine moieties, was observed. The characteristic signal of the carbon amidinate fragments in the  $^{13}C$  NMR spectrum resonates at 168.7 ppm. In addition, a  $^{119}Sn$  chemical shift at  $-494.5$  ppm was observed, consistent with reported stannoxane dimers [19]. The crystal structure of **1a** shows a dinuclear species with a central planar  $Sn_2O_2$  ring with two hexacoordinated tin atoms in a distorted pseudo-octahedral geometry (Figure 8). The O–Sn distances ( $\sim 2.000$  Å), Sn–Sn contact (2.9779(2) Å), and O–Sn–O angles ( $84.01^\circ$ ) are in the range of other dimeric tin amidinate-stannoxanes reported [19].

Scheme 6. Synthesis of **1a**.


**Figure 8.** Molecular structure of **1a**. Thermal ellipsoids are represented with a 30% probability. Hydrogens and solvent molecules have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Sn(1)-N(1) 2.2236(10); Sn(1)-N(2) 2.1644(11); Sn(1)-N(3) 2.1423(10); Sn(1)-N(4) 2.2742(10); Sn(1)-O(1) 2.0079(9); Sn(1)-O(1<sup>i</sup>) 1.9996(8); Sn(1)-Sn(1<sup>i</sup>) 2.97787(19); C(11)-N(1) 1.3285(16); C(11)-N(2) 1.3415(16); C(21)-N(4) 1.3113(17); C(21)-N(3) 1.3439(17); O(1)-Sn(1)-O(1<sup>i</sup>) 84.01(4); Sn(1)-O(1)-Sn(1<sup>i</sup>) 95.98(4); N(1)-C(11)-N(2) 110.50(11); N(3)-C(21)-N(4) 112.95(11).

Usually, stannylenes readily react with chalcogenides [6,20–25], and therefore we have investigated the reaction of **L<sub>1</sub>Sn** with elemental sulfur (Scheme 7). After the reaction mixture had been stirred at room temperature for 4 h, <sup>1</sup>H, and <sup>119</sup>Sn NMR spectroscopy showed a mixture of new signals and starting stannylene. In the <sup>119</sup>Sn NMR spectrum, next to the resonance of **L<sub>1</sub>Sn**, a new signal appears at −371 ppm, which evolves to a resonance singlet at −648.0 ppm after 24 h, corresponding to dimer **1b**. After purification, **1b** was isolated as pale-yellow crystals from a THF solution at −30 °C in 51% yield. The dimeric structure of **1b** has been determined by an X-ray diffraction analysis, revealing a central planar Sn<sub>2</sub>S<sub>2</sub> ring with the two tin centers in a distorted pseudo-octahedral geometry (Figure 9). The S–Sn distances are ~2.400 Å and Sn–Sn contact is 3.3847(10) Å [25], longer than that observed for the di-oxygen analog dimer **1a** due to the size of the sulfur atom.

In addition, compound **1b** was characterized by NMR spectroscopy. The <sup>1</sup>H NMR spectrum exhibits broadened signals. In the <sup>13</sup>C NMR spectrum, the characteristic signal of the NCN fragment appears at 167.4 ppm. In the <sup>119</sup>Sn NMR spectrum, a very high field chemical shift at −648.0 ppm is observed, a resonance signal that is more high-field shifted than those of previously reported Sn<sub>2</sub>S<sub>2</sub>-bridged dimeric complexes [26,27].



**Scheme 7.** Synthetic route to obtain **1b**.




**Figure 9.** Molecular structure of **1b**. Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms and solvent molecules have been omitted for clarity. The asymmetric unit contains two independent but very similar molecules; data for only one molecule are discussed. Selected bond distances [Å] and bond angles [deg]: Sn(1)-N(1) 2.130(4); Sn(1)-N(2) 2.374(3); Sn(1)-N(3) 2.240(3); Sn(1)-N(4) 2.198(4); Sn(1)-S(1) 2.4172(14); Sn(1)-S(1<sup>i</sup>) 2.4585(13); C(11)-N(1) 1.345(6); C(11)-N(2) 1.315(5); C(21)-N(3) 1.332(5); C(21)-N(4) 1.328(5); Sn(1)-S(1)-Sn(1<sup>i</sup>) 87.92(4); S(1)-Sn(1)-S(1<sup>i</sup>) 92.08(4); N(1)-C(11)-N(2) 112.8(4); N(3)-C(21)-N(4) 110.9(4).


**L**<sub>1</sub>**Sn** reacts with *p*-tolyldisulfide in THF at 70 °C overnight via an oxidative addition reaction with a tin insertion into the S–S bond (Scheme 8) to give a hexacoordinate Sn<sup>(IV)</sup> species **2a** [22]. This molecule presents a characteristic <sup>119</sup>Sn chemical shift at −456.7 ppm, consistent with a hexacoordinated tin center [24]. In addition, the X-ray diffraction analysis of **2a** reveals a monomeric Sn<sup>(IV)</sup> species in a distorted octahedral coordination sphere that included the four nitrogen atoms of the two chelating amidinates and the two *p*-tolyl sulfur fragments (Figure 10). The equivalence of C–N bond lengths within NCN frameworks and their magnitudes (1.301 to 1.361 Å) indicate that the π electrons within the ligands are delocalized.

3,5-di-*tert*-butyl-ortho-quinone also reacts with **L**<sub>1</sub>**Sn** to afford the corresponding [4 + 1]-cycloadduct **2b** (Scheme 9). The <sup>1</sup>H NMR spectrum reveals the characteristic signals of the quinone group in the aromatic region, doublets at 6.57 and 6.39 ppm (*J*<sub>HH</sub> = 2.4 Hz), and two singlets at 1.06 and 1.16 ppm, corresponding to the CH<sub>3</sub> groups of *t*butyl in the aliphatic area. Also, the signals corresponding to the amidine CH<sub>3</sub> groups' moieties integrating for 6 H as a singlet at 2.00 ppm and to the CH<sub>3</sub> groups of 2,6-dimethylphenyl moieties integrating for 12 H as a singlet at 2.08 ppm, respectively, are observed. The <sup>119</sup>Sn NMR spectrum shows a singlet at −512.2 ppm, in agreement with the cycloadduct structure.

The mass analysis (DCI/CH<sub>4</sub>) spectrum exhibits a peak at 786.29, which corresponds to [M]<sup>+</sup> of compound **2b**, evidencing its formation.





**Scheme 8.** Synthesis of **2a**.





**Figure 10.** Molecular structure of **2a**. Thermal ellipsoids are represented with a 30% probability. Hydrogen atoms have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Sn(1)-N(1) 2.132(3); Sn(1)-N(2) 2.411(3); Sn(1)-N(3) 2.230(3); Sn(1)-N(4) 2.180(3); Sn(1)-S(1) 2.4297(15); Sn(1)-S(2) 2.4483(16); C(11)-N(1) 1.361(4); C(11)-N(2) 1.301(4); C(21)-N(3) 1.334(4); C(21)-N(4) 1.337(4); N(1)-Sn(1)-N(4) 132.96(11); N(1)-Sn(1)-N(3) 78.01(11); N(3)-Sn(1)-N(4) 59.72(10); N(1)-Sn(1)-N(2) 58.27(11); N(2)-Sn(1)-N(4) 97.20(12); N(2)-Sn(1)-N(3) 86.71(11); N(1)-Sn(1)-S(1) 111.48(9); N(4)-Sn(1)-S(1) 102.83(8); N(3)-Sn(1)-S(1) 158.14(7); N(2)-Sn(1)-S(1) 82.44(9); N(1)-Sn(1)-S(2) 105.90(9); N(4)-Sn(1)-S(2) 97.26(9); N(3)-Sn(1)-S(2) 94.06(9); N(2)-Sn(1)-S(2) 163.67(8); S(1)-Sn(1)-S(2) 101.67(7); N(2)-C(11)-N(1) 113.5(3); N(3)-C(21)-N(4) 110.6(3).

As reported by So et al., an amidinatogermylene can act as a good ligand to form a germylene→ECl<sub>2</sub>-type adduct with GeCl<sub>2</sub> and SnCl<sub>2</sub> [28]. In contrast, **L<sub>1</sub>Sn** reacts with GeCl<sub>2</sub>(dioxane) in THF at room temperature via a transmetalation reaction to give a germylene dimer (**L<sub>1</sub>Ge**)<sub>2</sub> (Scheme 10). The <sup>119</sup>Sn NMR spectrum shows a resonance at −214.0 ppm, in agreement with the formation of SnCl<sub>2</sub> [29].

Scheme 9. Synthesis of **2b**.Scheme 10. Synthesis of **L<sub>1</sub>Ge** by transmetalation reaction.

Transmetalation reactions were also observed with  $\text{AlCl}_3$  and  $\text{GaCl}_3$ , affording the corresponding  $\text{Al}^{(\text{III})}$  **3a** and  $\text{Ga}^{(\text{III})}$ -chlorides **3b**, and in both cases, forming one equivalent of  $\text{SnCl}_2$  (Scheme 11). The modeling confirms an exergonic reaction ( $-18.92$  kcal/mol) during this “metal” exchange process. **L<sub>1</sub>Sn** reacts with  $\text{AlCl}_3$  quickly in THF at room temperature for 30 min to give **3a**, which was isolated as a white solid in 83% yield. The  $^1\text{H}$  NMR spectrum shows, in the aromatic region, the expected resonances and integrals, and in the aliphatic area, two singlets at 2.33 and 1.90 ppm, integrating for 12 H, corresponding to the methyl groups of 2,6-dimethylphenyl fragments and a singlet at 2.13 ppm (6 H) related to the  $-\text{CH}_3$  of the amidine groups. In the  $^{13}\text{C}$  NMR spectrum, the characteristic NCN signal is observed at 176.4 ppm. The symmetry of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra are in agreement with a pentacoordinated aluminum complex. Using the same experimental conditions, **L<sub>1</sub>Sn** also reacts with  $\text{GaCl}_3$  to give  $\text{Ga}^{(\text{III})}$ -chloride **3b**, which was isolated as colorless crystals from a pentane solution at  $-30$  °C in 88% yield. Gallium chloride **3b** was fully characterized by NMR spectroscopy (very similar to that of **3a**) and by X-ray diffraction analysis. The molecular structure of **3b** (Figure 11) shows a pentacoordinated gallium center with a distorted square-based pyramidal geometry. The Ga–Cl bond distance of 2.183 Å is in the range expected for such compounds [30].

Scheme 11. Synthesis of **3a** and **3b** via transmetalation reactions.



**Figure 11.** Molecular structure of **3b**. Thermal ellipsoids are represented with a 30% probability. Hydrogens have been omitted for clarity. Selected bond distances [Å] and bond angles [deg]: Ga(1)-N(1) 1.984(4); Ga(1)-N(2) 2.017(4); Ga(1)-N(3) 1.981(4); Ga(1)-N(4) 2.051(4); Ga(1)-Cl(1) 2.1828(14); N(1)-Ga(1)-N(3) 85.32(15); C(11)-N(1) 1.320(5); C(11)-N(2) 1.345(6); C(21)-N(3) 1.327(6); C(21)-N(4) 1.342(6); N(2)-Ga(1)-N(3) 139.06(16); N(1)-Ga(1)-N(2) 65.43(16); N(3)-Ga(1)-N(4) 65.69(15); N(1)-Ga(1)-N(4) 130.88(15); N(2)-Ga(1)-N(4) 112.06(16); N(3)-Ga(1)-Cl(1) 110.32(12); N(1)-Ga(1)-Cl(1) 115.30(11); N(2)-Ga(1)-Cl(1) 107.98(12); N(4)-Ga(1)-Cl(1) 111.79(12); N(1)-C(11)-N(2) 108.5(4); N(3)-C(21)-N(4) 110.1(4).

### 3. Materials and Methods

#### 3.1. General Comments

All manipulations were performed under an inert argon or nitrogen atmosphere using standard Schlenk-line and glovebox techniques. Dry, oxygen-free solvents were employed. Reagents were obtained from commercial suppliers unless otherwise stated. *N*-(2,6-dimethylphenyl)acetamide [31], *N*-(2,6-dimethylphenyl)acetimidoyl chloride [29], *N,N'*-(naphthalene-1,8-diyl)bis(2,2-dimethylpropanamide) [13], *N,N'*-(naphthalene-1,8-diyl)bis(2,2-dimethylpropanimidoyl chloride) [13], *L*<sub>1</sub> [12], and *L*<sub>2</sub> [13] were synthesized following reported procedures. The Lappert's germanium(II) and tin(II) derivatives were prepared according to the literature procedures [32]. The 1D and 2D NMR spectra were recorded with the following spectrometers for <sup>1</sup>H, <sup>13</sup>C, and <sup>119</sup>Sn: Bruker Avance II 300 MHz, Avance III HD 400 MHz, and Avance I and II 500 MHz spectrometers. The chemical shift has been counted positively versus the low field and expressed in part per million (ppm). The mass spectrometric analysis was performed using three techniques: direct chemical ionization (DCI-CH<sub>4</sub>) methods recorded on a GCT Premier Waters mass spectrometer (Headquarters, MN, USA); electrospray ionization (ESI, Los Angeles, CA, USA) recorded on a Waters Xevo G2 Q-TOF mass spectrometer; and a Maldi micro-MX micro-mass in a pyrene matrix (ratio of product/matrix 1/100). Melting points were measured with capillary Electrothermal Stuart SMP40 apparatus, and samples were prepared in the glovebox before the analysis. FT-IR spectra were measured on a Thermo Nicolet 6700 (Waltham, MA, USA) Nexus and recovered in a solid state (KBr). Single-crystal X-ray data were collected at a low temperature (193(2)K) on a Bruker APEX II Quazar (Billerica, MA, USA) diffractometer equipped with a 30W air-cooled microfocus source [(*L*<sub>2</sub>Sn)<sub>2</sub>, *L*<sub>2</sub>Ge, **2a** and **3b**] or on a Bruker D8 VENTURE diffractometer equipped with a PHOTON III detector [*L*<sub>3</sub>H<sub>2</sub>, *L*<sub>1</sub>Sn, (*L*<sub>1</sub>Ge)<sub>2</sub>, **1a** and **1b**] using MoK $\alpha$  radiation ( $\lambda = 0.71037$  Å). The structures were solved by the intrinsic phasing method (SHELXT) [33] and refined by the full-matrix least-squares method on F2 [34]. All non-H atoms were refined with anisotropic displacement parameters and all the hydrogen atoms were refined isotropically at calculated positions using a riding model. For **2a**, some solvent molecules were highly disordered and difficult to model correctly. Therefore, the SQUEEZE function of PLATON [35] was used to eliminate the contribution



of the electron density of those solvent molecules from the intensity data. Calculations were performed with the Gaussian 16 suite of programs [36] using the density functional method B3LYP with dispersion (D3) [37–40]. Tin atoms were treated with a Stuttgart–Dresden pseudopotential in combination with its adapted basis set [41]. All other atoms have been described with a 6–31G(d,p) basis set. Geometry optimizations were carried out without any symmetry restrictions. Frequency calculations were undertaken to confirm the nature of the stationary points, yielding one imaginary frequency for transition states (TS) and all of them were positive for minima. The connectivity of the transition states and their adjacent minima was confirmed by intrinsic reaction coordinate (IRC) calculations [42,43].

### 3.2. Synthesis

**Synthesis of L<sub>3</sub>H<sub>2</sub>.** *p*-toluidine (591 mg, 5.6 mmol) was added to a solution of *N,N'*-(naphthalene-1,8-diyl)bis(2,2-dimethylpropanimidoyl chloride) (1 g, 2.8 mmol) in dry toluene (50 mL). Then, Et<sub>3</sub>N (559 mg, 5.6 mmol) was added to the mixture. The reaction was stirred for 4 h under reflux. All volatiles were removed under reduced pressure. The solid residue was taken up in 40 mL of Et<sub>2</sub>O and washed with 25 mL of a saturated solution of Na<sub>2</sub>CO<sub>3</sub>. Then, the organic layer was washed with water (3 × 30 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. After that, the solvent was removed, and the crude product was recrystallized with CH<sub>2</sub>Cl<sub>2</sub>/pentane (1:2). Yellow crystals were obtained (63% yield). **Melting point:** 116–122 °C. **<sup>1</sup>H NMR** (DMSO-*d*<sub>6</sub>, 500 MHz): δ 9.43 (s, 1H, NH); 8.50 (s, 1H, NH); 7.26 (d, *J*<sub>HH</sub> = 6.5 Hz, 1H, aryl); 6.94 (d, *J*<sub>HH</sub> = 7.9 Hz, 1H, aryl); 6.88–6.83 (m, 3H, aryl); 6.80 (m, 3H, aryl); 6.66 (d, *J*<sub>HH</sub> = 8.1 Hz, 2H, aryl); 6.58–6.51 (m, 2H, aryl); 6.32–6.20 (m, 2H, aryl); 2.05 (s, 3H, CH<sub>3</sub>); 2.01 (s, 3H, CH<sub>3</sub>); 1.38 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>); 1.23 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (DMSO-*d*<sub>6</sub>, 125 MHz): δ 161.3 (C=N); 160.7 (C=N); 147.2 (aryl, C); 146.5 (aryl, C); 138.6 (aryl, C); 137.2 (aryl, C); 135.6 (aryl, C); 131.2 (aryl, C); 130.4 (aryl, C); 128.7 (aryl, CH); 128.0 (aryl, CH); 127.7 (aryl, CH); 124.8 (aryl, CH); 121.4 (aryl, CH); 120.9 (aryl, CH); 119.8 (aryl, CH); 116.5 (aryl, C); 114.8 (aryl, CH); 28.8 (C(CH<sub>3</sub>)<sub>3</sub>); 28.3 (C(CH<sub>3</sub>)<sub>3</sub>); 20.3 (CH<sub>3</sub>). **IR** (KBr, cm<sup>−1</sup>): 3373 (νNH); 1623 (νC=N). **ESI *m/z***: 504.33 ([M]<sup>+</sup>).

#### 3.2.1. General Synthetic Procedure of Metallylenes

THF (5 mL) was added to a mixture of one equiv. of MCl<sub>2</sub> (M = Sn or Ge) and two equiv. of K[N(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub> in a 20 mL vial inside the glovebox. The mixture was stirred for 30 min at room temperature. After this time, a white precipitate was obtained. The mixture was filtered, and the solution was added to one equiv. of the corresponding L<sub>1–3</sub>H<sub>2</sub>, stirring at 60 °C for 3 h. Then, all the volatiles were removed and the residual solid was washed with pentane (3 × 5 mL).

**L<sub>1</sub>Sn.** Colorless crystals were obtained by recrystallization from pentane at −30 °C (77% yield). **Melting point:** 202 °C (decomposition). **<sup>1</sup>H NMR** (THF-*d*<sub>8</sub>, 500 MHz): δ 7.42 (dd, *J*<sub>HH</sub> = 8.2, 1.1 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.31–7.27 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 7.07 (dd, *J*<sub>HH</sub> = 7.5, 1.1 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 6.96 (d, *J*<sub>HH</sub> = 7.4 Hz, 4H, C<sub>6</sub>H<sub>3</sub>); 6.83 (t, *J*<sub>HH</sub> = 7.5 Hz, 2H, C<sub>6</sub>H<sub>3</sub>); 2.18 (s, 12H, CH<sub>3</sub>); 1.97 (s, 6H, CH<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (THF-*d*<sub>8</sub>, 125 MHz): δ 169.0 (NCN); 145.3 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 144.5 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 138.6 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 134.4 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 128.8 (C<sub>6</sub>H<sub>3</sub>); 126.4 (C<sub>10</sub>H<sub>6</sub>); 124.8 (C<sub>6</sub>H<sub>3</sub>); 123.4 (C<sub>10</sub>H<sub>6</sub>); 119.8 (C<sub>10</sub>H<sub>6</sub>); 20.0 (CH<sub>3</sub>); 17.9 (CH<sub>3</sub>). **<sup>119</sup>Sn{<sup>1</sup>H} NMR** (THF-*d*<sub>8</sub>, 186 MHz): δ −276.7. **ESI *m/z***: 565.15 ([M]<sup>+</sup>); 449.27 ([M − Sn]<sup>+</sup>).

**(L<sub>2</sub>Sn)<sub>2</sub>.** Yellow crystals (44% yield). **Melting point:** 243 °C (decomposition). **MS** (Maldi-TOF) *m/z*: 649.3 ([M/2, monomer]<sup>+</sup>). Meaningful solution state spectroscopic data for the compound could not be obtained for the compound as it shows negligible solubility in normal non-coordinating deuterated solvents once crystallized.

**L<sub>3</sub>Sn.** Yellow solid (61% yield). **Melting point:** 189 °C (decomposition). **<sup>1</sup>H NMR** (THF-*d*<sub>8</sub>, 400 MHz): δ 7.38 (d, *J*<sub>HH</sub> = 7.5 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.24–7.19 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 7.09 (d, *J*<sub>HH</sub> = 7.1 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.05 (d, *J*<sub>HH</sub> = 8.0 Hz, 4H, C<sub>6</sub>H<sub>4</sub>); 6.97 (d, *J*<sub>HH</sub> = 8.2 Hz, 4H, C<sub>6</sub>H<sub>4</sub>); 2.29 (s, 6H, CH<sub>3</sub>); 1.30 (s, 18H, C(CH<sub>3</sub>)<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (THF-*d*<sub>8</sub>, 125 MHz): δ 177.5 (NCN); 146.7 (C<sub>6</sub>H<sub>4</sub><sub>ipso</sub>); 145.7 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 138.4 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 133.0 (C<sub>6</sub>H<sub>4</sub><sub>ipso</sub>); 130.4 (C<sub>6</sub>H<sub>4</sub>); 126.3 (C<sub>10</sub>H<sub>6</sub>); 124.9 (C<sub>6</sub>H<sub>4</sub>); 123.4 (C<sub>10</sub>H<sub>6</sub>); 121.3 (C<sub>10</sub>H<sub>6</sub>); 43.6 (C(CH<sub>3</sub>)<sub>3</sub>); 31.8 (C(CH<sub>3</sub>)<sub>3</sub>);



21.0 (CH<sub>3</sub>). <sup>119</sup>Sn{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 186 MHz): δ −254.9. MS (Maldi-TOF) *m/z*: 621.2 ([M]<sup>+</sup>); 503.4 ([M-Sn]<sup>+</sup>).

(L<sub>1</sub>Ge)<sub>2</sub>. Yellow crystals (42% yield). **Melting point**: 259 °C (decomposition). MS (Maldi-TOF) *m/z*: 519.14 ([M/2, monomer]<sup>+</sup>). Meaningful solution state spectroscopic data for the compound could not be obtained for the compound as it shows negligible solubility in normal non-coordinating deuterated solvents once crystallized.

L<sub>2</sub>Ge. Toluene was used instead of THF as solvent. Yellow crystals were obtained by recrystallization from pentane at −30 °C. (52% yield). **Melting point**: 199 °C (decomposition). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz): δ 6.98–6.91 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 6.89–6.81 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 6.62 (d, *J*<sub>HH</sub> = 7.7 Hz, 4H, C<sub>6</sub>H<sub>3</sub>); 6.47–6.40 (m, 2H, C<sub>6</sub>H<sub>3</sub>); 6.33 (d, *J*<sub>HH</sub> = 6.3 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 2.18 (s, 12H, CH<sub>3</sub>); 1.60 (s, 18H, C(CH<sub>3</sub>)<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, 125 MHz): δ 170.9 (NCN); 151.8 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 148.8 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 139.7 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 138.5 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 128.3 (C<sub>6</sub>H<sub>3</sub>); 126.3 (C<sub>10</sub>H<sub>6</sub>); 120.4 (C<sub>6</sub>H<sub>3</sub>); 114.9 (C<sub>10</sub>H<sub>6</sub>); 108.8 (C<sub>10</sub>H<sub>6</sub>); 41.9 (C(CH<sub>3</sub>)<sub>3</sub>); 30.6 (C(CH<sub>3</sub>)<sub>3</sub>); 19.7 (CH<sub>3</sub>).

**Synthesis of 1a.** In a J. Young valve NMR tube, a THF (0.4 mL) solution of L<sub>1</sub>Sn (30 mg, 0.053 mmol) was exposed to 5 bar of N<sub>2</sub>O. The reaction was monitored by NMR and proceeded quantitatively after 3 h at 70 °C. Colorless crystals were observed and separated from the solution by filtration. Then, the crystals were washed with pentane (3 × 2 mL). Colorless crystals were obtained (46% yield). **Melting point**: 204 °C. <sup>1</sup>H NMR (THF-d<sub>8</sub>, 500 MHz): δ 7.42 (d, *J*<sub>HH</sub> = 6.4 Hz, 4H, C<sub>10</sub>H<sub>6</sub>); 7.26 (t, *J*<sub>HH</sub> = 7.8 Hz; 4H, C<sub>10</sub>H<sub>6</sub>); 7.13 (d, *J*<sub>HH</sub> = 6.8 Hz, 4H, C<sub>10</sub>H<sub>6</sub>), 7.01 (s, 4H, C<sub>6</sub>H<sub>3</sub>); 6.92–6.83 (m, 8H, C<sub>6</sub>H<sub>3</sub>); 2.07 (s, 12H, CH<sub>3</sub>); 1.85 (s, 12H, CH<sub>3</sub>); 1.76 (s, 12H, CH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 125 MHz): δ 168.7 (NCN); 143.6 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 142.7 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 138.5 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 137.0 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 135.3 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 128.7 (C<sub>6</sub>H<sub>3</sub>); 128.3 (C<sub>6</sub>H<sub>3</sub>); 126.3 (C<sub>10</sub>H<sub>6</sub>); 125.7 (C<sub>6</sub>H<sub>3</sub>); 124.8 (C<sub>10</sub>H<sub>6</sub>); 120.6 (C<sub>10</sub>H<sub>6</sub>); 19.8 (CH<sub>3</sub>); 19.0 (CH<sub>3</sub>); 15.8 (CH<sub>3</sub>). <sup>119</sup>Sn{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 186 MHz): δ −494.5. MS (Maldi-TOF) *m/z*: 1161.26 ([M]<sup>+</sup>); 583.13 ([M/2, monomer]<sup>+</sup>).

### 3.2.2. General Reactivity Evaluation Procedure

THF was added to a mixture of one equiv. of L<sub>1</sub>Sn and one equiv. of S<sub>8</sub>, *p*-tolylidysulfide, 3,5-di-*tert*-butyl-*o*-benzoquinone, AlCl<sub>3</sub>, or GaCl<sub>3</sub>, accordingly. The mixture was stirred for the corresponding time and temperature. Then, the solvent was removed and the residual solid was washed with pentane (3 × 0.5 mL).

**1b.** Stirred overnight at room temperature. Pale-yellow crystals were obtained by recrystallization from THF at −30 °C (51% yield). **Melting point**: 229 °C (decomposition). <sup>1</sup>H NMR (THF-d<sub>8</sub>, 300 MHz): δ 7.40 (d, *J*<sub>HH</sub> = 7.5 Hz, 4H, C<sub>10</sub>H<sub>6</sub>); 7.22 (t, *J*<sub>HH</sub> = 7.8 Hz, 4H, C<sub>10</sub>H<sub>6</sub>); 7.11–7.02 (m, 4H, C<sub>10</sub>H<sub>6</sub>); 6.93 (bs, 12H, C<sub>6</sub>H<sub>3</sub>); 2.11 (br s, 24H, CH<sub>3</sub>); 1.75 (s, 12H, CH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 125 MHz): δ 167.4 (NCN); 142.7 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 138.2 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 135.9 (C<sub>6</sub>H<sub>3</sub><sub>ipso</sub>); 128.8 (C<sub>6</sub>H<sub>3</sub>); 126.4 (C<sub>10</sub>H<sub>6</sub>); 125.7 (C<sub>6</sub>H<sub>3</sub>); 124.8 (C<sub>10</sub>H<sub>6</sub>); 121.5 (C<sub>10</sub>H<sub>6</sub>); 19.7 (CH<sub>3</sub>); 16.4 (CH<sub>3</sub>). <sup>119</sup>Sn{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 186 MHz): δ −648.0. MS (Maldi-TOF) *m/z*: 1193.10 ([M]<sup>+</sup>); 597.04 ([M/2, monomer]<sup>+</sup>).

**2a.** Stirred overnight at 70 °C. Pale-brown crystals (51% yield). **Melting point**: 199 °C. <sup>1</sup>H NMR (THF-d<sub>8</sub>, 500 MHz): δ 7.40 (dd, *J*<sub>HH</sub> = 8.1, 1.1 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.25–7.21 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 7.17 (dd, *J*<sub>HH</sub> = 7.5, 1.2 Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.05–6.96 (m, 6H, C<sub>6</sub>H<sub>3</sub>); 6.63 (d, *J*<sub>HH</sub> = 8.1 Hz, 4H, C<sub>6</sub>H<sub>4</sub>); 6.44 (d, *J*<sub>HH</sub> = 7.9 Hz, 4H, C<sub>6</sub>H<sub>4</sub>); 2.19 (s, 12H, CH<sub>3</sub>); 2.09 (s, Sn satellite: *J*<sub>HSn</sub> = 7.3 Hz, 6H, CH<sub>3</sub>); 1.79 (s, Sn satellite: *J*<sub>HSn</sub> = 2.4 Hz, 6H, CH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 125 MHz): δ 168.8 (NCN); 143.8 (Sn satellite: *J*<sub>CSn</sub> = 5.2 Hz, C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 143.3 (Sn satellite: *J*<sub>CSn</sub> = 7.2 Hz, C<sub>6</sub>H<sub>3</sub>); 138.4 (Sn satellite: *J*<sub>CSn</sub> = 3.3 Hz, C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 136.1 (Sn satellite: *J*<sub>CSn</sub> = 12.8 Hz, C<sub>6</sub>H<sub>4</sub>); 135.6 (Sn satellite: *J*<sub>CSn</sub> = 3.1 Hz, C<sub>6</sub>H<sub>3</sub>); 135.5 (C<sub>6</sub>H<sub>4</sub><sub>ipso</sub>); 131.4 (C<sub>6</sub>H<sub>4</sub><sub>ipso</sub>); 129.2 (C<sub>6</sub>H<sub>3</sub>); 129.0 (Sn satellite: *J*<sub>CSn</sub> = 6.6 Hz, C<sub>6</sub>H<sub>4</sub>); 127.6 (C<sub>10</sub>H<sub>6</sub><sub>ipso</sub>); 126.4 (C<sub>10</sub>H<sub>6</sub>); 126.4 (C<sub>6</sub>H<sub>3</sub>); 125.2 (C<sub>10</sub>H<sub>6</sub>); 121.1 (Sn satellite: *J*<sub>CSn</sub> = 11.2 Hz, C<sub>10</sub>H<sub>6</sub>); 21.1 (CH<sub>3</sub>); 19.9 (CH<sub>3</sub>); 16.4 (Sn satellite: *J*<sub>CSn</sub> = 21.40 Hz, CH<sub>3</sub>). <sup>119</sup>Sn{<sup>1</sup>H} NMR (THF-d<sub>8</sub>, 186 MHz): δ −456.7. MS (Maldi-TOF) *m/z*: 687.13 ([M − S(tolyl)]<sup>+</sup>); 565.05 ([M − 2 S(tolyl)]<sup>+</sup>).

**2b.** Stirred for 15 min at room temperature. White solid (96% yield). **Melting point**: 240 °C (decomposition). <sup>1</sup>H NMR (THF-d<sub>8</sub>, 500 MHz): δ 7.65–7.62 (m, 2H, C<sub>10</sub>H<sub>6</sub>);

7.44–7.39 (m, 4H, C<sub>10</sub>H<sub>6</sub>); 6.98–6.92 (m, 6H, C<sub>6</sub>H<sub>3</sub>); 6.57 (d,  $J_{\text{HH}} = 2.4$  Hz, 1H, C<sub>6</sub>H<sub>2</sub>); 6.39 (d,  $J_{\text{HH}} = 2.4$  Hz, 1H, C<sub>6</sub>H<sub>2</sub>); 2.08 (s, 12H, CH<sub>3</sub>); 2.00 (s, Sn satellite:  $J_{\text{HSn}} = 3.5$  Hz, 6H, CH<sub>3</sub>); 1.16 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>); 1.06 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (THF-d<sub>8</sub>, 125 MHz):  $\delta$  171.6 (NCN); 151.3 (Sn satellite:  $J_{\text{CSn}} = 5.1$  Hz, C<sub>6</sub>H<sub>2ipso</sub>); 147.1 (Sn satellite:  $J_{\text{CSn}} = 3.3$  Hz, C<sub>6</sub>H<sub>2ipso</sub>); 142.0 (Sn satellite:  $J_{\text{CSn}} = 3.9$  Hz, C<sub>10</sub>H<sub>6ipso</sub>); 140.9 (Sn satellite:  $J_{\text{CSn}} = 7.5$  Hz, C<sub>6</sub>H<sub>3ipso</sub>); 138.7 (C<sub>6</sub>H<sub>2ipso</sub>); 138.5 (C<sub>10</sub>H<sub>6ipso</sub>); 136.1 (C<sub>6</sub>H<sub>3ipso</sub>); 134.1 (C<sub>6</sub>H<sub>2ipso</sub>); 129.1 (C<sub>6</sub>H<sub>3</sub>); 127.1 (C<sub>6</sub>H<sub>3</sub>); 126.7 (C<sub>10</sub>H<sub>6</sub>); 126.1 (C<sub>10</sub>H<sub>6ipso</sub>); 126.0 (C<sub>10</sub>H<sub>6</sub>); 121.2 (Sn satellite:  $J_{\text{CSn}} = 11.6$  Hz, C<sub>10</sub>H<sub>6</sub>); 111.5 (C<sub>6</sub>H<sub>2</sub>); 110.3 (Sn satellite:  $J_{\text{CSn}} = 40.2$  Hz, C<sub>6</sub>H<sub>2</sub>); 35.3 (C(CH<sub>3</sub>)<sub>3</sub>); 34.9 (C(CH<sub>3</sub>)<sub>3</sub>); 32.4 (C(CH<sub>3</sub>)<sub>3</sub>); 30.2 (C(CH<sub>3</sub>)<sub>3</sub>); 19.1 (CH<sub>3</sub>); 15.5 (Sn satellite:  $J_{\text{CSn}} = 33.7$  Hz, CH<sub>3</sub>). **<sup>119</sup>Sn{<sup>1</sup>H} NMR** (THF-d<sub>8</sub>, 186 MHz):  $\delta$  −512.2. **ESI  $m/z$** : 786.29 ([M]<sup>+</sup>).

**3a.** Stirred for 30 min at room temperature. White solid (85% yield). **Melting point:** 124 °C. **<sup>1</sup>H NMR** (THF-d<sub>8</sub>, 500 MHz):  $\delta$  7.46 (dd,  $J_{\text{HH}} = 8.3, 1.1$  Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.35–7.31 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 7.24 (dd,  $J_{\text{HH}} = 7.5, 1.2$  Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.02–6.90 (m, 6H, C<sub>6</sub>H<sub>3</sub>); 2.33 (s, 6H, CH<sub>3</sub>); 2.13 (s, 6H, CH<sub>3</sub>); 1.90 (s, 6H, CH<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (THF-d<sub>8</sub>, 125 MHz):  $\delta$  176.4 (NCN); 141.9 (C<sub>10</sub>H<sub>6ipso</sub>); 141.8 (C<sub>10</sub>H<sub>6ipso</sub>); 140.6 (C<sub>6</sub>H<sub>3ipso</sub>); 138.4 (C<sub>10</sub>H<sub>6ipso</sub>); 135.8 (C<sub>6</sub>H<sub>3ipso</sub>); 135.1 (C<sub>6</sub>H<sub>3ipso</sub>); 129.4 (C<sub>6</sub>H<sub>3</sub>); 128.8 (C<sub>6</sub>H<sub>3</sub>); 126.6 (C<sub>10</sub>H<sub>6</sub>); 126.2 (C<sub>6</sub>H<sub>3</sub>); 124.1 (C<sub>10</sub>H<sub>6</sub>); 123.9 (C<sub>10</sub>H<sub>6ipso</sub>); 117.8 (C<sub>10</sub>H<sub>6</sub>); 19.9 (CH<sub>3</sub>); 19.3 (CH<sub>3</sub>); 15.6 (CH<sub>3</sub>). **<sup>27</sup>Al{<sup>1</sup>H} NMR** (THF-d<sub>8</sub>, 130 MHz):  $\delta$  69.4.

**3b.** Stirred for 30 min at room temperature. Colorless crystals were obtained from recrystallization with pentane at −30 °C (88% yield). **Melting point:** 209 °C (decomposition). **<sup>1</sup>H NMR** (THF-d<sub>8</sub>, 500 MHz):  $\delta$  7.46 (dd,  $J_{\text{HH}} = 8.3, 1.1$  Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.34–7.30 (m, 2H, C<sub>10</sub>H<sub>6</sub>); 7.20 (dd,  $J_{\text{HH}} = 7.5, 1.2$  Hz, 2H, C<sub>10</sub>H<sub>6</sub>); 7.03–6.90 (m, 6H, C<sub>6</sub>H<sub>3</sub>); 2.34 (s, 6H, CH<sub>3</sub>); 2.12 (s, 6H, CH<sub>3</sub>); 1.98 (s, 6H, CH<sub>3</sub>). **<sup>13</sup>C{<sup>1</sup>H} NMR** (THF-d<sub>8</sub>, 125 MHz):  $\delta$  172.3 (NCN); 142.9 (C<sub>10</sub>H<sub>6ipso</sub>); 141.9 (C<sub>6</sub>H<sub>3ipso</sub>); 138.6 (C<sub>10</sub>H<sub>6ipso</sub>); 136.1 (C<sub>6</sub>H<sub>3ipso</sub>); 135.5 (C<sub>6</sub>H<sub>3ipso</sub>); 128.8 (C<sub>6</sub>H<sub>3</sub>); 126.4 (C<sub>10</sub>H<sub>6</sub>); 126.4 (C<sub>6</sub>H<sub>3</sub>); 124.3 (C<sub>10</sub>H<sub>6</sub>); 123.1 (C<sub>10</sub>H<sub>6ipso</sub>); 118.3 (C<sub>10</sub>H<sub>6</sub>); 19.8 (CH<sub>3</sub>); 15.1 (CH<sub>3</sub>). **MS (Maldi-TOF)  $m/z$** : 551.06 ([M]<sup>+</sup>); 517.04 ([M − Cl]<sup>+</sup>); 449.10 ([M − GaCl]<sup>+</sup>).

### 3.3. X-ray Data

CCDC-2303009 (L<sub>3</sub>H<sub>2</sub>), CCDC-2266345 (L<sub>1</sub>Sn), CCDC-2266346 [(L<sub>2</sub>Sn)<sub>2</sub>], CCDC-2266347 [(L<sub>1</sub>Ge)<sub>2</sub>], CCDC-2266348 (L<sub>2</sub>Ge), CCDC-2266349 (1a), CCDC-2266350 (1b), CCDC-2266351 (2a), and CCDC-2266352 (3b) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

## 4. Conclusions

Three tetradentate bis(amidine) ligands RNC(R′)N-linker-NC(R′)NR with a rigid naphthalene linker L<sub>1–3</sub> were successfully used for the stabilization of metallylenes. The corresponding stannylenes and germylenes (L<sub>1</sub>Sn and L<sub>1</sub>Ge) have been prepared by protonolysis reaction of Lappert's metallylenes [M(HMDS)<sub>2</sub>] (M = Ge or Sn) and were fully characterized by NMR spectroscopy and X-ray diffraction analysis. Structures in the solid state show either a monomer or a dimer depending on the different substituents of the amidine moiety, demonstrating that the nature of the bis(amidine) system can influence the stabilization and the reactivity of the corresponding tetrylenes. DFT calculations have been performed in order to define the electronic properties associated with tetradentate ligands. The reactivity of stannylene L<sub>1</sub>Sn was explored in oxidation, oxidative addition, and transmetalation reactions to form in particular the corresponding gallium and aluminum derivatives.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules29020325/s1>, NMR spectra, crystal structure refinements, and computational investigations (PDF).

**Author Contributions:** Conceptualization, R.S.R. and D.M.; investigation, A.A.; X-ray structural studies: S.M.-L.; DFT calculations, J.-M.S.; writing—original draft preparation, A.A., R.S.R. and D.M.; writing—review and editing, all authors; supervision, E.M., A.R.C., A.B., T.K., R.S.R. and D.M. project

administration, R.S.R. and D.M.; funding acquisition, T.K. and R.S.R. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by the Centre National de la Recherche Scientifique (CNRS), the Université de Toulouse (UPS), ECOS-Sud Chili (No. C19E04), and FONDECYT (project No. 1200748; 1230537). A. A. acknowledges funding from Ph.D. ANID 2019 fellowship No. 21190209.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available in article and Supplementary Materials.

**Conflicts of Interest:** The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## References

1. Aly, A.A.; Brâse, S.; Gomaa, M.A.-M. Amidines: Their synthesis, reactivity, and applications in heterocycle synthesis. *Arkivoc* **2018**, *vi*, 85–138. [[CrossRef](#)]
2. Aly, A.A.; El-Din, A.M.N. Functionality of Amidines and Amidrazones. *Arkivoc* **2008**, *i*, 153–194. [[CrossRef](#)]
3. Quek, J.Y.; Davis, T.P.; Lowe, A.B. Amidine Functionality as a Stimulus-Responsive Building Block. *Chem. Soc. Rev.* **2013**, *42*, 7326–7334. [[CrossRef](#)] [[PubMed](#)]
4. Coles, M.P. Application of Neutral Amidines and Guanidines in Coordination Chemistry. *Dalton Trans.* **2006**, 985–1001. [[CrossRef](#)] [[PubMed](#)]
5. Alvarez-Rodriguez, L.; Cabeza, J.-A.; Garcia-Alvarez, P.; Polo, D. The transition-metal chemistry of amidinatosilylenes, -germylenes and -stannylenes. *Coord. Chem. Rev.* **2015**, *300*, 1–28. [[CrossRef](#)]
6. Foley, S.R.; Bensimon, C.; Richeson, D.S. Facile Formation of Rare Terminal Chalcogenido Germanium Complexes with Alkylamidates as Supporting Ligands. *J. Am. Chem. Soc.* **1997**, *119*, 10359–10363. [[CrossRef](#)]
7. Karsch, H.H.; Schlüter, P.A.; Reisky, M. Bis(Amidinate) Complexes of Silicon and Germanium. *Eur. J. Inorg. Chem.* **1998**, 433–436. [[CrossRef](#)]
8. Aubrecht, K.B.; Hillmyer, M.A.; Tolman, W.B. Polymerization of Lactide by Monomeric Sn(II) Alkoxide Complexes. *Macromolecules* **2002**, *35*, 644–650. [[CrossRef](#)]
9. Garg, P.; Dange, D.; Jones, C. S- and p-Block Dinuclear Metal(Loid) Complexes Bearing 1,4-Phenylene and 1,4-Cyclohexylene Bridged Bis(Amidinate) Ligands. *Eur. J. Inorg. Chem.* **2020**, *2020*, 4037–4044. [[CrossRef](#)]
10. Garg, P.; Dange, D.; Jones, C. Bulky Arene-Bridged Bis(Amide) and Bis(Amidinate) Complexes of Germanium(II) and Tin(II). *Dalton Trans.* **2021**, *50*, 9118–9122. [[CrossRef](#)]
11. Dehmel, M.; Wünsche, M.A.; Görls, H.; Kretschmer, R. Dinuclear Chlorotetrylenes of Silicon, Germanium, and Tin Based on a Backbone-Bridged Bis(Amidine). *Eur. J. Inorg. Chem.* **2021**, *2021*, 4806–4811. [[CrossRef](#)]
12. Saltarini, S.; Villegas-Escobar, N.; Martínez, J.; Daniliuc, C.G.; Matute, R.A.; Gade, L.H.; Rojas, R.S. Toward a Neutral Single-Component Amidinate Iodide Aluminum Catalyst for the CO<sub>2</sub> Fixation into Cyclic Carbonates. *Inorg. Chem.* **2021**, *60*, 1172–1182. [[CrossRef](#)] [[PubMed](#)]
13. Yakovenko, M.V.; Cherkasov, A.V.; Fukin, G.K.; Cui, D.; Trifonov, A.A. Lanthanide Complexes Coordinated by a Dianionic Bis(Amidinate) Ligand with a Rigid Naphthalene Linker. *Eur. J. Inorg. Chem.* **2010**, *2010*, 3290–3298. [[CrossRef](#)]
14. Osorio Meléndez, D.; Castro-Osma, J.A.; Lara-Sánchez, A.; Rojas, R.S.; Otero, A. Ring-Opening Polymerization and Copolymerization of Cyclic Esters Catalyzed by Amidinate Aluminum Complexes. *J. Polym. Sci. Part A Polym. Chem.* **2017**, *55*, 2397–2407. [[CrossRef](#)]
15. Nimitsiriwat, N.; Gibson, V.C.; Marshall, E.L.; White, A.J.P.; Dale, S.H.; Elsegood, M.R.J. Tert-Butylamidinate Tin(II) Complexes: High Activity, Single-Site Initiators for the Controlled Production of Polylactide. *Dalton Trans.* **2007**, 4464. [[CrossRef](#)] [[PubMed](#)]
16. Phomphrai, K.; Pongchan-o, C.; Thumrongpatanaraks, W.; Sangtrirutnugul, P.; Kongsaree, P.; Pohmakotr, M. Synthesis of High-Molecular-Weight Poly( $\epsilon$ -Caprolactone) Catalyzed by Highly Active Bis(Amidinate) Tin(II) Complexes. *Dalton Trans.* **2011**, *40*, 2157–2159. [[CrossRef](#)] [[PubMed](#)]
17. Lentz, N.; Sodreau, A.; Acuña, A.; Mallet-Ladeira, S.; Maerten, E.; Sotiropoulos, J.-M.; Rojas Guerrero, R.S.; Madec, D. Synthesis and structures of homoleptic germylenes and stannylenes with sulfonimidamide ligands. *Dalton Trans.* **2023**, *52*, 6841–6846. [[CrossRef](#)]
18. Foley, S.R.; Zhou, Y.; Yap, G.P.A.; Richeson, D.S. Synthesis of M<sup>II</sup>[N(SiMe<sub>3</sub>)<sub>2</sub>][Me<sub>3</sub>SiNC(<sup>t</sup>Bu)NSiMe<sub>3</sub>] (M = Sn, Ge) from Amidinate Precursors: Active Catalysts for Phenyl Isocyanate Cyclization. *Inorg. Chem.* **2000**, *39*, 924–929. [[CrossRef](#)]
19. Chlupatý, T.; Růžičková, Z.; Horáček, M.; Alonso, M.; De Proft, F.; Kampová, H.; Brus, J.; Růžička, A. Oxidative Additions of Homoleptic Tin(II) Amidinate. *Organometallics* **2015**, *34*, 606–615. [[CrossRef](#)]

20. Nakata, N.; Hosoda, N.; Takahashi, S.; Ishii, A. Chlorogermynes and -Stannylenes Stabilized by Diimidodisulfinate Ligands: Synthesis, Structures, and Reactivity. *Dalton Trans.* **2018**, *47*, 481–490. [[CrossRef](#)]
21. Okazaki, R.; Tokitoh, N. Heavy Ketones, the heavier Element Congeners of a Ketone. *Acc. Chem. Res.* **2000**, *33*, 625–630. [[CrossRef](#)] [[PubMed](#)]
22. Foley, S.R.; Yap, G.P.A.; Richeson, D.S. Oxidative Addition to M(II) (M = Ge, Sn) Amidinate Complexes: Routes to Group 14 Chalcogenolates with Hypervalent Coordination Environments. *J. Chem. Soc. Dalton Trans.* **2000**, *10*, 1663–1668. [[CrossRef](#)]
23. Ahmet, I.Y.; Hill, M.S.; Raithby, P.R.; Johnson, A.L. Tin Guanidinato Complexes: Oxidative Control of Sn, SnS, SnSe and SnTe Thin Film Deposition. *Dalton Trans.* **2018**, *47*, 5031–5048. [[CrossRef](#)] [[PubMed](#)]
24. Ahmet, I.Y.; Thompson, J.R.; Johnson, A.L. Oxidative Addition to Sn(II) Guanidinate Complexes: Precursors to Tin(II) Chalcogenide Nanocrystals. *Eur. J. Inorg. Chem.* **2018**, *2018*, 1670–1678. [[CrossRef](#)]
25. Foley, S.R.; Yap, G.P.A.; Richeson, D.S. Formation of Novel Tetrasulfido Tin Complexes and Their Ability to Catalyze the Cyclotrimerization of Aryl Isocyanates. *Organometallics* **1999**, *18*, 4700–4705. [[CrossRef](#)]
26. Zhou, Y.; Richeson, D.S. Multiple Bonds between Sn and S: Synthesis and Structural Characterization of (CyNC(<sup>t</sup>Bu)NCy)<sub>2</sub>SnS and [(CyNC(Me)NCy)<sub>2</sub>Sn(μ-S)]<sub>2</sub>. *J. Am. Chem. Soc.* **1996**, *118*, 10850–10852. [[CrossRef](#)]
27. Sen, N.; Pal, S.; Khade, V.V.; Khan, S. Cyclic Four-Membered Stanna Thio and Seleno Compounds from 2-Aminopyridinato Stannylenes. *Eur. J. Inorg. Chem.* **2019**, *2019*, 4450–4454. [[CrossRef](#)]
28. Shan, Y.-L.; Leong, B.-X.; Xi, H.-W.; Ganguly, R.; Li, Y.; Lim, K.H.; So, C.-W. Reactivity of an Amidinato Silylene and Germylene toward Germanium(II), Tin(II) and Lead(II) Halides. *Dalton Trans.* **2017**, *46*, 3642–3648. [[CrossRef](#)]
29. Watson, I.C.; Ferguson, M.J.; Rivard, E. Zinc-Mediated Transmetalation as a Route to Anionic N-Heterocyclic Olefin Complexes in the p-Block. *Inorg. Chem.* **2021**, *60*, 18347–18359. [[CrossRef](#)]
30. Brazeau, A.L.; DiLabio, G.A.; Kreisel, K.A.; Monillas, W.; Yap, G.P.A.; Barry, S.T. Theoretical and experimental investigations of ligand exchange in guanidinate ligand systems for group 13 metals. *Dalton Trans.* **2007**, 3297–3304. [[CrossRef](#)]
31. Boeré, R.T.; Klassen, V.; Wolmershäuser, G. Synthesis of Some Very Bulky N,N'-Disubstituted Amidines and Initial Studies of Their Coordination Chemistry. *J. Chem. Soc. Dalton Trans.* **1998**, *24*, 4147–4154. [[CrossRef](#)]
32. Davidson, P.; Harris, D.; Lappert, M. Subvalent Group 4B metal alkyls and amides. Part I. The synthesis and physical properties of kinetically stable bis[bis(trimethylsilyl)methyl]- germanium(II), -tin(II), and -lead(II). *J. Chem. Soc. Chem. Dalton Trans.* **1976**, 2268–2274. [[CrossRef](#)]
33. Sheldrick, G.M. SHELXT—Integrated space-group and crystal-structure determination. *Acta Crystallogr. Sect. A* **2015**, *71*, 3–8. [[CrossRef](#)]
34. Sheldrick, G.M. Crystal structure refinement with SHELXL. *Acta Crystallogr. Sect. C* **2015**, *71*, 3–8. [[CrossRef](#)] [[PubMed](#)]
35. Van der Sluis, P.; Spek, A.L. BYPASS: An effective method for the refinement of crystal structures containing disordered solvent regions. *Acta Cryst. Sect. A* **1990**, *46*, 194–201. [[CrossRef](#)]
36. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R.; Scalmani, G.; Barone, V.; Petersson, G.A.; Nakatsuji, H.; et al. *Gaussian 16, Revision B.01*; Gaussian, Inc.: Wallingford, CT, USA, 2016.
37. Becke, A.D. Density-functional exchange-energy approximation with correct asymptotic behavior. *Phys. Rev. A* **1988**, *38*, 3098–3100. [[CrossRef](#)] [[PubMed](#)]
38. Becke, A.D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652. [[CrossRef](#)]
39. Lee, C.; Yang, W.; Parr, R.G. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B* **1988**, *37*, 785–789. [[CrossRef](#)]
40. Grimme, S. Semiempirical GGA-type density functional constructed with a long-range dispersion correction. *J. Comp. Chem.* **2006**, *27*, 1787–1799. [[CrossRef](#)]
41. Andrae, D.; Häussermann, U.; Dolg, M.; Stoll, H.; Preuss, H. Energy-adjusted ab initio pseudopotentials for the second and third row transition elements. *Theor. Chim. Acta* **1990**, *77*, 123–141. [[CrossRef](#)]
42. Fukui, K. The path of chemical reactions—The IRC approach. *Acc. Chem. Res.* **1981**, *14*, 363–368. [[CrossRef](#)]
43. Hratchian, H.P.; Schlegel, H.B. *Theory and Applications of Computational Chemistry: The First 40 Years*; Dykstra, D.C.E., Frenking, G., Kim, K.S., Scuseria, G., Eds.; Elsevier: Amsterdam, The Netherlands, 2005.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.