



**HAL**  
open science

# Towards experimental confirmation of quarkonia melting in Quark Gluon Plasma: A review of recent measurements of quarkonia production in relativistic heavy-ion collisions

Kara R Mattioli

► **To cite this version:**

Kara R Mattioli. Towards experimental confirmation of quarkonia melting in Quark Gluon Plasma: A review of recent measurements of quarkonia production in relativistic heavy-ion collisions. *Symmetry*, 2024, 16 (2), pp.225. 10.3390/sym16020225 . hal-04441033

**HAL Id: hal-04441033**

**<https://hal.science/hal-04441033>**

Submitted on 6 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Towards experimental confirmation of quarkonia melting in Quark Gluon Plasma: A review of recent measurements of quarkonia production in relativistic heavy-ion collisions

Kara R. Mattioli <sup>1</sup> 

<sup>1</sup> Laboratoire Leprince Ringuet, Palaiseau, France; kara.mattioli@llr.in2p3.fr

\* Correspondence: kara.mattioli@llr.in2p3.fr

**Abstract:** The dissociation, or “melting”, of heavy quarkonia states due to color charge screening is a predicted signature of the Quark Gluon Plasma (QGP) formation, with a quarkonium state predicted to dissociate when the temperature of the medium is higher than the binding energy of the quarkonium state. A conclusive experimental observation of quarkonium melting coupled with a detailed theoretical understanding of the melting mechanism would enable the use of quarkonia states as temperature probes of the QGP, a long-sought goal in the field of relativistic heavy-ion collisions. However, the interpretation of quarkonia suppression measurements in heavy-ion collisions is complicated by numerous other cold nuclear matter effects which also result in the dissociation of bound quarkonia states. A comprehensive understanding of these cold nuclear matter effects is therefore needed in order to correctly interpret quarkonia production measurements in heavy ion collisions and to observe the melting of quarkonium states experimentally. In this review, recent measurements of quarkonia production in  $pA$  and  $AA$  collisions and their state-of-the-art theoretical interpretations will be discussed, as well as the future measurements needed to further the knowledge of cold nuclear matter effects and realize a measurement of quarkonia melting in heavy ion collisions.

**Keywords:** quarkonia suppression; sequential suppression; cold nuclear matter effects

## 1. Introduction

Relativistic heavy-ion collisions provide the opportunity to study one of the rarest forms of matter ever created - the quark-gluon plasma (QGP). Quarks and gluons, which are normally tightly bound within hadronic bound states of protons and neutrons in nuclei, can be liberated in high-energy collisions of heavy nuclei and result in the formation of the dense partonic plasma phase [1]–[3]. The QGP is interesting for many reasons in addition to being a novel state of QCD matter; it is predicted to have existed shortly after the Big Bang, therefore studying it can provide insight into the early formation of the universe, and it also behaves as a nearly perfect hydrodynamic fluid [4][5]. Understanding the formation, dynamics, and behavior of the QGP, as with any unique state of matter, would result in significantly improving our understanding of the fundamental physics of the strong nuclear force and how it gives rise to the properties of nuclei.

A key property to understand the behavior of any medium is its temperature dependence. The idea to use heavy quarkonia, bound states of charm-anticharm ( $c\bar{c}$ , “charmonia”) or beauty-antibeauty ( $b\bar{b}$ , “bottomonia”) quarks, as temperature probes of the QGP came from a landmark paper by T. Matsui and H. Satz [6]. In their paper, Matsui and Satz proposed that quarkonia states would “melt”, or be prevented from forming, in a hot deconfined nuclear medium because the free QCD color charges would screen the potential between the two heavy quarks in a mechanism analogous to Debye screening. The screening was predicted to occur when the temperature of the medium exceeded that of the binding energy of the quarkonium state [6]. Since different quarkonium states have different binding energies, if one could experimentally observe the melting of quarkonia in

**Citation:** Mattioli, K. Towards experimental confirmation of quarkonia melting in Quark Gluon Plasma: A review of recent measurements of quarkonia production in relativistic heavy-ion collisions. *Symmetry* **2023**, *1*, 0. <https://doi.org/>

Received:

Revised:

Accepted:

Published:

**Copyright:** © 2024 by the authors. Submitted to *Symmetry* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

a hot nuclear medium, one could infer the temperature of the medium itself. The possibility of experimentally measuring the temperature of the QGP has made the observation of the “melting”, or dissociation of heavy quarkonia states due to color charge screening, a golden observable in the field of relativistic heavy-ion physics. The idea is that definitive experimental observation of quarkonia melting would enable the use of quarkonium states as temperature probes of the quark-gluon plasma.

The straightforward picture of quarkonia melting due to color-charge screening in high-energy nucleus-nucleus (“AA”) collisions has evolved over the years as experimental and theoretical studies have revealed that quarkonia production in high-energy collisions is much more complex than initially thought. From the theoretical side, quarkonium production in  $pp$  collisions is still not yet completely understood theoretically [7]-[9], and recent theoretical work revealing an imaginary component to the quarkonium potential complicates the picture of color-charge screened quarkonium dynamics proposed by Matsui and Satz [11]-[13]. From the experimental side, signatures previously thought to only occur in the presence of QGP have been observed even in small collision systems [14], raising questions about whether other QGP effects, including quarkonia melting, could be taking place in small systems as well. Additionally, other effects besides the proposed melting mechanism, commonly called Cold Nuclear Matter (CNM) effects, can cause the dissociation of quarkonia states, and these effects must be fully understood and quantified in order to isolate the signatures of quarkonia dissociation due to melting in the QGP [7]. Therefore, there are a few requirements that must be met before concluding that quarkonia melting has been experimentally observed. First, it must be proved that no other mechanisms (such as CNM effects) are responsible for the quarkonium dissociation. Second, it must be shown that the suppression depends on the binding energy of the measured quarkonia. The first criterion remains much more difficult to satisfy than the second. Disentangling dissociation by melting from dissociation due to other effects remains a challenge.

In this review, recent measurements of quarkonia production in  $pA$  and  $AA$  collisions will be discussed to highlight the current state of our experimental knowledge of quarkonium suppression in nuclear systems. Photoproduction of quarkonia and the production of quarkonia from beauty hadron decays will not be discussed, as these production mechanisms are different than that for prompt quarkonia and are also not subject to the predicted melting effects. For details on quarkonium production in vacuum ( $pp$  collisions) and comprehensive theoretical introductions, the reader is referred to excellent reviews on these topics [7]-[9] [15]-[17]. In Section 2, experimental methods for the reconstruction of quarkonia states and the common observables measured to probe quarkonia suppression are briefly reviewed. The discussion of recent experimental results begins in Section 3 with recent results in nucleus-nucleus collisions, where the clearest observations of quarkonium suppression are observed. Then in Section 4 recent results from proton-nucleus collisions are reviewed, which tend to complicate the interpretation of the results in nucleus-nucleus collisions.

## 2. Experimental methods and observables

### 2.1. Reconstruction of quarkonia states

In order to measure the predicted sequential suppression mechanism discussed in the previous section, one must measure both ground and excited states of the charmonia and bottomonia resonances. For charmonia, the states most experimentally accessible in heavy ion collisions are the  $J/\psi(1S)$ ,  $\chi_c(1P)$ , and  $\psi(2S)$ . For bottomonia, the corresponding most accessible states are the  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$ . With the exception of the  $\chi_c$ , the states are typically measured in the dielectron or dimuon decay channel, depending on the experiment. The dilepton decay channels produce clean experimental signals, as they produce two charged particle tracks in a detector which can be precisely measured. Electrons and muons are identified experimentally with dedicated particle identification or muon detectors, respectively.

Fewer experimental measurements exist for the  $\chi_c$  and  $Y(3S)$  states in heavy ion collisions as these states are particularly challenging to measure experimentally. The  $\chi_c$  state is typically reconstructed in the radiative decay  $\chi_{cn} \rightarrow J/\psi\gamma$ , which suffers from significant combinatorial background due to  $\pi^0 \rightarrow \gamma\gamma$  decays. The  $Y(3S)$  state is extremely heavy and therefore rarely produced. It is particularly difficult to measure in heavy ion collisions as the 3S state is predicted to be most affected by dissociation due to cold and hot nuclear matter effects. Therefore, large experimental luminosities are needed to precisely measure the  $Y(3S)$  in nuclear collisions.

The production of quarkonia states can be “prompt”, where the quarkonium state is produced directly in a hard QCD interaction, or “non-prompt”, where the quarkonium is produced in the decay of a beauty hadron. The production of a lower-energy quarkonium state from the decay of a higher-energy quarkonium state is included within the definition of prompt quarkonia, therefore the only production mechanism for bottomonia states is prompt production. For the charmonia states, however, measuring prompt or nonprompt production probes different physics [7]. Prompt charmonia are created directly in a hard QCD interaction in the early stages of a collision, and in the case of heavy-ion collisions are therefore thought to experience the full evolution of the QGP. Non-prompt charmonia are produced in the decays of beauty hadrons, and in this production mechanism the charmonia are believed to be produced outside the QGP medium [7]. In order to probe the dissociation of quarkonia states due to the presence of a nuclear medium, prompt charmonia are the preferred probes. Experimentally, quarkonia can be measured in inclusive, prompt, or non-prompt production. In inclusive production measurements, all candidates for a particular charmonium state in a particular decay channel are measured and no distinction is made between candidates produced promptly and non-promptly. Prompt and non-prompt charmonia production measurements are performed experimentally by fitting the pseudo-decay time distribution of the charmonia candidates in order to separate candidates produced from  $b$ -hadrons, which have larger decay times, from promptly produced charmonia.

## 2.2. Observables for quarkonia suppression

Measurements of quarkonia suppression begin by first measuring the cross section of a quarkonium state in  $pA$ ,  $AA$ , or even  $pp$  collisions. Then, one must choose a baseline for which the cross-section measurement will be compared against. Typically this baseline is either the cross-section of the quarkonium state in  $pp$  collisions, or for the case of a higher-resonance quarkonium state, the cross-section of the ground or lower-resonance state measured in the same collision system.

The classic observable for quarkonium suppression is the nuclear modification factor, defined as the ratio of the quarkonium production yield in a nuclear collision system ( $pA$ ,  $AA$ ) to the cross section in  $pp$  collisions scaled by the average nuclear overlap,  $\langle T_{AA} \rangle$ , determined from a Glauber model analysis [18]:

$$R_{AA} = \frac{dN/dx}{\langle T_{AA} \rangle \times d\sigma_{pp}/dx} \quad (1)$$

The nuclear modification factor is typically measured as a function of a variable  $x$ , where  $x$  is often a kinematic variable, such as the quarkonium transverse momentum  $p_T$  or rapidity  $y$ , or a variable describing the collision centrality, such as the number of participating nucleons in a collision,  $N_{part}$ .

Another observable used to probe suppression is the double ratio of quarkonia production in  $AA$  collisions relative to that in  $pp$  collisions. This ratio is sometimes denoted by  $\rho$ :

$$\rho = \frac{N_{\psi(2S)}^{AA} / N_{J/\psi}^{AA}}{N_{\psi(2S)}^{pp} / N_{J/\psi}^{pp}} \quad (2)$$

The double ratio of quarkonia production is not as powerful as the  $R_{AA}$  as it gives only information on the relative suppression between the excited state and the ground state, rather than the absolute suppression. However, it is easier to measure experimentally as many factors cancel in the ratio, and combined with an  $R_{AA}$  measurement of the ground state, it can be used to derive the excited state  $R_{AA}$ .

### 3. Recent results in Nucleus-Nucleus Collisions


#### 3.1. Bottomonia

The suppression of bottomonia states in PbPb collisions has been studied by ALICE, ATLAS, and CMS [19]–[27]. Measurements of elliptic flow of Y mesons are beyond the scope of this review but are included in the references for the interested reader [28][29]. The most precise measurement of bottomonia state suppression in AA collisions comes from CMS, which recently measured all three Y states in PbPb collisions at  $\sqrt{s} = 5.02$  TeV [26]. The measurements, shown in Figure 1, were performed as a function of centrality and  $p_T$  and compared to several different theoretical models. A clear sequential suppression is observed for the Y states, with the Y(3S) state the most suppressed, just barely more suppressed than the Y(2S) state, and both the 2S and 3S states being significantly suppressed compared to the Y(1S) state. The sequential suppression is observed to be stronger, with the higher Y states being more suppressed, in central collisions. The suppression measured as a function of  $p_T$  is constant within the experimental uncertainties [26].

The CMS data were compared to several theoretical models including a coupled Boltzmann model [30], an open quantum system (OQS) model [31], the Heidelberg model [32], the Comover Interaction Model (CIM) [33], and the TAMU transport model [34]. These models represent a wide variety of approaches to describing the dissociation of quarkonia states in nucleus-nucleus collisions and their key features will be briefly summarized here. The coupled Boltzmann and TAMU models are both transport models which describe the quarkonia production and dissociation in QGP using kinetic rate equations. Typically starting from a space- and momentum-integrated Boltzmann equation, the rate equation contains two transport coefficients that describe the quarkonia dynamics in a nuclear medium [35]. The first coefficient is the quarkonia dissociation rate and the second is the quarkonia yield in the equilibrium limit, which depends on the medium's evolution as a function of temperature. Different transport model calculations vary in their approach to obtaining the two transport coefficients and the medium temperature response. In the TAMU model that is compared to the CMS data in Figure 2, a microscopic model [36] is used to compute the modified quarkonium binding energies in the QGP, and a lattice QCD Equation of State is used to compute the medium evolution as a function of temperature [34]. Both transport models shown in Figure 2 include contributions from uncorrelated bottomonium recombination, where a  $b$ - and  $\bar{b}$ -quark pair produced in different initial partonic hard scattering events could recombine to form an Y meson. The coupled Boltzmann model instead uses two coupled Boltzmann equations, one for an unbound  $Q\bar{Q}$  pair ( $Q$  denoting a heavy quark) and one for bound  $Q\bar{Q}$  quarkonium states [30]. The coupled approach notably provides access to studying the effect of *correlated* quarkonium regeneration, in which a  $b$ - and  $\bar{b}$ -quark pair produced during the dissociation of a bottomonium state could once again recombine to form a new bottomonium state [30]. The Open Quantum System (OQS) approach treats the quarkonium as a quantum system that interacts with its surrounding environment, in this case the QGP [31][37]–[39]. The reduced density matrix of the quarkonium system is computed and its equation of motion, the so-called “Master equation”, is determined [9]. The OQS formalism provides a fully quantum description of the quarkonium system, including its dynamics out of thermal equilibrium with the QGP [31] [39] and accounting for quantum correlations between the quarkonium system and the QGP medium [9]. The OQS model compared to the CMS data in Figure 2 demonstrates a recent example of combining the OQS framework with an effective field theory that describes the non-relativistic nature of quarkonium, in this case potential non-relativistic QCD (NRQCD) [31] [39]. The Heidelberg model computes

the bottomonium wave functions by solving a temperature-dependent radial stationary Schrodinger equation with an effective potential that includes both real and imaginary components to account for the color screening and damping effects, respectively [32]. Finally, the Comover Interaction Model describes the dissociation of quarkonia states via their interactions with other “co-moving” particles with similar rapidities in a cold or hot nuclear medium [33]. A key parameter in this model is the comover density, which is obtained from measurements of the particle multiplicity as a function of rapidity [33].

The double ratio of the  $Y(3S)$  to the  $Y(2S)$  production yield in PbPb collisions to that in  $pp$  collisions provides an excellent testing ground for the theoretical models [26][40]. An example of this can be seen in Figure 2, where the measured double ratio and its theoretical predictions are shown as a function of the collision centrality and the quarkonium  $p_T$ . All models except for the Coupled Boltzmann model reasonably describe the data. The OQS predictions are closest to the measured data points, being directly on top of them in several cases. The Heidelberg model also does a good job but overpredicts the ratio at high  $p_T$ . The TAMU and CIM models are consistently slightly below the data points, and the Coupled Boltzmann model is significantly below the data. It would be interesting to see in the future if perhaps adjustments of the relative yields of correlated and uncorrelated bottomonium regeneration in the Coupled Boltzmann model could give a better agreement with the CMS data.




**Figure 1.** The  $R_{AA}$  of the  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$  states measured by CMS as a function of (a) centrality and (b) quarkonia  $p_T$ . Figures are by the CMS Collaboration, obtained from [26] under CERN copyright, and licensed under CC BY 4.0. Images are cropped from the original versions.

The  $R_{AA}$  of  $Y$  states has also recently been measured at RHIC by STAR [41]. Figure 3 shows the STAR results as a function of centrality compared to results from CMS and to two theoretical models already introduced above, the TAMU transport model [34] and the open quantum system model using potential non-relativistic QCD [37]-[39]. It is interesting to observe that the  $Y(1S)$  suppression at RHIC is consistent with that observed at the LHC, despite the significantly different center of mass energies. In contrast with the  $Y(1S)$ , the  $Y(2S)$  appears to be slightly more suppressed at RHIC than at the LHC, but the STAR measurement is still consistent with the CMS data within the experimental uncertainties. The theoretical models agree well with the CMS data and the STAR  $Y(2S)$  data, however both models predict a smaller  $Y(1S)$  suppression at RHIC than what is observed in the data. These recent measurements in the bottomonia sector confirm a sequential suppression pattern but do not yet single out a specific theoretical model that can describe the relative excited state suppression and the suppression dynamics as a function of collision energy.


### 3.2. Charmonia

#### 3.2.1. Ground state suppression


The ground state of the charmonia resonances, the  $J/\psi$ , has been studied extensively in relativistic heavy-ion collisions. Measurements of  $J/\psi$  hadroproduction have been performed in PbPb collisions at  $\sqrt{s} = 2.76$  TeV ([42]-[48]) and 5.02 TeV ([49] - [56]), in XeXe collisions at  $\sqrt{s} = 5.44$  TeV ([57]), in AuAu collisions at  $\sqrt{s} = 200$  GeV ([58]-[62]), and in



**Figure 2.** The  $R_{AA}$  of the  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$  states measured by CMS compared to several theoretical models, as a function of (a) centrality and (b) quarkonia  $p_T$ . See text for details of the theoretical models. Figures by the CMS Collaboration, obtained from [40] under CERN copyright. The size of the images is reduced with respect to the original versions.



**Figure 3.** The  $R_{AA}$  of  $Y(1S)$  (top) and  $Y(2S)$  (bottom) mesons measured by STAR and CMS, compared to theoretical models. See text for details of the theoretical models. Figure by the STAR collaboration, obtained from [41] under American Physical Society copyright and re-used with permission. Image is enlarged with respect to the original version.




**Figure 4.** The nuclear modification factor  $R_{AA}$  for prompt  $J/\psi$  production measured by ALICE [56], CMS [52], and ATLAS [51] in PbPb collisions at  $\sqrt{s} = 5.02$  TeV, measured in three different centrality bins: **a)** 0-10 %, **b)** 10-30 % (ATLAS 20-40 %), **c)** 30-50 % (ALICE), 30-100 % (CMS), 40-80 % (ATLAS). Figures are by the ALICE Collaboration, obtained from [56] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the images is reduced with respect to the original versions.

CuCu collisions at  $\sqrt{s} = 200$  GeV [59], to name the most recent measurements. Recent flow [63]–[68] and polarization [69] measurements are beyond the scope of this review but are included in the references for the interested reader. Several measurements in non-identical, or AB, nucleus-nucleus collisions have also been recently performed, in  $^3\text{HeAu}$  collisions by PHENIX [70] and in PbNe collisions by LHCb [71]. As previously mentioned in the preceding sections, the discussion here will focus on measurements of prompt charmonia production when available for the aforementioned datasets as prompt measurements are most sensitive to probing dissociation effects due to the nuclear medium. The most recent measurements from RHIC and the LHC, summarizing the current state of the art for  $J/\psi$  suppression measurements, will be reviewed and discussed along with some of the most common theoretical approaches used to interpret them.

The most recent measurement of prompt  $J/\psi$  production in high-energy nucleus-nucleus collisions was performed by ALICE in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV [56], where the ALICE results were also compared to measurements by CMS and ATLAS in PbPb collisions at the same collision energy [51] [52]. The  $R_{AA}$  results as a function of  $J/\psi$   $p_T$  at midrapidity for three different centrality bins are shown in Figure 4. The ATLAS and CMS results span the high- $p_T$  region, while the ALICE results cover the low- $p_T$  region down to 1.5 GeV/c. All three measurements are consistent in the  $p_T$  range where they overlap, resulting in a continuous description of the  $J/\psi$   $R_{AA}$  from low to high  $p_T$ . At low  $p_T$ , the  $R_{AA}$  rises dramatically, particularly in the most central collisions where it is even larger than one. In the next centrality bin from 10-30%, the central values in the two lowest  $p_T$  bins are slightly lower than in the 0-10% centrality bin but are still consistent with the first centrality bin within the uncertainties. The most noticeable difference is in the  $p_T$  range between 5-10 GeV/c, where the suppression in 10-30% centrality is slightly less than that seen in the 0-10% bin, consistent with previous measurements that found the  $J/\psi$  meson suppression to be strongest in the most central collisions. In the final centrality bin, interestingly the  $J/\psi$  suppression at low  $p_T$  is consistent within the uncertainties with the suppression observed by ATLAS and CMS at high  $p_T$ .




Moving to other collision systems besides PbPb, the inclusive  $J/\psi$   $R_{AA}$  has also been recently measured in XeXe collisions by ALICE [57] and in AuAu collisions by STAR [62]. Figure 5 shows the  $J/\psi$   $R_{AA}$  measured as a function of centrality in XeXe and PbPb collisions. The XeXe data set has much larger uncertainties and fewer measurement bins than the PbPb data due to the smaller integrated luminosity available. However, it is still interesting to see that the  $J/\psi$  suppression in XeXe is largely consistent with that observed in PbPb, despite the fact that the XeXe system is a much lighter collision system ( $A_{Xe} = 129$  and  $A_{Pb} = 208$ ). The data is compared to the TAMU transport model [72] [73] which was previously introduced for the  $Y$  states in Section 3.1. The transport model predictions describe the XeXe data more accurately than the PbPb data, which they slightly underpredict for the mid-centrality range.



**Figure 5.** The  $R_{AA}$  for  $J/\psi$  mesons in XeXe collisions compared to PbPb collisions, measured as a function of centrality. Figure by the ALICE Collaboration, obtained from [57] under CERN copyright, and licensed under CC BY 4.0. The size of the image is enlarged with respect to the original version.

In AuAu collisions, the most recent measurement of  $J/\psi$  suppression was performed by the STAR collaboration, which measured the  $J/\psi$   $R_{AA}$  as a function of  $p_T$  and centrality [62]. The results shown as a function of  $p_T$  and binned in five different centrality ranges are shown in Figure 6 and compared to previous results from RHIC and LHC experiments. The  $J/\psi$   $R_{AA}$  measured at RHIC is relatively constant as a function of  $p_T$  within the experimental uncertainties. The difference between the RHIC and LHC results is most prominent at low  $p_T$ , where the ALICE  $R_{AA}$  value is much larger than that measured by STAR. When comparing to the CMS measurements, the RHIC  $R_{AA}$  measurement is slightly above the CMS measurement for  $p_T$  between 8-10 GeV/c but is also consistent with the CMS result within the uncertainties. Comparisons with several theoretical models are presented. The Tsinghua [75] [76] and TAMU [72]-[74] models are both transport models employing a kinetic rate equation approach to describe the charmonium dissociation and regeneration. The transport models tend to agree with each other at low  $p_T$ , then diverge at high  $p_T$  but both are consistent with the data. The remaining collisional dissociation models are only shown in the most central AuAu bin (0-20%), and are based on NRQCD calculations computed with two different values of the charmonium formation time (" $t_{max}$ " and " $t_{min}$ ") [77]. The precision of the experimental data cannot yet distinguish between the two formation time scenarios, but this might be possible in the future [78]. All of the models also include cold nuclear matter effects on the  $J/\psi$  production.


The enhancement of the  $J/\psi$   $R_{AA}$  at low  $p_T$  has been attributed to the production of  $J/\psi$  mesons from the recombination of  $c$  and  $\bar{c}$  pairs in the extremely hot nuclear medium formed at the LHC [7][9]. At high  $p_T$ , the exact origin of the  $J/\psi$  suppression is not yet understood as multiple dissociation mechanisms could be involved, as will be discussed. The picture becomes murkier when moving away from very central collisions towards more peripheral collisions, where the extent of the produced QGP medium and possible interplay with other CNM effects is relatively unknown. The ATLAS and ALICE measurements of



**Figure 6.** The  $J/\psi$   $R_{AA}$  measured by STAR as a function of  $p_T$  and binned in centrality. AuAu results from RHIC are compared with PbPb results from the LHC. Figure by the STAR Collaboration, obtained from [62] and licensed under [CC BY 4.0](#). The figure is enlarged with respect to the original version.

the  $J/\psi$   $R_{AA}$  were compared to several theoretical predictions, shown in Figures 7 and 8. The ALICE measurements were compared to three different models: the Statistical Hadronization Model with charm quarks (SHMc) [79], the Boltzmann transport model (BT) [80][81] and a charmonium dissociation model [82] [83]. The Boltzmann transport model has already been described in the previous section. In the SHMc model, the charm quarks are assumed to thermalize in the QGP medium and the yields of the charm hadrons are computed with statistical mechanics using a grand canonical partition function [79] [84]. A key component of the SHMc model is that quarkonia states are only formed during hadronization at the phase transition of the QGP (at the chemical freeze-out temperature  $T_{CF}$ ) [84][85]. This approach differs from the other models shown in Figure 7, which assume that at least some quarkonia bound states remain intact and can interact with the QGP medium despite the existence of color screening effects [80]-[83]. The dissociation model shown in Figure 7 specifically focuses on describing the dynamics of high- $p_T$   $J/\psi$ s in a QGP, and includes  $J/\psi$  dissociation contributions from the expected color-charge screening and from collisions of the  $J/\psi$  with the medium constituents [82]. Despite the significantly different physical pictures described by the two models, both the Boltzmann transport model and the SHMc model describe the low  $p_T$  ALICE data particularly well, while at high  $p_T$  the dissociation model agrees extremely well with the ALICE data in very central collisions. In semi-central collisions, neither the Boltzmann transport or SHMc can fully describe the trend of the data.

The ATLAS measurements were compared to color-screening models [82][86] and models involving  $J/\psi$  suppression via parton energy loss [87][88]. The first color-screening model is the same model that is referred to as the “Dissociation” model in the ALICE plot in Figure 7, which accounts for  $J/\psi$  dissociation due to color screening and collisions in the medium [82]. The second color-screening model that is compared to the ATLAS data focuses particularly on dissociation due to color-exchanges between the  $Q\bar{Q}$  state and the QGP medium, which the authors describe as a nuclear absorption-like effect, in addition to the expected melting from color-charge screening [86]. Therefore, it is important to clarify that the “color-screening” models compared to the ATLAS data do not only include quarkonia dissociation due to color-screening, but they also include dissociation due to non-melting effects as well. The first energy loss model referred to in the ATLAS




**Figure 7.** The prompt  $J/\psi$   $R_{AA}$  measured by ALICE as a function of  $p_T$  and compared to predictions from several theoretical models. The data and theory comparisons are shown for two centrality bins: (a) 0–10% and (b) 30–50%. Figures are by the ALICE Collaboration, obtained from [56] under CERN copyright, and licensed under CC BY 4.0. The size of the images is reduced with respect to the original versions.

figure describes the quarkonia suppression in AA collisions as being predominantly due to radiative energy loss of the charmonia in the QGP [87]. The radiative energy loss model is based on the idea that quarkonia suppression in QGP could have the same underlying origin as the phenomenon of jet “quenching”, in which jets, high-energy sprays of particles produced from the hadronization of a single quark or gluon, are suppressed or “quenched” in the QGP compared to  $pp$  collisions [87]. Phenomenologically, the quenching (or suppression) depends on the original  $p_T$  of the quark or gluon (or charmonium state) and an effective color factor that accounts for the different radiation patterns between quarks and gluons [87]. It is indeed interesting that this quenching model gives almost identical predictions as the first color-screening model, despite the very different underlying theoretical mechanisms, and both are in very good agreement with the experimental data. The second energy loss model compared to the ATLAS data shares the idea that radiative energy loss is the dominant mechanism explaining the  $J/\psi$  suppression, but rather than comparing the  $J/\psi$  suppression with the quenching of jets, in this approach it is instead compared with the suppression of individual hadrons [88]. It is also found in this model that the  $J/\psi$  suppression and hadron suppression can be described by the same underlying mechanism. One caveat to the energy loss models is that they are expected to be most relevant at high  $p_T$  ( $> \approx 10$  GeV), therefore they are not expected to describe the observed  $J/\psi$  suppression at low  $p_T$  where melting could be playing an important role. However, it is certainly interestingly to observe that the ATLAS data at high  $p_T$  are consistent with both dissociation due to color screening and suppression due to energy loss, and that no single model in Figures 7 and 8 can describe the suppression over the full measurement range.

### 3.2.2. Excited state suppression

There are far fewer measurements of the charmonia excited states in relativistic heavy-ion collisions. Several measurements exist for the  $\psi(2S)$  in PbPb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV [45][48], and 5.02 TeV ([49][51][52][89]). The  $\chi_c$  state has never been measured in any nucleus-nucleus collision.

The most recent measurement comparing the suppression of  $\psi(2S)$  to  $J/\psi$  mesons was performed by the ALICE collaboration at  $\sqrt{s_{NN}} = 5.02$  TeV [89]. The suppression factor  $R_{AA}$  was measured as a function of  $p_T$  and centrality and compared to CMS results measured at the same center of mass energy [52]. Figure 9 shows the  $R_{AA}$  of  $J/\psi$  and  $\psi(2S)$  mesons measured by ALICE and CMS as a function of the quarkonia transverse momentum  $p_T$ . Note that the ALICE results are for inclusive  $J/\psi$  and  $\psi(2S)$ , while the CMS results are for prompt  $J/\psi$  and  $\psi(2S)$ . The ALICE and CMS results nicely complement each other, with the ALICE results covering the extremely low  $p_T$  region and the CMS results spanning




**Figure 8.** The prompt  $J/\psi$   $R_{AA}$  measured by ATLAS as a function of  $p_T$  and compared to predictions from theoretical models. Figure by the ATLAS Collaboration, obtained from [51] under CERN copyright, and licensed under [CC BY 4.0.](https://creativecommons.org/licenses/by/4.0/) Figure is enlarged with respect to original version.

the high  $p_T$  region up to  $p_T = 30$  GeV/c. The data clearly shows that the  $\psi(2S)$  has a lower  $R_{AA}$  across the entire  $p_T$  range measured compared to the  $J/\psi$ , indicating that the  $\psi(2S)$  is indeed more suppressed than the  $J/\psi$  in PbPb collisions. The data are compared to theoretical predictions based on the TAMU transport model, which was previously introduced in Section 3.1. The model calculations describe the data quite well, in particular accurately capturing the rise at low  $p_T$  attributed to charmonium recombination.

The  $R_{AA}$  of prompt  $J/\psi$  and  $\psi(2S)$  has also been measured as a function of the collision centrality by both ALICE and CMS [52][89]. The measured  $R_{AA}$  distributions are shown in Figure 10, with the CMS measurement shown on the left and the ALICE measurement shown on the right. When comparing the two measurements as a function of centrality, it is important to note that the experiments probe different charmonia  $p_T$  regions. This explains why the two measurements are consistent with each other when integrated over centrality and plotted as a function of  $p_T$ , but appear to disagree when integrated over  $p_T$  and plotted as a function of centrality. Nevertheless, although the absolute value of  $R_{AA}$  for a given value of  $N_{part}$  cannot be directly compared between the two measurements, the trends are still interesting. The ALICE measurement, which has a lower charmonia  $p_T$  range than the CMS measurement, flattens out towards the most central (highest  $\langle N_{part} \rangle$ ) collisions. The CMS results, however, show a generally decreasing trend of the  $R_{AA}$  towards the most central collisions for prompt  $J/\psi$ . The  $\psi(2S)$  results in the CMS measurement have larger uncertainties, so the trend with centrality is not as clear, but they could hint at a possible decrease with centrality as well.

The double ratio of  $\psi(2S)$  to  $J/\psi$  yields in PbPb collisions to the ratio in  $pp$  collisions has been measured by ALICE, ATLAS, and CMS at  $\sqrt{s} = 5.02$  TeV [49][51][89]. The ALICE measurements of the double ratio as a function of  $p_T$  indicate that the  $\psi(2S)$  is suppressed by roughly a factor of 40-50% in PbPb collisions compared to  $pp$  collisions [89]. The CMS measurement hints at a slightly larger suppression at midrapidity, but in the most forward rapidity bin probed,  $1.6 < |y| < 2.4$  the suppression is consistent with that observed by ALICE [49][89]. ATLAS measured the double ratio as a function of centrality for both prompt and non-prompt charmonia. The double ratio for the non-prompt charmonia was observed to be consistent with one, supporting the interpretation that the non-prompt charmonia, which originate from  $b$ -hadron decays, hadronize outside the medium and are therefore unaffected by the suppression mechanisms responsible for dissociating prompt charmonia in the medium [51].

A key missing component in the picture of sequential charmonium suppression in AA collisions is the  $\chi_c(1P)$  state. With a binding energy in between that of the  $J/\psi$  and




**Figure 9.** The nuclear modification factor  $R_{AA}$  for  $J/\psi$  and  $\psi(2S)$  mesons in PbPb collisions at  $\sqrt{s} = 5.02$  TeV, as a function of the quarkonium transverse momentum  $p_T$ . The ALICE data is inclusive mesons, while the CMS data is prompt mesons. The figure and ALICE data is from [89], while the CMS data is from [52]. Figure by the ALICE Collaboration, obtained from [89] under CERN copyright, and licensed under CC BY 4.0.. Figure is enlarged with respect to original version.

the  $\psi(2S)$ , a measurement of its suppression is needed to fully confirm the sequential suppression of charmonia states. As with the Y states, no single model can fully describe all of the experimental data on the charmonia, which likely indicates, as the variety of available models also suggests, that multiple mechanisms contribute simultaneously to charmonia dissociation in the QGP.

#### 4. Recent results in $pA$ collisions

##### 4.1. Bottomonia

Bottomonia production in  $pA$  collisions has been predominantly studied at the LHC, with measurements reported by all LHC experiments in  $pPb$  collisions [90]-[95]. The most recent measurement of the nuclear modification of bottomonia states at the LHC was performed by CMS [95]. Figure 11 shows the  $R_{pPb}$  for the  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$  states measured as a function of rapidity for  $p_T < 30$  GeV/c. A sequential suppression of the upsilon states is observed with the  $Y(3S)$  being more suppressed than the  $Y(2S)$ , which is more suppressed than the  $Y(1S)$ . The sequential suppression is flat within the experimental uncertainties as a function of the quarkonia  $p_T$  and  $y$ . The data in Figure 11 are compared to predictions from the Comover Interaction Model (CIM) [33] using two different sets of nuclear Parton Distribution Functions (nPDFs), nCTEQ15 [96] and EPS09 at leading order (LO) [97]. The CIM was previously introduced in Section 3.1 and describes the quarkonia dissociation via collisions with co-moving particles at similar rapidities as the quarkonium [33]. The use of different nPDFs is to test different descriptions of the initial nuclear state. Obtained from fits to a wide variety of data, nPDFs quantify the difference in the partonic structure of a bound nucleus compared to a hypothesis of the partonic structure in  $A$  non-interacting protons and neutrons, where  $A$  is the atomic mass number. If a bound nucleus were simply a superposition of non-interacting protons and neutrons, the nPDFs would not show significant modification relative to the  $A$ -scaled proton PDFs. However, indeed studies of vast amounts of experimental data have revealed that nuclei are not simple superpositions of non-interacting hadrons and have their own complex structure of which little is understood on a partonic level [96]. One such example of this structure is the existence of nuclear “shadowing”, in which the nuclear PDF is observed to be suppressed




**Figure 10.** The nuclear modification factor  $R_{AA}$  for prompt  $J/\psi$  and  $\psi(2S)$  mesons in PbPb collisions at  $\sqrt{s} = 5.02$  TeV as a function of the centrality (here defined with the average number of particles participating in the interaction,  $N_{part}$ ). (a) Measurement by CMS, from [52]. Figure by the CMS Collaboration, obtained from [52] under CERN copyright, and licensed under [CC BY 4.0](#). The figure size is reduced with respect to the original version. (b) Measurement by ALICE, from [89]. Figure by the ALICE Collaboration, obtained from [89] under CERN copyright, and licensed under [CC BY 4.0](#). The figure size is reduced with respect to the original version. Note that the color scheme for the  $J/\psi$  and  $\psi(2S)$  is switched between the two figures.


in certain partonic kinematic regions compared to the same regions in a proton PDF, for reasons not yet fully understood. Recent progress on nPDF interpretation and fitting is described in the excellent review in [98]. Both nPDF sets used in conjunction with the CIM model are fitted to Deep Inelastic Scattering and Drell-Yan data measured with nuclear targets, and also include inclusive  $\pi^0$  production data measured in  $dAu$  collisions at RHIC [96][97]. The nCTEQ15 nPDFs are fitted to fewer data points than the EPS09 nPDFs and the determination of the uncertainty approaches also varies between the two collaborations [96]. The CIM model with either nPDF set does a good job of describing the CMS  $Y$  data, with perhaps the only exception being a slight discrepancy with the  $Y(3S)$  data at negative rapidity.

The  $Y(1S)$   $R_{pPb}$  data were also compared to predictions involving an energy loss mechanism [99] as well as predictions that only included nuclear shadowing effects using the EPS09 nPDFs computed at next-to-leading-order [97][100] shown in Figure 12. The energy loss model depicted here specifically accounts for the fully coherent energy loss (FCEL) mechanism that occurs due to the interference of gluon emission amplitudes before and after a partonic hard scattering event [101][102]. The data are consistent with all of the compared theory models, indicating that the observed  $Y$  suppression is consistent with being produced as a result of CNM effects, but the exact effects responsible for the suppression remain unknown.


Recent measurements of  $Y$  production in  $\sqrt{s} = 8.16$  TeV  $pPb$  collisions were also performed by the ALICE experiment [94]. Figure 13 shows the ALICE and LHCb measurements of the  $Y(1S)$   $R_{pPb}$  compared to several theoretical models. The first is a model using the EPS09NLO nPDFs to describe the nuclear shadowing effects combined with the Color Evaporation Model (CEM) to describe the  $Y$  production [100]. In the CEM, the mass of a  $Q\bar{Q}$  pair is restricted to be less than the open heavy flavor meson threshold, and the total cross section for a quarkonium state given this mass constraint is computed from a fraction  $F$  of the total  $Q\bar{Q}$  pairs satisfying this criteria [100]. The next models were previously introduced during the discussion of Figure 12 and include fully coherent energy loss model predictions [103] with and without the additional inclusion of nuclear shadowing effects with the EPS09 nPDFs [104]. The EPPS16 reweighted and nCTEQ15 reweighted predictions are, respectively, the EPPS16 [105] and nCTEQ15 [96] nPDF sets reweighted with heavy flavor hadron production data from the LHC to study the effect of the heavy flavor data on constraining the gluon nuclear PDF [106]. The final model shown is the Comover model



**Figure 11.** The nuclear modification factors  $R_{pPb}$  measured for  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$  mesons as a function of  $y$  compared to predictions from the Comover Interaction Model (CIM). Figures are by the CMS Collaboration, obtained from [95] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figures is reduced with respect to the original version.



**Figure 12.** The nuclear modification factor  $R_{pPb}$  measured for  $Y(1S)$  mesons compared to (a) predictions involving nuclear shadowing effects and (b) predictions involving shadowing and energy loss effects. Figures are by the CMS Collaboration, obtained from [95] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figures is reduced with respect to the original version.



**Figure 13.** The nuclear modification factor  $R_{pPb}$  measured for  $Y(1S)$  states by ALICE and LHCb and compared to (a) predictions involving different CNM effects and (b) predictions from the Comover interaction model. Figures are by the ALICE Collaboration, obtained from [94] under CERN copyright, and licensed under CC BY 4.0. The size of the figures is reduced with respect to the original version.

with nCTEQ15 nPDFs for shadowing contributions [33]. As seen with the CMS results, the data from ALICE and LHCb are consistent with models involving several different CNM effects. The Comover model is particularly promising in that it can describe the suppression of all three  $Y$  states and it appears to do a slightly better job of describing the data at negative rapidity than the other models.


## 4.2. Charmonia

### 4.2.1. Ground state suppression

Following the strong suppression of  $J/\psi$  mesons observed in AA collisions, it is natural to ask what we currently know about  $J/\psi$  suppression in  $pA$  collisions. At the LHC, measurements of the  $J/\psi$   $R_{pPb}$  have been performed at  $\sqrt{s} = 5.02$  TeV and 8.16 TeV [93] [109]–[117]. As with the AA results, measurements of prompt  $J/\psi$  production will be highlighted here as these are most sensitive to probing  $J/\psi$  dissociation mechanisms. The LHC measurements will be briefly introduced and then their theoretical comparisons will be discussed, since the data are compared to many of the same theoretical models. Figure 14 shows the most recent prompt  $J/\psi$   $R_{pPb}$  results from ALICE [117], compared to previous LHC measurements at midrapidity as well as several theory predictions and plotted as a function of the  $J/\psi$   $p_T$ . While one rapidity bin of the CMS measurement is shown in Figure 14, the measurement was also performed in multiple rapidity bins to probe the nuclear effects on  $J/\psi$  production as a function of both  $p_T$  and rapidity, as shown in Figure 15. Due to its forward kinematic acceptance, LHCb has measured the prompt  $J/\psi$   $R_{pPb}$  in a complementary rapidity range to that of the midrapidity LHC experiments, probing more positive and negative rapidity values in the center of mass frame [110]. The LHCb  $R_{pPb}$  results are shown as a function of  $p_T$  for two different rapidity bins in Figure 16. The LHCb and ALICE  $R_{pPb}$  measurements are shown as a function of the center of mass rapidity in Figure 17.


All of the  $R_{pPb}$  results in Figures 14–17 are compared to theoretical predictions, most of which incorporate different cold nuclear matter effects. Several of the nuclear PDFs used in the models in Figures 14–17 have already been previously introduced, such as the EPS09 at LO and NLO [105] and the nCTEQ15 [96] and EPPS16 [105] nPDFs with a reweighting to include the impact of LHC heavy flavor data [106][107]. A new addition is the nCTEQ15HQ set [118], which included open heavy flavor and quarkonium measurements in  $pPb$  data from the LHC in a global nPDF analysis (the “HQ” in the name stands for Heavy Quark). The HELAC-Onia models compared to the LHCb data in Figure 16 are based on the HELAC-Onia package which computes the matrix elements involved in collinear NRQCD factorisation predictions of quarkonia production [119][120]. A Color Glass Condensate (CGC) approach is also introduced as a model comparison to the  $J/\psi$   $R_{pPb}$  data [121]–[123]. The CGC model is an effective field theory applicable when the






ALI-PUB-561246

**Figure 14.** The prompt  $J/\psi$   $R_{pPb}$  measured by ALICE, ATLAS, and CMS as a function of the  $J/\psi$   $p_T$  and compared to several theoretical predictions. See text for descriptions of the theory models. Figure by the ALICE Collaboration, obtained from [117] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The figure size is reduced with respect to the original version.



**Figure 15.** The  $J/\psi$   $R_{pPb}$  measured by CMS as a function of  $p_T$  and  $y$ . See text for descriptions of the theory models. Figure by the CMS Collaboration, obtained from [109] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figure is reduced with respect to the original version.




**Figure 16.** The prompt  $J/\psi$   $R_{pPb}$  measured by LHCb in two different rapidity bins corresponding to (a) the  $pPb$  beam configuration with the proton beam entering the detector and (b) the  $PbPb$  beam configuration with the Pb beam entering the detector. See text for details of the theoretical comparisons. Figures by the LHCb Collaboration, obtained from [110] under CERN copyright, and licensed under CC BY 4.0. The size of the figures is reduced with respect to the original version.

saturation scale, the scale at which the gluon density in a nucleus “saturates” and reaches its maximum limit, is much larger than the energy scale associated with nonperturbative QCD, denoted by  $\Lambda_{QCD}$  [121]. The  $c\bar{c}$  pair is produced from the radiation of a “probe” parton, and the pair proceeds to interact with the dense gluonic system in the target nucleus [122][124]. In contrast to the rest of the models, the transport model predictions [108] in Figure 14 involve a hot nuclear medium and the treatment of the quarkonia production and dissociation using a rate equation approach as described in Section 3.1.

Several interesting features can be noted from comparing the midrapidity results in Figures 14–15 with the forward and backward rapidity results in Figure 16. At midrapidity and high  $p_T$ , the  $R_{pPb}$  is slightly larger than one. This can be most clearly see in the CMS results shown in Figure 15, in which the  $R_{pPb}$  is observed to be larger than one in most of the measured  $p_T$  bins in  $-0.9 < y_{CM} < -1.93$ . It is when approaching the more forward rapidities that the value of  $R_{pPb}$  begins to drop below one, consistent with what is observed in the LHCb data. The  $R_{pPb}$  is smaller than one for almost the entire  $p_T$  range probed in the forward region, and for  $p_T < 3$  GeV at midrapidity. The  $R_{pPb}$  plotted as a function of rapidity shows a decreasing trend from negative to positive rapidity, as shown in Figure 17. Another observation is that the data cannot be described by a single nuclear PDF set, as can be seen most clearly with the LHCb data. The data are consistent with the models, which seem to suggest that nPDF modifications at a minimum are needed to describe the data but that other CNM effects such as energy loss or saturation effects could also be at play.

RHIC data is also extremely valuable for studying cold nuclear matter effects, as quarkonia measurements in many nuclear systems, collision energies, and phase space are needed to begin to disentangle the different effects and understand their underlying mechanisms. An example of this can be seen in the recent measurement from PHENIX of the  $J/\psi$   $R_{pAl}$ ,  $R_{pAu}$ , and  $R_{HeAu}$  [70]. The  $R_{AB}$  results from PHENIX are shown for each collision system as a function of  $p_T$  in Figure 18 and as a function of rapidity in Figure 19. The theoretical models, which have already been discussed previously, include different parameterizations of nuclear shadowing effects via the EPPS16 nPDFs [100][105], the EPS09 nPDFs [105], and the reweighted EPPS16 and nCTEQ15 nPDFs [106][107], and a transport model that includes hot medium effects on the charmonia production [74][108]. Contrary to the LHC results, the PHENIX results shown here begin to differentiate between theoretical models, as can be seen for the observed values of the  $R_{AB}$  at negative rapidity in  $p+Au$  and  $^3He+Au$ , which do not agree with the EPPS16 predictions from Vogt and Shao. These predictions did not include nuclear absorption effects, which is expected to be significant at RHIC energies [125]. When adding a nuclear absorption model [126] to the




**Figure 17.** The  $J/\psi$   $R_{pPb}$  measured as a function of rapidity and compared to several theoretical models. See text for details of the models. Figure by the ALICE Collaboration, obtained from [116] under CERN copyright, and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The figure size is reduced with respect to the original version.

same calculations (predictions labeled with “+ Abs” in the figure), the agreement between the data and predictions is greatly improved. This study presents a nice example of how a specific cold nuclear matter effect can begin to be isolated, or at least identified as the dominant effect, in different nuclear systems in order to better understand its underlying mechanism.


The STAR collaboration has also recently measured the  $J/\psi$   $R_{pAu}$ , as can be seen in Figure 20 [127]. The STAR  $p+Au$  results are compared to the  $J/\psi$   $R_{AA}$  measurement by STAR in Au+Au collisions, to  $d+Au$  results measured by PHENIX, and to several theoretical calculations. The  $p+Au$  and  $d+Au$  results are consistent with each other over the entire measured  $p_T$  range, which suggests that the measured nuclear modification is primarily due to nuclear effects associated with the Au nucleus. There are several similarities between the  $R_{pAu}$  measured at midrapidity and the  $R_{pPb}$  measured at midrapidity. For both cases, the nuclear modification factor is consistent with one starting around 3 GeV/c. The small enhancement above one at high  $p_T$  that was observed in the ATLAS and CMS data is not observed in the ALICE or STAR data, possibly due to the larger statistical uncertainty for the latter results. The theoretical comparisons include effects due to nuclear shadowing [97][106][107][119][120], a CGC model [128][129], a transport model including a hot nuclear medium effect [73][108], an energy loss model [130], and a comover model [131]. Notably, the STAR data are consistent with predictions involving cold nuclear matter effects and the TAMU transport model which involves the formation of a QGP medium. The Comover model predictions seem to do particularly well at describing the data at low  $p_T$ , but the other models do a better job of describing the data over the entire measured  $p_T$  range. It is worth mentioning that particularly for the  $J/\psi$  results discussed in this section, the data can be described by many models that describe individual CNM effects, usually in combination with nuclear shadowing. However it is likely the case that multiple CNM effects occurring simultaneously are responsible for the observed  $J/\psi$  dissociation. Understanding the relative contributions of various CNM effects to the observed suppression remains a challenging but important goal.

#### 4.2.2. Excited state suppression


Proton-nucleus systems are ideal for studying excited-state charmonium production as these states are generally statistically-limited in nucleus-nucleus collisions. Recent measurements in  $pA$  collisions have studied  $\chi_c$  production, which is particularly interesting as it has never been studied in AA collisions. As previously discussed in Section 2, the prompt  $J/\psi$  yield measured experimentally includes  $J/\psi$  production via the decays of



**Figure 18.** The  $J/\psi$  nuclear modification factor as a function of  $p_T$  measured by PHENIX in three different collision systems:  $p+Al$  (left plots),  $p+Au$  (middle plots), and  ${}^3He+Au$  (right plots). The top row shows the results at negative rapidity, while the bottom row shows the results at positive rapidity. See text for details of the theoretical predictions. Figures by the PHENIX Collaboration, obtained from [70] and licensed under [CC BY 4.0](#). The size of the figures is reduced with respect to the original version.



**Figure 19.** The  $J/\psi$  nuclear modification factor as a function of  $y$  measured by PHENIX in three different collision systems:  $p+Al$  (left),  $p+Au$  (middle), and  ${}^3He+Au$  (right). See text for details of the theoretical predictions. Figure by the PHENIX Collaboration, obtained from [70] and licensed under [CC BY 4.0](#). The size of the figure is reduced with respect to the original version.




**Figure 20.** The nuclear modification factor  $R_{pAu}$  for  $J/\psi$  mesons measured as a function of  $p_T$  by STAR and compared to several theoretical models. Figure by the STAR Collaboration, obtained from [127] and licensed under CC BY 4.0. The size of the figure is reduced with respect to the original version.


higher charmonium states, including the  $\chi_c$ . LHCb measured for the first time at the LHC the fraction of  $J/\psi$  mesons produced from  $\chi_c$  decays [132], where the total  $\chi_c$  yield included the  $\chi_{c1}$  and  $\chi_{c2}$  states. The result is shown in Figure 21 as a function of the  $J/\psi$   $p_T$ . The  $pPb$  measurement is consistent with the fraction measured in  $pp$  collisions except in the lowest  $p_T$  bins at negative rapidity, however the increased fraction in these bins is compatible with the observed  $\psi(2S)$  suppression in  $pPb$  collisions [132].

The production of  $\psi(2S)$  mesons in  $pPb$  collisions has been measured by all four LHC experiments [93], [133]-[139]. Figure 22 shows the  $\psi(2S)$   $R_{pPb}$  measured by CMS as a function of  $y$  in three different  $p_T$  bins [136]. The  $\psi(2S)$  suppression is most evident in the lowest  $p_T$  bin probed, from 4 to 6.5 GeV/c, where the  $\psi(2S)$  is significantly suppressed with respect to the  $J/\psi$  particularly at negative rapidity. In the middle  $p_T$  bin, from 6.5 to 10 GeV/c, the  $\psi(2S)$  appears significantly suppressed at negative rapidity but is consistent with the  $J/\psi$ , and no suppression, for  $y > 1$ . In the highest  $p_T$  bin probed, the  $\psi(2S)$  suppression is consistent with one although it is systematically lower than the  $J/\psi$   $R_{pPb}$  values. It is interesting that the  $R_{pPb}$  values for the  $J/\psi$  are consistently slightly larger than one, while the values for the  $\psi(2S)$  tend to be smaller than one except in the highest  $p_T$  bin.


LHCb has also measured the prompt  $\psi(2S)$   $R_{pPb}$  and compared it to several theoretical models [134]. The LHCb measurement as a function of rapidity is shown in Figure 23. The LHCb results clearly show that the  $\psi(2S)$  is more suppressed than the  $J/\psi$  at negative rapidity than at positive rapidity, which is also consistent with the trend observed by CMS in the  $p_T$  bins where the measurements overlap. The LHCb data are compared to models involving shadowing via the EPS09 PDFs at LO [140][141] and NLO [142], and to energy loss predictions with and without nuclear shadowing effects included [103]. Interestingly, while the  $J/\psi$  suppression can be described by the theoretical models, none of them successfully describe the  $\psi(2S)$  suppression. The same difficulty of theoretical models to describe both the  $J/\psi$  and  $\psi(2S)$  suppression was observed by ALICE when they measured the inclusive  $\psi(2S)$   $R_{pPb}$  [137]. Figure 24(a) shows the ALICE measurements compared to several theoretical models, employing some of the same cold nuclear matter effects as were included in the models compared to the LHCb data. The models include shadowing effects [106][142], CGC models [123][143], an energy loss model [99] and the Comover model [131]. The Comover model compared to the data in Figure 24(b) appears



**Figure 21.** The fraction of prompt  $J/\psi$  mesons produced from  $\chi_c$  decays, measured in  $pp$ ,  $p\text{Pb}$ , and  $\text{Pb}p$  collisions as a function of the  $J/\psi$   $p_T$ . Figure by the LHCb Collaboration, obtained from [132] under CERN copyright and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figure is reduced with respect to the original version.



**Figure 22.** The nuclear modification factor  $R_{p\text{Pb}}$  for prompt  $J/\psi$  and  $\psi(2\text{S})$  mesons measured as a function of rapidity and  $p_T$ . The top left figure shows the results in the lowest  $p_T$  bin probed, 4–6.5 GeV/c, the top right figure shows the results in  $p_T = 6.5$ –10 GeV/c, and the bottom figure shows the results in  $p_T = 10$ –30 GeV/c. Figures by the CMS Collaboration, obtained from [136] under CERN copyright and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figures is reduced with respect to the original version.




**Figure 23.** The nuclear modification factor  $R_{pPb}$  for prompt  $J/\psi$  and  $\psi(2S)$  mesons, measured as a function of rapidity and compared to several theoretical models. Figure by the LHCb Collaboration, obtained from [134] under CERN copyright and licensed under CC BY 4.0. The size of the figure is reduced with respect to the original version.

to be the only model able to describe both the  $J/\psi$  and  $\psi(2S)$  suppression, at least getting the approximate order of magnitude for the  $\psi(2S)$  suppression correct. 593


Recent studies of  $\psi(2S)$  suppression have also been performed at RHIC by PHENIX in  $p+Al$  and  $p+Au$  collisions [144]. The PHENIX results also show a significant  $\psi(2S)$  suppression at negative rapidity that cannot be explained by models using different nuclear PDFs, specifically the reweighted EPPS16 and nCTEQ15 nPDFs [106][107] as shown in Figure 25. The authors note that nuclear absorption cannot explain the observed suppression as the initial-state CNM effects should be similar for the  $J/\psi$  and  $\psi(2S)$ . It would however still be interesting to confirm this by comparing to models including a nuclear absorption component, as was done in [70]. It would also be interesting to see if the Comover model can reproduce the observed  $\psi(2S)$  suppression in the backward region, as it appears to be able to do for the ALICE data shown in Figure 24, which could also not be described by shadowing alone. The PHENIX nuclear modification factor measurements were also compared to recent measurements from ALICE [115] as a function of centrality and transport model predictions [108], which are shown in Figure 26. Note that in Figure 26, the ALICE nuclear modification factor as a function of centrality is denoted by  $Q_{pPb}$  due to potential biases in the determination of centrality in  $pPb$  collisions. The transport model predictions describe the data better in the forward rapidity region than in the backward rapidity region, although the backward predictions are still consistent with the data points within the large uncertainties. It is important to note here that the ALICE data shown in Figure 26 were also well described by Comover model predictions, again showing that the observed suppression is consistent with predictions involving hot or cold nuclear matter effects. 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615

## 5. Conclusions 616


Significant progress has been made both experimentally and theoretically to better understand quarkonium production and modification in nuclear systems. The measurements presented here show that the production of quarkonia in nuclear systems are indeed suppressed with respect to  $pp$  collisions, and the suppression in nucleus-nucleus collisions where QGP is expected to be present is indeed stronger than that observed in smaller proton-nucleus systems. There is experimental evidence of the sequential suppression of quarkonia states as a function of their binding energy for both the bottomonia and charmonia systems. However, the experimental measurements and theoretical descriptions 617 618 619 620 621 622 623 624



**Figure 24.** The nuclear modification factor of inclusive  $J/\psi$  and  $\psi(2S)$  mesons as a function of rapidity and compared to several theoretical models. **(a)** depicts comparisons to models using different nPDFs as well as an energy loss model and **(b)** depicts comparisons to CGC and Comover model predictions. See text for details of the theoretical models. Figures by the ALICE Collaboration, obtained from [137] under CERN copyright and licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). The size of the figures is reduced with respect to the original version.



**Figure 25.** The  $J/\psi$  and  $\psi(2S)$  nuclear modification factors measured as a function of rapidity by PHENIX in **(a)** p+Al and **(b)** p+Au collisions. Figures by the PHENIX collaboration, obtained from [144] under American Physical Society copyright and re-used with permission. The size of the figures is reduced with respect to the original version.



**Figure 26.** The  $\psi(2S)$  nuclear modification factor measured as a function of centrality ( $R_{pAu}$ ) by PHENIX ( $R_{pAu}$ ) and ALICE ( $Q_{pPb}$ ). Figure by the PHENIX collaboration, obtained from [144] under American Physical Society copyright and re-used with permission. The size of the figure is reduced with respect to the original version.



cannot yet definitively conclude that the observed suppression in AA collisions is due to quarkonium melting. Other effects, such as parton energy loss and comover interactions can also describe the data within the current experimental uncertainties. In order to better understand the complex quarkonium dynamics in AA collisions, the cold nuclear matter effects that modify quarkonium production in  $pA$  collisions and the quarkonium production mechanisms in  $pp$  collisions, the baseline for comparisons to AA collisions, must be more thoroughly understood. Future measurements in a variety of collision systems, combined with the increased luminosities expected with future RHIC and LHC data, will enable precise and differential measurements that will help to improve our knowledge of quarkonia production and dissociation in nuclear systems.

**Funding:** This research was funded by Agence Nationale de la Recherche grant number ANR-21-CE31-0006. The APC was funded by MDPI.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. J. C. Collins and M. J. Perry. Superdense matter: Neutrons or asymptotically free quarks? *Phys. Rev. Lett.* **1975**, *34*, 1353.
2. N. Cabibbo and G. Parisi. Exponential Hadronic Spectrum and Quark Liberation. *Phys. Lett. B* **1975**, *59*, 67-69.
3. E. V. Shuryak. Theory of Hadronic Plasma. *Sov. Phys. JETP* **1978**, *47*, 212-219.
4. P. Braun-Munzinger, V. Koch, T. Schäfer, and J. Stachel. Properties of hot and dense matter from relativistic heavy ion collisions. *Phys. Rep.* **2016**, *621*, 76-126.
5. W. Busza, K. Rajagopal, and W. van der Schee. Heavy Ion Collisions: The Big Picture, and the Big Questions. *Ann. Rev. Nucl. Part. Phys.* **2018**, *68*, 339-376.
6. Matsui, T. and Satz, H.  $J/\psi$  suppression by quark gluon plasma formation. *Phys. Lett. B* **1986**, *178*, 416.
7. A. Andronic et al. Heavy-flavour and quarkonium production in the LHC era: from proton-proton to heavy-ion collisions. *Eur. Phys. J. C* **2016**, *76*, 107.
8. J.-P. Lansberg. New observables in inclusive production of quarkonia. *Phys. Rep.* **2020**, *889*, 1-106.
9. A. Rothkopf. Heavy quarkonium in extreme conditions. *Phys. Rep.* **2020**, *858*, 1-117.
14. J. L. Nagle and W. A. Zajc. Small System Collectivity in Relativistic Hadronic and Nuclear Collisions. *Ann. Rev. Nucl. Part. Sci.* **2018**, *68*, 211-235.
11. M. Laine, O. Philipsen, M. Tassler and P. Romatschke. Real-time static potential in hot QCD. *JHEP* **2007**, *03*, 054.
12. A. Beraudo, J. P. Blaizot, C. Ratti. Real and imaginary-time  $Q\bar{Q}$  correlators in a thermal medium. *Nucl. Phys. A* **2008**, *806*, 312-338.
13. N. Brambilla, J. Ghiglieri, A. Vairo, and P. Petreczky. Static quark-antiquark pairs at finite temperature. *Phys. Rev. D* **2008**, *78*, 014017.
14. J. L. Nagle and W. A. Zajc. Small System Collectivity in Relativistic Hadronic and Nuclear Collisions. *Ann. Rev. Nucl. Part. Sci.* **2018**, *68*, 211-235.
15. N. Brambilla et al. Heavy Quarkonium Physics. CERN Yellow Report, CERN-2005-005 (2005).
16. A. Mocsy, P. Petreczky, and M. Strickland. Quarkonia in the Quark Gluon Plasma. *Int. J. Mod. Phys. A* **2013**, *28*, 1340012.
17. L. Apolinário, Y.-J. Lee, and M. Winn. Heavy quarks and jets as probes of the QGP. *Prog. Part. Nucl. Phys.* **2022**, *127*, 103990.
18. M. L. Miller, K. Reygers, S. J. Sanders, P. Steinberg. Glauber Modeling in High Energy Nuclear Collisions. *Ann. Rev. Nucl. Part. Sci.* **2007**, *57*, 205-243.
19. CMS Collaboration. Observation of sequential Y suppression in PbPb collisions. *Phys. Rev. Lett.* **2012**, *109*, 222301.
20. ALICE Collaboration. Suppression of Y(1S) at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *Phys. Lett. B* **2014**, *738*, 361-372.
21. CMS Collaboration. Suppression of Y(1S), Y(2S), and Y(3S) quarkonium states in PbPb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *Phys. Lett. B* **2017**, *770*, 357.
22. CMS Collaboration. Suppression of excited Y states relative to the ground state in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Rev. Lett.* **2018**, *120*, 142301.
23. CMS Collaboration. Measurement of nuclear modification factors of Y(1S), Y(2S), and Y(3S) mesons in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2019**, *790*, 270.
24. ALICE Collaboration. Y suppression at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2019**, *790*, 89.
25. ALICE Collaboration. Y production and nuclear modification at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B*, **2021**, *822*, 136579.
26. CMS Collaboration. Observation of the Y(3S) meson and suppression of Y states in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Rev. Lett.* **2023**, submitted. DOI: <https://doi.org/10.48550/arXiv.2303.17026>.
27. ATLAS Collaboration. Production of Y( $nS$ ) mesons in Pb+Pb and  $pp$  collisions at 5.02 TeV. *Phys. Rev. C*, **2023**, *107*, 054912.
28. ALICE Collaboration. Measurement of Y(1S) Elliptic Flow at Forward Rapidity in Pb-Pb Collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Rev. Lett.* **2019**, *123*, 192301.

29. CMS Collaboration. Measurement of the azimuthal anisotropy of  $Y(1S)$  and  $Y(2S)$  mesons in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2021**, *819*, 136385. 681
30. Yao, X. et al. Coupled Boltzmann transport equations of heavy quarks and quarkonia in quark-gluon plasma. *JHEP*, **2021**, *01*, 046. 682
31. N. Brambilla, M. A. Escobedo, A. Islam, M. Strickland, A. Tiwari, A. Vairo, and P. V. Griend. Regeneration of bottomonia in an open quantum systems approach. *Phys. Rev. D*, **2023**, *108*, L011502. 683
32. G. Wolschin. Bottomonium spectroscopy in the quark-gluon plasma. *Int. Jour. Mod. Phys. A*, **2020**, *35*, 2030016. 684
33. E. G. Ferreira and J. P. Lansberg. Is bottomonium suppression in proton-nucleus and nucleus-nucleus collisions at LHC energies due to the same effects? *JHEP*, **2018**, *10*, 094. 685
34. X. Du, M. He, and R. Rapp. Color screening and regeneration of bottomonia in high-energy heavy-ion collisions. *Phys. Rev. C* **2017**, *96*, 054901. 686
35. L. Grandchamp, R. Rapp, and G. E. Brown. In-Medium Effects on Charmonium Production in Heavy-Ion Collisions. *Phys. Rev. Lett.* **2004** *92*, 212301. 687
36. F. Riek and R. Rapp. Quarkonia and heavy-quark relaxation times in the quark-gluon plasma. *Phys. Rev. C* **2010**, *82*, 035201. 688
37. N. Brambilla, M. A. Escobedo, M. Strickland, A. Vairo, P. Vander Griend, J.H. Weber. Bottomonium production in heavy-ion collisions using quantum trajectories: Differential observables and momentum anisotropy. *Phys. Rev. D*, **2021**, *104*, 094049. 689
38. N. Brambilla, M. A. Escobedo, M. Strickland, A. Vairo, P. Vander Griend, J.H. Weber. Bottomonium suppression in an open quantum system using the quantum trajectories method. *JHEP* **2021**, *05*, 136. 690
39. N. Brambilla, M. A. Escobedo, A. Islam, M. Strickland, A. Tiwari, A. Vairo, P. Vander Griend. Heavy quarkonium dynamics at next-to-leading order in the binding energy over temperature. *JHEP* **2022**, *08*, 303. 691
40. Observation of the  $Y(3S)$  meson and suppression of  $Y$  states in PbPb collisions at  $\sqrt{s} = 5$  TeV. CMS Public Results Website, CERN, <https://cms-results.web.cern.ch/cms-results/public-results/publications/HIN-21-007/>. 692
41. STAR Collaboration. Measurement of sequential  $Y$  suppression in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV with the STAR experiment. *Phys. Rev. Lett.* **2023**, *130*, 112301. DOI: <https://doi.org/10.1103/PhysRevLett.130.112301> 693
42. ATLAS Collaboration. Measurement of the centrality dependence of  $J/\psi$  yields and observation of  $Z$  production in lead-lead collisions with the ATLAS detector at the LHC. *Phys. Lett. B* **2011**, *697*, 294-312. 694
43. ALICE Collaboration.  $J/\psi$  suppression at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *Phys. Rev. Lett.* **2012**, *109*, 072301. 695
44. CMS Collaboration. Suppression of non-prompt  $J/\psi$ , prompt  $J/\psi$ , and  $Y(1S)$  in PbPb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *JHEP* **2012**, *05*, 063. 696
45. CMS Collaboration. Measurement of prompt  $\psi(2S)$  to  $J/\psi$  yield ratios in PbPb and pp collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *Phys. Rev. Lett.* **2014**, *113*, 262301. 697
46. ALICE Collaboration. Centrality, rapidity, and transverse momentum dependence of  $J/\psi$  suppression in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *Phys. Rev. B* **2014**, *734*, 314-327. 698
47. ALICE Collaboration. Inclusive, prompt and non-prompt  $J/\psi$  production at mid-rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *JHEP* **2015**, *07*, 051. 699
48. ALICE Collaboration. Differential studies of inclusive  $J/\psi$  and  $\psi(2S)$  production at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV. *JHEP* **2016**, *05*, 179. 700
49. CMS Collaboration. Relative modification of prompt  $\psi(2S)$  and  $J/\psi$  yields from pp to PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Rev. Lett.* **2017**, *118*, 162301. 701
50. ALICE Collaboration.  $J/\psi$  suppression at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2017**, *766*, 212-224. 702
51. ATLAS Collaboration. Prompt and non-prompt  $J/\psi$  and  $\psi(2S)$  suppression at high transverse momentum in 5.02 TeV Pb+Pb collisions with the ATLAS experiment. *Eur. Phys. J. C* **2018**, *78*, 762. DOI: <https://doi.org/10.1140/epjc/s10052-018-6219-9> 703
52. CMS Collaboration. Measurement of prompt and nonprompt charmonium suppression in PbPb collisions at 5.02 TeV. *Eur. Phys. J. C* **2018**, *78*, 509. DOI: <https://doi.org/10.1140/epjc/s10052-018-5950-6> 704
53. ALICE Collaboration. Studies of  $J/\psi$  production at forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *JHEP* **2020**, *02*, 041. 705
54. ALICE Collaboration. Centrality and transverse momentum dependence of inclusive  $J/\psi$  production at midrapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2020**, *805*, 135434. 706
55. ALICE Collaboration. Measurements of inclusive  $J/\psi$  production at midrapidity and forward rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2023**, *submitted*. 707
56. ALICE Collaboration. Prompt and non-prompt  $J/\psi$  production at midrapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *JHEP* **2023**, *submitted*. DOI: <https://doi.org/10.48550/arXiv.2308.16125> 708
57. ALICE Collaboration. Inclusive  $J/\psi$  production in Xe-Xe collisions at  $\sqrt{s_{NN}} = 5.44$  TeV. *Phys. Lett. B* **2018**, *785*, 419-428. DOI: <https://doi.org/10.1016/j.physletb.2018.08.047> 709
58. PHENIX Collaboration.  $J/\psi$  Production vs Centrality, Transverse Momentum, and Rapidity in Au+Au Collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Rev. Lett.* **2007**, *98*, 249902. 710
59. STAR Collaboration.  $J/\psi$  production at low  $p_T$  in Au+Au and Cu+Cu collisions at  $\sqrt{s_{NN}} = 200$  GeV at STAR. *Phys. Rev. C* **2014**, *90*, 024906. 711
60. STAR Collaboration.  $J/\psi$  production at high transverse momenta in p+p and Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Lett. B* **2013**, *722*, 55-62. 712

681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739

61. STAR Collaboration. Energy dependence of  $J/\psi$  production in Au+Au collisions at  $\sqrt{s_{NN}} = 39, 62.4, \text{ and } 200 \text{ GeV}$ . *Phys. Lett. B* **2017**, *771*, 13-20. 740
62. STAR Collaboration. Measurement of inclusive  $J/\psi$  suppression in Au+Au collisions at  $\sqrt{s_{NN}} = 200 \text{ GeV}$  through the dimuon channel at STAR. *Phys. Lett. B* **2019**, *797*, 134917. DOI: <https://doi.org/10.1016/j.physletb.2019.134917> 741
63. ALICE Collaboration.  $J/\psi$  Elliptic Flow in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76 \text{ TeV}$ . *Phys. Rev. Lett.* **2013**, *111*, 162301. 742
64. CMS Collaboration. Suppression and azimuthal anisotropy of prompt and nonprompt  $J/\psi$  production in PbPb collisions at  $\sqrt{s_{NN}} = 2.76 \text{ TeV}$ . *Eur. Phys. J. C* **2017**, *77*, 252. 743
65. ALICE Collaboration.  $J/\psi$  Elliptic Flow in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *Phys. Rev. Lett.* **2017**, *119*, 242301. 744
66. ATLAS Collaboration. Prompt and non-prompt  $J/\psi$  elliptic flow in Pb+Pb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$  with the ATLAS experiment. *Eur. Phys. J. C* **2018**, *78*, 784. 745
67. ALICE Collaboration.  $J/\psi$  elliptic and triangular flow in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *JHEP* **2020**, *10*, 141 746
68. CMS Collaboration. Measurements of the azimuthal anisotropy of prompt and nonprompt charmonia in PbPb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *JHEP* **2023**, *10*, 115. 747
69. ALICE Collaboration. Measurement of the  $J/\psi$  polarization with respect to the event plane in Pb-Pb collisions at the LHC. *Phys. Rev. Lett.* **2023**, *131*, 042303. 748
70. PHENIX Collaboration. Measurement of  $J/\psi$  at forward and backward rapidity in  $p+p, p+\text{Al}, p+\text{Au}, \text{ and } ^3\text{He}+\text{Au}$  collisions at  $\sqrt{s_{NN}} = 200 \text{ GeV}$ . *Phys. Rev. C*, **2020**, *102*, 014902. DOI: <https://doi.org/10.1103/PhysRevC.102.014902> 749
71. LHCb Collaboration.  $J/\psi$  and  $D^0$  production in  $\sqrt{s_{NN}} = 68.5 \text{ GeV}$  PbNe collisions. *Eur. Phys. J. C* **2023**, *83*, 658. 750
72. X. Zhao and R. Rapp. Medium modifications and production of charmonia at LHC. *Nucl. Phys. A* **2011**, *859*, 114-125. 751
73. X. Du and R. Rapp. Sequential Regeneration of Charmonia in Heavy-Ion Collisions. *Nucl. Phys. A* **2015**, *943*, 147-158. 752
74. X. Zhao and R. Rapp. Charmonium in Medium: From Correlators to Experiment. *Phys. Rev. C* **2010**, *82*, 064905. 753
75. L. Yan, P. Zhuang, and N. Xu.  $J/\psi$  Production in Quark-Gluon Plasma. *Phys. Rev. Lett* **2006**, *97*, 232301. 754
76. Y. Liu, Z. Qu, N. Xu, and P. Zhuang.  $J/\psi$  transverse momentum distribution in high energy nuclear collisions. *Phys. Lett. B* **2009**, *678*, 72-76. 755
77. R. Sharma and I. Vitev. High transverse momentum quarkonium production and dissociation in heavy ion collisions. *Phys. Rev. C* **2013**, *87*, 044905. 756
78. E. G. Ferreira, F. Fleuret, and E. Maurice. Towards quarkonium formation time determination. *Eur. Phys. J. C*, **2022** *82*, 201. 757
79. A. Andronic, P. Braun-Munzinger, M. K. Köhler, K. Redlich, and J. Stachel. Transverse momentum distributions of charmonium states with the statistical hadronization model. *Phys. Lett. B* **2019**, *797*, 134836. 758
80. K. Zhou, N. Xu, Z. Xu, and P. Zhuang. Medium effects on charmonium production at ultrarelativistic energies available at the CERN Large Hadron Collider. *Phys. Rev. C* **2014**, *89*, 054911. 759
81. B. Chen. Thermal production of charmonia in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *Chin. Phys. C* **2019**, *43*, 124101. 760
82. S. Aronson, E. Borrás, B. Odegard, R. Sharma, and I. Vitev. Collisional and thermal dissociation of  $J/\psi$  and  $Y$  states at the LHC. *Phys. Lett. B* **2018**, *778*, 384-391. 761
83. Y. Makris and I. Vitev. An Effective Theory of Quarkonia in QCD Matter. *JHEP* **2019**, *10*, 111. 762
84. A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel. Statistical hadronization of heavy quarks in ultra-relativistic nucleus-nucleus collisions. *Nucl. Phys. A* **2007**, *89*, 334-356. 763
85. A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel. Evidence for charmonium generation at the phase boundary in ultra-relativistic nuclear collisions. *Phys. Lett. B* **2007**, *652*, 259-261. 764
86. B. Z. Kopeliovich, I. K. Potashnikova, I. Schmidt, and M. Siddikov. Survival of charmonia in a hot environment. *Phys. Rev. C* **2015**, *91*, 024911. 765
87. M. Spousta. On similarity of jet quenching and charmonia suppression. *Phys. Lett. B*, **2017**, *767*, 10. 766
88. F. Arleo. Quenching of Hadron Spectra in Heavy Ion Collisions at the LHC. *Phys. Rev. Lett.* **2017**, *119*, 062302. 767
89. ALICE Collaboration.  $\psi(2S)$  suppression in Pb-Pb collisions at the LHC. *Phys. Rev. Lett.* **2023**, *Accepted*. DOI: <https://doi.org/10.48550/arXiv.2210.08893> 768
90. LHCb Collaboration. Study of  $Y$  production and cold nuclear matter effects in pPb collisions at  $\sqrt{s_{NN}} = 5 \text{ TeV}$ . *JHEP* **2014**, *07*, 094. 769
91. ALICE Collaboration. Production of inclusive  $Y(1S)$  and  $Y(2S)$  in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *Phys. Lett. B* **2015**, *740*, 105-117. 770
92. LHCb Collaboration. Study of  $Y$  production in pPb collisions at  $\sqrt{s_{NN}} = 8.16 \text{ TeV}$ . *JHEP* **2018**, *11*, 194. 771
93. ATLAS Collaboration. Measurement of quarkonium production in proton-lead and proton-proton collisions at 5.02 TeV with the ATLAS detector. *Eur. Phys. J. C* **2018**, *78*, 171. 772
94. ALICE Collaboration.  $Y$  production in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16 \text{ TeV}$ . *Phys. Lett. B* **2020**, *806*, 135486. DOI: <https://doi.org/10.1016/j.physletb.2020.135486> 773
95. CMS Collaboration. Nuclear modification of  $Y$  states in pPb collisions at  $\sqrt{s_{NN}} = 5.02 \text{ TeV}$ . *Phys. Lett. B* **2022**, *835*, 137397. DOI: <https://doi.org/10.1016/j.physletb.2022.137397> 774
96. K. Kovarik et al. nCTEQ15 - Global analysis of nuclear parton distributions with uncertainties in the CTEQ framework. *Phys. Rev. D* **2016**, *93*, 085037. 775

97. K. J. Eskola, H. Paukkunen, C. A. Salgado. EPS09 - a New Generation of NLO and LO Nuclear Parton Distribution Functions. *JHEP* **2009**, *04*, 065. 798
98. M. Klasen and H. Paukkunen. Nuclear PDFs After the First Decade of LHC Data. *Ann. Rev. Nucl. Part. Sci.* **2024**, *74*, 1-41. 799
99. F. Arleo and S. Peigné. Quarkonium suppression in heavy-ion collisions from coherent energy loss in cold nuclear matter. *JHEP* **2014**, *10*, 073. 800
100. R. Vogt. Shadowing Effects on  $J/\psi$  and  $Y$  Production at the LHC. *Phys. Rev. C* **2015**, *92*, 034909. 801
101. F. Arleo, S. Peigné, T. Sami. Revisiting scaling properties of medium-induced gluon radiation. *Phys. Rev. D* **2011**, *83*, 114036. 802
102. S. Peigné, F. Arleo, and R. Kolevatov. Coherent medium-induced gluon radiation in hard forward  $1 \rightarrow 1$  partonic processes. *Phys. Rev. D* **2016**, *93*, 014006. 803
103. F. Arleo and S. Peigné. Heavy-quarkonium suppression in p-A collisions from parton energy loss in cold QCD matter. *JHEP* **2013**, *03*, 122. 804
104. J. L. Albacete et al. Predictions for Cold Nuclear Matter Effects in  $p$ +Pb Collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *Nucl. Phys. A* **2018**, *972*, 18-85. 805
105. K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado. EPPS16: Nuclear parton distributions with LHC data. *Eur. Phys. J. C* **2017**, *77*, 163. 806
106. A. Kusina, J. Lansberg, I. Schienbein, and H. Shao. Gluon Shadowing in Heavy-Flavor Production at the LHC. *Phys. Rev. Lett.* **2018**, *121*, 052004. 807
107. J. Lansberg and H. Shao. Towards an automated tool to evaluate the impact of the nuclear modification of the gluon density on quarkonium,  $D$  and  $B$  meson production in proton-nucleus collisions. *Eur. Phys. J. C*, **2017**, *77*, 1. 808
108. X. Du and R. Rapp. In-Medium Charmonium Production in Proton-Nucleus Collisions. *JHEP* **2019**, *03*, 015. 809
109. CMS Collaboration. Measurement of prompt and non-prompt  $J/\psi$  production in pp and pPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Eur. Phys. J. C* **2017**, *77*, 269. DOI: <https://doi.org/10.1140/epjc/s10052-017-4828-3> 810
110. LHCb Collaboration. Prompt and nonprompt  $J/\psi$  production and nuclear modification in pPb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *Phys. Lett. B* **2017**, *774*, 159. DOI: <https://doi.org/10.1016/j.physletb.2017.09.058> 811
111. ALICE Collaboration. Inclusive  $J/\psi$  production at forward and backward rapidity in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP* **2018**, *07*, 160. 812
112. ALICE Collaboration. Prompt and non-prompt  $J/\psi$  production and nuclear modification at mid-rapidity in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Eur. Phys. J. C* **2018**, *78*, 466. 813
113. ALICE Collaboration.  $J/\psi$  production as a function of charged-particle pseudorapidity density in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2018**, *776*, 91. 814
114. ALICE Collaboration.  $J/\psi$  production as a function of charged-particle multiplicity in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP* **2020**, *2009*, 162. 815
115. ALICE Collaboration. Centrality dependence of  $J/\psi$  and  $\psi(2S)$  production and nuclear modification in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP* **2021**, *02*, 002. 816
116. ALICE Collaboration. Inclusive, prompt, and non-prompt  $J/\psi$  production at midrapidity in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *JHEP* **2022**, *06*, 011. DOI: <https://doi.org/10.1007/JHEP06%282022%29011> 817
117. ALICE Collaboration.  $J/\psi$  production at midrapidity in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP* **2023**, *07*, 137. DOI: [https://doi.org/10.1007/JHEP07\(2023\)137](https://doi.org/10.1007/JHEP07(2023)137) 818
118. P. Duwentäster et al. Impact of heavy quark and quarkonium data on nuclear gluon PDFs. *Phys. Rev. D* **2022**, *105*, 114043. 819
119. H.-S. Shao. HELAC-Onia: an automatic matrix element generator for heavy quarkonium physics. *Computer Physics Communications* **2013**, *184*, 2562-2570. 820
120. H.-S. Shao. HELAC-Onia 2.0: An upgraded matrix-element and event generator for heavy quarkonium physics. *Computer Physics Communications* **2016**, *198*, 238. 821
121. Y.-Q. Ma, R. Venugopalan, H.-F. Zhang.  $J/\psi$  production and suppression in high energy proton-nucleus collisions. *Phys. Rev. D* **2015**, *92*, 071901. 822
122. B. Ducloué, T. Lappi and H. Mäntysaari. Forward  $J/\psi$  production in proton-nucleus collisions at high energy. *Phys. Rev. D* **2015**, *91*, 114005. 823
123. B. Ducloué, T. Lappi and H. Mäntysaari. Forward  $J/\psi$  production at high energy: centrality dependence and mean transverse momentum. *Phys. Rev. D* **2016**, *94*, 074031. 824
124. F. Gelis, E. Iancu, J. Jalilian-Marian, R. Venugopalan. The Color Glass Condensate. *Ann. Rev. Nucl. Part. Sci.* **2010**, *60*, 463-489. 825
125. D. McGlinchey, A. D. Frawley, and R. Vogt. Impact-parameter dependence of the nuclear modification of  $J/\psi$  production in  $d$ +Au collisions at  $\sqrt{s} = 200$  GeV. *Phys. Rev. C*, **2013**, *87*, 054910. 826
126. F. Arleo, P. B. Gossiaux, T. Gousset, and J. Aichelin. Charmonium suppression in  $pA$  collisions. *Phys. Rev. C*, **2000**, *61*, 054906. 827
127. STAR Collaboration. Measurement of cold nuclear matter effects for inclusive  $J/\psi$  in p+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Lett. B*, **2022**, *825*, 136865. DOI: <https://doi.org/10.1016/j.physletb.2021.136865> 828
128. Y.-Q. Ma and R. Vogt. Quarkonium production in an improved color evaporation model. *Phys. Rev. D*, **2016**, *94*, 114029. 829
129. Y.-Q. Ma, R. Venugopalan, K. Watanabe and H.-F. Zhang.  $\psi(2S)$  versus  $J/\psi$  suppression in proton-nucleus collisions from factorization violating soft color exchanges. *Phys. Rev. C* **2018**, *97*, 014909. 830

130. F. Arleo, R. Kolevatov, S. Peigné, and M. Rustamova. Centrality and  $p_T$  dependence of  $J/\psi$  suppression in proton-nucleus collisions from parton energy loss. *JHEP* **2013**, 05, 155. 856
131. E. G. Ferreiro. Excited charmonium suppression in proton-nucleus collisions as a consequence of comovers. *Phys. Lett. B*, **2015**, 749, 98-103. 857
132. LHCb Collaboration. Fraction of  $\chi_c$  decays in prompt  $J/\psi$  production measured in  $p$ Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *Phys. Rev. Lett.* **2023**, submitted. DOI: <https://doi.org/10.48550/arXiv.2311.01562> 858
133. ALICE Collaboration. Suppression of  $\psi(2S)$  production in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *JHEP*, **2014**, 12, 073. 859
134. LHCb Collaboration. Study of  $\psi(2S)$  production and cold nuclear matter effects in  $p$ Pb collisions at  $\sqrt{s_{NN}} = 5$  TeV. *JHEP* **2016**, 03, 133. DOI: <https://doi.org/10.1007/JHEP03%282016%29133> 860
135. ALICE Collaboration. Centrality dependence of  $\psi(2S)$  suppression in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *JHEP*, **2016**, 06, 50. 861
136. CMS Collaboration. Measurement of prompt  $\psi(2S)$  production cross sections in proton-lead and proton-proton collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. *Phys. Lett. B* **2019**, 790, 509. DOI: <https://doi.org/10.1016/j.physletb.2019.01.058> 862
137. ALICE Collaboration. Measurement of nuclear effects on  $\psi(2S)$  production in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP*, **2020**, 07, 237. DOI: <https://doi.org/10.1007/JHEP07%282020%29237> 863
138. ALICE Collaboration. Centrality dependence of  $J/\psi$  and  $\psi(2S)$  production and nuclear modification in p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV. *JHEP*, **2021**, Jem 02, 002. 864
139. ALICE Collaboration. Measurement of  $\psi(2S)$  production as a function of charged-particle pseudorapidity density in pp collisions at  $\sqrt{s} = 13$  TeV and p-Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV with ALICE at the LHC. *JHEP* **2023**, 06, 147. 865
140. E. G. Ferreiro, F. Fleuret, J. P. Lansberg, and A. Rakotozafindrabe. Impact of the nuclear modification of the gluon densities on  $J/\psi$  production in  $p$ Pb collisions at  $\sqrt{s_{NN}} = 5$  TeV. *Phys. Rev. C* **2013**, 88, 047901. 866
141. Z. Conesa del Valle, E. G. Ferreiro, F. Fleuret, J. P. Lansberg, and A. Rakotozafindrabe. Open-beauty production in  $p$ Pb collisions at  $\sqrt{s_{NN}} = 5$  TeV: effect of the gluon nuclear densities. *Nuclear Physics A* **2014**, 926, 236-241. 867
142. J. L. Albacete et al. Predictions for  $p$ +Pb Collisions at  $\sqrt{s} = 5$  TeV. *Int. J. Mod. Phys. E* **2013**, 22, 1330007. 868
143. Y.-Q. Ma, R. Venugopalan, K. Watanabe, and H.-F. Zhang.  $\psi(2S)$  versus  $J/\psi$  suppression in proton-nucleus collisions from factorization violating soft color exchanges. *Phys. Rev. C* **2018**, 97, 014909. 869
144. PHENIX Collaboration. Measurement of  $\psi(2S)$  nuclear modification at backward and forward rapidity in  $p+p$ ,  $p+Al$ , and  $p+Au$  collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Rev. C*, **2022**, 105, 064912. DOI: <https://doi.org/10.1103/PhysRevC.105.064912> 870

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content. 871