



HAL
open science

Thermotropic Ionic Liquid Crystals: Structure/ion transport correlation within stimuli-responsive electrolytes for energy

Manuel Maréchal, Hélène Pung, Celso Yassuo Imp Okada-Junior, Mirella Simoes Santos, Sébastien Livi, Jannick Duchet-Rumeau, Agílio Pádua, Patrice Rannou

► To cite this version:

Manuel Maréchal, Hélène Pung, Celso Yassuo Imp Okada-Junior, Mirella Simoes Santos, Sébastien Livi, et al.. Thermotropic Ionic Liquid Crystals: Structure/ion transport correlation within stimuli-responsive electrolytes for energy. International Society of Electrochemistry, International Society of Electrochemistry, Sep 2023, Lyon, France. hal-04442311

HAL Id: hal-04442311

<https://hal.science/hal-04442311v1>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Thermotropic Ionic Liquid Crystals: Structure/ion transport correlation within stimuli-responsive electrolytes for energy

Hélène PUNG¹, Celso Yassuo OKADA-JUNIOR², Mirella SIMOES SANTOS³, Sébastien LIVI², Jannick DUCHET-RUMEAU², Agilio PADUA³, Patrice RANNOU⁴, Manuel MARÉCHAL^{1,*}

¹Univ. Grenoble Alpes, CNRS, CEA, IRIG-SyMMES, 38000 Grenoble, France

²Université de Lyon, CNRS, Univ. Claude Bernard Lyon 1, INSA Lyon, Univ. Jean Monnet, UMR 5223, Ingénierie des Matériaux Polymères, 69621 Lyon, France

³Laboratoire de Chimie, Ecole Normale Supérieure de Lyon, CNRS, Lyon, 69342 France

⁴Univ. Grenoble Alpes, Univ. Savoie Mont-Blanc, CNRS, Grenoble INP, LEPMI, Grenoble, 38000 France

*manuel.marechal@univ-grenoble-alpes.fr

Thermotropic Ionic Liquid Crystals (TILCs)¹ have recently emerged as a smart class of organic electrolytes encoding the unique hierarchical self-assembling properties and stimuli-responsive mosaicity of mesophases featured by thermotropic liquid crystals² with the efficient ion transport of ionic liquids. As a result, TILCs stand out as a promising electrolytic platform for energy storage (batteries³/supercapacitors) and conversion (fuel cells/DSSCs) as they use industry-compatible and simple-to-implement manufacturing processes. The multi-scale correlation relating their structures and (an/cat)ionic transport properties is crucial to pave understanding of their ionic transport. A fundamental question of interest is here the influence of the dynamical mosaicity related to the average size of their self-assembled domains, with or without (w vs. w/o) dynamic grain boundaries. Remarkably, stimuli such as electric⁴ or magnetic field can be leveraged to sample (dis)order till ultimately reaching monodomains (eradication of mosaicity) to reveal inherent (*i.e.* defect-free) features of their ionic transport.

Exploring this research endeavour, in this communication, the case study of an anionic conductive TILC (A-TILC) composed of an imidazolium-based cation with symmetrical *n*-alkyl chains (C₁₈C₁₈Im⁺) and a N(CN)₂⁻ anion featuring a Smectic A (SmA⁵) mesophase (lamellar organization) will be deciphered by combining *state-of-the-art* (SoA) theoretical and experimental approaches. First, coarse-grained simulations derived from Martinelli⁶ will be presented to unravel its set of structural and dynamic descriptive parameters. Second, an unique *operando* experiment combining (synchrotron-based) SoA structural studies (SAXS+WAXS) with ionic transport measurements (probed by EIS) w vs. w/o a tuneable (0 up to 1T) magnetic field (**Fig. 1**) will be introduced, allowing in-depth study of the interplay linking its structure with its ion transport properties. In closing and beyond the case study of an A-TILC encoding a nanoconfined 2D transport within a SmA mesophase, we will illustrate the potential offered by TILCs to impart next generation (conversion/storage) energy devices with self-healing functionalities.




Fig.1. A. Chemical structure of the C₁₈C₁₈Im⁺/N(CN)₂⁻ TILC, LC cell (through-plane configuration) and mesomorphic behaviours (phase transition and texture) of its SmA mesophase. B. 1D SAXS/WAXS scattering profiles (first heating) as a function of temperature. C. vs. D. Temperature dependent evolution of the correlation length (average domain size) and ionic conductivity w vs. w/o 1T magnetic field.

Acknowledgments: We are indebted to the French National Agency of Research (ANR) for the funding of the collaborative research project CITADEL under the grant ID # [ANR-19-CE05-0028](#). The coauthors acknowledge [ESRF-EBS](#) for beamtime allocation at [ID31](#) under the SC-5292 proposal (DOI: 10.15151/ESRF-ES-894751246).

References: ¹K. Goossens *et al.*, *Chem. Rev.* **2016**, *116*, 4643. DOI: [10.1021/cr400334b](#) ²J. Uchida *et al.*, *Adv. Mater.* **2022**, *34*, 2109063. DOI: [10.1002/adma.202109063](#) ³Z. Ahmad *et al.*, *Proc. Natl. Acad. Sci. (USA)* **2020**, *117*, 26672. DOI: [10.1073/pnas.2008841117](#) ⁴H. Shimura *et al.*, *Adv. Mater.* **2009**, *21*, 1591. DOI: [10.1002/adma.200802252](#) ⁵I.W. Hamley, *Soft Matter* **2022**, *18*, 711. DOI: [10.1039/d1sm01758f](#) ⁶G.D. Barbosa. *et al.*, *Macromolecules* **2022**, *55*, 26. DOI: [10.1021/acs.macromol.1c01932](#)