



Electron contribution to photon-induced desorption in molecular ices

D Torres-Díaz, R Basalgète, A Hacquard, J.-H Fillion, X Michaut, L Philippe,
G Feraud, L Amiaud, M Bertin, A Lafosse

► To cite this version:

D Torres-Díaz, R Basalgète, A Hacquard, J.-H Fillion, X Michaut, et al.. Electron contribution to photon-induced desorption in molecular ices. POSMOL 2023 - XXI International Workshop on Low-Energy Positron and Positronium Physics & XXIII International Symposium on Electron-Molecule Collisions and Swarms, Aug 2023, Notre Dame, IN, United States. hal-04442689

HAL Id: hal-04442689

<https://hal.science/hal-04442689v1>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electron contribution to photon-induced desorption in molecular ices

D. Torres-Díaz^{1,2}, R. Basalgète,² A. Hacquard,² J.-H. Fillion,² X. Michaut,² L. Philippe,² G. Feraud,² L. Amiaud¹, M. Bertin², A. Lafosse¹


¹ISMO, CNRS, Université Paris-Saclay, France

²LERMA, CNRS, Sorbonne Université, Observatoire de Paris, Université PSL, France
CNRS, Université Paris-Saclay, France

anne.lafosse@universite-paris-saclay.fr

Chemistry in the interstellar medium (ISM) takes place both in the gas phase and on the surface of interstellar dust grains, where films of physisorbed molecules accumulate. These ices are processed by photons, cosmic rays and the large number of secondary electrons produced inside the ice. Desorption is the key step in this interaction between gas phase, solid phase and radiation, and is at the heart of the interpretation of the most recent observational data obtained by the latest generation of telescopes (ALMA, NOEMA, JWST).

Our aim is to assess the contribution of electron-induced processes to the non-thermal desorption of cryogenic molecular films subjected to photon irradiation, and to better understand the complex surface chemistry in icy mantles. We carry on comparative measurements of electron- and photon-induced desorption (ESD and PSD) using two setups, the E/SOLID setup at ISMO and the SPICES setup in the LERMA group. Model molecular ices of methanol (CH_3OH) [1,2] and ammonia (NH_3) [3,4] are irradiated either in the core-shell or in the valence-shell energy range, and the desorption of neutral species is monitored by mass spectrometry. Desorption efficiencies per incident/absorbed particle are compared for electron and photon irradiations, either as a function of incident energy in the low-dose regime (spectroscopic study), or as a function of cumulative dose at a fixed irradiation energy (kinetic study).



[1] D. Torres-Díaz et al., ChemPhysChem (2023) e202200912(1-7)

[2] R. Balsagete et al., A&A 647, A35 (2021)

[3] L.A. Sala, PhD Thesis, Univ. Paris-Sud, 2018 <https://theses.hal.science/tel-02073621>

[4] R. Dupuy, PhD Thesis, Sorbonne Univ., 2019, <https://tel.archives-ouvertes.fr/tel-02354689v2>