



HAL
open science

On certain partial sums involving squares of Hecke eigenvalues

Winfried Kohnen, Florian Luca

► **To cite this version:**

Winfried Kohnen, Florian Luca. On certain partial sums involving squares of Hecke eigenvalues. Hardy-Ramanujan Journal, 2024, 46, pp.1 – 5. 10.46298/hrj.2024.13031 . hal-04442692

HAL Id: hal-04442692

<https://hal.science/hal-04442692v1>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

On certain partial sums involving squares of Hecke eigenvalues

Winfried Kohnen and Florian Luca

Abstract. Let $a(n)$ be the n th Fourier coefficient of a cuspidal Hecke eigenform of even integral weight $k \geq 2$ and trivial character that is a normalized new form for some level N . We show that the partial sums

$$H_n = \sum_{m=1}^n a(m)^2/m^k$$

are not integral for $n \geq n_0$.

Keywords. Fourier coefficients of automorphic forms.

2010 Mathematics Subject Classification. 11F30, 11F11, 11N36

1. Introduction and Results

Let

$$h_n := 1 + \frac{1}{2} + \cdots + \frac{1}{n} \quad (n \in \mathbf{N})$$

be the n -th harmonic number. Harmonic numbers have been studied since antiquity and are important in various branches of number theory. They also have interesting arithmetic properties. For example, a result of Theisinger [The1915] asserts that h_n for $n \geq 2$ never is an integer. Except for the use of “Bertrand’s postulate” (cf. below) the proof is completely elementary.

If one seeks for a generalization in the context of modular forms one naturally is led to the partial sums formally attached to the Rankin-Selberg L -series at $s = k$ where k is the weight of the form in question. (For properties of the Rankin-Selberg zeta function we refer, e.g. to [Bu98, sect. 1.6].) Indeed, more precisely let

$$f(z) = \sum_{n \geq 1} a(n)e^{2\pi inz}$$

($z \in \mathcal{H} =$ complex upper half-plane) be a cuspidal Hecke eigenform of even integral weight $k \geq 2$ and trivial character that is a normalized new form of some level N . Recall that $a(1) = 1$ and $a(n)$ is the eigenvalue of the n -th Hecke operator $T(n)$. It is well-known that the $a(n)$ are algebraic integers and generate a totally real number field K . Let

$$D_f(s) := \sum_{n \geq 1} a(n)^2 n^{-s} \quad (\Re(s) \gg 1)$$

be the Rankin-Selberg L -series attached to f . Recall that $\sigma_0 = k$ is the abscissa of convergence of $D_f(s)$. We define

$$H_n = H_n(f) := \sum_{m=1}^n a(m)^2 m^{-k} \quad (n \in \mathbf{N}). \quad (1.1)$$

Then $H_n \in K$ and the H_n could be viewed as (at least) formal generalizations of the numbers h_n .

We thank episciences.org for providing open access hosting of the electronic journal *Hardy-Ramanujan Journal*

We prove:

Theorem 1. *Assume that $n \geq 8$ is an integer such that $a(p) \neq 0$ for some prime $p \in (n/2, n]$. Then H_n is not an algebraic integer.*

The proof of Theorem 1 works in a similar way as the proof in [The1915] with an additional use of Deligne's deep result

$$|a(p)| \leq 2p^{\frac{k-1}{2}}. \quad (1.2)$$

The assumption $n \geq 8$ (which implies that $p \geq 5$) comes from the fact that seemingly we do not have good information about the exact p -power dividing $a(p)$ for $p = 2, 3$, as will be clear from the proof.

2. Proof of Theorem 1

Assume that $n \geq 8$. Let p be any prime number in the interval $(n/2, n]$ which is guaranteed to exist by the hypothesis of Theorem 1. Further, $p \geq 5$. We denote by $\mathcal{O} \subset K$ the ring of algebraic integers of K . Suppose that $H_n \in \mathcal{O}$.

Let $C(n, p)$ be the n -th factorial, with the factor p omitted; i.e.:

$$C(n, p) := (p-1)! \cdot \prod_{\nu=1}^{n-p} (p+\nu).$$

We multiply both sides of (1.1) with $C(n, p)^k$. Our assumption implies that $C(n, p)^k H_n \in \mathcal{O}$. Hence, we infer that

$$C(n, p)^k \cdot \frac{a(p)^2}{p^k} \in \mathcal{O}$$

since the denominators of all other terms of the resulting sum on the right-hand side cancel. Therefore also

$$C(n, p)^{k/2} \cdot \frac{a(p)}{p^{k/2}} \in \mathcal{O}. \quad (2.3)$$

Now note that p does not divide any of the numbers $p+1, p+2, \dots, n$, because the next multiple of p , namely $2p$ is larger than n .

By the unique factorization of integral ideals into products of prime ideals in the Dedekind domain \mathcal{O} , applied to the principal ideals in question and using that p does not divide $C(n, p)$, we deduce from (2.3) that $p^{k/2} | a(p)$; i.e., $a(p) = p^{k/2} \alpha_p$ with $\alpha_p \in \mathcal{O}$.

From (1.2) we therefore find that

$$|\alpha_p| \leq \frac{2}{\sqrt{p}} < 1 \quad (2.4)$$

where for the last inequality in (2.4) we have used that $p \geq 5$.

Let σ be one of the finitely many embeddings of K over \mathbf{Q} into an algebraic closure of \mathbf{Q} . Then as is well-known

$$f^\sigma(z) := \sum_{n \geq 1} a(n)^\sigma e^{2\pi i n z} \quad (z \in \mathcal{H})$$

is a normalized cuspidal Hecke eigenform of weight k , trivial character and level N . We have $a(p)^\sigma = p^{k/2} \alpha_p^\sigma$ and

$$|\alpha_p^\sigma| \leq \frac{2}{\sqrt{p}} < 1.$$

From the above we therefore find that the norm of α_p satisfies

$$|N(\alpha_p)| = \left| \prod_{\sigma} \alpha_p^{\sigma} \right| < 1.$$

However, $N(\alpha_p)$ is a rational integer. Hence, we should have that $a(p) = 0$, which contradicts our assumption. This proves Theorem 1.

Remarks. i) For the Ramanujan τ -function, which are the Fourier coefficients of the discriminant function Δ of weight $k = 12$ (see [Leh47]), Lehmer conjectured that $a(n) \neq 0$ for all n . If this were true, then Theorem 1 together with a well-known theorem of Tchebyshev (also known under the name of Bertrand's postulate, see [HaWr59, p. 373, notes on chap. XXII]) would apply immediately to show that H_n is not algebraic integer for all $n \geq 8$.

ii) Theorems about the vanishing of $a(p)$ for primes p were proved by Serre in his seminal paper [Ser81]. Theorem 15 on Page 174 and Proposition 18 on page 180 there, give the following:

Serre's Theorem *Assume that $k \geq 2$ and $x \geq 2$.*

i) In case f is non-CM (there is no complex quadratic field \mathbf{L} such that $a(p) = 0$ for all primes p which are inert in \mathbf{L}), we have that

$$\#\{p \leq x : a(p) = 0\} = O\left(\frac{x}{(\log x)^{4/3}}\right).$$

ii) In case f is CM, we have that

$$\#\{p \leq x : a(p) = 0\} = \frac{x}{2 \log x} + O\left(\frac{x}{(\log x)^2}\right).$$

In either case, with n large and $x = n$, we conclude that the interval $(x/2, x]$ contains

$$\begin{aligned} \pi(x) - \pi(x/2) - \#\{p \in (x/2, x] : a(p) = 0\} &\geq \frac{x}{\log x} - \frac{x/2}{\log(x/2)} - \left(\frac{x/2}{\log(x/2)} - \frac{x/4}{\log(x/4)}\right) \\ &+ O\left(\frac{x}{(\log x)^{4/3}}\right) \geq \frac{x}{4 \log x} + O\left(\frac{x}{(\log x)^{4/3}}\right) \end{aligned} \quad (2.5)$$

primes p such that $a(p) \neq 0$. In the above, we used the Prime Number Theorem with the error term

$$\pi(x) = \frac{x}{\log x} \left(1 + O\left(\frac{1}{\log x}\right)\right) \text{ as } x \rightarrow \infty. \quad (2.6)$$

Since the last quantity indicated in the right-hand side of (2.5) is positive for large x , we conclude that for large n , the interval $(n/2, n]$ contains a prime p with $a(p) \neq 0$ and thanks to Theorem 1, we have that H_n is not an algebraic integer. Thus, we may write our conclusion as:

Theorem 2. *There exists $n_0 = n_0(f)$ such that H_n is not an algebraic integer for $n \geq n_0$.*

3. Comments and a challenge problem

We would like to suggest the following challenge question. Let ℓ, j be integers and put

$$H_{n,\ell,j} = \sum_{m=1}^n a(m)^\ell m^j.$$

Our paper discussed the case $\ell = 2$ and $j = -k$. We ask under what conditions on ℓ, j one may guarantee that $H_{n,\ell,j}$ is an algebraic integer for only finitely many n . When j is negative and large in absolute value (say $-j > \ell(k-1)/2$) the above argument applies. Indeed, the argument used in Theorem 1 creates an algebraic integer all whose conjugates have absolute values at most $2/p^{-j-\ell(p-1)/2}$ and since the exponent in the denominator is positive, this bounds p ; hence, n . If $\ell \geq 1$ and $j \geq 0$, then $H_{n,\ell,j}$ is an algebraic integer for all n , and the case when $\ell < 0$ is difficult since $a(m)$ might be zero (although we may decide to sum only over the values of $a(m)$ which are nonzero). The case $\ell \geq 1$ and $j \in [-\ell(k-1)/2, -1]$ is different since the argument of Theorem 1 creates an algebraic integer with large conjugates so it does not lead to a contradiction. As a toy example, we took $a(m) = \tau(m)$, the Ramanujan function of m and studied the borderline case $\ell = 2, j = -1$. Let n be large and assume $s \geq 1$ is such that 13^s is the largest power of 13 smaller than or equal to n . Then the only numbers $m \leq n$ divisible by 13^s are of the form $\ell \times 13^s$ where $\ell \in \{1, 2, \dots, K\}$ and $K \leq 12$ is some integer. Separating those ones out and multiplying everything by $A = 13^{s-1} \cdot \text{lcm}[p^t \leq n, p \neq 13]$, and using the multiplicativity of $\tau(m)^2$, we get that

$$AH(n, 2, -1) = \frac{(A/13^{s-1})\tau(13^s)^2}{13} \sum_{\ell=1}^K \frac{\tau(\ell)^2}{\ell} \pmod{\mathbf{Z}}.$$

But $13 \nmid \tau(13^s)$, $13 \nmid A/13^{s-1}$ and a small calculation with Mathematica reveals that

$$\sum_{\ell=1}^K \frac{\tau(\ell)^2}{\ell}$$

is not a rational number whose numerator is divisible by 13 for any $K \in \{1, \dots, 12\}$. Thus, $H(n, 2, -1) \notin \mathbf{Z}$ for any $n \geq 13$. Presumably there is an integer n_0 such that $H_{n,\ell,j}$ is not an integer for any $n \geq n_0, \ell \geq 1$ and $j < 0$. Note that since $p \mid \tau(p)$ for $p \in \{2, 3, 5, 7\}$, we have that $H_{n,\ell,j}$ is an integer for all $n \leq 10, \ell \geq 1$ and j negative and small in absolute value. Knowledge of non-ordinary primes p , namely primes p such that $p \mid \tau(p)$ would be useful. It has been checked that in addition to the above four, 2411, 7758337633 are the only ones below 10^{10} . See [MoSm13] for some results on this problem.

Acknowledgement. This work was done when the second author was visiting the Max Planck Institute for Software Systems in Saarbücken, Germany in Spring 2023. This author thanks Professor J. Ouaknine for the invitation and the Institute for financial support.

References

- [Bu98] D. Bump, *Automorphic forms and representations*, Cambridge University Press 1998.
- [HaWr59] G.H. Hardy and E.M. Wright, *An introduction to the theory of numbers*, 4th edition, Oxford 1959
- [Leh47] D.H. Lehmer, The vanishing of Ramanujan's function $\tau(n)$, *Duke Math. J.*, **14** (1947), 429–433.
- [MoSm13] B.K. Moriya and C.J. Smyth, Index-dependent divisors of coefficients of modular forms, *Int. J. Number Theory*, **9** (2013), 1841–1853.
- [Ser81] J.P. Serre, Quelques applications du théorème de densité de Chebotarev, *Inst. Hautes Études Sci. Publ. Math.*, **54** (1981), 123–201.

[The1915] L. Theisinger, Bemerkung über die harmonische Reihe, *Monatshefte für Mathematik und Physik*, **26** (1915), 132–134.

Winfried Kohlen

Mathematisches Institut der Universität Heidelberg

INF 205

69120 Heidelberg, Germany

e-mail: winfried@mathi.uni-heidelberg.de

Florian Luca

School of Maths, Wits University

South Africa

and

Centro de Ciencias Matemáticas UNAM

Morelia, Mexico

e-mail: florian.luca@wits.ac.za