



HAL
open science

Temporal Dynamics in an Attractor Model of the Songbird's Premotor Nucleus

Fjola Hyseni, Arthur Leblois, Nicolas P. Rougier

► **To cite this version:**

Fjola Hyseni, Arthur Leblois, Nicolas P. Rougier. Temporal Dynamics in an Attractor Model of the Songbird's Premotor Nucleus. COSYNE 2022 - Computational and Systems Neuroscience, Mar 2022, Lisbon, Portugal. hal-04443688

HAL Id: hal-04443688

<https://hal.science/hal-04443688>

Submitted on 7 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Temporal Dynamics in an Attractor Model of the Songbird's Premotor Nucleus

Fjola Hyseni, Arthur Leblois, Nicolas P. Rougier
 Univ. Bordeaux, CNRS, IMN, UMR 5293, F-33000 Bordeaux, France
 fjola.hyseni@u-bordeaux.fr

What do songbirds tell us about timing?





Synfire chains are however not robust to noise and function for a narrow range of feedforward weights (W , see figure below), requiring fine tuning during learning.



Attractors dynamics and timing


Here, we propose that HVC neuronal dynamics may be modelled by attractor dynamics, where recurrent connections allow the formation of a travelling activity bump with robust functional properties.

Net Connectivity Profile Feature space (x : preferred timing during song)



1. Activity profile in HVC

The width of the activity bump, and thus the duration of transient neuronal activation, can be decreased to reproduce the brief activity bursts of HVC neurons.





2. Modifying syllable duration

Relying on a reward covariance rule, we show that the duration of a syllable can be modified in response to a perturbed reward profile, as implemented in a widely used lab conditioning paradigm.

$$\Delta W_{ij} = \gamma \int_0^{t^{(k)}} R(t) e_{ij} dt$$

γ : learning rate parameter, $R(t)$: the reward, η : noise at the postsynaptic neuron, r_j : rate of the presynaptic neuron.


$$e_{ij} = \int_0^t \frac{dt'}{\tau_e} e^{-(t-t')/\tau_e} \eta_i(t') r_j(t')$$



3. Local inhibition in HVC

Local inhibition administered to the HVC.

$$\text{Inhibition} = \alpha \frac{1}{\sigma \sqrt{2\pi}} \exp \left(-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right)$$



Conclusion


We propose that HVC neuronal dynamics may be modelled using the ring model.

The duration of a syllable can be modified in response to a perturbed reward profile and consistent with behavioral data, the change in duration is specific to the target syllable.

We derive a new prediction from the model. Following local muscimol (GABA_A agonist) injection in HVC, in contrast to the synfire chain, a perturbation in the initial phase of the pattern formation would delay song initiation, but singing would be sustained.

Future perspectives:

A spiking network (with adapting LIF neurons) to replicate the behaviours acquired from the rate model, i.e 10ms bump width, learning and prediction.



- References
- Hansel, D., & Sompolinsky, H. (1998). 13 Modeling Feature Selectivity in Local Cortical Circuits. 69
 - Ali F. Pehlevan C and Olveczky. Nat Commun 9, 977, 2018.
 - Williams, R. J. (1992). Machine Learning, 8(3).
 - Zhang, K. (1996). Journal of Neuroscience, 16(6).
 - Mauk, M. D., & Buonomano, D. V. (2004). Annual Review of Neuroscience, 27(1).
 - Hahnloser, R., Kozhevnikov, A. & Fee, M. Nature 419,(2002)
 - Brainard, M. S., & Doupe, A. J. (2002). Nature, 417(6886).