

The detection of vulnerable road users by younger and older drivers

Maud Ranchet, Roland Brémond, Prashant Pala, Michèle Colomb, Viola

Cavallo

► To cite this version:

Maud Ranchet, Roland Brémond, Prashant Pala, Michèle Colomb, Viola Cavallo. The detection of vulnerable road users by younger and older drivers. Transportation Research Part F: Traffic Psychology and Behaviour, 2022, 91, pp.357-367. 10.1016/j.trf.2022.10.018 . hal-04444293

HAL Id: hal-04444293 https://hal.science/hal-04444293

Submitted on 12 Feb 2024 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Detection of Vulnerable Road Users by Younger and Older Drivers

Maud Ranchet^{1*}, Roland Brémond², Prashant Pala², Michèle Colomb³, and Viola Cavallo² ¹ TS2-LESCOT, Univ Gustave Eiffel, IFSTTAR, Univ Lyon, F-69675 Lyon, France ² COSYS-PICS-L, Univ Gustave Eiffel, IFSTTAR, F-77454 Marne-la-Vallée Cedex 2, France ³ CEREMA, 8-10, rue Bernard Palissy – 63017 Clermont-Ferrand Cedex 2, France

* Corresponding author:

Maud Ranchet, PhD Université Gustave Eiffel – TS2-LESCOT Cité des Mobilités 25, Avenue François Mitterrand Case 24 F-69675 Bron Cedex, France phone: +33 (0) 4 72 14 25 99 email: maud.ranchet@univ-eiffel.fr

Highlights

- The aim was to investigate the effect of aging on the detection, under time constraints, of vulnerable road users
- The detection of vulnerable road users decreased with aging, even in the younger-old group
- Older-old adults compensated for their visual-field decline by doing more visual exploration than the other two groups did to correctly detect vulnerable road users
- Further studies should investigate the detection of vulnerable road users, including those on personal mobility devices and using different exploration times.

Abstract

The detection of vulnerable road users (VRUs), especially under time constraints, may be impaired in elderly drivers, due to their visual and cognitive decline. This represents a major concern for road safety. The objective of the present study was to investigate the effect of aging on the detection of VRUs. A further aim was to investigate the impact of external factors on VRU detection. Twenty-two young adults, 20 younger-old adults, and 32 older-old adults were included in the study. A series of photographs were displayed for 500 ms. Participants were asked to detect the VRU, which could be a motorcyclist, a cyclist, or a pedestrian. The VRU was located at one of two distances (near/far), one of two locations (off-centered/centered), and in two car-DRL (daytime running lights) environments (on/off). The ability to correctly detect a VRU was measured. An eve-tracker was used to record eve movements. The main findings showed that VRU-detection performance decreased with aging, even in the younger-old group (ages 55 to 68). The ability to correctly detect a VRU by older-old adults was poorer particularly when the VRU was a motorcyclist or a cyclist. As a whole, the older-old adults made more fixations to correctly detect the VRU than the other two age groups did. Moreover, the visual angle between the gaze of the participant's last fixation and the target in the older-old group was lower than in the other two groups, particularly when the VRU was off-centered. This finding suggests that older-old adults compensated for their visual-field decline by doing more visual exploration than the other two groups did to correctly detect the VRU. The results are discussed with regards to age-related cognitive and visual deficits.

Word count: 279

Keywords: older drivers; vulnerable road users; detection performance; visual exploration

1 **1.** Introduction

2 The safety of vulnerable road users (VRUs), including pedestrians, cyclists, and 3 motorcyclists, is a major concern in many countries (WHO, 2018) where they have a significantly higher risk of being killed or seriously injured than do car drivers. VRU 4 fatalities currently represent 42.5% of all fatalities in France (ONISR, 2019), 46.5% in 5 6 Europe (ERSO, 2018), and 56% in the world (WHO, 2018). With the COVID-19 7 pandemic situation, daily mobility of individuals has considerably changed (Li et al., 8 2022): the number of VRUs, especially in the big cities is constantly growing. A further 9 increase in the number of VRUs can be expected due to a rise in petrol prices but 10 also due to ecological and health considerations.

11 Main factors that could explain poor VRU detection

In order to safely interact with vulnerable road users, the drivers of motorized vehicles 12 13 must detect them quickly. In fact, late detection is the basic driver error that leads to 14 collisions (Rumar, 1990). Besides expectation errors (i.e., failure to scan for a specific 15 class of road users), Rumar et al. (1990) mentioned perceptual difficulties, especially 16 when low-contrast conditions and/or the peripheral visual field of the observer are involved. VRUs are often more difficult to detect than cars because of their smaller 17 18 size and irregular contours (Cavallo & Pinto, 2012). The fact that they are not very 19 conspicuous visually and fail to be perceived by other vehicle drivers are frequent 20 causes of accidents (Van Elslande et al., 2012). A previous study showed that drivers 21 detect pedestrians less often when they appear in urban areas and more often when 22 they appear in residential areas (Borowsky et al., 2012). In urban areas, drivers are 23 not necessarily attuned to pedestrians, whereas in residential areas, their mindset is 24 such that they expect and search for pedestrians. Furthermore, findings have

25 demonstrated that driving experience is also a dominant factor that affects the 26 identification of pedestrians. Regarding motorcycle accidents, for instance, it has 27 been shown that poor motorcycle detectability is one of the main risk factors and 28 leads to right-of-way violations (ACEM, 2009; Hurt et al., 1981). While motorcycle 29 conspicuity has improved since the use of daytime running lights (DRLs, which 30 became compulsory in the seventies in many countries), this conspiculty advantage 31 has been jeopardized by the increasing use of DRLs on cars, which became 32 compulsory in Europe in 2011. Previous experimental studies have shown that car 33 DRLs, which act as a kind of visual noise, hamper motorcycle detection (Brendicke 34 et al., 1994; Cavallo & Pinto, 2012), and also have a detrimental effect on the 35 detection of bicyclists and pedestrians (Cavallo & Pinto, 2012). Although rarely 36 studied, VRU detectability, especially in a car-DRL environment, represents a major 37 concern for road safety.

38 The impact of age-related declines on VRU detection

39 Perceptual difficulties in detecting VRUs can also be caused by factors internal to 40 drivers who are visually impaired and/or exhibit age-related declines. It has been shown that elderly drivers are over-represented in accidents involving pedestrians 41 42 (Bromberg et al., 2012) and motorcyclists (Pai et al., 2009). While speed/distance judgment errors are often mentioned (e.g., Hancock & Caird, 1993; Oxley et al., 2006; 43 44 Pai, 2011), detection failures in elderly drivers, together with longer perception and 45 reaction times at intersections, have also been observed (e.g., Staplin, 1995). Using a video observation technique, the Bromberg et al. study (2012) showed that elderly 46 47 drivers had longer response times on hazard detection, especially for pedestrianrelated hazardous events. 48

Late detection and failure to detect among elderly people are classically explained in terms of an age-related deterioration of visual capacities, as reflected by declines in visuo-attentional abilities, and also in terms of less effective visual exploration strategies.

53 Among the age-related sensory changes, we find impaired visual acuity in both 54 central and peripheral vision (Collins et al., 1989; Derefeldt et al., 1979; Harwood, 55 2001; Ivers et al., 1998; Owsley et al., 1983), low motion sensitivity (Sekuler et al., 56 2000; Snowden & Kavanagh, 2006), and poor adaptation to the dark (McMurdo & 57 Gaskell, 1991). The visual field, as measured in optometry, is also subject to an age effect (Coleman et al., 2007; Freeman et al., 2007; Jaffe et al., 1986), with a greater 58 59 decrease in peripheral than in central vision. Contrast sensitivity also deteriorates 60 with age (Elliott, 1987), particularly at high spatial frequencies, i.e., the ability to see 61 details.

62 The useful field of view test (Ball et al., 1988) refers to the ability, in one glance, to 63 detect a peripheral visual target and to identify a central visual target. Findings from a computer-screen task (UFOV^R), have consistently shown that the ability to detect 64 peripheral targets in this dual task deteriorates with age, and that the difference 65 66 between younger and older people becomes greater as the eccentricity of the peripheral target increases (Ball et al., 1988; Scialfa et al., 1987; Sekuler & Ball, 67 68 1986). This is partly due to the lowering of processing speed with ageing, as demonstrated by Salthouse (Salthouse, 1996). 69

Similar age effects have also been observed when participants performed a task of
detecting visual targets in a road scene while driving on a simulator (Rogé et al., 2003,
2004; Rogé & Pébayle, 2009). It has been shown that the accident risk was much

73 higher for elderly drivers when their useful field of view was reduced by 40% or more (Ball & Owsley, 1993; Owsley, 1994; Sims et al., 2000). The poor UFOV performance 74 75 of participants referred for fitness-to-drive assessment was found to be associated 76 with a large number of driving errors during real-world driving (Myers et al., 2000). 77 Similar results have been found on a driving simulator (Henderson et al., 2013). With 78 respect to the detection of VRUs, Rogé and Pébayle's (2009) simulator study showed 79 that drivers with a limited useful field of view detected fewer motorcycles in the road scene, especially in the distant periphery, than those who had a more extensive useful 80 81 field of view. Although that study only involved younger drivers, it can be assumed that such motorcycle-detection difficulties are more pronounced for older drivers, 82 83 whose UFOV performance is generally impaired.

It is also well known in vision science that age impacts eye movements. While age does not seem to affect eye-movement accuracy (Moschner & Baloh, 1994), it has been shown to impact saccadic speed (Abel et al., 1983), saccadic latency, and saccadic reaction time (SRT) (Moschner & Baloh, 1994; Munoz et al., 1998). More specifically, an effect of age on SRT has been observed for persons over 40.

89 <u>Objectives</u>

The number of older drivers is increasing and their age-related declines may negatively impact the detection of VRUs, which may represent a major concern for road safety. It is surprising to notice that the detection of VRUs by older drivers has been very little investigated (Borowsky et al., 2012; Bromberg et al., 2012). The first objective of the present study was therefore to investigate the effect of aging on the detection of VRUs. Three age groups (young adults, younger-old adults, older-old adults) participated in the study. We expected an age-related decrease in the

97 detection of VRUs. We also hypothesized that a limited useful field of view, as
98 measured by the UFOV test, would be associated with poor detection of VRUs.

Another objective was to investigate the extent to which the detection of VRUs is affected by external factors such as the presence/absence of daytime running lights on cars, the kind of VRU to be detected (pedestrian, cyclist, motorcyclist), their distance from the observer, and their location in the visual scene. The exploratory aim was to better understand the visual strategies used by older drivers to correctly detect a VRU.

105 **2. Method**

106 2.1 Participants

107 A total of 74 participants were included in the study. In the group of young adults 108 (N=22), there were 10 males and 12 females. Their median age was 29 years (range 25-46) and their median number of years of driving experience was 5 (range 1-25). 109 110 In the group of younger-old adults (N=20), there were 16 males and 4 females. Their 111 median age was 61.5 years (range 55-68) and their median number of years of driving 112 experience was 42 (age 15-50). As cognitive decline often occurs at older ages (70 113 or more) (Aartsen et al., 2002), we added a third group of older-old adults (N = 32) whose median age was 73 years (range 70-87) and whose median number of years 114 of driving experience was 51 (range 4-70). There were 19 males and 13 females. 115

The inclusion criteria were: (1) having normal or corrected-to-normal vision (at least
6/10), (2) being a driving-license holder, (3) being an active driver (driving more than
2500 km/year), and (4) not declaring any neurological disorders.

None of the participants were professional drivers. Participants were recruited by
flyers or social networks. Written informed consent was obtained from all participants
before the experiment.

122 2.2 Task and experimental design

123 A series of photographs were displayed for 500 ms (for an example, see Figure 1). The participants had to detect the presence of a VRU and determine whether it was 124 125 a motorcyclist, bicyclist, or pedestrian. They had to choose between four response 126 types displayed on the screen: bicyclists, motorcyclists, pedestrians, or none of the above. They indicated their response to the experimenter orally. Practice trials were 127 performed before the experiment. A mixed experimental design was employed, with 128 group (young, younger-old, older-old) as a between-subject variable. There were four 129 within-subject variables: the targeted VRU (motorcyclist, bicyclist, or pedestrian), the 130 distance of the VRU in the scene (far vs. near), the DRL environment of the cars 131 (DRLs on vs. off), the VRU's *location* (centered vs. off-centered). Each experimental 132 133 condition was repeated three times (VRU (3) × Distance (2) × Car-DRL (2) × Location (2)), making for a total of 72 experimental trials. Eighteen distractors containing a 134 VRU (located at different distances from those used in the experimental trials) were 135 136 added. In order to obtain a balanced design, 90 distractors with no target were added, 137 making a total of 180 trials. The stimuli were presented in four blocks of 45 trials each.



Figure 1. Example of a trial involving a motorcyclist (a scooter) centered at a neardistance, among cars with their daytime running lights off

141

142 2.3 Apparatus and stimuli

143 The stimuli were photographs of urban traffic at intersections in overcast weather. The photographs were presented on a 47" flat-panel LCD display (width: 1.02 m; 144 145 height: 0.57 m). Participants were seated 1.60 m from the screen (angular size of the 146 screen: 32.5° × 19.6°). The eye-tracking system was SMI (SensoMotoric 147 Instruments), and the sampling rate was 60 Hz. The data was processed using BeGaze (SMI software). Participants performed Part 3 of the UFOV test, considered 148 to be a measure of selective attention (Ball et al., 1993). In this part of the test, an 149 150 object (truck or car) was presented in the center of the screen. At the same time, another object (truck or car) was presented in the periphery among 47 distractor 151 triangles of the same size, contrast, and luminance. Following the identification of the 152 153 central target object, participants had to indicate at which location the peripheral object was displayed. 154

156 2.4 Procedure

All participants first took a visual acuity test (Ergovision) and Part 3 of the UFOV test. 157 158 Then the participants read the instructions of the experiment. The order of the four blocks was counterbalanced across participants. Each trial included the display of a 159 fixation cross for 1500 ms, followed by the photographs for 500 ms. The screen turned 160 161 gray at the end of the trial. Then the participants had to choose between four response types displayed on the screen. No feedback on response accuracy was given. 162 163 Participants were allowed breaks during the blocks and between the blocks. The experimental session lasted approximately one hour and 30 minutes. 164

165 2.5 Dependent variables and statistical analyses

166 2.5.1 Dependent variables

167 For each experimental trial, we determined the number of correct detections, 168 identification errors, and misses. A correct detection was counted when the right VRU category was recognized. An identification error was counted when a VRU was 169 170 detected but not correctly identified, and a miss was counted when the VRU was not detected at all. Performance on distractor items was not analyzed. The number of 171 fixations, known to be an indicator of visual exploration, was analyzed. Given that 172 visual-field size decreases and peripheral vision declines with aging, the visual angle 173 between the gaze of the last fixation and the target was also analyzed. To better 174 175 understand the visual strategies used by older drivers to correctly detect a VRU, the 176 number of fixations and the visual angle were analyzed specifically on correct 177 detections. However, differences in visual strategies between correct detections and 178 no detections (including misses and identification errors) were also explored for all participants and for each age group. For the UFOV test, the presentation time inmilliseconds (ms), necessary for correct identification, was measured.

181 2.5.2 Statistical analyses

A binary logistic regression was performed on the ability to correctly detect a VRU. This was done in the framework of the generalized linear mixed model (GLMM) in order to take into account the individual random effect, using the SPSS v.26s Genlinmixed procedure (see Agresti, 2013, for more details). The binary response on each trial (yes = 1, no = 0, 0 being the reference category), indicating the ability to correctly detect a VRU, was chosen as the dependent variable. A binomial distribution and a logit link function were assumed.

The number of fixations during a trial, as well as the visual angle between the gaze (of the last fixation) and the target, were analyzed for correct detections. To analyze the number of fixations on correct detections, a Poisson regression in the framework of the GLMM model was conducted. To analyze the visual angle between the gaze (of the last fixation) and the target on correct detections, a linear regression was performed in the framework of the GLMM model.

195 All of these models (binary logistic regression, Poisson regression, and linear regression) included five predictors that were categorical variables. The four within-196 subject variables were VRU type (pedestrian, motorcyclist, cyclist), VRU distance (far, 197 198 near), car-DRL environment (on, off), and VRU location (centered, off-centered). The 199 single between-subject variable was age group. For the purpose of this study, only 200 the interactions between age group and VRU type, age group and distance, age 201 group and location, and age group and car-DRL environment were examined. Posthoc comparisons for the GLMMs were performed using t-tests. Significance levels 202

were adjusted for multiple comparisons using the sequential Bonferroni correction (or 203 Holm's sequential Bonferroni procedure; Holm, 1979). Successive models were 204 205 tested by removing non-significant terms one by one until the best fit was found using the corrected Akaike Information Criterion (cAIC), which decreases with increasing 206 207 model fit. Given the small number of identification errors and misses, we did not 208 examine the effect of within-subject variables on errors and misses. However, paired t-tests for all participants and for each group were performed between correct 209 detections and no detections for the number of fixations and the degree of visual 210 211 angle to determine whether there are any significant differences in visual strategies 212 between correct detections and no detections.

For identification errors, misses, and UFOV performance, the data were not normally distributed. Values are expressed as medians (Q1-Q3). Between-group differences were examined using Kruskal-Wallis tests (KW). Post-hoc analyses used Wilcoxon rank-sum tests, with Holm's correction for multiple comparisons (Holm, 1979).

Pearson correlations between UFOV performance and VRU detection performancewere conducted for all participants.

All statistical analyses were performed using SPSS, version 26.0.

220 **3. Results**

3.1 Detection task

3.1.1 Ability to correctly detect a VRU

Table 1 presents the findings for the GLMM. A significant main group effect was found on the ability to correctly detect a VRU. Post hoc tests showed that the older-old participants detected the VRUs significantly less well than both the younger-old and young groups did (p < 0.001), and the younger-old adults detected the VRUs significantly less well than the young adults did (p < 0.001). The mean (\pm standard deviation) proportion of correct detections was 0.55 \pm 0.01 for the older-old adults, 0.67 \pm 0.01 for the younger-old adults, and 0.82 \pm 0.01 for the young adults.

For all within-group variables, significant main effects on the ability to correctly detect 230 231 a VRU were also found (Table 1). Post-hoc comparisons showed that the mean proportion of correct detections was significantly lower in the far condition (0.57 ± 232 233 0.01) than in near conditions (0.79 \pm 0.001). The proportion of correct detections was also significantly lower in off-centered than in centered conditions. For location, the 234 values were 0.78 ± 0.001 for centered VRUs and 0.59 ± 0.01 for off-centered VRUs. 235 The proportion of correct detections was also lower for the DRLs-on than for the 236 237 DRLs-off condition, but this difference was not significant. For the car-DRL 238 environment, the values were 0.71 ± 0.01 for DRLs off and 0.67 ± 0.01 for DRLs on.


239 Table 1. Fixed effects using GLMM

Age group	<i>F</i> _{2,4842} = 115.17, p < 0.001	
Within-group variables		
Type of VRU	<i>F</i> _{2,4842} =43.02, p < 0.001	
VRU distance	<i>F</i> _{1,4842} = 219.28, p < 0.001	
VRU location	<i>F</i> _{1, 4842} = 172.98, p < 0.001	
Car-DRL environment	<i>F</i> _{1, 4842} = 5.00, p = 0.02	

240

Figure 2 shows the estimated mean proportion of correct detections per age group 241 for each VRU type. The older-old adults detected significantly fewer pedestrians, 242 243 motorcyclists, and cyclists than the young adults did (p < 0.001); the older-old adults detected significantly fewer motorcyclists and cyclists than the younger-old adults did 244 (p < 0.001); and the younger-old adults detected significantly fewer motorcyclists, 245 246 cyclists, and pedestrians than the young adults did (p < 0.001). Pairwise comparisons also revealed that older-old adults detected significantly more pedestrians than 247 248 cyclists and motorcyclists (p < 0.001). Younger-old adults detected significantly more pedestrians and motorcyclists than cyclists (p < 0.001). Young adults detected more 249 motorcyclists than cyclists and pedestrians (p < 0.001). They also detected more 250 251 pedestrians than cyclists (p < 0.001).

No significant interactions between age group and VRU distance, age group and VRU
location, or age group and car-DRL environment were found.



255 *** p < 0.001

Figure 2. Estimated proportion of correct detections as a function of age group and VRU type. Only significant between-group differences are shown in this figure. Error bars represent the standard error.

259

260 3.1.2 Identification errors and misses

Older-old adults made more identification errors than young adults (p < 0.01). The median percentage of identification errors for older-old adults was 5.59 (4.17 – 10.14) and for young adults, it was 2.78 (1.41 – 4.17). Both older-old adults and younger-old adults missed more VRUs than young adults did (p < 0.01). The median percentage of misses for older-old adults was 37.50 (29.13 – 42.85). For younger-old adults, this value was 30.07 (22.22 – 36.11), and for young adults, it was 19.44 (11.94 – 23.61).

267 3.2 Eye-related variables for correct detections


For the number of fixations, the final model included the significant main effects of distance ($F_{1,3157} = 18.47$, p < 0.001), location ($F_{1,3157} = 8.28$, p = 0.004), and age group ($F_{2,3157} = 204.93$, p < 0.001). For the visual angle, the final model included the significant main effects of distance ($F_{1,3153} = 8.78$, p = 0.003), location ($F_{2,3153} =$ 2507.81, p < 0.001), age group ($F_{2,3153} = 16.09$, p < 0.001), and a significant distance by age group interaction ($F_{2,3153} = 3.86$, p = 0.02) as well as a significant location by age group interaction ($F_{2,3153} = 59.64$, p < 0.001).

There were a greater number of fixations in far VRU locations than in near ones. The mean number of fixations in a far location ($M \pm SD$) was 1.77 ± 0.02; in a near location, it was 1.68 ± 0.02, p < 0.001. The findings also revealed a significant main effect of location, with a significantly higher average number of fixations in the off-centered condition than in centered conditions. The mean number of fixations ($M \pm SD$) in off-centered condition was 1.76 ± 0.02 . In the centered condition, it was 1.69 ± 0.02 , p = 0.004.

There was a significant main effect of age group on the number of fixations for correct 282 283 detections, and on the visual angle between the gaze and the target. Post-hoc tests showed that the older-old group had a significantly higher number of fixations than 284 285 either younger groups. The mean number of fixations for the older-old adults was 286 2.05 ± 0.02 . For the younger-old adults, it was 1.51 ± 0.02 , and for the young adults, it was 1.61 ± 0.02 (p < 0.001 on all pairwise comparisons). The older-old group had 287 288 smaller visual angle than the younger-old group (p < 0.001) and the young group (p= 0.001). The visual angle in degrees for the older-old adults was 5.05 ± 0.09 . For the 289 290 younger-old adults, it was 5.85 ± 0.11 , and for the young adults, it was 5.49 ± 0.09 (p 291 \leq 0.01 on all pairwise comparisons).

292 Interaction between age group and distance

The older-old adults had a significantly smaller visual angle than younger-old group did in both the near and far locations (p < 0.001, p = 0.009, respectively). In the far location, the older-old adults had smaller visual angle than did the young group (p = 0.003) (Figure 3). Only the younger group had a visual angle greater in the far location than in the near location (p < 0.001).



299 **p<0.01


300 ***p<0.001

Figure 3. Visual angle by age group, for each distance. This represents the estimated values from the
 final model. Only significant between-group differences are shown in this figure. Error bars represent
 the standard error.

304

305 Interaction between age group and location

The older-old group had a significantly smaller visual angle than did the younger-old group and the young group in the off-centered condition (p < 0.001 on all pairwise comparisons) (Figure 4). The younger-old group also had a greater visual angle than the young group did (p < 0.001) in the off-centered condition. In the centered condition, the older-old group had a greater visual angle than the younger groups did (p = 0.001 on all pairwise comparisons). For all age groups, the visual angle was greater when the VRU was off-centered than when it was centered (p < 0.001).



314 **p<0.01




Figure 4. Visual angle by age group, for each location. This represents the estimated values from the final model. Only significant between-group differences are shown in this figure. Error bars represent the standard error.

319 3.3 Differences in visual strategies between correct detections and no detections

For the number of fixations, there was no significant difference between correct 320 detections and no detections (correct detections = 1.76 ± 0.30 versus no detections 321 = 1.77 ± 0.29 , t = 0.99, p = 0.33). Furthermore, no significant differences between 322 323 correct detections and no detections were found for each age group (P > 0.05). However, a significant difference was found between correct detections and no 324 325 detections for the visual angle (Visual angle for correct detections = $4.96^{\circ} \pm 0.98$ versus Visual angle for no detections = $8.24^{\circ} \pm 1.48$, t = 0.99, p < 0.001). These 326 327 differences remain significant for each age group (Younger adults: Visual angle for correct detections = $5.21^{\circ} \pm 0.67$ versus Visual angle for no detections = $7.31^{\circ} \pm 1.32$, 328 329 t = 6.66, p < 0.001; younger-old adults: Visual angle for correct detections = $5.25^{\circ} \pm$ 0.59 versus Visual angle for no detections = $7.93^{\circ} \pm 1.17$, t = 8.04, p < 0.001; Older-330 16

old adults: Visual angle for correct detections = $4.64^{\circ} \pm 1.23$ versus Visual angle for no detections = 9.05 ± 1.32 , t = 14.29, p < 0.001).

333

334 3.3 UFOV test

The younger-old and older-old groups performed significantly less well on Part 3 of

the UFOV test than the young group did (p < 0.001). The young adults had a median

337 (Q1 - Q3) of 103 ms (80 - 140). The younger-old adults' median (Q1 - Q3) was 300

ms (257 - 407) and the older-old adults' median (Q1 - Q3) was 335 ms (253 - 500).

339 3.4 Correlation between Part 3 of the UFOV test and detection performance

For all participants, poor performance on Part 3 of the UFOV test was strongly correlated with poor VRU detection (r = -0.67, p<0.001).

342 **4. Discussion**

343 The aim of this study was to investigate the effects of aging on the detection of VRUs. The VRU could be a pedestrian, a motorcyclist, or a cyclist, positioned in two different 344 locations and at two different distances in urban traffic. The cars could have their 345 DRLs turned on or off. The contribution of this study is that VRU detection 346 performance decreased with age, even in the younger-old group (ages 55 to 68). 347 348 Furthermore, older-old adults performed poorly, particularly when the VRU was a motorcyclist or a cyclist. Finally, this study suggests that older participants used 349 350 compensatory mechanisms for their poorer visual field by increasing visual exploration, without reaching the detection performance of the younger group. 351

As a whole, the older-old adults made more fixations to correctly detect the VRU than the other two age groups did. As expected, there were a greater number of fixations

to far VRU locations than to near ones. The mean number of fixations to detect the VRU in a far location was 1.77 whereas in a near location, it was 1.68, meaning that more individuals needed 2 fixations to detect far VRUs as compared to near VRUs. Moreover, the visual angle between the detected VRU and the last fixation was smaller in the older-old group than in the younger-old group, whether the VRU was in the far or near location. The visual angle of the older-old group was also smaller than that of the other two groups when the VRU was off-centered.

361 Importantly, detection performance decreased with aging, not only in the older-old participants (ages 70 to 87) but even in the younger-old group (ages 55 to 68). These 362 findings are consistent with previous driving studies showing that age has an impact 363 on the ability to detect visual signals (Rogé et al., 2003, 2004). Poor detection 364 365 performance may be explained by a deterioration of selective attention with aging. In 366 the present study, lower detection performance was strongly associated with a poorer measure of selective attention (Part 3 of the UFOV). Previous findings have shown 367 368 that a decrease in selective attention appears early in the course of aging, between the ages of 40 and 50 (Sekuler et al., 2000). Another explanation could be the slowing 369 of processing speed with aging (Owsley, 2011; Salthouse, 1996). In our study, 370 371 participants were under time constraints when detecting a VRU since the 372 photographs were displayed for only 500 ms. This could explain why even the younger-old adults could not compensate for their slowing in processing speed and 373 exhibited a decrease in VRU detection performance. Older-old adults made more 374 identification errors and both older groups (older-old and younger-old) missed more 375 376 VRUs than young adults did, suggesting that they did not have time to see and identify 377 the VRU.

Unexpectedly, the interaction between age group and centeredness on the ability to 378 379 detect a VRU was not significant. Yet, for correct detections, the visual angle between 380 the gaze and a VRU located in the periphery was narrower for older-old adults than 381 for younger-old and young adults. This finding indicates that older-old adults' eyes 382 need to land closer to the target to correctly detect a VRU in the periphery. It suggests 383 that the older-old adults used mechanisms that compensate for visual-field decline by doing more visual exploration than the other two groups to correctly detect the VRU 384 (Coeckelbergh et al., 2004). Our findings on the detection of VRUs located in the 385 386 periphery differ from those of previous studies (Rogé et al., 2003, 2004; Rogé & 387 Pébayle, 2009), which showed that older drivers detected fewer signals than younger 388 drivers, particularly when the signal was farther away from the fixation point. 389 Discrepancies across studies may be due to differences in methodology (screen size) 390 or the type of task (observation versus driving). Interestingly, the number of fixations did not differ between correct detections and no detections whereas the degree of 391 392 visual angle was larger for no detections, for all age groups. This result suggests that 393 visual strategies of participants differ between correct detections and no detections: 394 all participants (regardless of the age) need to land closer to the target to correctly 395 detect a VRU.

Similarly, no age-related deterioration in the ability to detect a VRU among cars with their DRLs on was found in our study. Yet, the use of DRLs had a negative impact on detection performance in all three age groups. Previous studies have shown that the use of DRLs by cars may create "visual noise" that hinders the perception of other VRUs, such as pedestrians, cyclists, or motorcyclists (Cavallo & Pinto, 2012), which makes them less conspicuous and less noticeable (Brouwer et al., 2004). Our results

thus suggest that the use of DRLs by cars affects the detection of VRUs for allindividuals, irrespective of their age.

Another interesting finding is that older-old adults and younger-old adults detected pedestrians better than they did motorcyclists and/or cyclists, which was not the case for our young adults. One explanation could be that older adults identified more with pedestrians than with motorcyclists and cyclists.

408 **5. Limitations and future research avenues**

409 Methodological considerations include the use of static images, i.e., photographs of urban traffic instead of dynamic images in videos or during simulated driving. The 410 411 decision to use static images in the present study allowed us to investigate attentional aspects of road-user behavior. Photographs were displayed for 500 ms, which is quite 412 413 short. In real life, drivers need to quickly detect relevant information in the road scene, 414 and detection failures are more likely to occur under time pressure and when visual 415 exploration time is short. Also, the age gap between our younger-old and older-old 416 groups was quite close (65-68 vs. 70-87). However, our findings indicated some 417 significant differences in VRU detection performance and visual exploration. 418 Furthermore, the age gap between the older groups and the younger group was 419 relatively large, which can explain significant differences between these two groups. 420 Additional studies that include an intermediate age group with a mean age around 50 421 should be conducted to better understand the time course of visual and attentional 422 aging in VRU detection. Time constraints play a major role in detection performance 423 in older adults. In the present study, photographs were displayed for 500 ms, which 424 may be not enough to correctly detect the VRU, especially for older adults. Future

research should vary presentation times of photographs in order to determine the roleof exploration times in relation with compensation strategies in older adults.

It would be interesting in future studies on older adults to investigate VRU detection using a driving simulator. The use of a driving simulator enables one to assess VRU detection during driving, and to better understand the compensatory strategies of older adults. A previous simulator study showed that elderly drivers drove about 20% slower than experienced non-elderly drivers (Bromberg et al., 2020), which suggests that as a whole, elderly drivers are aware of their deficiencies in detecting hazards.

A valuable contribution for improving road safety would be to investigate different 433 434 types of interventions aimed at helping older drivers better detect VRUs, particularly motorcyclists and cyclists. Based on the results of this study, we can contend that 435 older individuals detect fewer motorcyclists and cyclists than pedestrians. There are 436 437 several ways to improve VRU detection performance. A variety of training methods 438 have been employed to improve VRU detection by car drivers. One such method is 439 useful-visual-field training which could improve elderly car drivers' VRU-detection 440 (Rogé et al., 2014). It has been shown that drivers who had undergone training for 441 perceiving motorcycles were better able to identify motorcyclists than ones who had 442 not been trained (Crundall et al., 2017). Also, safety messages have been developed 443 to improve the perception of VRU vulnerability during driving (Rogé et al., 2015).

Additional research needs to be conducted to explore the benefits of automated vehicles for detecting VRUs, particularly among older adults. The introduction of automated vehicles with external human-machine interfaces, and the use of augmented reality in automated vehicles, could help drivers to better detect VRUs and may also enhance future interaction between VRUs and autonomous vehicles 21 (Tabone et al., 2021). Several safety systems such as pedestrian alerts could inform
drivers of a potential hazard (Large et al., 2019). However, the additional cognitive
load induced by these systems should be considered with caution in future studies,
especially for older adults.

As a complement to studies investigating VRU detection by drivers, eye-tracking
studies should also explore the interactions between drivers and pedestrians from the
pedestrian's point of view (Lévêque et al., 2020).

Further research should also help improve the visibility of VRUs. Previous studies have shown that the use of specific conspicuous clothes (for example, retroreflective and yellow clothes) by cyclists, or having specific headlights on motorcycles could improve their detection (e.g., Olson et al., 1981; Pinto et al., 459

2014; Ranchet et al., 2016).

461 **6. Conclusion**

462 The main findings of this study are that VRU detection performance decreased with 463 aging, even in the group of younger-old adults (after age 55). This may be partly due to age-related deficits in selective attention and visual exploration. With the increasing 464 465 number of personal mobility devices (PMD) (e.g., electric scooters, unicycles, segways, hoverboards), particularly in cities, the number of accidents involving both 466 467 PMD riders and other road users is increasing (Ang et al., 2021). The detectability of 468 VRUs on PMDs by drivers or other road users, particularly older adults, has become a major challenge for road safety. Further studies should investigate the detection of 469 VRUs - including those on PMDs - by other road users (e.g., drivers, pedestrians, 470 471 cyclists).

472 Declarations of interest: none

Acknowledgments:

photographs.

The authors are grateful to Vivian Waltz and James Thomson for the English

- editing. We would also like to thank Jean-Philippe Cavallo for taking and editing the
- 175

- 476
- 477

478 References

Aartsen, M. J., Smits, C. H. M., van Tilburg, T., Knipscheer, K. C. P. M., & Deeg, D.

- 480 J. H. (2002). Activity in older adults : Cause or consequence of cognitive
- 481 functioning? A longitudinal study on everyday activities and cognitive performance

in older adults. *The Journals of Gerontology. Series B, Psychological Sciences and*

- 483 Social Sciences, 57(2), P153-162. https://doi.org/10.1093/geronb/57.2.p153
- Abel, L. A., Troost, B. T., & Dell'Osso, L. F. (1983). The effects of age on normal
 saccadic characteristics and their variability. *Vision Research*, *23*(1), 33-37.
 https://doi.org/10.1016/0042-6989(83)90038-X
- 487 ACEM. (2009). *Maids—In-Depth investigation of motorcycle accidents*.
 488 http://www.maids-study.eu/
- 489 Agresti, A. (2013). *Categorical Data Analysis* (3rd éd.). Wiley.
- 490 Ang, K. X. M., Chandrakumara, S. D., Kon Kam King, C., & Loh, S. Y. J. (2021).
- The Orthopedic Injury Burden of Personal Mobility Devices in Singapore Our
- 492 Experience in the East Coast. Journal of Clinical Orthopaedics and Trauma, 13,
- 493 66-69. https://doi.org/10.1016/j.jcot.2020.08.015
- Ball, K., Beard, B. L., Roenker, D. L., Miller, R. L., & Griggs, D. S. (1988). Age and
- 495 visual search : Expanding the useful field of view. Journal of the Optical Society of
- 496 America. A, Optics and Image Science, 5(12), 2210-2219.
- 497 https://doi.org/10.1364/josaa.5.002210
- Ball, K., & Owsley, C. (1993). The useful field of view test : A new technique for
- evaluating age-related declines in visual function. *Journal of the American Optometric Association*, *64*(1), 71-79.
- Borowsky, A., Oron-Gilad, T., Meir, A., & Parmet, Y. (2012). Drivers' perception of
- 502 vulnerable road users : A hazard perception approach. Accident Analysis &
- 503 *Prevention*, 44(1), 160-166. https://doi.org/10.1016/j.aap.2010.11.029
- 504 Brendicke, R., Forke, E., & Schaefer, D. (1994). Auswirkungen einer allgemeinen
- tageslichtpflicht auf die sicherheit motorisierter zweiraeder. *VDI-Berichte*, *1159*.
 https://trid.trb.org/view/1000302
- Bromberg, S., Oron-Gilad, T., Ronen, A., Borowsky, A., & Parmet, Y. (2012). The
- 508 perception of pedestrians from the perspective of elderly experienced and
- 509 experienced drivers. Accident Analysis & Prevention, 44(1), 48-55.
- 510 https://doi.org/10.1016/j.aap.2010.12.028
- Brouwer, R. F. T., Jansen, W. H., Theeuwes, J., Duistermaat, M., & Alferdinck, J.
- 512 W. A. M. (2004). Do other road users suffer from the presence of cars that have
- their daytime running lights on ? *TNO*.

- 514 Cavallo, V., & Pinto, M. (2012). Are car daytime running lights detrimental to
- 515 motorcycle conspicuity? *Accident; Analysis and Prevention, 49,* 78-85.
- 516 https://doi.org/10.1016/j.aap.2011.09.013
- 517 Coeckelbergh, T. R. M., Cornelissen, F. W., Brouwer, W. H., & Kooijman, A. C.
- 518 (2004). Age-Related Changes in the Functional Visual Field : Further Evidence for
- an Inverse Age × Eccentricity Effect. The Journals of Gerontology: Series B, 59(1),
- 520 P11-P18. https://doi.org/10.1093/geronb/59.1.P11
- 521 Coleman, A. L., Cummings, S. R., Yu, F., Kodjebacheva, G., Ensrud, K. E.,
- 522 Gutierrez, P., Stone, K. L., Cauley, J. A., Pedula, K. L., Hochberg, M. C., Mangione,
- 523 C. M., & Study Group of Osteoporotic Fractures. (2007). Binocular visual-field loss
- increases the risk of future falls in older white women. *Journal of the American*
- 525 Geriatrics Society, 55(3), 357-364. https://doi.org/10.1111/j.1532-
- 526 5415.2007.01094.x
- 527 Collins, M. J., Brown, B., & Bowman, K. J. (1989). Peripheral visual acuity and age.
- 528 Ophthalmic & Physiological Optics: The Journal of the British College of Ophthalmic
- 529 Opticians (Optometrists), 9(3), 314-316. https://doi.org/10.1111/j.1475-
- 530 1313.1989.tb00914.x
- 531 Crundall, D., Howard, A., & Young, A. (2017). Perceptual training to increase
- drivers' ability to spot motorcycles at T-junctions. *Transportation Research Part F:*
- 533 *Traffic Psychology and Behaviour, 48*, 1-12. https://doi.org/10.1016/j.trf.2017.05.003
- 534 Derefeldt, G., Lennerstrand, G., & Lundh, B. (1979). Age variations in normal
- human contrast sensitivity. *Acta Ophthalmologica*, *57*(4), 679-690.
- 536 https://doi.org/10.1111/j.1755-3768.1979.tb00517.x
- 537 Elliott, D. B. (1987). Contrast sensitivity decline with ageing : A neural or optical
- 538 phenomenon? *Ophthalmic and Physiological Optics*, 7(4), 415-419.
- 539 https://doi.org/10.1111/j.1475-1313.1987.tb00771.x
- 540 ERSO. (2018). Annual Accident Report 2018 (European Road Safety Observatory).
- 541 Freeman, E. E., Muñoz, B., Rubin, G., & West, S. K. (2007). Visual field loss
- 542 increases the risk of falls in older adults : The Salisbury eye evaluation. Investigative
- 543 *Ophthalmology & Visual Science*, *48*(10), 4445-4450.
- 544 https://doi.org/10.1167/iovs.07-0326
- Hancock, P. A., & Caird, J. K. (1993). Experimental evaluation of a model of mental
- 546 workload. *Human Factors*, *35*(3), 413-429.
- 547 https://doi.org/10.1177/001872089303500303
- 548 Harwood, R. H. (2001). Visual problems and falls. Age and Ageing, 30 Suppl 4,
- 549 13-18. https://doi.org/10.1093/ageing/30.suppl_4.13
- Henderson, S., Gagnon, S., Collin, C., Tabone, R., & Stinchcombe, A. (2013). Near
- 551 peripheral motion contrast threshold predicts older drivers' simulator performance.

- 552 Accident; Analysis and Prevention, 50, 103-109.
- 553 https://doi.org/10.1016/j.aap.2012.03.035
- Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure.
- 555 Scandinavian Journal of Statistics, 6(2), 65-70.
- Hurt, H. H. J., Ouellet, J. V., & Thom, D. R. (1981). *Motorcycle accident cause*
- 557 factors and identification of countermeasures (NHTSA report N° 81-1-VOL-1 Final
- 558 Rpt.). Department of Transportation. https://trid.trb.org/view/170348
- Ivers, R. Q., Cumming, R. G., Mitchell, P., & Attebo, K. (1998). Visual impairment
- and falls in older adults : The Blue Mountains Eye Study. *Journal of the American Geriatrics Society*, *46*(1), 58-64. https://doi.org/10.1111/j.1532-5415.1998.tb01014.x
- Jaffe, G. J., Alvarado, J. A., & Juster, R. P. (1986). Age-Related Changes of the
- Normal Visual Field. Archives of Ophthalmology, 104(7), 1021-1025.
- 564 https://doi.org/10.1001/archopht.1986.01050190079043
- Large, D. R., Kim, H., Merenda, C., Leong, S., Harvey, C., Burnett, G., & Gabbard,
- J. (2019). Investigating the effect of urgency and modality of pedestrian alert
- 567 warnings on driver acceptance and performance. *Transportation Research Part F:*
- 568 Traffic Psychology and Behaviour, 60, 11-24.
- 569 https://doi.org/10.1016/j.trf.2018.09.028
- Li, X., Farrukh, M., Lee, C., Khreis, H., Sarda, S., Sohrabi, S., Zhang, Z., &
- 571 Dadashova, B. (2022). COVID-19 impacts on mobility, environment, and health of
- active transportation users. *Cities*, 103886.
- 573 https://doi.org/10.1016/j.cities.2022.103886
- 574 McMurdo, M. E., & Gaskell, A. (1991). Dark adaptation and falls in the elderly.
- 575 Gerontology, 37(4), 221-224. https://doi.org/10.1159/000213264
- 576 Moschner, C., & Baloh, R. W. (1994). Age-Related Changes in Visual Tracking.
- 577 Journal of Gerontology, 49(5), M235-M238.
- 578 https://doi.org/10.1093/geronj/49.5.M235
- 579 Munoz, D. P., Broughton, J. R., Goldring, J. E., & Armstrong, I. T. (1998). Age-
- related performance of human subjects on saccadic eye movement tasks.
- 581 Experimental Brain Research.
- 582 Myers, R. S., Ball, K. K., Kalina, T. D., Roth, D. L., & Goode, K. T. (2000). Relation
- of Useful Field of View and other Screening Tests to on-Road Driving Performance.
- 584 *Perceptual and Motor Skills*, *91*(1), 279-290.
- 585 https://doi.org/10.2466/pms.2000.91.1.279
- 586 ONISR. (2019). Bilan 2019 de la sécurité routière | Observatoire national
- 587 *interministériel de la sécurité routière*. https://www.onisr.securite-
- routiere.gouv.fr/etat-de-l-insecurite-routiere/bilans-annuels-de-la-securite-
- routiere/bilan-2019-de-la-securite-routiere

- 590 Owsley, C. (1994). Vision and driving in the elderly. Optometry and Vision Science:
- ⁵⁹¹ Official Publication of the American Academy of Optometry, 71(12), 727-735.
- 592 https://doi.org/10.1097/00006324-199412000-00002
- 593 Owsley, C. (2011). Aging and vision. *Vision Research*, *51*(13), 1610-1622. 594 https://doi.org/10.1016/j.visres.2010.10.020
- 595 Owsley, C., Sekuler, R., & Siemsen, D. (1983). Contrast sensitivity throughout
- adulthood. Vision Research, 23(7), 689-699. https://doi.org/10.1016/00426989(83)90210-9
- Oxley, J., Fildes, B., Corben, B., & Langford, J. (2006). Intersection design for older
 drivers. *Transportation Research Part F: Traffic Psychology and Behaviour*, *9*(5),
 335-346. https://doi.org/10.1016/j.trf.2006.06.005
- Pai, C.-W. (2011). Motorcycle right-of-way accidents—A literature review. Accident
 Analysis & Prevention, 43(3), 971-982. https://doi.org/10.1016/j.aap.2010.11.024
- Pai, C.-W., Hwang, K. P., & Saleh, W. (2009). A mixed logit analysis of motorists'
 right-of-way violation in motorcycle accidents at priority T-junctions. *Accident; Analysis and Prevention*, *41*(3), 565-573. https://doi.org/10.1016/j.aap.2009.02.007
- Ranchet, M., Cavallo, V., Dang, N.-T., & Vienne, F. (2016). Improving motorcycle
 conspicuity through innovative headlight configurations. *Accident; Analysis and Prevention*, *94*, 119-126. https://doi.org/10.1016/j.aap.2016.05.011
- 609 Roenker, D. L., Cissell, G. M., Ball, K. K., Wadley, V. G., & Edwards, J. D. (2003).
- 610 Speed-of-processing and driving simulator training result in improved driving
- 611 performance. *Human Factors*, *45*(2), 218-233.
- 612 https://doi.org/10.1518/hfes.45.2.218.27241
- Rogé, J., El Zufari, V., Vienne, F., & Ndiaye, D. (2015). Safety messages and
- visibility of vulnerable road users for drivers. *Safety Science*, *79*, 29-38.
- 615 https://doi.org/10.1016/j.ssci.2015.05.002
- Rogé, J., Ndiaye, D., & Vienne, F. (2014). Useful visual field training : A way to
- 617 improve elderly car drivers' ability to detect vulnerable road users. *Transportation*
- 618 Research Part F: Traffic Psychology and Behaviour, 26, 246-257.
- 619 https://doi.org/10.1016/j.trf.2014.08.005
- Rogé, J., & Pébayle, T. (2009). Deterioration of the useful visual field with ageing
- during simulated driving in traffic and its possible consequences for road safety.
- 622 Safety Science, 47(9), 1271-1276. https://doi.org/10.1016/j.ssci.2009.03.012
- Rogé, J., Pébayle, T., El Hannachi, S., & Muzet, A. (2003). Effect of sleep
- 624 deprivation and driving duration on the useful visual field in younger and older
- subjects during simulator driving. *Vision Research*, *43*(13), 1465-1472.
- 626 https://doi.org/10.1016/s0042-6989(03)00143-3
- Rogé, J., Pébayle, T., Lambilliotte, E., Spitzenstetter, F., Giselbrecht, D., & Muzet,
- A. (2004). Influence of age, speed and duration of monotonous driving task in traffic

- on the driver's useful visual field. *Vision Research*, *44*(23), 2737-2744.
- 630 https://doi.org/10.1016/j.visres.2004.05.026
- Rumar, K. (1990). The basic driver error : Late detection. *Ergonomics*, *33*(10-11),
 1281-1290. https://doi.org/10.1080/00140139008925332
- 633 Salthouse, T. A. (1996). The processing-speed theory of adult age differences in
- 634 cognition. *Psychological Review*, *103*(3), 403-428. https://doi.org/10.1037/0033-635 295X.103.3.403
- 636 Scialfa, C. T., Kline, D. W., & Lyman, B. J. (1987). Age differences in target
- 637 identification as a function of retinal location and noise level : Examination of the
- useful field of view. *Psychology and Aging*, 2(1), 14-19.
- 639 https://doi.org/10.1037//0882-7974.2.1.14
- 640 Sekuler, A. B., Bennett, P. J., & Mamelak, M. (2000). Effects of aging on the useful
- 641 field of view. *Experimental Aging Research*, *26*(2), 103-120.
- 642 https://doi.org/10.1080/036107300243588
- 643 Sekuler, R., & Ball, K. (1986). Visual localization : Age and practice. Journal of the
- 644 Optical Society of America. A, Optics and Image Science, 3(6), 864-867.
- 645 https://doi.org/10.1364/josaa.3.000864
- Sims, R. V., McGwin, G., Allman, R. M., Ball, K., & Owsley, C. (2000). Exploratory
- 647 study of incident vehicle crashes among older drivers. *The Journals of Gerontology.*
- 648 Series A, Biological Sciences and Medical Sciences, 55(1), M22-27.
- 649 https://doi.org/10.1093/gerona/55.1.m22
- Snowden, R. J., & Kavanagh, E. (2006). Motion perception in the ageing visual
- 651 system : Minimum motion, motion coherence, and speed discrimination thresholds.
- 652 *Perception*, 35(1), 9-24. https://doi.org/10.1068/p5399
- 653 Staplin, L. (1995). Simulator and field measures of driver age differences in left-turn
- 654 gap judgments. *Transportation Research Record*, 1485.
- 655 https://trid.trb.org/view/451819
- Tabone, W., de Winter, J., Ackermann, C., Bärgman, J., Baumann, M., Deb, S.,
- 657 Emmenegger, C., Habibovic, A., Hagenzieker, M., Hancock, P. A., Happee, R.,
- Krems, J., Lee, J. D., Martens, M., Merat, N., Norman, D., Sheridan, T. B., &
- 659 Stanton, N. A. (2021). Vulnerable road users and the coming wave of automated
- 660 vehicles : Expert perspectives. *Transportation Research Interdisciplinary*
- 661 *Perspectives*, *9*, 100293. https://doi.org/10.1016/j.trip.2020.100293
- Van Elslande, P., Fournier, J.-Y., & Jaffard, M. (2012). In-depth analysis of road
- 663 crashes involving powered two-wheelers vehicles : Typical human functional failures
- and conditions of their production. Work (Reading, Mass.), 41 Suppl 1, 5871-5873.
- 665 https://doi.org/10.3233/WOR-2012-0978-5871
- 666 WHO. (2018). *Global status report on road safety 2018 : Summary* (World Health 667 Organization).