

The detection of vulnerable road users by younger and older drivers

Maud Ranchet, Roland Brémond, Prashant Pala, Michèle Colomb, Viola Cavallo

▶ To cite this version:

Maud Ranchet, Roland Brémond, Prashant Pala, Michèle Colomb, Viola Cavallo. The detection of vulnerable road users by younger and older drivers. Transportation Research Part F: Traffic Psychology and Behaviour, 2022, 91, pp.357-367. 10.1016/j.trf.2022.10.018. hal-04444293

HAL Id: hal-04444293 https://hal.science/hal-04444293v1

Submitted on 12 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Detection of Vulnerable Road Users by Younger and Older Drivers

Maud Ranchet^{1*}, Roland Brémond², Prashant Pala², Michèle Colomb³, and Viola Cavallo²

¹ TS2-LESCOT, Univ Gustave Eiffel, IFSTTAR, Univ Lyon, F-69675 Lyon, France

²COSYS-PICS-L, Univ Gustave Eiffel, IFSTTAR, F-77454 Marne-la-Vallée Cedex 2, France

³ CEREMA, 8-10, rue Bernard Palissy – 63017 Clermont-Ferrand Cedex 2, France

* Corresponding author:

Maud Ranchet, PhD Université Gustave Eiffel – TS2-LESCOT Cité des Mobilités 25, Avenue François Mitterrand Case 24 F-69675 Bron Cedex, France phone: +33 (0) 4 72 14 25 99

phone: +33 (0) 4 72 14 25 99 email: maud.ranchet@univ-eiffel.fr

Highlights

- The aim was to investigate the effect of aging on the detection, under time constraints, of vulnerable road users
- The detection of vulnerable road users decreased with aging, even in the younger-old group
- Older-old adults compensated for their visual-field decline by doing more visual exploration than the other two groups did to correctly detect vulnerable road users
- Further studies should investigate the detection of vulnerable road users, including those on personal mobility devices and using different exploration times.

Abstract

The detection of vulnerable road users (VRUs), especially under time constraints, may be impaired in elderly drivers, due to their visual and cognitive decline. This represents a major concern for road safety. The objective of the present study was to investigate the effect of aging on the detection of VRUs. A further aim was to investigate the impact of external factors on VRU detection. Twenty-two young adults, 20 younger-old adults, and 32 older-old adults were included in the study. A series of photographs were displayed for 500 ms. Participants were asked to detect the VRU, which could be a motorcyclist, a cyclist, or a pedestrian. The VRU was located at one of two distances (near/far), one of two locations (off-centered/centered), and in two car-DRL (daytime running lights) environments (on/off). The ability to correctly detect a VRU was measured. An eye-tracker was used to record eye movements. The main findings showed that VRU-detection performance decreased with aging, even in the younger-old group (ages 55 to 68). The ability to correctly detect a VRU by older-old adults was poorer particularly when the VRU was a motorcyclist or a cyclist. As a whole, the older-old adults made more fixations to correctly detect the VRU than the other two age groups did. Moreover, the visual angle between the gaze of the participant's last fixation and the target in the older-old group was lower than in the other two groups, particularly when the VRU was off-centered. This finding suggests that older-old adults compensated for their visual-field decline by doing more visual exploration than the other two groups did to correctly detect the VRU. The results are discussed with regards to age-related cognitive and visual deficits.

Word count: 279

Keywords: older drivers; vulnerable road users; detection performance; visual exploration

1. Introduction

The safety of vulnerable road users (VRUs), including pedestrians, cyclists, and motorcyclists, is a major concern in many countries (WHO, 2018) where they have a significantly higher risk of being killed or seriously injured than do car drivers. VRU fatalities currently represent 42.5% of all fatalities in France (ONISR, 2019), 46.5% in Europe (ERSO, 2018), and 56% in the world (WHO, 2018). With the COVID-19 pandemic situation, daily mobility of individuals has considerably changed (Li et al., 2022): the number of VRUs, especially in the big cities is constantly growing. A further increase in the number of VRUs can be expected due to a rise in petrol prices but also due to ecological and health considerations.

Main factors that could explain poor VRU detection

In order to safely interact with vulnerable road users, the drivers of motorized vehicles must detect them quickly. In fact, late detection is the basic driver error that leads to collisions (Rumar, 1990). Besides expectation errors (i.e., failure to scan for a specific class of road users), Rumar et al. (1990) mentioned perceptual difficulties, especially when low-contrast conditions and/or the peripheral visual field of the observer are involved. VRUs are often more difficult to detect than cars because of their smaller size and irregular contours (Cavallo & Pinto, 2012). The fact that they are not very conspicuous visually and fail to be perceived by other vehicle drivers are frequent causes of accidents (Van Elslande et al., 2012). A previous study showed that drivers detect pedestrians less often when they appear in urban areas and more often when they appear in residential areas (Borowsky et al., 2012). In urban areas, drivers are not necessarily attuned to pedestrians, whereas in residential areas, their mindset is such that they expect and search for pedestrians. Furthermore, findings have

demonstrated that driving experience is also a dominant factor that affects the identification of pedestrians. Regarding motorcycle accidents, for instance, it has been shown that poor motorcycle detectability is one of the main risk factors and leads to right-of-way violations (ACEM, 2009; Hurt et al., 1981). While motorcycle conspicuity has improved since the use of daytime running lights (DRLs, which became compulsory in the seventies in many countries), this conspicuity advantage has been jeopardized by the increasing use of DRLs on cars, which became compulsory in Europe in 2011. Previous experimental studies have shown that car DRLs, which act as a kind of visual noise, hamper motorcycle detection (Brendicke et al., 1994; Cavallo & Pinto, 2012), and also have a detrimental effect on the detection of bicyclists and pedestrians (Cavallo & Pinto, 2012). Although rarely studied, VRU detectability, especially in a car-DRL environment, represents a major concern for road safety.

The impact of age-related declines on VRU detection

Perceptual difficulties in detecting VRUs can also be caused by factors internal to drivers who are visually impaired and/or exhibit age-related declines. It has been shown that elderly drivers are over-represented in accidents involving pedestrians (Bromberg et al., 2012) and motorcyclists (Pai et al., 2009). While speed/distance judgment errors are often mentioned (e.g., Hancock & Caird, 1993; Oxley et al., 2006; Pai, 2011), detection failures in elderly drivers, together with longer perception and reaction times at intersections, have also been observed (e.g., Staplin, 1995). Using a video observation technique, the Bromberg et al. study (2012) showed that elderly drivers had longer response times on hazard detection, especially for pedestrian-related hazardous events.

- Late detection and failure to detect among elderly people are classically explained in terms of an age-related deterioration of visual capacities, as reflected by declines in visuo-attentional abilities, and also in terms of less effective visual exploration strategies.
- 53 Among the age-related sensory changes, we find impaired visual acuity in both 54 central and peripheral vision (Collins et al., 1989; Derefeldt et al., 1979; Harwood, 55 2001; Ivers et al., 1998; Owsley et al., 1983), low motion sensitivity (Sekuler et al., 56 2000; Snowden & Kavanagh, 2006), and poor adaptation to the dark (McMurdo & 57 Gaskell, 1991). The visual field, as measured in optometry, is also subject to an age effect (Coleman et al., 2007; Freeman et al., 2007; Jaffe et al., 1986), with a greater 58 59 decrease in peripheral than in central vision. Contrast sensitivity also deteriorates 60 with age (Elliott, 1987), particularly at high spatial frequencies, i.e., the ability to see 61 details.

62

63

64

65

66

67

68

- The useful field of view test (Ball et al., 1988) refers to the ability, in one glance, to detect a peripheral visual target and to identify a central visual target. Findings from a computer-screen task (UFOV^R), have consistently shown that the ability to detect peripheral targets in this dual task deteriorates with age, and that the difference between younger and older people becomes greater as the eccentricity of the peripheral target increases (Ball et al., 1988; Scialfa et al., 1987; Sekuler & Ball, 1986). This is partly due to the lowering of processing speed with ageing, as demonstrated by Salthouse (Salthouse, 1996).
- Similar age effects have also been observed when participants performed a task of detecting visual targets in a road scene while driving on a simulator (Rogé et al., 2003, 2004; Rogé & Pébayle, 2009). It has been shown that the accident risk was much

higher for elderly drivers when their useful field of view was reduced by 40% or more (Ball & Owsley, 1993; Owsley, 1994; Sims et al., 2000). The poor UFOV performance of participants referred for fitness-to-drive assessment was found to be associated with a large number of driving errors during real-world driving (Myers et al., 2000). Similar results have been found on a driving simulator (Henderson et al., 2013). With respect to the detection of VRUs, Rogé and Pébayle's (2009) simulator study showed that drivers with a limited useful field of view detected fewer motorcycles in the road scene, especially in the distant periphery, than those who had a more extensive useful field of view. Although that study only involved younger drivers, it can be assumed that such motorcycle-detection difficulties are more pronounced for older drivers, whose UFOV performance is generally impaired.

It is also well known in vision science that age impacts eye movements. While age does not seem to affect eye-movement accuracy (Moschner & Baloh, 1994), it has been shown to impact saccadic speed (Abel et al., 1983), saccadic latency, and saccadic reaction time (SRT) (Moschner & Baloh, 1994; Munoz et al., 1998). More specifically, an effect of age on SRT has been observed for persons over 40.

Objectives

The number of older drivers is increasing and their age-related declines may negatively impact the detection of VRUs, which may represent a major concern for road safety. It is surprising to notice that the detection of VRUs by older drivers has been very little investigated (Borowsky et al., 2012; Bromberg et al., 2012). The first objective of the present study was therefore to investigate the effect of aging on the detection of VRUs. Three age groups (young adults, younger-old adults, older-old adults) participated in the study. We expected an age-related decrease in the

97 detection of VRUs. We also hypothesized that a limited useful field of view, as 98 measured by the UFOV test, would be associated with poor detection of VRUs.

Another objective was to investigate the extent to which the detection of VRUs is affected by external factors such as the presence/absence of daytime running lights on cars, the kind of VRU to be detected (pedestrian, cyclist, motorcyclist), their distance from the observer, and their location in the visual scene. The exploratory aim was to better understand the visual strategies used by older drivers to correctly detect a VRU.

2. Method

2.1 Participants

A total of 74 participants were included in the study. In the group of young adults (N=22), there were 10 males and 12 females. Their median age was 29 years (range 25-46) and their median number of years of driving experience was 5 (range 1-25). In the group of younger-old adults (N=20), there were 16 males and 4 females. Their median age was 61.5 years (range 55-68) and their median number of years of driving experience was 42 (age 15-50). As cognitive decline often occurs at older ages (70 or more) (Aartsen et al., 2002), we added a third group of older-old adults (N = 32) whose median age was 73 years (range 70-87) and whose median number of years of driving experience was 51 (range 4-70). There were 19 males and 13 females.

The inclusion criteria were: (1) having normal or corrected-to-normal vision (at least 6/10), (2) being a driving-license holder, (3) being an active driver (driving more than 2500 km/year), and (4) not declaring any neurological disorders.

None of the participants were professional drivers. Participants were recruited by flyers or social networks. Written informed consent was obtained from all participants before the experiment.

2.2 Task and experimental design

A series of photographs were displayed for 500 ms (for an example, see Figure 1). The participants had to detect the presence of a VRU and determine whether it was a motorcyclist, bicyclist, or pedestrian. They had to choose between four response types displayed on the screen: bicyclists, motorcyclists, pedestrians, or none of the above. They indicated their response to the experimenter orally. Practice trials were performed before the experiment. A mixed experimental design was employed, with group (young, younger-old, older-old) as a between-subject variable. There were four within-subject variables: the targeted VRU (motorcyclist, bicyclist, or pedestrian), the distance of the VRU in the scene (far vs. near), the DRL environment of the cars (DRLs on vs. off), the VRU's location (centered vs. off-centered). Each experimental condition was repeated three times (VRU (3) × Distance (2) × Car-DRL (2) × Location (2)), making for a total of 72 experimental trials. Eighteen distractors containing a VRU (located at different distances from those used in the experimental trials) were added. In order to obtain a balanced design, 90 distractors with no target were added, making a total of 180 trials. The stimuli were presented in four blocks of 45 trials each.



Figure 1. Example of a trial involving a motorcyclist (a scooter) centered at a near distance, among cars with their daytime running lights off

2.3 Apparatus and stimuli

The stimuli were photographs of urban traffic at intersections in overcast weather. The photographs were presented on a 47" flat-panel LCD display (width: 1.02 m; height: 0.57 m). Participants were seated 1.60 m from the screen (angular size of the screen: 32.5° × 19.6°). The eye-tracking system was SMI (SensoMotoric Instruments), and the sampling rate was 60 Hz. The data was processed using BeGaze (SMI software). Participants performed Part 3 of the UFOV test, considered to be a measure of selective attention (Ball et al., 1993). In this part of the test, an object (truck or car) was presented in the center of the screen. At the same time, another object (truck or car) was presented in the periphery among 47 distractor triangles of the same size, contrast, and luminance. Following the identification of the central target object, participants had to indicate at which location the peripheral object was displayed.

2.4 Procedure

All participants first took a visual acuity test (Ergovision) and Part 3 of the UFOV test. Then the participants read the instructions of the experiment. The order of the four blocks was counterbalanced across participants. Each trial included the display of a fixation cross for 1500 ms, followed by the photographs for 500 ms. The screen turned gray at the end of the trial. Then the participants had to choose between four response types displayed on the screen. No feedback on response accuracy was given. Participants were allowed breaks during the blocks and between the blocks. The experimental session lasted approximately one hour and 30 minutes.

2.5 Dependent variables and statistical analyses

2.5.1 Dependent variables

For each experimental trial, we determined the number of correct detections, identification errors, and misses. A correct detection was counted when the right VRU category was recognized. An identification error was counted when a VRU was detected but not correctly identified, and a miss was counted when the VRU was not detected at all. Performance on distractor items was not analyzed. The number of fixations, known to be an indicator of visual exploration, was analyzed. Given that visual-field size decreases and peripheral vision declines with aging, the visual angle between the gaze of the last fixation and the target was also analyzed. To better understand the visual strategies used by older drivers to correctly detect a VRU, the number of fixations and the visual angle were analyzed specifically on correct detections. However, differences in visual strategies between correct detections and no detections (including misses and identification errors) were also explored for all

participants and for each age group. For the UFOV test, the presentation time in milliseconds (ms), necessary for correct identification, was measured.

2.5.2 Statistical analyses

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

A binary logistic regression was performed on the ability to correctly detect a VRU. This was done in the framework of the generalized linear mixed model (GLMM) in order to take into account the individual random effect, using the SPSS v.26s Genlinmixed procedure (see Agresti, 2013, for more details). The binary response on each trial (yes = 1, no = 0, 0 being the reference category), indicating the ability to correctly detect a VRU, was chosen as the dependent variable. A binomial distribution and a logit link function were assumed. The number of fixations during a trial, as well as the visual angle between the gaze (of the last fixation) and the target, were analyzed for correct detections. To analyze the number of fixations on correct detections, a Poisson regression in the framework of the GLMM model was conducted. To analyze the visual angle between the gaze (of the last fixation) and the target on correct detections, a linear regression was performed in the framework of the GLMM model. All of these models (binary logistic regression, Poisson regression, and linear regression) included five predictors that were categorical variables. The four withinsubject variables were VRU type (pedestrian, motorcyclist, cyclist), VRU distance (far, near), car-DRL environment (on, off), and VRU location (centered, off-centered). The single between-subject variable was age group. For the purpose of this study, only the interactions between age group and VRU type, age group and distance, age group and location, and age group and car-DRL environment were examined. Post-

hoc comparisons for the GLMMs were performed using t-tests. Significance levels

were adjusted for multiple comparisons using the sequential Bonferroni correction (or Holm's sequential Bonferroni procedure; Holm, 1979). Successive models were tested by removing non-significant terms one by one until the best fit was found using the corrected Akaike Information Criterion (cAIC), which decreases with increasing model fit. Given the small number of identification errors and misses, we did not examine the effect of within-subject variables on errors and misses. However, paired t-tests for all participants and for each group were performed between correct detections and no detections for the number of fixations and the degree of visual angle to determine whether there are any significant differences in visual strategies between correct detections and no detections.

For identification errors, misses, and UFOV performance, the data were not normally distributed. Values are expressed as medians (Q1-Q3). Between-group differences were examined using Kruskal-Wallis tests (KW). Post-hoc analyses used Wilcoxon rank-sum tests, with Holm's correction for multiple comparisons (Holm, 1979).

- Pearson correlations between UFOV performance and VRU detection performance were conducted for all participants.
- All statistical analyses were performed using SPSS, version 26.0.

3. Results

- 221 3.1 Detection task
- 3.1.1 Ability to correctly detect a VRU
- Table 1 presents the findings for the GLMM. A significant main group effect was found on the ability to correctly detect a VRU. Post hoc tests showed that the older-old participants detected the VRUs significantly less well than both the younger-old and

young groups did (p < 0.001), and the younger-old adults detected the VRUs significantly less well than the young adults did (p < 0.001). The mean (\pm standard deviation) proportion of correct detections was 0.55 \pm 0.01 for the older-old adults, 0.67 \pm 0.01 for the younger-old adults, and 0.82 \pm 0.01 for the young adults.


For all within-group variables, significant main effects on the ability to correctly detect a VRU were also found (Table 1). Post-hoc comparisons showed that the mean proportion of correct detections was significantly lower in the far condition (0.57 \pm 0.01) than in near conditions (0.79 \pm 0.001). The proportion of correct detections was also significantly lower in off-centered than in centered conditions. For location, the values were 0.78 \pm 0.001 for centered VRUs and 0.59 \pm 0.01 for off-centered VRUs. The proportion of correct detections was also lower for the DRLs-on than for the DRLs-off condition, but this difference was not significant. For the car-DRL environment, the values were 0.71 \pm 0.01 for DRLs off and 0.67 \pm 0.01 for DRLs on.

Table 1. Fixed effects using GLMM

Between-group variable		
Age group	F _{2,4842} = 115.17, p < 0.001	
Within-group variables		
Type of VRU	<i>F</i> _{2,4842} =43.02, p < 0.001	
VRU distance	<i>F</i> _{1,4842} = 219.28, p < 0.001	
VRU location	$F_{1, 4842}$ = 172.98, p < 0.001	
Car-DRL environment	$F_{1, 4842} = 5.00, p = 0.02$	

Figure 2 shows the estimated mean proportion of correct detections per age group for each VRU type. The older-old adults detected significantly fewer pedestrians, motorcyclists, and cyclists than the young adults did (p < 0.001); the older-old adults detected significantly fewer motorcyclists and cyclists than the younger-old adults did (p < 0.001); and the younger-old adults detected significantly fewer motorcyclists, cyclists, and pedestrians than the young adults did (p < 0.001). Pairwise comparisons also revealed that older-old adults detected significantly more pedestrians than cyclists and motorcyclists (p < 0.001). Younger-old adults detected significantly more pedestrians and motorcyclists than cyclists (p < 0.001). Young adults detected more motorcyclists than cyclists and pedestrians (p < 0.001). They also detected more pedestrians than cyclists (p < 0.001).

No significant interactions between age group and VRU distance, age group and VRU location, or age group and car-DRL environment were found.



- 255 *** p < 0.001
- 256 Figure 2. Estimated proportion of correct detections as a function of age group and VRU type. Only
- 257 significant between-group differences are shown in this figure. Error bars represent the standard
- 258 error.


- 260 3.1.2 Identification errors and misses
- Older-old adults made more identification errors than young adults (p < 0.01). The
- median percentage of identification errors for older-old adults was 5.59 (4.17 10.14)
- and for young adults, it was 2.78 (1.41 4.17). Both older-old adults and younger-old
- 264 adults missed more VRUs than young adults did (p < 0.01). The median percentage
- of misses for older-old adults was 37.50 (29.13 42.85). For younger-old adults, this
- value was 30.07 (22.22 36.11), and for young adults, it was 19.44 (11.94 23.61).
- 3.2 Eye-related variables for correct detections
- 268 For the number of fixations, the final model included the significant main effects of
- distance ($F_{1,3157} = 18.47$, p < 0.001), location ($F_{1,3157} = 8.28$, p = 0.004), and age group
- $(F_{2.3157} = 204.93, p < 0.001)$. For the visual angle, the final model included the
- significant main effects of distance ($F_{1,3153} = 8.78$, p = 0.003), location ($F_{2,3153} =$
- 272 2507.81, p < 0.001), age group ($F_{2,3153}$ = 16.09, p < 0.001), and a significant distance
- by age group interaction ($F_{2,3153} = 3.86$, p = 0.02) as well as a significant location by
- age group interaction ($F_{2,3153} = 59.64$, p < 0.001).
- There were a greater number of fixations in far VRU locations than in near ones. The
- mean number of fixations in a far location (M \pm SD) was 1.77 \pm 0.02; in a near location,
- it was 1.68 ± 0.02 , p < 0.001.

The findings also revealed a significant main effect of location, with a significantly higher average number of fixations in the off-centered condition than in centered conditions. The mean number of fixations (M \pm SD) in off-centered condition was 1.76 \pm 0.02. In the centered condition, it was 1.69 \pm 0.02, p = 0.004.

There was a significant main effect of age group on the number of fixations for correct detections, and on the visual angle between the gaze and the target. Post-hoc tests showed that the older-old group had a significantly higher number of fixations than either younger groups. The mean number of fixations for the older-old adults was 2.05 ± 0.02 . For the younger-old adults, it was 1.51 ± 0.02 , and for the young adults, it was 1.61 ± 0.02 (p < 0.001 on all pairwise comparisons). The older-old group had smaller visual angle than the younger-old group (p < 0.001) and the young group (p = 0.001). The visual angle in degrees for the older-old adults was 5.05 ± 0.09 . For the younger-old adults, it was 5.85 ± 0.11 , and for the young adults, it was 5.49 ± 0.09 (p ≤ 0.01 on all pairwise comparisons).

Interaction between age group and distance

The older-old adults had a significantly smaller visual angle than younger-old group did in both the near and far locations (p < 0.001, p = 0.009, respectively). In the far location, the older-old adults had smaller visual angle than did the young group (p = 0.003) (Figure 3). Only the younger group had a visual angle greater in the far location than in the near location (p < 0.001).




299 **p<0.01

***p<0.001

Figure 3. Visual angle by age group, for each distance. This represents the estimated values from the final model. Only significant between-group differences are shown in this figure. Error bars represent the standard error.

Interaction between age group and location

The older-old group had a significantly smaller visual angle than did the younger-old group and the young group in the off-centered condition (p < 0.001 on all pairwise comparisons) (Figure 4). The younger-old group also had a greater visual angle than the young group did (p < 0.001) in the off-centered condition. In the centered condition, the older-old group had a greater visual angle than the younger groups did (p = 0.001 on all pairwise comparisons). For all age groups, the visual angle was greater when the VRU was off-centered than when it was centered (p < 0.001).



314 **p<0.01

315 ***p<0.001

Figure 4. Visual angle by age group, for each location. This represents the estimated values from the final model. Only significant between-group differences are shown in this figure. Error bars represent the standard error.

3.3 Differences in visual strategies between correct detections and no detections

For the number of fixations, there was no significant difference between correct detections and no detections (correct detections = 1.76 ± 0.30 versus no detections = 1.77 ± 0.29 , t = 0.99, p = 0.33). Furthermore, no significant differences between correct detections and no detections were found for each age group (P > 0.05). However, a significant difference was found between correct detections and no detections for the visual angle (Visual angle for correct detections = $4.96^{\circ} \pm 0.98$ versus Visual angle for no detections = $8.24^{\circ} \pm 1.48$, t = 0.99, p < 0.001). These differences remain significant for each age group (Younger adults: Visual angle for correct detections = $5.21^{\circ} \pm 0.67$ versus Visual angle for no detections = $7.31^{\circ} \pm 1.32$, t = 6.66, p < 0.001; younger-old adults: Visual angle for correct detections = $5.25^{\circ} \pm 0.59$ versus Visual angle for no detections = $7.93^{\circ} \pm 1.17$, t = 8.04, p < 0.001; Older-

old adults: Visual angle for correct detections = $4.64^{\circ} \pm 1.23$ versus Visual angle for no detections = 9.05 ± 1.32 , t = 14.29, p < 0.001).

333

334

342

343

344

345

346

347

348

349

350

351

352

353

3.3 UFOV test

- The younger-old and older-old groups performed significantly less well on Part 3 of the UFOV test than the young group did (p < 0.001). The young adults had a median (Q1 - Q3) of 103 ms (80 - 140). The younger-old adults' median (Q1 - Q3) was 300 ms (257 - 407) and the older-old adults' median (Q1 - Q3) was 335 ms (253 - 500).
- 3.4 Correlation between Part 3 of the UFOV test and detection performance
- For all participants, poor performance on Part 3 of the UFOV test was strongly correlated with poor VRU detection (r = -0.67, p<0.001).

4. Discussion

- The aim of this study was to investigate the effects of aging on the detection of VRUs. The VRU could be a pedestrian, a motorcyclist, or a cyclist, positioned in two different locations and at two different distances in urban traffic. The cars could have their DRLs turned on or off. The contribution of this study is that VRU detection performance decreased with age, even in the younger-old group (ages 55 to 68). Furthermore, older-old adults performed poorly, particularly when the VRU was a motorcyclist or a cyclist. Finally, this study suggests that older participants used compensatory mechanisms for their poorer visual field by increasing visual exploration, without reaching the detection performance of the younger group.
 - As a whole, the older-old adults made more fixations to correctly detect the VRU than the other two age groups did. As expected, there were a greater number of fixations

to far VRU locations than to near ones. The mean number of fixations to detect the VRU in a far location was 1.77 whereas in a near location, it was 1.68, meaning that more individuals needed 2 fixations to detect far VRUs as compared to near VRUs. Moreover, the visual angle between the detected VRU and the last fixation was smaller in the older-old group than in the younger-old group, whether the VRU was in the far or near location. The visual angle of the older-old group was also smaller than that of the other two groups when the VRU was off-centered.

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

Importantly, detection performance decreased with aging, not only in the older-old participants (ages 70 to 87) but even in the younger-old group (ages 55 to 68). These findings are consistent with previous driving studies showing that age has an impact on the ability to detect visual signals (Rogé et al., 2003, 2004). Poor detection performance may be explained by a deterioration of selective attention with aging. In the present study, lower detection performance was strongly associated with a poorer measure of selective attention (Part 3 of the UFOV). Previous findings have shown that a decrease in selective attention appears early in the course of aging, between the ages of 40 and 50 (Sekuler et al., 2000). Another explanation could be the slowing of processing speed with aging (Owsley, 2011; Salthouse, 1996). In our study, participants were under time constraints when detecting a VRU since the photographs were displayed for only 500 ms. This could explain why even the younger-old adults could not compensate for their slowing in processing speed and exhibited a decrease in VRU detection performance. Older-old adults made more identification errors and both older groups (older-old and younger-old) missed more VRUs than young adults did, suggesting that they did not have time to see and identify the VRU.

Unexpectedly, the interaction between age group and centeredness on the ability to detect a VRU was not significant. Yet, for correct detections, the visual angle between the gaze and a VRU located in the periphery was narrower for older-old adults than for younger-old and young adults. This finding indicates that older-old adults' eyes need to land closer to the target to correctly detect a VRU in the periphery. It suggests that the older-old adults used mechanisms that compensate for visual-field decline by doing more visual exploration than the other two groups to correctly detect the VRU (Coeckelbergh et al., 2004). Our findings on the detection of VRUs located in the periphery differ from those of previous studies (Rogé et al., 2003, 2004; Rogé & Pébayle, 2009), which showed that older drivers detected fewer signals than younger drivers, particularly when the signal was farther away from the fixation point. Discrepancies across studies may be due to differences in methodology (screen size) or the type of task (observation versus driving). Interestingly, the number of fixations did not differ between correct detections and no detections whereas the degree of visual angle was larger for no detections, for all age groups. This result suggests that visual strategies of participants differ between correct detections and no detections: all participants (regardless of the age) need to land closer to the target to correctly detect a VRU.

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

Similarly, no age-related deterioration in the ability to detect a VRU among cars with their DRLs on was found in our study. Yet, the use of DRLs had a negative impact on detection performance in all three age groups. Previous studies have shown that the use of DRLs by cars may create "visual noise" that hinders the perception of other VRUs, such as pedestrians, cyclists, or motorcyclists (Cavallo & Pinto, 2012), which makes them less conspicuous and less noticeable (Brouwer et al., 2004). Our results

thus suggest that the use of DRLs by cars affects the detection of VRUs for all individuals, irrespective of their age.

Another interesting finding is that older-old adults and younger-old adults detected pedestrians better than they did motorcyclists and/or cyclists, which was not the case for our young adults. One explanation could be that older adults identified more with pedestrians than with motorcyclists and cyclists.

5. Limitations and future research avenues

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

Methodological considerations include the use of static images, i.e., photographs of urban traffic instead of dynamic images in videos or during simulated driving. The decision to use static images in the present study allowed us to investigate attentional aspects of road-user behavior. Photographs were displayed for 500 ms, which is quite short. In real life, drivers need to quickly detect relevant information in the road scene, and detection failures are more likely to occur under time pressure and when visual exploration time is short. Also, the age gap between our younger-old and older-old groups was quite close (65-68 vs. 70-87). However, our findings indicated some significant differences in VRU detection performance and visual exploration. Furthermore, the age gap between the older groups and the younger group was relatively large, which can explain significant differences between these two groups. Additional studies that include an intermediate age group with a mean age around 50 should be conducted to better understand the time course of visual and attentional aging in VRU detection. Time constraints play a major role in detection performance in older adults. In the present study, photographs were displayed for 500 ms, which may be not enough to correctly detect the VRU, especially for older adults. Future

research should vary presentation times of photographs in order to determine the role
of exploration times in relation with compensation strategies in older adults.

It would be interesting in future studies on older adults to investigate VRU detection using a driving simulator. The use of a driving simulator enables one to assess VRU detection during driving, and to better understand the compensatory strategies of older adults. A previous simulator study showed that elderly drivers drove about 20% slower than experienced non-elderly drivers (Bromberg et al., 2020), which suggests that as a whole, elderly drivers are aware of their deficiencies in detecting hazards.

A valuable contribution for improving road safety would be to investigate different types of interventions aimed at helping older drivers better detect VRUs, particularly motorcyclists and cyclists. Based on the results of this study, we can contend that older individuals detect fewer motorcyclists and cyclists than pedestrians. There are several ways to improve VRU detection performance. A variety of training methods have been employed to improve VRU detection by car drivers. One such method is useful-visual-field training which could improve elderly car drivers' VRU-detection (Rogé et al., 2014). It has been shown that drivers who had undergone training for perceiving motorcycles were better able to identify motorcyclists than ones who had not been trained (Crundall et al., 2017). Also, safety messages have been developed to improve the perception of VRU vulnerability during driving (Rogé et al., 2015).

Additional research needs to be conducted to explore the benefits of automated vehicles for detecting VRUs, particularly among older adults. The introduction of automated vehicles with external human-machine interfaces, and the use of augmented reality in automated vehicles, could help drivers to better detect VRUs and may also enhance future interaction between VRUs and autonomous vehicles

(Tabone et al., 2021). Several safety systems such as pedestrian alerts could inform drivers of a potential hazard (Large et al., 2019). However, the additional cognitive load induced by these systems should be considered with caution in future studies, especially for older adults.

As a complement to studies investigating VRU detection by drivers, eye-tracking studies should also explore the interactions between drivers and pedestrians from the pedestrian's point of view (Lévêque et al., 2020).

Further research should also help improve the visibility of VRUs. Previous studies have shown that the use of specific conspicuous clothes (for example, retroreflective and yellow clothes) by cyclists, or having specific headlights on motorcycles could improve their detection (e.g., Olson et al., 1981; Pinto et al., 2014; Ranchet et al., 2016).

6. Conclusion

The main findings of this study are that VRU detection performance decreased with aging, even in the group of younger-old adults (after age 55). This may be partly due to age-related deficits in selective attention and visual exploration. With the increasing number of personal mobility devices (PMD) (e.g., electric scooters, unicycles, segways, hoverboards), particularly in cities, the number of accidents involving both PMD riders and other road users is increasing (Ang et al., 2021). The detectability of VRUs on PMDs by drivers or other road users, particularly older adults, has become a major challenge for road safety. Further studies should investigate the detection of VRUs - including those on PMDs - by other road users (e.g., drivers, pedestrians, cyclists).

472	Declarations of interest: none
473	Acknowledgments:
474	The authors are grateful to Vivian Waltz and James Thomson for the English
475	editing. We would also like to thank Jean-Philippe Cavallo for taking and editing the
476	photographs.
477	

478 References

- Aartsen, M. J., Smits, C. H. M., van Tilburg, T., Knipscheer, K. C. P. M., & Deeg, D.
- J. H. (2002). Activity in older adults: Cause or consequence of cognitive
- functioning? A longitudinal study on everyday activities and cognitive performance
- in older adults. The Journals of Gerontology. Series B, Psychological Sciences and
- 483 Social Sciences, 57(2), P153-162. https://doi.org/10.1093/geronb/57.2.p153
- Abel, L. A., Troost, B. T., & Dell'Osso, L. F. (1983). The effects of age on normal
- saccadic characteristics and their variability. Vision Research, 23(1), 33-37.
- 486 https://doi.org/10.1016/0042-6989(83)90038-X
- 487 ACEM. (2009). Maids—In-Depth investigation of motorcycle accidents.
- 488 http://www.maids-study.eu/
- 489 Agresti, A. (2013). Categorical Data Analysis (3rd éd.). Wiley.
- 490 Ang, K. X. M., Chandrakumara, S. D., Kon Kam King, C., & Loh, S. Y. J. (2021).
- The Orthopedic Injury Burden of Personal Mobility Devices in Singapore Our
- Experience in the East Coast. Journal of Clinical Orthopaedics and Trauma, 13,
- 493 66-69. https://doi.org/10.1016/j.jcot.2020.08.015
- 494 Ball, K., Beard, B. L., Roenker, D. L., Miller, R. L., & Griggs, D. S. (1988). Age and
- visual search: Expanding the useful field of view. Journal of the Optical Society of
- 496 America. A, Optics and Image Science, 5(12), 2210-2219.
- 497 https://doi.org/10.1364/josaa.5.002210
- Ball, K., & Owsley, C. (1993). The useful field of view test: A new technique for
- evaluating age-related declines in visual function. *Journal of the American*
- 500 *Optometric Association*, *64*(1), 71-79.
- Borowsky, A., Oron-Gilad, T., Meir, A., & Parmet, Y. (2012). Drivers' perception of
- vulnerable road users: A hazard perception approach. Accident Analysis &
- 503 Prevention, 44(1), 160-166. https://doi.org/10.1016/j.aap.2010.11.029
- Brendicke, R., Forke, E., & Schaefer, D. (1994). Auswirkungen einer allgemeinen
- tageslichtpflicht auf die sicherheit motorisierter zweiraeder. VDI-Berichte, 1159.
- 506 https://trid.trb.org/view/1000302
- Bromberg, S., Oron-Gilad, T., Ronen, A., Borowsky, A., & Parmet, Y. (2012). The
- 508 perception of pedestrians from the perspective of elderly experienced and
- experienced drivers. Accident Analysis & Prevention, 44(1), 48-55.
- 510 https://doi.org/10.1016/j.aap.2010.12.028
- Brouwer, R. F. T., Jansen, W. H., Theeuwes, J., Duistermaat, M., & Alferdinck, J.
- W. A. M. (2004). Do other road users suffer from the presence of cars that have
- their daytime running lights on ? *TNO*.

- Cavallo, V., & Pinto, M. (2012). Are car daytime running lights detrimental to
- motorcycle conspicuity? Accident; Analysis and Prevention, 49, 78-85.
- 516 https://doi.org/10.1016/j.aap.2011.09.013
- Coeckelbergh, T. R. M., Cornelissen, F. W., Brouwer, W. H., & Kooijman, A. C.
- 518 (2004). Age-Related Changes in the Functional Visual Field: Further Evidence for
- an Inverse Age × Eccentricity Effect. The Journals of Gerontology: Series B, 59(1),
- 520 P11-P18. https://doi.org/10.1093/geronb/59.1.P11
- 521 Coleman, A. L., Cummings, S. R., Yu, F., Kodjebacheva, G., Ensrud, K. E.,
- Gutierrez, P., Stone, K. L., Cauley, J. A., Pedula, K. L., Hochberg, M. C., Mangione,
- 523 C. M., & Study Group of Osteoporotic Fractures. (2007). Binocular visual-field loss
- increases the risk of future falls in older white women. *Journal of the American*
- 525 Geriatrics Society, 55(3), 357-364. https://doi.org/10.1111/j.1532-
- 526 5415.2007.01094.x
- 527 Collins, M. J., Brown, B., & Bowman, K. J. (1989). Peripheral visual acuity and age.
- 528 Ophthalmic & Physiological Optics: The Journal of the British College of Ophthalmic
- 529 Opticians (Optometrists), 9(3), 314-316. https://doi.org/10.1111/j.1475-
- 530 1313.1989.tb00914.x
- 531 Crundall, D., Howard, A., & Young, A. (2017). Perceptual training to increase
- drivers' ability to spot motorcycles at T-junctions. Transportation Research Part F:
- 533 *Traffic Psychology and Behaviour, 48*, 1-12. https://doi.org/10.1016/j.trf.2017.05.003
- Derefeldt, G., Lennerstrand, G., & Lundh, B. (1979). Age variations in normal
- 535 human contrast sensitivity. *Acta Ophthalmologica*, *57*(4), 679-690.
- 536 https://doi.org/10.1111/j.1755-3768.1979.tb00517.x
- Elliott, D. B. (1987). Contrast sensitivity decline with ageing: A neural or optical
- 538 phenomenon? Ophthalmic and Physiological Optics, 7(4), 415-419.
- 539 https://doi.org/10.1111/j.1475-1313.1987.tb00771.x
- 540 ERSO. (2018). Annual Accident Report 2018 (European Road Safety Observatory).
- Freeman, E. E., Muñoz, B., Rubin, G., & West, S. K. (2007). Visual field loss
- increases the risk of falls in older adults: The Salisbury eye evaluation. *Investigative*
- 543 Ophthalmology & Visual Science, 48(10), 4445-4450.
- 544 https://doi.org/10.1167/iovs.07-0326
- Hancock, P. A., & Caird, J. K. (1993). Experimental evaluation of a model of mental
- 546 workload. *Human Factors*, 35(3), 413-429.
- 547 https://doi.org/10.1177/001872089303500303
- Harwood, R. H. (2001). Visual problems and falls. Age and Ageing, 30 Suppl 4,
- 13-18. https://doi.org/10.1093/ageing/30.suppl 4.13
- Henderson, S., Gagnon, S., Collin, C., Tabone, R., & Stinchcombe, A. (2013). Near
- peripheral motion contrast threshold predicts older drivers' simulator performance.

- 552 Accident; Analysis and Prevention, 50, 103-109.
- 553 https://doi.org/10.1016/j.aap.2012.03.035
- Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure.
- 555 Scandinavian Journal of Statistics, 6(2), 65-70.
- Hurt, H. H. J., Ouellet, J. V., & Thom, D. R. (1981). *Motorcycle accident cause*
- 557 factors and identification of countermeasures (NHTSA report N° 81-1-VOL-1 Final
- Rpt.). Department of Transportation. https://trid.trb.org/view/170348
- Ivers, R. Q., Cumming, R. G., Mitchell, P., & Attebo, K. (1998). Visual impairment
- and falls in older adults: The Blue Mountains Eye Study. *Journal of the American*
- 561 Geriatrics Society, 46(1), 58-64. https://doi.org/10.1111/j.1532-5415.1998.tb01014.x
- Jaffe, G. J., Alvarado, J. A., & Juster, R. P. (1986). Age-Related Changes of the
- Normal Visual Field. *Archives of Ophthalmology*, *104*(7), 1021-1025.
- https://doi.org/10.1001/archopht.1986.01050190079043
- Large, D. R., Kim, H., Merenda, C., Leong, S., Harvey, C., Burnett, G., & Gabbard,
- J. (2019). Investigating the effect of urgency and modality of pedestrian alert
- warnings on driver acceptance and performance. *Transportation Research Part F:*
- 568 Traffic Psychology and Behaviour, 60, 11-24.
- 569 https://doi.org/10.1016/j.trf.2018.09.028
- Li, X., Farrukh, M., Lee, C., Khreis, H., Sarda, S., Sohrabi, S., Zhang, Z., &
- Dadashova, B. (2022). COVID-19 impacts on mobility, environment, and health of
- active transportation users. *Cities*, 103886.
- 573 https://doi.org/10.1016/j.cities.2022.103886
- McMurdo, M. E., & Gaskell, A. (1991). Dark adaptation and falls in the elderly.
- 575 Gerontology, 37(4), 221-224. https://doi.org/10.1159/000213264
- 576 Moschner, C., & Baloh, R. W. (1994). Age-Related Changes in Visual Tracking.
- 577 *Journal of Gerontology*, *49*(5), M235-M238.
- 578 https://doi.org/10.1093/geronj/49.5.M235
- Munoz, D. P., Broughton, J. R., Goldring, J. E., & Armstrong, I. T. (1998). Age-
- related performance of human subjects on saccadic eye movement tasks.
- 581 Experimental Brain Research.
- 582 Myers, R. S., Ball, K. K., Kalina, T. D., Roth, D. L., & Goode, K. T. (2000). Relation
- of Useful Field of View and other Screening Tests to on-Road Driving Performance.
- 584 Perceptual and Motor Skills, 91(1), 279-290.
- 585 https://doi.org/10.2466/pms.2000.91.1.279
- 586 ONISR. (2019). Bilan 2019 de la sécurité routière | Observatoire national
- interministériel de la sécurité routière. https://www.onisr.securite-
- routiere.gouv.fr/etat-de-l-insecurite-routiere/bilans-annuels-de-la-securite-
- routiere/bilan-2019-de-la-securite-routiere

- 590 Owsley, C. (1994). Vision and driving in the elderly. *Optometry and Vision Science:*
- Official Publication of the American Academy of Optometry, 71(12), 727-735.
- 592 https://doi.org/10.1097/00006324-199412000-00002
- 593 Owsley, C. (2011). Aging and vision. *Vision Research*, *51*(13), 1610-1622.
- 594 https://doi.org/10.1016/j.visres.2010.10.020
- Owsley, C., Sekuler, R., & Siemsen, D. (1983). Contrast sensitivity throughout
- adulthood. Vision Research, 23(7), 689-699. https://doi.org/10.1016/0042-
- 597 6989(83)90210-9
- Oxley, J., Fildes, B., Corben, B., & Langford, J. (2006). Intersection design for older
- 599 drivers. Transportation Research Part F: Traffic Psychology and Behaviour, 9(5),
- 335-346. https://doi.org/10.1016/j.trf.2006.06.005
- Pai, C.-W. (2011). Motorcycle right-of-way accidents—A literature review. Accident
- 602 Analysis & Prevention, 43(3), 971-982. https://doi.org/10.1016/j.aap.2010.11.024
- Pai, C.-W., Hwang, K. P., & Saleh, W. (2009). A mixed logit analysis of motorists'
- right-of-way violation in motorcycle accidents at priority T-junctions. *Accident;*
- Analysis and Prevention, 41(3), 565-573. https://doi.org/10.1016/j.aap.2009.02.007
- Ranchet, M., Cavallo, V., Dang, N.-T., & Vienne, F. (2016). Improving motorcycle
- conspicuity through innovative headlight configurations. Accident; Analysis and
- 608 Prevention, 94, 119-126. https://doi.org/10.1016/j.aap.2016.05.011
- 609 Roenker, D. L., Cissell, G. M., Ball, K. K., Wadley, V. G., & Edwards, J. D. (2003).
- Speed-of-processing and driving simulator training result in improved driving
- 611 performance. *Human Factors*, *45*(2), 218-233.
- 612 https://doi.org/10.1518/hfes.45.2.218.27241
- Rogé, J., El Zufari, V., Vienne, F., & Ndiaye, D. (2015). Safety messages and
- visibility of vulnerable road users for drivers. Safety Science, 79, 29-38.
- 615 https://doi.org/10.1016/j.ssci.2015.05.002
- Rogé, J., Ndiaye, D., & Vienne, F. (2014). Useful visual field training: A way to
- 617 improve elderly car drivers' ability to detect vulnerable road users. *Transportation*
- Research Part F: Traffic Psychology and Behaviour, 26, 246-257.
- 619 https://doi.org/10.1016/j.trf.2014.08.005
- Rogé, J., & Pébayle, T. (2009). Deterioration of the useful visual field with ageing
- during simulated driving in traffic and its possible consequences for road safety.
- Safety Science, 47(9), 1271-1276. https://doi.org/10.1016/j.ssci.2009.03.012
- Rogé, J., Pébayle, T., El Hannachi, S., & Muzet, A. (2003). Effect of sleep
- deprivation and driving duration on the useful visual field in younger and older
- subjects during simulator driving. *Vision Research*, *43*(13), 1465-1472.
- 626 https://doi.org/10.1016/s0042-6989(03)00143-3
- Rogé, J., Pébayle, T., Lambilliotte, E., Spitzenstetter, F., Giselbrecht, D., & Muzet,
- A. (2004). Influence of age, speed and duration of monotonous driving task in traffic

- on the driver's useful visual field. Vision Research, 44(23), 2737-2744.
- 630 https://doi.org/10.1016/j.visres.2004.05.026
- Rumar, K. (1990). The basic driver error: Late detection. *Ergonomics*, 33(10-11),
- 632 1281-1290. https://doi.org/10.1080/00140139008925332
- 633 Salthouse, T. A. (1996). The processing-speed theory of adult age differences in
- 634 cognition. Psychological Review, 103(3), 403-428. https://doi.org/10.1037/0033-
- 635 295X.103.3.403
- 636 Scialfa, C. T., Kline, D. W., & Lyman, B. J. (1987). Age differences in target
- identification as a function of retinal location and noise level: Examination of the
- useful field of view. Psychology and Aging, 2(1), 14-19.
- 639 https://doi.org/10.1037//0882-7974.2.1.14
- Sekuler, A. B., Bennett, P. J., & Mamelak, M. (2000). Effects of aging on the useful
- 641 field of view. Experimental Aging Research, 26(2), 103-120.
- 642 https://doi.org/10.1080/036107300243588
- 643 Sekuler, R., & Ball, K. (1986). Visual localization: Age and practice. Journal of the
- Optical Society of America. A, Optics and Image Science, 3(6), 864-867.
- 645 https://doi.org/10.1364/josaa.3.000864
- Sims, R. V., McGwin, G., Allman, R. M., Ball, K., & Owsley, C. (2000). Exploratory
- study of incident vehicle crashes among older drivers. *The Journals of Gerontology.*
- Series A, Biological Sciences and Medical Sciences, 55(1), M22-27.
- 649 https://doi.org/10.1093/gerona/55.1.m22
- Snowden, R. J., & Kavanagh, E. (2006). Motion perception in the ageing visual
- system: Minimum motion, motion coherence, and speed discrimination thresholds.
- 652 *Perception*, 35(1), 9-24. https://doi.org/10.1068/p5399
- Staplin, L. (1995). Simulator and field measures of driver age differences in left-turn
- 654 gap judgments. Transportation Research Record, 1485.
- 655 https://trid.trb.org/view/451819
- Tabone, W., de Winter, J., Ackermann, C., Bärgman, J., Baumann, M., Deb, S.,
- Emmenegger, C., Habibovic, A., Hagenzieker, M., Hancock, P. A., Happee, R.,
- Krems, J., Lee, J. D., Martens, M., Merat, N., Norman, D., Sheridan, T. B., &
- 659 Stanton, N. A. (2021). Vulnerable road users and the coming wave of automated
- vehicles: Expert perspectives. *Transportation Research Interdisciplinary*
- 661 Perspectives, 9, 100293. https://doi.org/10.1016/j.trip.2020.100293
- Van Elslande, P., Fournier, J.-Y., & Jaffard, M. (2012). In-depth analysis of road
- crashes involving powered two-wheelers vehicles: Typical human functional failures
- and conditions of their production. Work (Reading, Mass.), 41 Suppl 1, 5871-5873.
- 665 https://doi.org/10.3233/WOR-2012-0978-5871
- 666 WHO. (2018). Global status report on road safety 2018: Summary (World Health
- 667 Organization).