



Indirect X-ray photodesorption from ices

Romain Basalgète, G Féraud, Daniela Torres-Diaz, A Lafosse, Lionel Amiaud, Pascal Jeseck, L Philippe, Xavier Michaut, Jean-Hugues Fillion, Mathieu Bertin

► To cite this version:

Romain Basalgète, G Féraud, Daniela Torres-Diaz, A Lafosse, Lionel Amiaud, et al.. Indirect X-ray photodesorption from ices. 15th International Conference on Electronic Spectroscopy and Structure, Aug 2023, Oulu, Finland. hal-04444570

HAL Id: hal-04444570

<https://hal.science/hal-04444570>

Submitted on 7 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Indirect X-ray photodesorption from ices

R. Basalgète¹, G. Féraud¹, D. Torres-Díaz^{1,2}, A. Lafosse², L. Amiaud², P. Jeseck¹, L. Philippe¹, X. Michaut¹, J-H. Fillion¹, and M. Bertin¹

¹ Sorbonne Université, Observatoire de Paris, Université PSL, CNRS, LERMA, F-75005, Paris, France

² Univ. Paris Saclay, CNRS, ISMO, 91405 Orsay, France

*Contact: geraldine.feraud@sorbonne-universite.fr

Keywords: X-ray processes, molecular ices, energy transfer, solid-to-gas interface

In the colder parts of the interstellar medium, dust grains are covered with molecular ices composed of H₂O, CO,... Among the many radiation-induced processes that occur in these icy mantles, the ones induced by X-rays may play an important role in star and planet formation regions [1]. In particular, X-rays interacting with interstellar ices can release molecules or radicals into the gas phase, by so-called photodesorption processes, and thus have an important role in the gas phase abundances measured from observations by the most recent telescopes (ALMA, NOEMA). This X-ray induced desorption process is however rarely considered because of a lack of experimental quantitative data and because the processes at play are poorly understood [2-5].

We irradiated ices containing ¹⁵N₂ and ¹³CO with soft X-rays using the SPICES 2 set-up connected to the SEXTANTS beamline at the SOLEIL synchrotron [6]. The release of ¹³CO or of ¹⁵N₂ is recorded by mass spectrometry as a function of the incident photon energy near the N (~ 400 eV) and O K-edge (~ 500 eV). Several ices were studied including pure CO ices, pure ¹⁵N₂ ices, and mixed or layered ¹³CO/¹⁵N₂ ices. Photodesorption from a layered ¹³CO/¹⁵N₂ ice reveals an indirect desorption mechanism for which the desorption of ¹³CO is triggered by the photo-absorption of ¹⁵N₂ at the N-edge. This enables to quantify the relevant depth involved in the indirect desorption process, which is found to be 30 - 40 Monolayers in that case. This value is further related to the energy transport of Auger electrons emitted from the photo-absorbing ¹⁵N₂ molecules that scatter towards the ice surface, inducing the desorption of ¹³CO. We show that X-ray induced electron stimulated desorption (XESD), mediated by Auger scattering and subsequent secondary electrons, is the dominant process explaining the desorption from the ices studied in this work [6].




Figure 1: Left: scheme of a simple model to estimate the energy deposited to the top ¹³CO layer of a layered ¹³CO/¹⁵N₂ ice by scattering of Auger electrons emitted after photo-absorption by ¹⁵N₂ molecules. Right: in green is displayed the estimation of this energy, compared with the X-ray photodesorption yields of ¹³CO (orange squares) [7].

- Walsh, C., Nomura H., Millar T. J., et Aikawa Y. ApJ 747, 2 (2012): 114.
- Ciaravella,A.; Caro,G. M.; Escobar,A. J.; Cecchi-Pestellini,C.; Giarrusso,S.; Barbera,M.; Collura,A. ApJ 2010, 722, L45–L48.
- Dupuy, R., M. Bertin, G. Féraud, M. Hassenfratz, X. Michaut, et al. Nat. Astron. 2, 10 (2018): 796-80
- Basalgète, R.; Dupuy, R.; Féraud, G.; Romanzin, C.; Philippe, L.; Michaut, X.; Michoud, J.; Amiaud, L.; Lafosse, A.; Fillion, J.-H.; Bertin, M. A&A 2021, 647, A35.
- Basalgète, R.; Dupuy, R.; Féraud, G.; Romanzin, C.; Philippe, L.; Michaut, X.; Michoud, J.; Amiaud, L.; Lafosse, A.; Fillion, J.-H.; Bertin, M. A&A 2021, 647, A36.
- Basalgète, R.; Torres-Díaz, D.; Lafosse, A.; Amiaud, L.; Féraud, G.; Jeseck, P.; Philippe, L.; Michaut, X.; Fillion, J.-H.; Bertin, M. J. Chem. Phys. 2022, 157 (8), 084308.