

14th International Conference on Atmospheric Sciences and Application to Air Quality 29-31 May 2017, Strasbourg, France

Urban compaction vs city sprawl : impact of road traffic on air quality in the greater Paris

Arthur Elessa Etuman⁽¹⁾, Isabelle Coll⁽¹⁾, Vincent Viguie⁽²⁾, Nicolas Coulombel⁽³⁾, and Julie Prud'homme

(1) LISA (Laboratoire Interuniversitaire des Systèmes Atmosphériques), (2) CIRED (Centre International de Recherche sur l'Environnement et le Développement), (3) LVMT (Laboratoire Ville Mobilité Transport)











Plan




















liso

Energy transition

- o Global priority
- Environmentals policies to reduce GHG and air pollutant emissions
- Limitations of abatement technologies (Internal combustion engine)

Energy use in the future ?

New reflexions on reducing energy use (Urban planning)




liso

Energy transition

- o Global priority
- Environmentals policies to reduce GHG and air pollutant emissions
- Limitations of abatement technologies (Internal combustion engine)

Energy use in the future ?

New reflexions on reducing energy use (Urban planning)



Source: INSEE, population projections 2005-2050


liso

Energy transition


- o Global priority
- Environmentals policies to reduce GHG and air pollutant emissions
- Limitations of abatement technologies (Internal combustion engine)

Energy use in the future ?

New reflexions on reducing energy use (Urban planning)




Source: INSEE, population projections 2005-2050




Source: INSEE, population projections 2005-2050


High density urban settlement
 Proximity and mixity of services
 Reducing commuting distances





High density urban settlement
 Proximity and mixity of services
 Reducing commuting distances








liso


High density urban settlement
 Proximity and mixity of services
 Reducing commuting distances




liso

□ High density urban settlement □ Proximity and mixity of services □ Reducing commuting distances Reduction of energy use/ GHG /air pollutant Organizing energy networks (heating, services) 0 rightarrow travel by car <math>rightarrow walking / cycling0 HOME HOME **WORK** WØRK Leisure Walking / cycling distance Distance we need car Leisure

liso




liso





Integrated modeling of the analysis of impacts of urban policies on air quality






The modeling chain






The modeling chain : NEDUM






NEDUM : Land-Use model

- how the inhabitants choose to locate in a city ?
- how policies can influence their choice ?
- what are the socio-economic effects associated to these policies ?





liso

NEDUM - Methodology

NEDUM

Locate households of a zone in terms

- based on budgetary constraint of households (cost of transport, housing, and size of dwellings)
- **Q** Real estate development depend on the demand
- out of the city


70 65

60 55


<section-header> NEDUM - Validation NEDUM Calibrated on the greater Paris Image: Calibrated on the greater Paris Image: Calibrated on the greater Paris


Mumber and size of houses

Figure : Simulated urbanized area compared to actual urbanized area. Actual urban area appears in black (Source : Corinne Land Cover and IAU), whereas model simulation appears in transparent green. -and-Use Model











The modeling chain : GAIA


GAIA (GenerAtion of synthetic populAtion)

- Generate a synthetic population based on population density (at district level)
- Provides several data (age, sex, job, driving licence, car...)






Greater Paris population pyramid (GAIA)






GAIA - Methodology







GAIA - Validation

0 a 14 ans

Ensemble

GAIA

12


10 -

8 -

6 =

4 =

2 -




30 a 44 ans

15 a 29 ans

population distribution (greater Paris)

45 a 59 ans

60 a 74 ans 75 ans ou plus







The modelling chain : Moirai



- Generation of agent timetable
- Modal split
- Activity duration












MOIRAI - Methodology





Calculation of the attractiveness of jobs center per zone Probabiblity_{Attractivity} (*i*, *j*, *distance*) = K × Attractivity Attractivity(*i*, *j*, *distance*) = Emploi_i × Emploi_j × $\frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{1}{2}(\frac{x-distance_{moyenne}}{\sigma})}$ $\sigma = \frac{distance_{moyenne}}{2}$ $-K = \frac{Attractivity(ij)}{\sum_{j=zone_{0}}^{j=zone_{0}}Attractivity(ij)}$ The attractivities of the activities vary according to the average distance calibrated using the EGT (French Census)







8000000

MOIRAI - Validation

MOIRAI


- **Calibrated on the greater Paris**
- Modal split (work in progress...)

0

2000000


4000000

6000000




1000000 1200000 1400000 1600000 1800000 2000000

(Omnil)











The modeling chain : HERMES


HERMES : Spatial modeling of individualized travel path


- Generation of car fleet
- Generation of the path (Dijsktra algorithm of shortest path Graph theory)





liso





liso

HERMES - Validation


HERMES


- Calibrated on the greater Paris
- □ Validation in progress

The modeling chain : IFRIT

IFRIT : Spatial modeling of individualized travel emissions

□ Modelling the **the exhaust emissions** for passenger car








IFRIT - Methodology

IFRIT : Spatial modeling of individualized travel emissions

 Calculate hot and cold start emissions NOx, NMVOC, CO,PM for every trip (COPERT methodology)



lisa

_copert[@]

Emission factor

100

120


IFRIT - Validation




IFRIT

Comparison with french Air quality monitoring network

in progress...






The modeling chain : HESTIA

HESTIA : Spatial modeling of small combustion emissions

- □ Modelling the **small combustion** installations for every household in residential sector
 - heating
 - provision of hot water






HESTIA - Methodology

HESTIA

Modelling the small combustion installations for every household in residential sector




liso


HESTIA - Validation

HESTIA

- □ Validation in progress
- **Comparison with french Air quality monitoring network**









The modeling chain : CHIMERE

Eulerian off-line chemistry-transport model (CTM)

- \Box Calculate at each timestep ΔC in each box
- □ Forecasting and analysis of air quality
- Driven by weather outputs WRF





lisa

□ triple nested simulation with :

- Global domain (LMDZ-INCA) 1.
- 2. Continental domain (CHIMERE)

Longitude

CHIMERE – simulations

National domain (CHIMERE) 3.

20

atit

46

44


₽

Global domain 1° X 1°

Continental domain 0.5° X 0.5°

National domain 15km X 15km

Regional domain 1km X 1km







Continental domain

0.5° X 0.5°


CHIMERE – validation

CHIMERE


□ Validation in progress

R 0.27

Comparison with french Air quality monitoring network



R and RMSE for NO2 for rural Air-Base stations on the first week of January



Concentrations of NO2 in ppb (City station Paris 13^{ème})

Weather fields Emissions CHIMERE




Regional domain

1km X 1km


Scenarios






Scenarios









liso

Generation of land use scenarios


Study on past urbanization







lisa


Scenarios parametrization




Scenarios - Land-use transport outputs










seholds 17500 15000 12500 - 10000 7500 Nur - 5000 2500






Scenarios - Road emissions outputs








Scenarios – CTM outputs





Exposure

Scenarios – CTM outputs time series




Conclusion

- Operational modeling chain
- □ Sensitivity in terms of air quality
- □ In the center of the city, pollutants concentration appears higher, in average, in the compact scenario
- □ Simplied modeling of the city Change modal split sensibility

Ongoing

Downscaling

Modelling at a smaller scale








Downscaling

Extrapolation of CTM mesoscale to micro-environments **Chimere** Work in progress...

Learning Capacity / Extrapolation

D Potential for transposition to other sites / Standard micro-environments



Conclusion

- Operational modeling chain
- □ Sensitivity in terms of air quality
- □ In the center of the city, pollutants concentration appears higher, in average, in the compact scenario
- □ Simplied modeling of the city Change modal split sensibility


Ongoing

Downscaling

Modelling at a smaller scale

Individual exposure

□ Time geography exposure (Work in progess...)







Individual exposure

Time geography exposure (Work in progess...)

□ MOIRAI + CHIMERE (Downscaling)







Individual daily movement trajectory

Conclusion

Outlook

- Operational modeling chain
- □ Sensitivity in terms of air quality
- □ In the center of the city, pollutants concentration appears higher, in average, in the compact scenario
- □ Simplied modeling of the city Change modal split sensibility

Ongoing

Downscaling

Modelling at a smaller scale

Individual exposure

□ Time geography exposure (Work in progess...)

Interaction on the LUTI

In progress

Deposition

In progress

