



**HAL**  
open science

# Remarks on variable Lebesgue spaces and fractional Navier-Stokes equations

Gastón Vergara-Hermosilla

► **To cite this version:**

Gastón Vergara-Hermosilla. Remarks on variable Lebesgue spaces and fractional Navier-Stokes equations. 2024. hal-04447573v1

**HAL Id: hal-04447573**

**<https://hal.science/hal-04447573v1>**

Preprint submitted on 8 Feb 2024 (v1), last revised 10 Jul 2024 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Remarks on variable Lebesgue spaces and fractional Navier-Stokes equations

Gastón Vergara-Hermosilla\*

LaMME, Univ. Evry, CNRS, Université Paris-Saclay, 91025, Evry, France.

February 8, 2024

## Abstract

In this work we study the 3D Navier-Stokes equations, under the action of an external force and with the fractional Laplacian operator  $(-\Delta)^\alpha$  in the diffusion term, from the point of view of variable Lebesgue spaces. Based on decay estimates of the fractional heat kernel we prove the existence and uniqueness of mild solutions on this functional setting. Thus, in a first theorem we obtain an unique local-in-time solution in the space  $L^{p(\cdot)}([0, T], L^q(\mathbb{R}^3))$ . As a bi-product, in a second theorem we prove the existence of an unique global-in-time solution in the mixed-space  $\mathcal{L}^{\frac{p(\cdot)}{2\alpha-1}}(\mathbb{R}^3, L^\infty([0, T]))$ .

## 1 Introduction

### 1.1 General setting

In this paper we consider the fractional incompressible Navier-Stokes equations defined in  $\mathbb{R}^3$ , namely the system

$$\begin{cases} \partial_t \vec{u} = -(-\Delta)^\alpha \vec{u} - (\vec{u} \cdot \nabla) \vec{u} + \nabla P + \vec{f}, \\ \operatorname{div}(\vec{u}) = 0, \\ \vec{u}(0, \mathbf{x}) = \vec{u}_0(\mathbf{x}), \quad \operatorname{div}(\vec{u}_0) = 0, \quad \mathbf{x} \in \mathbb{R}^3, \end{cases} \quad (1.1)$$

with  $\alpha \in ]1/2, 1]$  and  $(-\Delta)^\alpha$  is the fractional Laplacian operator, which is defined at the Fourier level by the symbol  $|\xi|^{2\alpha}$ . Considering the traditional notation, the vector field  $\vec{u} : [0, +\infty[ \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$  denotes the velocity of a viscous, incompressible and homogeneous fluid,  $P : [0, +\infty[ \times \mathbb{R}^3 \rightarrow \mathbb{R}$  is its pressure and  $\vec{u}_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ ,  $\vec{f} : [0, +\infty[ \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$  stand for a given initial data and a given external force, respectively.

This system is of interest by several reasons. For instance, it can be seen as a natural generalization of the classical incompressible 3D Navier-Stokes equations. In fact, equations (1.1) are the equations resulting from replacing the Laplacian  $(-\Delta)$  in the Navier-Stokes equation by  $(-\Delta)^\alpha$ . On the other hand, system (1.1) has similar energy estimates and scaling properties as the classical Navier-Stokes

---

\*[gaston.vergarahermosilla@univ-evry.fr](mailto:gaston.vergarahermosilla@univ-evry.fr)

equations. In this case, the existence and uniqueness of mild solutions was pioneered by the works of Fujita and Kato [6, 11], where they reformulated the classical Navier-Stokes equation into an integral equation and proved the local well-posedness on appropriate Sobolev and Lebesgue spaces. Since [6, 11], many other functional spaces had been considered in the literature, such as Besov spaces [1, 13], Morrey spaces [7, 10], Fourier-Herz spaces [2] and the  $BMO^{-1}$  space [12]. For a rigorous review about these and other possible functional spaces, see the book [14].

In this work we are concerned by the existence of *mild* solutions of (1.1) which are obtained via a fixed-point method. To develop this theory, the choice of a *appropriate* functional setting is crucial. Lions in [17] proved that in the case that  $\alpha \geq 5/4$ , the fractional 3D Navier-Stokes equations (1.1) have a global and unique regular solution on appropriate Lebesgue and Sobolev spaces (for a simple proof of this fact, see the appendix of [19]). Thus, since the seminal work of Lions many other functional spaces have been considered to obtain global solutions for (1.1), for instance Q-type spaces [16, 15], Besov spaces [20, 22], Triebel-Lizorki type spaces [21], the  $BMO^{1-2\alpha}$  space [23], etc.

In this paper we focus our efforts on the exploration of some existence and uniqueness results for equations (1.1) considering as functional framework the Lebesgue spaces of variable exponent  $L^{p(\cdot)}(\mathbb{R}^3)$ . Intuitively, these spaces are a natural generalization of the classical Lebesgue spaces  $L^p$ , however, in the construction of them there exist particular issues that make these spaces quite dissimilar. Being more precise, in our functional setting the (classical) constant parameter  $p \in [1, +\infty[$  is replaced by a measurable function  $p(\cdot) : \mathbb{R}^3 \rightarrow [1, +\infty[$ . Thus, in order to construct the spaces  $L^{p(\cdot)}$ , we start by considering a measurable function  $\vec{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , and we define the *modular function*  $\rho_{p(\cdot)}$  associated to  $p(\cdot)$  by the formula

$$\rho_{p(\cdot)}(\vec{f}) = \int_{\mathbb{R}^3} |\vec{f}(x)|^{p(x)} dx. \quad (1.2)$$

In this point, we should remark that in the case that  $p(\cdot) \equiv p \in [1, +\infty[$ , it is possible to recover the classical Lebesgue spaces  $L^p$ . In fact, the  $L^p$ -norm can be obtained by considering the modular function as follows,

$$\|\vec{f}\|_{L^p(\mathbb{R}^3)} = \left( \rho_{p(\cdot)}(\vec{f}) \right)^{\frac{1}{p}}.$$

At this stage a relevant difference arises; if we consider the measurable function  $p(\cdot)$ , we cannot simply consider the variable exponent  $\frac{1}{p(\cdot)}$  instead of the constant exponent  $\frac{1}{p}$  in the previous expression. In order to bypass this issue, it is classical to equip the space  $L^{p(\cdot)}$  with the *Luxemburg norm* associated to the modular function  $\rho_{p(\cdot)}$  (see [4], [5] and [9]), which is given by the expression

$$\|\vec{f}\|_{L^{p(\cdot)}(\mathbb{R}^3)} = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^3} \left| \frac{\vec{f}(x)}{\lambda} \right|^{p(x)} dx \leq 1 \right\}. \quad (1.3)$$

Then, we define the variable Lebesgue spaces  $L^{p(\cdot)}(\mathbb{R}^3)$  as the set of measurable functions such that the quantity  $\|\cdot\|_{L^{p(\cdot)}(\mathbb{R}^3)}$  above is finite. A comprehensive presentation on the theory of variable Lebesgue spaces can be consulted in the books [4], [5] and [9].

To the best of our knowledge, these functional spaces have not been considered before in the analysis of the fractional Navier-Stokes equations. Thus, our aim in this article is to present a first application

of the variable Lebesgue spaces to the study of this system. More precisely, here we study mild solutions for the fractional Navier-Stokes equations considering variable Lebesgue spaces  $L^{p(\cdot)}(\mathbb{R}^3)$  as functional setting. The presentation of these results motivate the next subsection.

## 1.2 Presentation of the results

In this subsection we state our main results about the well-posedness of the 3D fractional Navier-Stokes equations on  $L^{p(\cdot)}$ -spaces. In a first theorem, we will study the existence and uniqueness of a mild solution by considering a Lebesgue space of variable exponent in the time variable  $t > 0$ , and by setting a classical  $L^q$ -space in the space variable  $x \in \mathbb{R}^3$ . This first result reads as follows.

**Theorem 1** *Let  $\alpha \in ]\frac{1}{2}, 1]$ ,  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^2)$  with  $p^- > 2$  and fix a parameter  $q > \frac{3}{2\alpha-1}$  by the relationship  $\frac{\alpha}{p(\cdot)} + \frac{3}{2q} < \alpha - \frac{1}{2}$ . If  $\vec{f} \in L^1([0, +\infty[, L^q(\mathbb{R}^3))$  is an exterior force and if is a divergence free initial data  $\vec{u}_0 \in L^q(\mathbb{R}^2)$ , then there exist a time  $0 < T < +\infty$  and an unique mild solution of the fractional Navier-Stokes equations (1.1) in the space  $L^{p(\cdot)}([0, T], L^q(\mathbb{R}^3))$ .*

We should stress the fact that, in this theorem we first measure the information in the space variable and then we measure the information in the time variable. In our next result, we will study a variant of the variable Lebesgue spaces for the space variable  $x \in \mathbb{R}^3$  and we will set the classical space  $L^\infty(\mathbb{R}^3)$  as the functional framework for the time variable  $t \in \mathbb{R}_+$ . The theorem reads as follows.

**Theorem 2** *Consider  $\alpha \in ]\frac{1}{2}, 1]$ , a variable exponent  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  such that  $p^- > 1$ , an initial data  $\vec{u}_0 \in \mathcal{L}^{\frac{p(\cdot)}{3}}(\mathbb{R}^3)$  such that  $\operatorname{div}(\vec{u}_0) = 0$ , and let  $\vec{f}$  be a divergence free external force such that  $\vec{f} = \operatorname{div}(\mathcal{F})$  where  $\mathcal{F}$  is a tensor in  $\mathcal{L}^{\frac{p(\cdot)}{2}}(\mathbb{R}^3, L^\infty([0, T]))$ . If  $\|\vec{u}_0\|_{\mathcal{L}^{\frac{p(\cdot)}{3}}} + \|\mathcal{F}\|_{\mathcal{L}^{\frac{p(\cdot)}{2(2\alpha-1)}, x}(L_t^\infty)}$  is small enough, then the fractional Navier-Stokes equations (1.1) admits a unique, global mild solution in the space  $\mathcal{L}^{\frac{p(\cdot)}{3}}(\mathbb{R}^3, L^\infty([0, T]))$ .*

Note that, in this theorem we have considered the *mixed variable Lebesgue spaces*  $\mathcal{L}_p^{p(\cdot)}$  (see Subsection 2.1 for the definition of these spaces) by merely technical reasons and it is motivated by the lack of flexibility in the parameters that intervene in the boundedness of the Riesz transforms involved in its proof. See Subsection 2.1 and Remark 3.4 below for more details on this particular issue.

## Organization of the paper

The present paper is structured as follows. In Section 2 we present a concise review of the main definitions and properties of the Variable Lebesgue spaces  $L^{p(\cdot)}$  and fractional heats kernels. Section 3 is devoted to the proof of the theorems.

## 2 Preliminaries

To keep this paper reasonably self-contained, several results and definitions on variable Lebesgue spaces and fractional heat kernels are recalled. Thus, we begin by presenting a brief summary on the key elements of variable Lebesgue spaces theory.

## 2.1 Variable Lebesgue spaces

We start this subsection by formalising the notion of variable exponent, set of variable exponents and limit exponents.

**Definition 2.1** *Let consider  $n \in \mathbb{N}$ , and a function  $p : \mathbb{R}^n \rightarrow [1, +\infty[$ . Then,  $p$  will be called a variable exponent if  $p(\cdot)$  is a measurable function and we define  $\mathcal{P}(\mathbb{R}^n)$  to be the set of variable exponents. Moreover, we define the limit exponents  $p^- = \inf \text{ess}_{x \in \mathbb{R}^n} \{p(x)\}$  and  $p^+ = \sup \text{ess}_{x \in \mathbb{R}^n} \{p(x)\}$ .*

For the sake of simplicity in the rest of the paper we will assume  $1 < p^- \leq p^+ < +\infty$ , and in order of differentiate between variable and constant exponents, we will always denote variable exponent by  $p(\cdot)$ . Note that, the spaces  $L^{p(\cdot)}(\mathbb{R}^n)$  are Banach function spaces and they have very interesting features. Thus, we start by presenting the generalization of the Hölder inequalities in this setting.

**Lemma 2.1** *Let  $p(\cdot), q(\cdot), r(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  be functions such that we have the pointwise relationship  $\frac{1}{p(x)} = \frac{1}{q(x)} + \frac{1}{r(x)}$ ,  $x \in \mathbb{R}^n$ . Then there exists a constant  $C > 0$  such that for all  $f \in L^{q(\cdot)}(\mathbb{R}^n)$  and  $g \in L^{r(\cdot)}(\mathbb{R}^n)$ , the pointwise product  $fg$  belongs to the space  $L^{p(\cdot)}(\mathbb{R}^n)$  and we have the estimate*

$$\|fg\|_{L^{p(\cdot)}} \leq C \|f\|_{L^{q(\cdot)}} \|g\|_{L^{r(\cdot)}}. \quad (2.1)$$

As is natural, this estimate can be easily generalized to vector fields  $\vec{f}, \vec{g} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  and to the product  $\vec{f} \cdot \vec{g}$ . For a proof of this result we recommend to the interested reader to [4, Section 2.4] or [5, Section 3.2].

Note that the quantity  $\|\cdot\|_{L^{p(\cdot)}}$  satisfies the *Norm conjugate formula* given in [5, Corollary 3.2.14]:

**Proposition 2.1** *Let  $p(\cdot), q(\cdot), r(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  be functions such that we have the pointwise relationship  $1 = \frac{1}{p(x)} + \frac{1}{q(x)}$ ,  $x \in \mathbb{R}^n$ . Then, for all  $f \in L^{p(\cdot)}$  we have*

$$\|f\|_{L^{p(\cdot)}} \leq 2 \sup_{\|g\|_{L^{q(\cdot)}} \leq 1} \int_{\mathbb{R}^n} |f(x)| |g(x)| dx. \quad (2.2)$$

**Remark 2.1** *Note that in the previous notions the space  $\mathbb{R}^n$  can be replaced by an interval  $[0, T]$ .*

To continue we present the following embedding result [4, Corollary 2.48].

**Lemma 2.2** *Let consider  $n \geq 1$ , a bounded domain  $\mathcal{X} \subset \mathbb{R}^n$  and two variable exponents  $q_1(\cdot), q_2(\cdot) \in \mathcal{P}(\mathcal{X})$  such that  $1 < q_1^+, q_2^+ < +\infty$ . Then,  $L^{q_2(\cdot)}(\mathcal{X}) \subset L^{q_1(\cdot)}(\mathcal{X})$  if and only if  $q_1(x) \leq q_2(x)$  almost everywhere. Furthermore, in this case we have that*

$$\|f\|_{L^{q_1(\cdot)}} \leq (1 + |\mathcal{X}|) \|f\|_{L^{q_2(\cdot)}}.$$

It is important to emphasize that the convolution product  $f * g$  is not well adapted to the structure of the  $L^{p(\cdot)}$  spaces, in particular the Young inequalities for convolution are not valid anymore (see [5, Section 3.6]) and thus many of the usual operators that appear in PDEs must be treated very carefully. Note also that Fourier-based methods are not so easy to use as we lack of an alternative for the Plancherel formula.

In order to study the boundedness of such operators is classical to impose some conditions on the variable exponents  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ . To this end, we introduce the following definition.

**Definition 2.2** Let  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  and consider the limit value  $\frac{1}{p_\infty} = \lim_{|x| \rightarrow +\infty} \frac{1}{p(x)}$ . We say that  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  belongs to the class  $\mathcal{P}^{\log}(\mathbb{R}^n)$  if we have:

1.  $\left| \frac{1}{p(x)} - \frac{1}{p(y)} \right| \leq \frac{C}{\log(e+1/|x-y|)}$  for all  $x, y \in \mathbb{R}^n$ , and
2.  $\left| \frac{1}{p(x)} - \frac{1}{p_\infty} \right| \leq \frac{C}{\log(e+|x|)}$  for all  $x \in \mathbb{R}^n$ .

With the condition  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  we can state the following definition and theorem.

**Definition 2.3** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  a locally integrable function. The Hardy-Littlewood maximal function  $\mathcal{M}$  is defined by

$$\mathcal{M}(f)(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy$$

where  $B$  is an open ball of  $\mathbb{R}^n$ .

**Theorem 3** Let  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  with  $p^- > 1$ . Then, there exist a constant  $C > 0$  such that

$$\|\mathcal{M}(f)\|_{L^{p(\cdot)}} \leq C \|f\|_{L^{p(\cdot)}}. \quad (2.3)$$

A proof of this result can be consulted in [5, Section 4.3]. At this point, we recall a classical result about the Hardy-Littlewood maximal function (see [8, Section 2.1]):

**Lemma 2.3** If  $\varphi$  is a radially decreasing function on  $\mathbb{R}^3$  and  $\vec{f}$  is a locally integrable function, then

$$|(\varphi * \vec{f})(x)| \leq \|\varphi\|_{L^1} \mathcal{M}(\vec{f})(x),$$

where  $\mathcal{M}$  is the Hardy-Littlewood maximal function.

We remark that the usual Riesz transforms  $(\mathcal{R}_j)_{1 \leq j \leq n}$  defined formally in the Fourier level by  $\widehat{\mathcal{R}_j(f)}(\xi) = -\frac{i\xi_j}{|\xi|} \widehat{f}(\xi)$  are also bounded in Lebesgue spaces of variable exponent. This fact is stated in the following Lemma.

**Lemma 2.4** Let  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  and  $1 < p^- \leq p^+ < +\infty$ . Then, given  $f \in L^{p(\cdot)}$ , there exist a constant  $C > 0$  such that

$$\|\mathcal{R}_j(f)\|_{L^{p(\cdot)}} \leq C \|f\|_{L^{p(\cdot)}}, \quad (2.4)$$

A proof of this result can be consulted in [5, Sections 6.3 and 12.4]. To continue, we introduce the following classical operator.

**Definition 2.4** Let  $0 < \beta < n$ . Then, given a measurable function  $f$ , we define the Riesz potential operator  $\mathcal{J}_\beta(f) : \mathbb{R}^n \rightarrow [0, +\infty]$  by

$$\mathcal{J}_\beta(f)(x) := \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^{n-\beta}} dy. \quad (2.5)$$

The Riesz potential operator is bounded in variable Lebesgue spaces if we consider appropriate variable exponents in the class  $\mathcal{P}^{\log}(\mathbb{R}^n)$ . A precise statement of this result reads as follows.

**Theorem 4** Let  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  and  $0 < \beta < n/p^+$ . Then, there exist  $C > 0$  such that

$$\|J_\beta(f)\|_{L^{q(\cdot)}} \leq C \|f\|_{L^{p(\cdot)}}, \quad \text{with} \quad \frac{1}{q(\cdot)} = \frac{1}{p(\cdot)} - \frac{\beta}{n}. \quad (2.6)$$

A proof of this theorem can be consulted in [5, Section 6.1].

Note that the estimate (2.6) introduces a very strong relationship between the variable exponents  $p(\cdot)$  and  $q(\cdot)$ . Thus, in order to obtain more flexibility on these parameters (see Remark 3.4 below) we will consider the following spaces introduced in [3].

**Definition 2.5** Let  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  a variable exponent and  $1 < p < +\infty$  a constant. Then, the mixed Lebesgue space  $\mathcal{L}_p^{p(\cdot)}(\mathbb{R}^n)$  is defined by

$$\mathcal{L}_p^{p(\cdot)}(\mathbb{R}^n) = L^{p(\cdot)}(\mathbb{R}^n) \cap L^p(\mathbb{R}^n),$$

which can be normed by the quantity

$$\|\cdot\|_{\mathcal{L}_p^{p(\cdot)}} = \max\{\|\cdot\|_{L^{p(\cdot)}}, \|\cdot\|_{L^p}\}. \quad (2.7)$$

With the help of these spaces we have the following result.

**Proposition 2.2** Let  $1 < p < +\infty$  be a constant exponent,  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$  a variable exponent and fix a parameter  $0 < \beta < \min\{n/p^+, n/p\}$ . Given,  $f \in \mathcal{L}_p^{p(\cdot)}(\mathbb{R}^n)$  and a function  $\rho(\cdot)$  satisfying the following condition

$$\rho(\cdot) = \frac{np(\cdot)}{n - sp}, \quad (2.8)$$

then, there exist a constant  $C > 0$  such that

$$\|J_\beta(f)\|_{L^{\rho(\cdot)}} \leq C \|f\|_{\mathcal{L}_p^{p(\cdot)}}. \quad (2.9)$$

A proof of this result can be consulted in [3].

**Remark 2.2** We emphasize in particular the fact that the index  $p$  is not related to  $p^-$  or  $p^+$  nor to  $p(\cdot)$  and this inequality gives more flexibility in the exponents than the conditions stated in Theorem 4.

**Remark 2.3** Note that, by construction, the mixed spaces  $\mathcal{L}_p^{p(\cdot)}$  inherit the properties of the spaces  $L^{p(\cdot)}$  and  $L^p$ . In particular we have the Hölder inequality  $\|\varphi\|_{\mathcal{L}_p^{p(\cdot)}} \leq \|\varphi\|_{\mathcal{L}_q^{q(\cdot)}} \|\varphi\|_{\mathcal{L}_r^{r(\cdot)}}$  with  $\frac{1}{p(\cdot)} = \frac{1}{q(\cdot)} + \frac{1}{r(\cdot)}$  and  $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$  and of course the Riesz transforms are also bounded in these spaces.

For more details about the Lebesgue spaces of variable exponent, their inner structure as well as many other properties, see the books [4], [5] and [9].

## 2.2 Some preliminary estimates on the fractional heat kernel

In this subsection we recall some estimates related to the fractional heat kernel involved in the integral formulation of the system (1.1).

**Remark 2.4** Consider  $x \in \mathbb{R}^n$  and the fractional heat kernel  $\mathfrak{g}_t^\alpha(x)$ . Then, there exist a constant  $C > 0$  such that the following pointwise estimate follows

$$|\nabla \mathfrak{g}_t^\alpha(x)| \leq C \frac{1}{(t^{\frac{1}{2\alpha}} + |x|)^{n+1}}. \quad (2.10)$$

More details about this remark can be consulted in [18, Remark 2.1]. Another useful result for our purposes is the following.

**Lemma 2.5** Consider  $x \in \mathbb{R}^n$ , the fractional heat kernel  $\mathfrak{g}_t^\alpha(x)$ , and the parameters  $1 \leq r \leq p \leq +\infty$ . Then, for  $\alpha, \nu > 0$ , there exist  $C > 0$  such that

1.  $\|\mathfrak{g}_t^\alpha * \varphi(x)\|_{L^p} \leq C t^{-\frac{n}{2\alpha}(\frac{1}{r}-\frac{1}{p})} \|\varphi\|_{L^r},$
2.  $\|(-\Delta)^{\nu/2} \mathfrak{g}_t^\alpha * \varphi(x)\|_{L^p} \leq C t^{-\frac{\nu}{2\alpha} - \frac{n}{2\alpha}(\frac{1}{r}-\frac{1}{p})} \|\varphi\|_{L^r}.$

A proof of this result can be consulted in [18, Lemma 3.1].

## 3 Mild solutions in variable Lebesgue spaces

We present here a general approach to mild solutions for the fractional Navier-Stokes equations (1.1) in the setting of variable Lebesgue spaces. These mild solutions are obtained via the following classical result:

**Theorem 5 (Banach-Picard principle)** Consider a Banach space  $(E, \|\cdot\|_E)$  and a bounded bilinear application  $\mathcal{B} : E \times E \rightarrow E$ :

$$\|\mathcal{B}(e, e)\|_E \leq C_{\mathcal{B}} \|e\|_E \|e\|_E.$$

Given  $e_0 \in E$  such that  $\|e_0\|_E \leq \delta$  with  $0 < \delta < \frac{1}{4C_{\mathcal{B}}}$ , then the equation

$$e = e_0 - \mathcal{B}(e, e),$$

admits an unique solution  $e \in E$  which satisfies  $\|e\|_E \leq 2\delta$ .

Note that the divergence-free condition allows us to recover the pressure  $P$  from  $\vec{u}$  through the identity

$$-\Delta P = \operatorname{div}(\vec{u} \cdot \vec{\nabla} \vec{u}),$$

thus, in the following we will focus our analysis on the velocity field  $\vec{u}$ .

In order to apply this result to the fractional Navier-Stokes equations (1.1) we need to get rid of the pressure  $P$  and for this we apply to this system the Leray projector  $\mathbb{P}$  defined by  $\mathbb{P} := \operatorname{Id} + \vec{\nabla}(-\Delta)^{-1} \operatorname{div}$ .

**Remark 3.1** We emphasise the fact that the Leray projector  $\mathbb{P}$  can be defined equivalently by considering Riesz transforms. In fact, given a function  $\vec{g}$  in a functional space  $\mathcal{X}$ , the Leray projector is defined as follows:  $\mathbb{P}(\vec{g}) = (\operatorname{Id}_{3 \times 3} - \vec{R} \otimes \vec{R})(\vec{g})$  where  $\vec{R} = (R_1, R_2, R_3)$ , with  $R_j$  denoting the  $j$ -th Riesz transform. Thus, if the Riesz transforms are bounded in  $\mathcal{X}$ , then the Leray's projector  $\mathbb{P}$  is also bounded on  $\mathcal{X}$ , and then we can write  $\|\mathbb{P}(\vec{g})\|_{\mathcal{X}} \leq C \|\vec{g}\|_{\mathcal{X}}$ .



As we anticipated above, considering the pressure term  $\mathbb{P}$  in (1.1) we have  $\mathbb{P}(\vec{\nabla}\mathbb{P}) \equiv 0$ . Moreover, given a vector field is divergence free we have the identity  $\mathbb{P}(\vec{g}) = \vec{g}$ . A proof of these facts and many others properties of  $\mathbb{P}$ , can be consulted in the book [14].

Since  $\vec{u}$  and  $\vec{f}$  are divergence free, by applying Leray's projector to 1.1 we obtain the equation

$$\begin{cases} \partial_t \vec{u} = -(-\Delta)^\alpha \vec{u} - \mathbb{P}(\operatorname{div}(\vec{u} \otimes \vec{u})) + \vec{f}, & \operatorname{div}(\vec{u}) = 0, \\ \vec{u}(0, \mathbf{x}) = \vec{u}_0(\mathbf{x}), & \mathbf{x} \in \mathbb{R}^3. \end{cases}$$

Now, due to the Dumahel formula, we can write this equation in the following form

$$\vec{u}(t, \mathbf{x}) = \mathbf{g}_t^\alpha * \vec{u}_0(\mathbf{x}) + \int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \mathbf{x}) ds - \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{u} \otimes \vec{u}))(s, \mathbf{x}) ds, \quad (3.1)$$

where  $\mathbf{g}_t^\alpha$  is the usual fractional heat kernel. In the following we will consider the integro-differential equation above in order to apply the Banach-Picard principle. To this end, we set the bilinear application as

$$\mathcal{B}(\vec{u}, \vec{v}) = \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{u} \otimes \vec{v}))(s, \mathbf{x}), \quad (3.2)$$

and we consider the term  $\mathbf{g}_t^\alpha * \vec{u}_0(\mathbf{x}) + \int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \mathbf{x}) ds$  as  $\mathbf{e}_0$  (in Theorem 5).

### 3.1 Proof of Theorem 1

In the following we consider a variable exponent  $\mathbf{p}(\cdot) \in \mathcal{P}^{\log}([0, +\infty[)$ , and the functional space

$$\mathfrak{E}_T = L^{\mathbf{p}(\cdot)}([0, T], L^q(\mathbb{R}^2)),$$

with  $T \in ]0, +\infty[$  to be precised later. The space  $\mathfrak{E}_T$  is endowed with a Luxemburg norm as follows

$$\|\vec{\varphi}\|_{\mathfrak{E}_T} = \inf \left\{ \lambda > 0 : \int_0^T \left| \frac{\|\vec{\varphi}(t, \cdot)\|_{L^q}}{\lambda} \right|^{\mathbf{p}(t)} dt \leq 1 \right\}. \quad (3.3)$$

Under this functional setting we will consider the Banach-Picard principle to construct mild solutions for the integral equation (3.1). More precisely, in the following we will prove 3 propositions which will provide the core of the hypotheses of Theorem 5.

To this end, we star by proving the following result regarding a control to the initial data.

**Proposition 3.1** *Let  $\alpha \in ]\frac{1}{2}, 1]$ ,  $\mathbf{p}(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^2)$  with  $\mathbf{p}^- > 2$  and fix an index  $q > \frac{3}{2\alpha-1}$  by the relationship  $\frac{\alpha}{\mathbf{p}(\cdot)} + \frac{3}{2q} < \alpha - \frac{1}{2}$ . Consider a function  $\vec{u}_0 \in L^q(\mathbb{R}^2)$ . Then, there exist a constant  $C_1 > 0$  such that*

$$\|\mathbf{g}_t^\alpha * \vec{u}_0\|_{\mathfrak{E}_T} \leq C_1 \|\vec{u}_0\|_{L^q}. \quad (3.4)$$

**Proof:** In order to conclude the inequality (3.4), we start by considering a Young inequality to obtain

$$\|\mathbf{g}_t^\alpha * \vec{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)} \leq \|\mathbf{g}_t^\alpha\|_{L^1(\mathbb{R}^2)} \|\vec{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)} = \|\vec{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)}.$$

Before continue, let consider the following classical result in the context of Variable Lebesgue spaces (see[5, Lemma 3.2.12, Section 3.2]).

**Lemma 3.1** *Let  $p(\cdot) \in \mathcal{P}([0, +\infty[)$  such that  $1 < p^- \leq p^+ < +\infty$ . Then, there exist  $C > 0$  such that*

$$\|1\|_{L^{p(\cdot)}([0, T])} \leq C \max \left\{ T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}} \right\}.$$

Thus, by taking the  $L_t^{p(\cdot)}$ -norm we obtain

$$\|\mathbf{g}_t^\alpha * \vec{\mathbf{u}}_0\|_{L_t^{p(\cdot)} L_x^q} \leq C \|\vec{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)} \|1\|_{L^{p(\cdot)}([0, T])} \quad (3.5)$$

$$\leq C \|\vec{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)} \max \left\{ T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}} \right\}. \quad (3.6)$$

Thus, by considering  $C_1 = C \max \left\{ T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}} \right\}$  we deduce (3.4) and we conclude the proof of Proposition 3.1.  $\square$

**Proposition 3.2** *Let  $\alpha \in ]\frac{1}{2}, 1]$ ,  $p(\cdot) \in \mathcal{P}^{log}(\mathbb{R}^2)$  with  $p^- > 2$  and fix an index  $q > \frac{3}{2\alpha-1}$  by the relationship  $\frac{\alpha}{p(\cdot)} + \frac{3}{2q} < \alpha - \frac{1}{2}$ . Then, given a function  $\vec{f} \in L^{p(\cdot)}([0, +\infty[, L^q(\mathbb{R}^3))$ , there exist a numerical constant  $C_2 > 0$  such that*

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \cdot) ds \right\|_{\mathfrak{E}_T} \leq C_2 \|\vec{f}\|_{L_t^1(L_x^q)}. \quad (3.7)$$

**Proof:** To conclude the inequality (3.7), we start by considering the  $L_x^q$ -norm to the term  $\int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \cdot) ds$  in order to get

$$\begin{aligned} \left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \cdot) ds \right\|_{L^q} &\leq \int_0^t \|\mathbf{g}_{t-s}^\alpha\|_{L^1} \|\vec{f}(s, \cdot)\|_{L^q} ds \\ &\leq C \|\vec{f}\|_{L_t^1(L_x^q)}. \end{aligned}$$

Then, by taking the  $L_t^{p(\cdot)}$ -norm in the time variable and considering Lemma 3.1 we deduce the estimates

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \vec{f}(s, \cdot) ds \right\|_{L_t^{p(\cdot)}(L_x^q)} \leq C \left\| \|\vec{f}\|_{L_t^1(L_x^q)} \right\|_{L^{p(\cdot)}} \quad (3.8)$$

$$\begin{aligned} &\leq C \|\vec{f}\|_{L_t^1(L_x^q)} \|1\|_{L^{p(\cdot)}([0, T])} \\ &\leq C \|\vec{f}\|_{L_t^1(L_x^q)} \max \left\{ T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}} \right\}. \end{aligned} \quad (3.9)$$

Thus, by considering  $C_2 = C \max \left\{ T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}} \right\}$  we deduce the inequality (3.7) and then we conclude the proof of Proposition 3.2.  $\square$

**Proposition 3.3** *Let  $\alpha \in ]\frac{1}{2}, 1]$ ,  $p(\cdot) \in \mathcal{P}^{log}(\mathbb{R}^2)$  with  $p^- > 2$  and fix an index  $q > \frac{3}{2\alpha-1}$  by the relationship  $\frac{\alpha}{p(\cdot)} + \frac{3}{2q} < \alpha - \frac{1}{2}$ . Then, there exist a constant  $C_B > 0$  such that*

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) ds \right\|_{\mathfrak{E}_T} \leq C_B \|\vec{\mathbf{u}}\|_{\mathfrak{E}_T} \|\vec{\mathbf{u}}\|_{\mathfrak{E}_T}. \quad (3.10)$$

**Proof:** To conclude the inequality (3.22), we start by considering the  $L_x^q$ -norm to the term  $\int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) ds$  to get

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) ds \right\|_{L^q} \leq C \int_0^t \left\| \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) \right\|_{L^q} ds.$$

Considering the properties of the Leray projector  $\mathbb{P}$  and the fact that the Riesz transform which define it are bounded on classical Lebesgue spaces  $L^q$ , with  $1 < q < +\infty$ , we conclude

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) ds \right\|_{L^q} \leq C \int_0^t \left\| \vec{\nabla} \mathbf{g}_{t-s}^\alpha * \vec{\mathbf{u}} \otimes \vec{\mathbf{u}}(s, \cdot) \right\|_{L^q} ds.$$

Now, note that a Young inequality, with  $1 + \frac{1}{q} = \frac{q-1}{q} + \frac{2}{q}$ , and Lemma 2.5, with parameters  $\nu = 1$ ,  $n = 3$ ,  $p = p$  and  $r = \frac{p}{2}$ , yield

$$\int_0^t \left\| \vec{\nabla} \mathbf{g}_{t-s}^\alpha * \vec{\mathbf{u}} \otimes \vec{\mathbf{u}}(s, \cdot) \right\|_{L^q} ds \leq C \int_0^t \frac{1}{(t-s)^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q} ds.$$

To continue, we take  $L_t^{p(\cdot)}$ -norm and we consider the conjugate exponent  $p'(\cdot)$  defined by  $1 = \frac{1}{p(\cdot)} + \frac{1}{p'(\cdot)}$ . Then, by considering the norm conjugate formula (2.2) we can write

$$\begin{aligned} \left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(s, \cdot) ds \right\|_{L_t^{p(\cdot)}(L_x^q)} &\leq \left\| \int_0^t \frac{1}{(t-s)^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds \right\|_{L_t^{p'(\cdot)}([0, T])} \\ &\leq \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \int_0^t \frac{|\psi(t)|}{|t-s|^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds dt. \end{aligned} \quad (3.11)$$

The Fubini Theorem yields

$$\sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \int_0^t \frac{|\psi(t)|}{|t-s|^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds dt = \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \int_0^T \frac{1_{\{0 < s < t\}} |\psi(t)|}{|t-s|^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} dt \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds,$$

thus, by extending the function  $\psi(t)$  by zero on  $\mathbb{R} \setminus [0, T]$ , we obtain

$$\begin{aligned} \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \int_0^t \frac{|\psi(t)|}{|t-s|^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds dt &= \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \left( \int_{-\infty}^{+\infty} \frac{|\psi(t)|}{|t-s|^{\frac{1}{2\alpha} + \frac{3}{2\alpha q}}} dt \right) \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds \\ &= \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \mathcal{J}_\beta(|\psi|)(s) \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds, \end{aligned}$$

where  $\mathcal{J}_\beta$  is the 1D Riesz potential with  $\beta = 1 - \frac{1}{2\alpha} - \frac{3}{2\alpha q} < 1$  (see Definition 2.4).

**Remark 3.2** We emphasise the fact that the constraints  $\frac{3}{2\alpha-1} < q$  and  $\alpha \in ]\frac{1}{2}, 1]$  imply  $0 < 1 - \frac{1}{2\alpha} - \frac{3}{2\alpha q} < 1$ , and thus the Riesz potential considered is well defined.

An Hölder inequality with  $1 = \frac{1}{p(\cdot)} + \frac{1}{p(\cdot)} + \frac{1}{\tilde{p}(\cdot)}$  yields

$$\sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \int_0^T \mathcal{J}_\beta(|\psi|)(s) \|\vec{\mathbf{u}}(s, \cdot)\|_{L^q}^2 ds \leq C \sup_{\|\psi\|_{L^{p'(\cdot)}} \leq 1} \|\mathcal{J}_\beta(|\psi|)\|_{L_t^{\tilde{p}(\cdot)}} \left\| \|\vec{\mathbf{u}}(\cdot, \cdot)\|_{L_x^q} \right\|_{L_t^{p(\cdot)}} \left\| \|\vec{\mathbf{u}}(\cdot, \cdot)\|_{L_x^q} \right\|_{L_t^{p(\cdot)}}.$$

**Remark 3.3** Note that the condition  $p^- > 2$  in the statement of the proposition become from the relationship  $1 = \frac{2}{p(\cdot)} + \frac{1}{\tilde{p}(\cdot)}$ .

Thus by considering the indexes defined by the relationship

$$\frac{1}{\tilde{p}(\cdot)} = \frac{1}{r(\cdot)} - \left(1 - \frac{1}{2\alpha} - \frac{3}{2\alpha q}\right), \quad (3.12)$$

on Theorem 4, we obtain

$$\begin{aligned} \sup_{\|\Psi\|_{L^{p'(\cdot)}} \leq 1} \|\mathcal{J}_\beta(|\Psi|)\|_{L^{\tilde{p}(\cdot)}} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \\ \leq C \sup_{\|\Psi\|_{L^{p'(\cdot)}} \leq 1} \|\Psi\|_{L_t^{r(\cdot)}} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)}. \end{aligned} \quad (3.13)$$

Gathering the hypothesis  $\frac{\alpha}{p(\cdot)} + \frac{3}{2q} < \alpha - \frac{1}{2}$  and the relationship (3.12) we conclude that

$$\frac{1}{\tilde{p}(\cdot)} = 1 - \frac{2}{p(\cdot)} \quad \text{and} \quad \frac{1}{p'(\cdot)} = 1 - \frac{1}{p(\cdot)},$$

and then, we deduce  $r(\cdot) < p'(\cdot)$ . Considering Lemma 2.2 with  $r(\cdot) < p'(\cdot)$  and  $\mathcal{X} = [0, T]$  in (3.13) we get

$$\begin{aligned} \sup_{\|\Psi\|_{L^{p'(\cdot)}} \leq 1} \|\Psi\|_{L_t^{r(\cdot)}} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \\ \leq \sup_{\|\Psi\|_{L^{p'(\cdot)}} \leq 1} (1 + T) \|\Psi\|_{L_t^{p'(\cdot)}} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \\ \leq (1 + T) \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)}. \end{aligned}$$

Thus, by gathering these estimates with (3.11) we deduce

$$\left\| \int_0^t \mathfrak{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\tilde{\mathbf{u}} \otimes \tilde{\mathbf{u}}))(\cdot, \cdot) ds \right\|_{L_t^{p(\cdot)}(L_x^q)} \leq C(1 + T) \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)} \|\tilde{\mathbf{u}}(\cdot, \cdot)\|_{L_t^{p(\cdot)}(L_x^q)}. \quad (3.14)$$

Thus, by considering  $C_B = C(1 + T)$  we deduce the inequality (3.22) and then we conclude the proof of Proposition 3.3.  $\square$

## End of the proof of Theorem 1

Gathering together the hypothesis assumed in the statement of the theorem and the estimates obtained in Propositions 3.1, 3.2 and 3.3, we conclude that there exist  $0 < T < +\infty$  such that

$$\|\tilde{\mathbf{u}}_0\|_{L^q(\mathbb{R}^2)} + \|\vec{f}\|_{L_t^1(L_x^q)} \leq \frac{C}{(1 + T) \max\left\{T^{\frac{1}{p^-}}, T^{\frac{1}{p^+}}\right\}}.$$

Thus, by applying the Banach-Picard principle we obtain the desired solution. With this we conclude the proof of Theorem 1.

### 3.2 Proof of Theorem 2

In the following we consider a variable exponent  $p(\cdot) \in \mathcal{P}^{\log}([0, +\infty[)$ , and the functional space

$$\mathcal{E} = \mathcal{L}^{\frac{p(\cdot)}{2\alpha-1}}(\mathbb{R}^3, L^\infty([0, T])).$$

The space  $\mathcal{E}$  is endowed with a Luxemburg norm as follows

$$\|\cdot\|_{\mathcal{E}} = \max\{\|\cdot\|_{L_x^{p(\cdot)}(L_t^\infty)}, \|\cdot\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)}\}, \quad (3.15)$$

Under this functional setting we will consider the Banach-Picard principle to construct mild solutions for the integral equation (3.1). More precisely, in the following we will prove 3 propositions which will provide the core of the hypotheses of Theorem 5.

To this end, we start by proving the following result regarding a control to the initial data.

**Proposition 3.4** *Consider  $\alpha \in ]\frac{1}{2}, 1]$ , consider a variable exponent  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  such that  $p^- > 1$ , and a divergence free function  $\vec{u}_0 \in \mathcal{L}^{\frac{p(\cdot)}{2\alpha-1}}(\mathbb{R}^3)$ . Then, there exist a constant  $C > 0$  such that*

$$\|\mathfrak{g}_t^\alpha * \vec{u}_0\|_{\mathcal{E}} \leq C \|\vec{u}_0\|_{\mathcal{L}^{\frac{p(\cdot)}{2\alpha-1}}}. \quad (3.16)$$

**Proof:** Since  $\vec{u}_0 \in L^{\frac{3}{2\alpha-1}} \subset L_{\text{loc}}^1$ , we have that  $\vec{u}_0$  is a locally integrable function. Now, considering that the fractional heat kernel  $\mathfrak{g}_t^\alpha$  is a radially decreasing function, by Lemma 2.3 we can write

$$\|\mathfrak{g}_t^\alpha * \vec{u}_0(x)\|_{L_t^\infty} \leq C \mathcal{M}(\vec{u}_0)(x).$$

Thus, by recalling the norm defined in (3.15), we obtain the estimate

$$\|\mathfrak{g}_t^\alpha * \vec{u}_0\|_{\mathcal{E}} \leq C \max\{\|\mathcal{M}(\vec{u}_0)\|_{L^{p(\cdot)}}, \|\mathcal{M}(\vec{u}_0)\|_{L^{\frac{3}{2\alpha-1}}}\}.$$

Now, since  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  with  $p^- > 1$ , by Theorem 3 we conclude that the maximal function  $\mathcal{M}$  is bounded in the Lebesgue space  $L^{p(\cdot)}(\mathbb{R}^3)$ . Considering this, and the fact that  $\mathcal{M}$  is also bounded in  $L^{\frac{3}{2\alpha-1}}$ , we obtain

$$\|\mathfrak{g}_t^\alpha * \vec{u}_0\|_{\mathcal{E}} \leq C \max\{\|\vec{u}_0\|_{L^{p(\cdot)}}, \|\vec{u}_0\|_{L^{\frac{3}{2\alpha-1}}}\} \leq C \|\vec{u}_0\|_{\mathcal{L}^{\frac{p(\cdot)}{2\alpha-1}}}.$$

With this we conclude the proof.  $\square$

**Proposition 3.5** *Consider  $\alpha \in ]\frac{1}{2}, 1]$ , a variable exponent  $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  such that  $p^- > 1$ , and a function  $\mathcal{F} \in \mathcal{L}^{\frac{p(\cdot)}{2(2\alpha-1)}}(\mathbb{R}^3, L^\infty([0, T]))$ . Then, there exist a numerical constant  $C > 0$  such that*

$$\left\| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(\cdot, \cdot) ds \right\|_{\mathcal{E}} \leq C \|\mathcal{F}\|_{\mathcal{L}^{\frac{p(\cdot)}{2(2\alpha-1)}, x}(L_t^\infty)}. \quad (3.17)$$

**Proof:** By the Minkowski's integral inequality, we can write

$$\left| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right| \leq C \int_0^t \int_{\mathbb{R}^3} |\vec{\nabla} \mathbf{g}_{t-s}^\alpha(\mathbf{x} - \mathbf{y})| |\mathcal{F}(s, \mathbf{y})| d\mathbf{y} ds.$$

Then, by the decay properties of the fractional heat kernel in Remark 2.4, and the Fubini theorem, we obtain

$$\left| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right| \leq C \int_{\mathbb{R}^3} \int_0^t \frac{1}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} |\mathcal{F}(s, \mathbf{y})| ds d\mathbf{y}.$$

Now, by considering the  $L_t^\infty$  norm on  $\mathcal{F}$ , we get

$$\left| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right| \leq C \int_{\mathbb{R}^3} \int_0^t \frac{1}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} ds \|\mathcal{F}(\cdot, \mathbf{y})\|_{L_t^\infty} d\mathbf{y}.$$

Then, considering the Riesz potential defined in (2.4), the fact that  $\alpha \in ]\frac{1}{2}, 1]$ , and the estimate

$$\begin{aligned} \int_0^t \frac{ds}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} &\leq \int_0^{+\infty} \frac{ds}{(s^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} \\ &= \int_0^{+\infty} \frac{|\mathbf{x} - \mathbf{y}|^{2\alpha} d\beta}{((|\mathbf{x} - \mathbf{y}|^{2\alpha} \beta)^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} = \frac{1}{|\mathbf{x} - \mathbf{y}|^{4-2\alpha}} \int_0^{+\infty} \frac{d\beta}{(1 + \beta^{\frac{1}{2\alpha}})^4}, \end{aligned} \quad (3.18)$$

we obtain

$$\left| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right| \leq C \int_{\mathbb{R}^3} \frac{1}{|\mathbf{x} - \mathbf{y}|^{4-2\alpha}} \|\mathcal{F}(\cdot, \mathbf{y})\|_{L_t^\infty} d\mathbf{y} = C \mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\mathbf{x}).$$

Note that, this last estimates implies

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right\|_{L_t^\infty} \leq C \mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\mathbf{x}). \quad (3.19)$$

Then, to obtain the  $\mathcal{L}_{\frac{3}{2\alpha-1}}^{\mathbf{p}(\cdot)}$ -norm given in (3.15), from the estimate (3.19) we get

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right\|_{\mathcal{L}_x^{\mathbf{p}(\cdot)}(L_t^\infty)} \leq C \|\mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\cdot)\|_{\mathcal{L}_x^{\mathbf{p}(\cdot)}},$$

and

$$\left\| \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right\|_{\mathcal{L}_x^{\frac{3}{2\alpha-1}}(L_t^\infty)} \leq C \|\mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\cdot)\|_{\mathcal{L}_x^{\frac{3}{2\alpha-1}}}.$$

Thus, considering Proposition 2.2 we can write

$$\|\mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\cdot)\|_{\mathcal{L}_x^{\mathbf{p}(\cdot)}} \leq C \|\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty}\|_{\mathcal{L}_{\frac{3}{2(2\alpha-1)}, x}^{\mathbf{p}(\cdot)}} = \|\mathcal{F}\|_{\mathcal{L}_{\frac{3}{2(2\alpha-1)}, x}^{\mathbf{p}(\cdot)}(L_t^\infty)}. \quad (3.20)$$

On the other hand, since the Riesz potentials are bounded on the classical Lebesgue space  $L^{\frac{3}{2\alpha-1}}$  we obtain

$$\left\| \mathcal{J}_{2\alpha-1}(\|\mathcal{F}(\cdot, \cdot)\|_{L_t^\infty})(\cdot) \right\|_{L_x^{\frac{3}{2\alpha-1}}} \leq C \|\mathcal{F}\|_{L_t^\infty} \Big\|_{L_x^{\frac{3}{2(2\alpha-1)}}} = \|\mathcal{F}\|_{L_x^{\frac{3}{2(2\alpha-1)}}(L_t^\infty)}. \quad (3.21)$$

Then, gathering together the norm  $\|\cdot\|_{\mathcal{E}}$  given in (3.15) with the estimates (3.20) and (3.21), we get

$$\left\| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\mathcal{F})(s, \mathbf{x}) ds \right\|_{\mathcal{E}} \leq C \|\mathcal{F}\|_{\mathcal{L}^{\frac{\mathfrak{p}(\cdot)}{2}}_{\frac{3}{2(2\alpha-1)}, \mathbf{x}}(L_t^\infty)} < +\infty.$$

With this we conclude the proof.  $\square$

**Proposition 3.6** *Consider  $\alpha \in ]\frac{1}{2}, 1]$  and a variable exponent  $\mathfrak{p}(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  such that  $\mathfrak{p}^- > 1$ . Then, there exist a constant  $C_{\mathcal{B}} > 0$  such that*

$$\left\| \int_0^t \mathfrak{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}))(\cdot, \cdot) ds \right\|_{\mathcal{E}} \leq C_{\mathcal{B}} \|\vec{\mathbf{u}}\|_{\mathcal{E}} \|\vec{\mathbf{u}}\|_{\mathcal{E}}. \quad (3.22)$$

**Proof:** We begin by noticing that, due to the properties of the Leray projector, we can write

$$\int_0^t \mathfrak{g}_{t-s}^\alpha * \mathbb{P}(\operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}})) ds = \int_0^t \mathbb{P}(\mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}})) ds = \mathbb{P} \left( \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}) ds \right).$$

Thus, by the Minkowski's integral inequality and the decay properties of the fractional heat kernel (see Remark 2.4), we obtain

$$\begin{aligned} \left| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}) ds \right| &\leq C \int_0^t \int_{\mathbb{R}^3} |\vec{\nabla} \mathfrak{g}_{t-s}^\alpha(\mathbf{x} - \mathbf{y})| |\vec{\mathbf{u}}(s, \mathbf{y})| |\vec{\mathbf{u}}(s, \mathbf{y})| d\mathbf{y} ds \\ &\leq C \int_{\mathbb{R}^3} \int_0^t \frac{1}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} |\vec{\mathbf{u}}(s, \mathbf{y})| |\vec{\mathbf{u}}(s, \mathbf{y})| ds d\mathbf{y}. \end{aligned}$$

Then, considering the  $L_t^\infty$ -norm we can write

$$\left| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}) ds \right| \leq C \int_{\mathbb{R}^3} \left( \int_0^t \frac{1}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} ds \right) \|\vec{\mathbf{u}}(\cdot, \mathbf{y})\|_{L_t^\infty} \|\vec{\mathbf{u}}(\cdot, \mathbf{y})\|_{L_t^\infty} d\mathbf{y}.$$

Now, remark that similarly to in the proof of Proposition 3.5, we have

$$\int_0^t \frac{ds}{(|t-s|^{\frac{1}{2\alpha}} + |\mathbf{x} - \mathbf{y}|)^4} = C \frac{1}{|\mathbf{x} - \mathbf{y}|^{4-2\alpha}} \quad (3.23)$$

and then, we conclude the estimate

$$\left| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}}) ds \right| \leq C \int_{\mathbb{R}^3} \frac{1}{|\mathbf{x} - \mathbf{y}|^{4-2\alpha}} \|\vec{\mathbf{u}}(\cdot, \mathbf{y})\|_{L_t^\infty} \|\vec{\mathbf{u}}(\cdot, \mathbf{y})\|_{L_t^\infty} d\mathbf{y}. \quad (3.24)$$

Thus, by considering Definition 2.4, the last inequality can be recast as

$$\left| \int_0^t \mathfrak{g}_{t-s}^\alpha * \operatorname{div}(\vec{\mathbf{u}} \otimes \vec{\mathbf{u}})(s, \mathbf{x}) ds \right| \leq C \mathcal{J}_{2\alpha-1}(\|\vec{\mathbf{u}}\|_{L_t^\infty} \|\vec{\mathbf{u}}\|_{L_t^\infty})(\mathbf{x}),$$

from where get the estimate

$$\mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}})(s, \mathbf{x}) ds \right) \leq C \mathbb{P} \left( \mathcal{J}_{2\alpha-1}(\|\bar{\mathbf{u}}\|_{L_t^\infty} \|\bar{\mathbf{u}}\|_{L_t^\infty})(\mathbf{x}) \right).$$

Now, to obtain the  $\mathcal{L}^{\mathbf{p}(\cdot)}_{\frac{3}{2\alpha-1}}$ -norm given in (3.15) note that the Riesz transforms are bounded in the spaces  $L^{\frac{3}{2\alpha-1}}$  and  $L^{\mathbf{p}(\cdot)}$  (recall that we have supposed that the variable exponent  $\mathbf{p}(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^3)$  with  $\mathbf{p}^- > 1$ ). Thus, the Leray projector  $\mathbb{P}$  is bounded in such spaces and we obtain the estimates

$$\left\| \mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}}) ds \right) \right\|_{L_x^{\mathbf{p}(\cdot)}(L_t^\infty)} \leq C \|\mathcal{J}_{2\alpha-1}(\|\bar{\mathbf{u}}\|_{L_t^\infty} \|\bar{\mathbf{u}}\|_{L_t^\infty})\|_{L_x^{\mathbf{p}(\cdot)}(L_t^\infty)},$$

and

$$\left\| \mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}}) ds \right) \right\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)} \leq C \|\mathcal{J}_{2\alpha-1}(\|\bar{\mathbf{u}}\|_{L_t^\infty} \|\bar{\mathbf{u}}\|_{L_t^\infty})\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)}.$$

Now, considering that the Riesz potential  $\mathcal{J}_{2\alpha-1}$  satisfies  $\|\mathcal{J}_{2\alpha-1}(\varphi)\|_{L^{\frac{3}{2\alpha-1}}} \leq C \|\varphi\|_{L^{\frac{3}{2\alpha-1}}}$ , an Hölder inequality (see Remark 2.3) and Proposition 2.2, we get the estimates

$$\left\| \mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}}) ds \right) \right\|_{L_x^{\mathbf{p}(\cdot)}(L_t^\infty)} \leq C \|\|\bar{\mathbf{u}}\|_{L_t^\infty} \|\bar{\mathbf{u}}\|_{L_t^\infty}\|_{\mathcal{L}^{\frac{\mathbf{p}(\cdot)}{2}}_{\frac{3}{2(2\alpha-1)}}} \leq C \|\bar{\mathbf{u}}\|_{\mathcal{L}^{\frac{\mathbf{p}(\cdot)}{2\alpha-1}, x}(L_x^\infty)} \|\bar{\mathbf{u}}\|_{\mathcal{L}^{\frac{\mathbf{p}(\cdot)}{2\alpha-1}, x}(L_x^\infty)},$$

and

$$\left\| \mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}}) ds \right) \right\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)} \leq C \|\|\bar{\mathbf{u}}\|_{L_t^\infty} \|\bar{\mathbf{u}}\|_{L_t^\infty}\|_{L_x^{\frac{3}{2(2\alpha-1)}}} \leq C \|\bar{\mathbf{u}}\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)} \|\bar{\mathbf{u}}\|_{L_x^{\frac{3}{2\alpha-1}}(L_t^\infty)}.$$

**Remark 3.4** Note that, in the case that we had considered Theorem 4 instead of Proposition 2.2, we had obtained an estimate of the form  $\|\mathcal{J}_{2\alpha-1}(\varphi)\|_{L^{\mathbf{p}(\cdot)}} \leq \|\varphi\|_{L^{\frac{\mathbf{p}(\cdot)}{2}}}$ , which, due the strong relationship between the the variable exponents involved, yields the constant exponent  $\mathbf{p}(\cdot) \equiv \frac{3}{2\alpha-1}$ .

Gathering together these last estimates, and using the definition of the norm  $\|\cdot\|_{\mathcal{E}}$  given in (3.15) we obtain

$$\left\| \mathbb{P} \left( \int_0^t \mathbf{g}_{t-s}^\alpha * \operatorname{div}(\bar{\mathbf{u}} \otimes \bar{\mathbf{u}}) ds \right) \right\|_{\mathcal{E}} \leq C_{\mathcal{B}} \|\bar{\mathbf{u}}\|_{\mathcal{E}} \|\bar{\mathbf{u}}\|_{\mathcal{E}}.$$

Thus Proposition 3.3 is proven.  $\square$

## End of the proof of Theorem 2

Considering the hypothesis assumed and the estimates obtained in Propositions 3.4, 3.5 and 3.6, we get the condition

$$C \left( \|\bar{\mathbf{u}}_0\|_{\mathcal{L}^{\frac{\mathbf{p}(\cdot)}{2\alpha-1}}} + \|\mathcal{F}\|_{\mathcal{L}^{\frac{\mathbf{p}(\cdot)}{2}}_{\frac{3}{2(2\alpha-1)}, x}(L_t^\infty)} \right) < \frac{1}{4C_{\mathcal{B}}}.$$

Thus, by applying the Banach-Picard principle stated in Theorem 5, we obtain the desired solution. With this we conclude the proof of Theorem 2.



**Acknowledgements.** The author warmly thanks Pierre-Gilles Lemarié-Rieusset and Diego Chamorro for their helpful comments and advises. The author is supported by the ANID postdoctoral program BCH 2022 grant No. 74220003.

## References

- [1] M. CANNONE, *Ondelettes, paraproducts et Navier-Stokes*, Diderot Editeur, Paris, 1995. With a preface by Yves Meyer.
- [2] M. CANNONE AND G. WU, *Global well-posedness for Navier-Stokes equations in critical Fourier-Herz spaces*, *Nonlinear Anal.*, 75 (2012), pp. 3754–3760.
- [3] D. CHAMORRO, *Mixed Sobolev-like inequalities in Lebesgue spaces of variable exponents and in Orlicz spaces*, *Positivity*, 26 (2022), pp. Paper No. 5, 21.
- [4] D. V. CRUZ-URIBE AND A. FIORENZA, *Variable Lebesgue spaces: Foundations and harmonic analysis*, Springer Science & Business Media, 2013.
- [5] L. DIENING, P. HARJULEHTO, P. HÄSTÖ, AND M. RUZICKA, *Lebesgue and Sobolev spaces with variable exponents*, Springer, 2011.
- [6] H. FUJITA AND T. KATO, *On the Navier-Stokes initial value problem. I*, *Arch. Rational Mech. Anal.*, 16 (1964), pp. 269–315.
- [7] Y. GIGA AND T. MIYAKAWA, *Navier-Stokes flow in  $\mathbf{R}^3$  with measures as initial vorticity and Morrey spaces*, *Comm. Partial Differential Equations*, 14 (1989), pp. 577–618.
- [8] L. GRAFAKOS, *Classical fourier analysis*, vol. 2, Springer, 2008.
- [9] A. KALTENBACH, *Pseudo-monotone operator theory for unsteady problems with variable exponents*, vol. 2329 of *Lecture Notes in Mathematics*, Springer, Cham, [2023] ©2023.
- [10] T. KATO, *Strong solutions of the Navier-Stokes equation in Morrey spaces*, *Bol. Soc. Brasil. Mat. (N.S.)*, 22 (1992), pp. 127–155.
- [11] T. KATO AND H. FUJITA, *On the nonstationary Navier-Stokes system*, *Rend. Sem. Mat. Univ. Padova*, 32 (1962), pp. 243–260.
- [12] H. KOCH AND D. TATARU, *Well-posedness for the Navier-Stokes equations*, *Adv. Math.*, 157 (2001), pp. 22–35.
- [13] P. G. LEMARIÉ-RIEUSSET, *Recent developments in the Navier-Stokes problem*, vol. 431 of *Chapman & Hall/CRC Research Notes in Mathematics*, Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [14] P. G. LEMARIÉ-RIEUSSET, *The Navier-Stokes problem in the 21st century*, CRC Press, Boca Raton, FL, 2016.
- [15] P. LI AND Z. ZHAI, *Generalized Navier-Stokes equations with initial data in local Q-type spaces*, *J. Math. Anal. Appl.*, 369 (2010), pp. 595–609.

- [16] ———, *Well-posedness and regularity of generalized Navier-Stokes equations in some critical Q-spaces*, J. Funct. Anal., 259 (2010), pp. 2457–2519.
- [17] J.-L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris; Gauthier-Villars, Paris, 1969.
- [18] C. MIAO, B. YUAN, AND B. ZHANG, *Well-posedness of the Cauchy problem for the fractional power dissipative equations*, Nonlinear Analysis: Theory, Methods & Applications, 68 (2008), pp. 461–484.
- [19] J. WU, *Generalized MHD equations*, J. Differential Equations, 195 (2003), pp. 284–312.
- [20] ———, *The generalized incompressible Navier-Stokes equations in Besov spaces*, Dynamics of Partial Differential Equations, 1 (2004), pp. 381–400.
- [21] Q. YANG AND P. LI, *Regular wavelets, heat semigroup and application to the magneto-hydrodynamic equations with data in critical Triebel-Lizorkin type oscillation spaces*, Taiwanese J. Math., 20 (2016), pp. 1335–1376.
- [22] X. YU AND Z. ZHAI, *Well-posedness for fractional Navier–Stokes equations in the largest critical spaces*, Mathematical Methods in the Applied Sciences, 35 (2012), pp. 676–683.
- [23] Z. ZHAI, *Well-posedness for fractional Navier-Stokes equations in critical spaces close to  $\dot{B}_{\infty,\infty}^{-(2\beta-1)}(\mathbb{R}^n)$* , Dyn. Partial Differ. Equ., 7 (2010), pp. 25–44.