



HAL
open science

Importance and roles of arbuscular mycorrhizal fungi in New Caledonian ultramafic soils

Hamid Amir, Yvon Cavaloc, Thomas Crossay, Alexandre Bourles, Simon Gensous, Alexandre Lagrange, Valérie Burtet-Sarramegna, Linda Guentas

► **To cite this version:**

Hamid Amir, Yvon Cavaloc, Thomas Crossay, Alexandre Bourles, Simon Gensous, et al.. Importance and roles of arbuscular mycorrhizal fungi in New Caledonian ultramafic soils. *Botany Letters*, 2022, 170, pp.449 - 458. 10.1080/23818107.2022.2160808 . hal-04448048

HAL Id: hal-04448048

<https://hal.science/hal-04448048>

Submitted on 5 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Importance and roles of arbuscular mycorrhizal fungi in New Caledonian**
2 **ultramafic soils**

3

4 Hamid Amir, Yvon Cavaloc, Thomas Crossay, Alexandre Bourles, Simon Gensous, Alexandre
5 Lagrange, Valérie Burtet-Sarramegna, and Linda Guentas

6

7 Institut des Sciences Exactes et Appliquées (ISEA), Université de la Nouvelle-Calédonie, Noumea
8 98851, New Caledonia

9

10 Contact Hamid Amir : hamid.amir@unc.nc; <https://orcid.org/0000-0001-9777-4050>

11

12

13 **ABSTRACT**

14

15 Our knowledge about New Caledonian serpentine ecosystems has increased greatly during the past
16 half-century, mainly thanks to Jaffré's group. However, research on soil microflora and plant
17 symbionts started only in the nineties and was mainly published during the last two decades. We
18 aimed to synthesize these studies, focusing particularly on arbuscular mycorrhizal fungi (AMF).
19 Research on AMF consisted firstly of a global and inventory approach aiming to produce basic but
20 essential lacking knowledge. These studies showed that AMF are abundant in ultramafic soils and
21 concerned nearly all plant species of these ecosystems. Even Ni-hyperaccumulator plants and
22 sedges, generally considered non-mycorrhizal, were found to be functionally colonized by AMF in
23 New Caledonian ultramafic soils. The adaptation of AMF communities to the extreme conditions of
24 these soils led to high levels of metal tolerance (particularly to Ni) and noticeable originality of the
25 taxa. The influence of these symbionts on plant growth and adaptation was assessed in greenhouse
26 and field conditions. An accurate selection of AMF isolates that improve plant growth, and plant
27 metal tolerance was performed. It was demonstrated that combinations of AMF isolates with
28 complementary functional traits showed high synergistic effects on plant development. Finally, a
29 partnership with a biotechnological company led to the production of an efficient commercial
30 inoculant now used in the ecological restoration of mine-degraded areas. Today studies are focused
31 mainly on the additive effects of AMF and mycorrhiza-helper bacteria.

32

33 **KEYWORDS**

34 AMF; bacteria; serpentine ecosystems; abiotic stress; toxic metals; restoration ecology; New
35 Caledonia

36

37 **Introduction**

38

39 Arbuscular mycorrhiza is among the most frequent plant-microbe mutualistic symbiosis, established
40 with 80% of vascular plants (Wang and Qiu 2006; Smith and Read 2008; Strullu-Derrien et al.
41 2017). This symbiosis is also one of the most ancient, initiated probably when plants have colonized
42 terrestrial areas more than 400 million years ago (Karatygin et al. 2006; Humphreys et al. 2010;
43 Strullu-Derrien et al. 2014). Arbuscular mycorrhizal fungi (AMF) act fundamentally on different
44 mechanisms of plant development, including mineral nutrition, drought resistance and toxic metal
45 tolerance (Smith and Smith 2011; Amir et al. 2014; Augé et al. 2015; Ferrol et al. 2016). Their use
46 in environmental remediation and ecological restoration is then particularly appropriate (Göhre and
47 Paszkowski 2006; Amir et al. 2019; Valliere et al. 2020). Moreover, these targeted environmental
48 applications are among the most advanced for large-scale mycorrhizal inoculation practices today
49 (Hart et al. 2017).

50 Serpentine ecosystems developed in ultramafic soils are generally characterized by extreme
51 conditions: macronutrient deficiency, unbalanced Ca/Mg ratio, metal toxicity and frequent drought
52 stress. These conditions have contributed to promoting original flora (Brooks 1987; Proctor 2003;
53 Jaffré and L'Huillier 2010; Isnard et al 2016). It is particularly the case in New Caledonia, a Pacific
54 island biodiversity hot spot (Myers et al. 2000, Marchese 2015), characterized by endemism of 96.7
55 % of the ultramafic obligate flora (Isnard et al. 2016). This precious biodiversity is threatened by
56 mining activities, fires, and other anthropogenic influences (Jaffré et al. 1998; L'Huillier et al.
57 2010).

58 The origin and pedological classification of new Caledonian ultramafic soils and their
59 physicochemical features have been subject of many studies from the seventies (Jaffré et al. 1971;
60 Jaffré and Latham 1974; Latham 1975; Latham et al. 1978; Jaffré and L'Huillier 2010). Metal
61 concentrations and their bioavailability in these soils were also investigated in relation to their
62 potential toxicity on plants (L'Huillier and Edighoffer 1996, Quantin et al. 1997; L'Huillier et al
63 1998). However, their microbiology has not been investigated until the nineties, and this was also
64 true for other countries, with rare exceptions (Lebedeva et al. 1978). In particular, AMF survival
65 and activity, and their relationships with plants in these extreme conditions, with very low

66 phosphorus availability (Dubus and Becquer 2001; Jaffré and L’Huillier 2010) and high potentially
67 toxic metal concentrations constituted an original unexplored research subject.

68 The first study on mycorrhizal associations in ultramafic soils worldwide concerned AMF
69 spore densities and mycorrhizal colonization of 13 endemic plant species in different New
70 Caledonian sites (Amir et al. 1997). From the year 2000, several publications showed the high
71 interest of these extreme soils to investigate the role of mycorrhizal symbiosis in the adaptation to
72 edaphic and environmental stresses of plants (Boulet et al. 2005; Jourand et al. 2010a; Lagrange et
73 al. 2011; Orłowska et al. 2011; Amir et al. 2013). The adaptive traits of fungal symbionts were also
74 highlighted (Turnau and Mesjasz-Przybyłowics 2003; Amir et al. 2008; Jourand et al. 2010b;
75 Branco and Ree 2010; Majorel et al. 2012).

76 Studies on ectomycorrhizas (EM) in New Caledonian ultramafic ecosystems were not detailed
77 here. EM are important particularly in forests (Carriconde et al. 2019) and in maquis dominated by
78 *Tristaniopsis* and *Sannantha* (Myrtaceae) species or *Acacia spirorbis* (Fabaceae) (Pillon et al.
79 2021). They are present on trees belonging to Myrtaceae, Fabaceae, Casuarinaceae and
80 Nothofagaceae families (Perrier 2005; Amir and Ducouso 2010; Carriconde et al. 2019). A recent
81 list of EM plants was reported in Pillon et al. (2021). Their role is then complementary with AMF,
82 root system of some plants (*Tristaniopsis* and *Nothofagus* (Nothofagaceae) species for example)
83 being simultaneously colonized by the two symbiont groups (Perrier et al. 2006). EM have also a
84 good potential for their use in ecological restoration of mined areas (Houlès et al. 2022).

85 In New Caledonia, studies on AMF were taken over mainly by the University of New
86 Caledonia. This review synthesizes those focused on ultramafic soils, particularly the ecological
87 aspects and their practical consequences for the ecological restoration of mine-degraded areas. Are
88 these symbionts able to maintain their spores and colonize root systems in these extreme soils, and
89 how AMF diversity is shaped by these conditions? Are the most frequent plants in ultramafic
90 maquis, particularly those used in restoration programs, arbuscular mycorrhizal? What are the main
91 functions of AMF in plant nutrition, and how could they participate to plant adaptation to highly
92 stressed conditions? These questions are relevant not only for science, but also for their potential
93 use in ecological restoration of mined areas.

94

95 **Exploration and inventory studies**

96

97 When research on ultramafic soil microbiology started in New Caledonia, the abiotic
98 characterization of these substrates in relation to ecosystem functions was relatively advanced (Jaffré

99 et al. 1971; Jaffré and Latham 1974; Lee et al. 1977; Becquer et al. 1995). Microbiological studies
100 were then focused first on a global approach that can help to explain some ecological specificities of
101 the studied ecosystems. Quantitative and qualitative analyses of soil microbiota and its activity in
102 relation with plants and soil metal concentrations were assessed (Amir and Pineau 1998; Saintpierre
103 2001; Amir and Pineau 2003a, 2003b). These publications showed different specificities of
104 ultramafic soil microbiology, with low level of activity but high metal tolerance of the microbiota,
105 part of which being responsible of the release of different metals, including Ni and Co. *Frankia*
106 symbiosis with Casuarinaceae is frequent in New Caledonian ultramafic ecosystems (Navarro et al.
107 1999; Gauthier et al. 1999) and may play an important role in nitrogen enrichment of these soils. A
108 global approach on plant growth promoting rhizobacteria was initiated (Mercky 1999). More
109 recently, it was shown that ultramafic soil properties do not limit their microbial diversity (Bordez et
110 al. 2016; Carriconde et al. 2019). However, this rich diversity showed wide variations in relation to
111 environmental conditions (Bordez et al. 2016; Gourmelon et al. 2016). The vegetation cover,
112 particularly its density, seemed to play a major role. Among soil properties, pH, nitrogen and
113 calcium influenced the distribution patterns of microbial communities.


114 After exploratory works on mycorrhizas (Amir et Pineau 1997; Perrier 2005; Perrier et al.
115 2006), it was clear that AMF are abundant in New Caledonian ultramafic soils. Most of the plant
116 species tested were found to be arbuscular mycorrhizal, and AMF spore densities were globally
117 similar to those of non-ultramafic soils (Perrier et al. 2006; Amir and Ducouso 2010). If AMF root
118 colonization does not generally induce changes in root morphology, Duhoux et al. (2001) showed
119 that *Gymnostoma* trees (Casuarinaceae) produce small root nodules called myconodules, colonized
120 by AMF related to *Glomus*. Similar small nodules rich in AMF structures were found in roots of
121 *Podocarpus* trees (Podocarpaceae) in New Zealand (Russel et al. 2002), and were confirmed to be
122 present with the same characteristics in *Araucaria* trees in New Caledonia (Amir, unpublished
123 data). A correlation between AMF spore abundance and soil global activity was detected (Perrier et
124 al. 2006) probably due to negative effect on AMF development of high phosphorus deficiency of
125 the poorest soils (Lagrange et al. 2013; Amir et al. 2021). A list of over 60-studied plant species,
126 particularly those used in ecological restoration, presenting their affinity with AMF or
127 ectomycorrhizal fungi was then published (Amir and Ducouso 2010).

128 **Adaptation of arbuscular mycorrhizal symbiosis to ultramafic soils**

129

130 In the past, Ni-hyperaccumulating plants were generally assumed to be non-mycorrhizal (Leyval et
131 al. 1997; Pawlowska et al. 2000). Amir et al. (2007) tested nine Ni-accumulating New Caledonian

132 plants ranging from weak accumulators to hypernickelophore species. All species were more or less
 133 mycorrhized, but mycorrhizal colonization was reduced by high levels of Ni-hyperaccumulation
 134 (Figure 1). Indeed, for the same species, the level of mycorrhizal colonization was negatively
 135 correlated with Ni-concentration in plant leaves. The main reason for classically considering Ni-
 136 hyperaccumulators as non-mycorrhizal was that the plant species generally studied for this property
 137 in Europe and USA belonged to non-mycorrhizal plant families (Brassicaceae, Caryophyllaceae).



138

139 **Figure 1.** Relationships between leaf Ni content and mycorrhizal intensity (M) of 9 Ni-
 140 accumulating plant species (means of 8-10 samples for each value); *G* = *Geissos*. The
 141 coefficient r (a-b) indicates the total linear correlation between the two parameters. From
 142 Amir et al. (2007), reprinted with the permission of Plant and Soil.

143

144 Some studies on metal-polluted soils have revealed the high tolerance to Zn and Cd of AMF
 145 isolated from this environment (Weissenhorn et al. 1994; Tullio et al. 2003). When tested for Ni
 146 tolerance, AMF from ultramafic soils were more tolerant than those from other origins (Amir et al.

147 2008). Moreover, isolates from roots of Ni-hyperaccumulating plants tolerated the highest Ni
148 concentrations (Amir et al. 2007; 2008). This last study also reported that Ni-tolerance of AMF
149 isolates could be inducted by the presence of this metal in the substrate solution during host plant
150 growth.

151 In New Caledonian ultramafic areas, the herbaceous stratum is mainly represented by
152 Cyperaceae (sedges), a family generally considered non-mycorrhizal (Brundrett, 2009). However,
153 several endemic Cyperaceae species highly adapted to ultramafic soils were colonized at low levels
154 by AMF (Perrier et al. 2006; Lagrange et al. 2011). Tested in greenhouse conditions and field
155 experiment, *Tetraria comosa*, one of the most frequent and pioneer sedge species in New
156 Caledonian maquis, was shown to be functionally mycorrhizal, even at low levels of root
157 colonization, with positive effects on plant survival rate and plant growth (Lagrange et al. 2011;
158 2013).

159

160 **Importance and role of AMF on plant adaptation to ultramafic soils**

161

162 AMF are obligate symbionts so their cultivation needs a host plant. Isolates from New Caledonian
163 ultramafic soils were obtained in a greenhouse in association with sorghum. This grass was chosen
164 not only because it has a good affinity with AMF, but also because it can grow in difficult
165 conditions with a relatively high rate, contrary to native species that grow slowly. The effects of
166 AMF isolates on plant development have then been tested with sorghum and two endemic shrubs,
167 *Alphitonia neocaledonica* (Rhamnaceae) and *Cloezia artensis* (Myrtaceae), currently used in
168 restoration programs (Amir et al. 2013). The study showed that AMF not only stimulated the
169 growth and phosphorus nutrition of these plants, but also reduced nickel absorption and its toxicity
170 (Figure 2). Similar but less important effects were obtained on *T. comosa* (Lagrange et al. 2011).




171

172 Figure 2. Aspect of *Alphitonia neocaledonica* plants in pots treated or not with nickel and
 173 inoculated or not with SFONL arbuscular mycorrhizal fungus isolate, showing Ni-toxicity
 174 symptoms. *A. neocaledonica* plants without Ni are not shown because they have the same
 175 aspect than SFONL 30 $\mu\text{g g}^{-1}$ Ni. NM: non mycorrhizal plants. From Amir et al. (2013),
 176 reprinted with the permission of Mycorrhiza.
 177

178 Ultramafic soils are often highly deficient in phosphorus (P), this element being moreover
 179 strongly adsorbed to iron oxides (Dubus and Becquer 2001; Jaffré and L'Huillier 2010). It is well
 180 known that an optimal level of available P in soil generally reduces AMF root colonization
 181 (Marschner 1995; Smith and Read 2008), but the effects of high P deficiency has received less
 182 attention (Abbott et al. 1984; Lambers et al. 2009). Lagrange et al. (2011) suggested that P
 183 deficiency could explain low levels of AMF root colonization in New Caledonian ultramafic soils.
 184 A field experiment with *T. comosa* in an ultramafic area confirmed this hypothesis: it showed that
 185 AMF root colonization and arbuscules' density of non-inoculated plants were clearly enhanced by
 186 phosphorus supply, with correlatively a better mineral nutrition (Lagrange et al. 2013).

187 In the latter cited study, plants were not inoculated with AMF. More recently, a field
 188 experiment with *Metrosideros laurifolia* (Myrtaceae) plants inoculated in greenhouse with a mix of
 189 three AMF isolates was assessed in an ultramafic area in Goro (south of New Caledonia). Stored
 190 topsoil spread in this area a few years before, and characterized by a low density of AMF spores,
 191 was amended or not with sewage sludge and planted with *M. laurifolia* plantlets (Amir et al. 2019).
 192 After 528 days, the aerial dry biomass of AMF-inoculated plants was four times greater than the

193 control ones (Figure 3). Mycorrhizal colonization was increased by sewage sludge inducing a
 194 complementary effect with AMF inoculation. This resulted in better plant growth and mineral
 195 nutrition, with plant biomass 7.7 times greater than control ones. Ca/Mg ratio was also enhanced by
 196 AMF inoculation, and the translocation to aerial parts of potentially toxic metals (particularly Mn,
 197 Ni, and Co) was strongly reduced. The analysis of AMF diversity in *M. laurifolia* roots showed a
 198 higher number of operational taxonomic units (OTU) in AMF-inoculated plants grown on non-
 199 amended plots compared to non-inoculated control. However, the introduced AMF isolates
 200 constituted at most, only 9% of the detected OTUs. These results were explained by the strong
 201 increase in plant biomass of AMF-inoculated plants. This increase allowed more carbon allocation
 202 to AMF in roots, the competition between AMF communities, then inducing a relative reduction of
 203 inoculated isolates. However, the relative decrease of the introduced isolates did not weaken the
 204 positive effects of mycorrhizal inoculation on plant nutrition and growth, even after 3.5 years (Amir
 205 et al. 2019). This latter point is interesting for AMF biotechnologies, knowing that inoculum
 206 reduction after a few months to a few years is one of the drawbacks of this practice (Pellegrino et al.
 207 2012; Janoušková et al. 2013).



208

209 Figure 3. Influence of AMF inoculation, sewage sludge and their combination on growth
 210 (plant dry weight of aerial parts) and mycorrhizal colonization (M%) of *Metrosideros*
 211 *laurifolia* plants, 528 days after plantation (mean values with standard error). Bars with the
 212 same letters are non-significantly different (post-hoc Bonferroni corrected Mann-Whitney
 213 test). From Amir et al. (2019), reprinted with the permission of Science of the Total
 214 Environment.

215

216 Studies on AMF diversity of New Caledonian soils, quite recently initiated, revealed the
 217 originality and high richness of AMF communities of ultramafic soils (Gensous 2014; Amir et al.
 218 2015; Cavaloc and Amir 2019). A comparison between AMF communities associated with roots of
 219 *Psychotria* (Rubiaceae) and *Phyllanthus* (Phyllanthaceae) species from tropical rain forests

220 developed on ultramafic and volcano-sedimentary soils, in four regions, was performed (Gensous
221 2014). AMF diversity richness did not differ significantly between the two types of forests, but
222 community composition was different. Ultramafic plants were associated mostly with the genera
223 *Rhizophagus*, *Glomus*, *Diversispora* and *Archaeospora*, whereas plants from volcano-sedimentary
224 soils were associated more frequently with *Acaulospora*, *Claroideoglomus*, and an unknown
225 Glomeraceae. Moreover, in ultramafic forest, AMF community composition differed significantly
226 between Ni-hyperaccumulating and non-accumulating plants. Cavaloc and Amir (2019) analyzed
227 the diversity of AMF in soils of 4 ultramafic open, dense and bushy maquis in south New
228 Caledonia. They reported total number of 188 Amplicon Sequence Variants (ASVs), varying from
229 72 to 104 ASVs per type of maquis, increasing with the vegetation density. Each analyzed plot
230 contained about 10 to 30 ASVs. Shannon-Wiener AMF diversity index varied from 1.5 to 3.5. Amir
231 et al. (2015) estimated AMF diversity inside roots of 4 endemic plant species collected from a dense
232 maquis in Goro (south New Caledonia). They detected 427 sequences corresponding to 54
233 Operational Taxonomic Units (OTUs), varying from 13 to 20 per plant species. These AMF
234 communities were dominated by Glomeraceae family, the genera *Rhizophagus* and *Glomus*
235 generally constituting more than 60% of the sequences (Gensous 2014; Amir et al. 2019; Cavaloc
236 and Amir 2019). The other most detected genera were *Sclerocystis*, *Acaulospora*, *Scutellospora*,
237 *Claroideoglomus*, *Diversispora*, *Redeckera*, and *Paraglomus*. Five AMF new species were
238 published (Blaszkowski et al. 2017; Crossay et al. 2018).

239 This diversity has also been the object of functional experiments. With a larger collection of
240 fungal isolates, greenhouse studies on the effects of different AMF mixes using the endemic plant
241 *M. laurifolia* grown on ultramafic soil were assessed, assuming synergistic effects would be found.
242 Indeed, combinations of co-inoculated isolates belonging to different families and having different
243 functional traits were more efficient than single ones (Crossay et al. 2019). The best mix improved
244 not only mineral nutrition with better absorption of major elements, particularly P and K, compared
245 to single isolates, but also increased Ca/Mg ratio and reduced Ni and Co translocation more clearly.
246 These effects resulted in a great growth improvement, with an aerial dry biomass 21.6 times bigger
247 than the non-inoculated plants. In contrast, the best single isolate induced a 12-fold increase of this
248 biomass. These results comfort the hypothesis stating that synergistic effects of AMF on plant
249 development are optimal when the phylogenetic distance between the mixed isolates is maximized
250 (Maherali and Klironomos, 2007), thus complementing biological niches and reducing functional
251 redundancy. Mathematical modeling of these AMF synergistic effects was produced (Crossay et al.
252 2019) explaining the experimental results with a high concordance ($R^2 = 0.90$, $p < 0.00001$).

253

254 **AMF and mycorrhiza helper bacteria**

255

256 Studies on bacteria and their communities in New Caledonian ultramafic soils have received less
257 attention than fungal symbionts (Héry et al. 2003; Herrera et al. 2007; Bordez et al. 2016;
258 Gourmelon et al. 2016). A few articles reported some of their growth promoting effects on plants
259 (Mercky 1999; Gonin et al. 2013; Bourles et al. 2019; Bourles et al. 2020a). Complementary effects
260 between bacteria and AMF have been addressed only recently (Bourles et al. 2020b). The latter
261 studied the strain BE isolated from the rhizosphere of *Tetraria* plants and screened by Gonin et al.
262 (2013) for its metal tolerance and plant growth promotion properties : phosphorus solubilization,
263 and production of indole acetic acid (IAA), siderophores, ammonium and 1-aminocyclopropane-1-
264 carboxylate deaminase (ACC deaminase). This bacterium belonging to the species *Curtobacterium*
265 *citreum* was able to form biofilm and produce an exopolysaccharide inducing Ni adsorption and
266 inactivation capacities (Bourles 2019; Bourles et al. 2019). The strain reduced Ni, Co and Cr
267 absorption by sorghum plants (Bourles et al. 2019) and strongly enhanced mycorrhizal colonization
268 (Bourles et al. 2020b). The latter study reported complementary effects on the growth and nutrition
269 parameters of *T. comosa* plants when the bacterium was co-inoculated with AMF in greenhouse
270 conditions. A field experiment testing these effects is underway.

271

272 **AMF and ecological restoration of ultramafic degraded areas**

273

274 A part of AMF studies in New Caledonia concerned directly the ecological restoration of ultramafic
275 mine-degraded areas. It was shown that AMF communities of revegetated subsoils on mine-
276 degraded areas are different and functionally non-equivalent with AMF communities of non-
277 degraded related ecosystems (Cavaloc et al. unpublished data). AMF diversity seemed to adapt to
278 the new conditions of the restored ecosystem. The reuse of topsoil, generally not recuperated by
279 mine companies until the nineties, was strongly recommended (Amir et al. 2003; Bordez et al.
280 2018), but in practice, this substrate was generally stored for years before being spread for
281 revegetation. Studies showed that topsoil storage not only reduces seed bank, soil activity and
282 nutrients' reserve, but also AMF spore density and AMF diversity (Amir et al. 2015; Amir et al.
283 2022). It was then recommended to reuse the topsoil for the revegetation as soon as possible,
284 preferably not more than 6 months after its stripping (Bordez et al. 2018; Amir et al. 2022).

285 Considering the important role of AMF in plant development on ultramafic areas, a company
286 partner of the University of New Caledonia developed recently a commercial inoculum, containing
287 synergistic isolates, based on Crossay et al. (2019) results, after innovation of an aseptic method for
288 AMF culture. Today, this inoculum is used to enhance growth and adaptation of plants for the
289 revegetation of mined areas in ecological restoration perspective. It is also envisaged to develop
290 inoculants with isolates that are specific to particular lateritic soils to avoid reduction of native
291 AMF biodiversity.

292 On-going greenhouse and field experiments aim to adapt the inoculation technic to seeding
293 methods, by incorporating AMF inoculum into seed balls. These experiments also test the co-
294 inoculation of AMF and MHB. Three to ten seeds of different native species, depending on their
295 size, are incorporated with AMF powder inoculum into 8 mm balls. The balls made with a mix of
296 clay and organic matter are then sowed and followed-up for seed germination, plant growth,
297 mycorrhizal colonization and inoculum behavior.

298

299 **Conclusion**

300

301 Research on mycorrhizas' role in New Caledonian ultramafic soils during the last two decades
302 followed a progression dictated by the local context. The protection of the biodiversity of valuable
303 ecosystems geared the studies toward the ecological restoration perspective. Yet, the originality of
304 ultramafic environment and the absence of knowledge worldwide on AMF symbiosis of these soils
305 have made this subject particularly attractive to investigate plant adaptation in extreme
306 environments.

307 The abundance and importance of these symbionts in New Caledonian ultramafic ecosystems
308 were shown, and their diversity and relative originality in relation to the high plant endemism of
309 this environment was then highlighted. The ability of AMF to improve plant mineral nutrition and
310 metal tolerance in harsh conditions pointed out the central role of these symbionts in plant
311 adaptation to multi-stress environments. Greenhouse and field experiments then revealed practical
312 perspectives.

313 If an efficient AMF inoculant adapted to ultramafic mined areas is commercialized today, more
314 research on microbial ecology are needed to improve restoration technics in such extreme
315 conditions. New microbe consortia containing AMF and bacteria constitute particularly an
316 interesting target for deeper investigations.

317

318 Acknowledgments

319 The authors gratefully acknowledge for their contribution all of the research team, particularly,
320 Laurent Bordez, Valerie Médevielle, Audrey Laurent, Monika Lemestre, Véronique Gaudichon,
321 Clément Richert, Vincent Véron, Cyril Antheaume and all the trainees that worked with us from
322 1995 to 2020.

323

324 Author contributions

325 HA managed the research, designed the experiments and wrote the article; YC designed and
326 followed up biomolecular studies; TC, AB, AL and SG contributed as PhD students to significant
327 parts of the studies; TC worked as co-manager of a private company in partnership with the
328 research team for the commercial use of selected AMF; VBS and LG followed up and designed
329 biochemical and bacterial studies; all reviewed the article text.

330

331 Disclosure statement

332 No potential conflict of interest was reported by the authors.

333

334 Funding

335 The research synthesized here were supported by different financial funds, particularly, Agence
336 Nationale de la Recherche scientifique (ANR), CNRT Nickel et son Environnement, the South
337 Province of New Caledonia, Koniambo Nickel Society (KNS), and the University of New
338 Caledonia.

339

340 References

341

342 Abbott LK, AD Robson, G de Boer. 1984. The effect of phosphorus on the formation of hyphae in
343 soil by the vesicular-arbuscular mycorrhizal fungus, *Glomus fasciculatum*. *New Phytol.*
344 97:437–446.

345 Amir H, R Pineau, Z Violette. 1997. Premiers résultats sur les endomycorhizes des plantes de
346 maquis miniers de Nouvelle-Caledonie [first results on the endomycorrhizas of plants of New
347 Caledonian ultramafic areas]. In T Jaffré, RD Reeves, T Becquer, editors. *Ecologie des milieux*

- 348 sur roches ultramafiques et sur sols métallifères [The ecology of ultramafic and metalliferous
349 areas]. Noumea : ORSTOM ed.; p. 79–85. French.
- 350 Amir H, R Pineau. 1998. Influence of plants and cropping on microbiological characteristics of
351 some New Caledonian ultramafic soils. *Aust J Soil Res.* 361:457–471.
- 352 Amir H, N Barre, J Boulegue, F Colin, B Dreyfus, JF Marini, T Mennesson, B Pelletier, C Quantin,
353 R Pineau, J Tassin. 2003. Résumé et recommandations des travaux du colloque sur la Préservation
354 et restauration écologique en milieu tropical minier [Abstract and recommendations of the
355 International conference on Preservation and ecological restoration in tropical mining
356 environments]. Noumea, july 15-20, 2003. French.
- 357 Amir H, R Pineau. 2003a. Relationships between extractable Ni, Co and other metals and some
358 microbiological characteristics of different ultramafic soils from New Caledonia. *Aust J Soil
359 Res.* 41:215–228.
- 360 Amir H, R Pineau. 2003b. Release of Ni and Co by microbial activity in New Caledonian
361 ultramafic soils. *Can J Microbiol.* 49:288–293.
- 362 Amir H, N Perrier, F Rigault, T Jaffré. 2007. Relationships between Ni-hyperaccumulation and
363 mycorrhizal status of endemic plant species from New Caledonian ultramafic soils. *Plant Soil.*
364 293:23–35.
- 365 Amir H, DA Jasper, LK Abbott. 2008. Tolerance and induction of tolerance to Ni of arbuscular
366 mycorrhizal fungi from New Caledonian ultramafic soils. *Mycorrhiza.* 19:1–6.
- 367 Amir H, M Ducouso. 2010. Les bactéries et les champignons du sol sur roches ultramafiques
368 [bacteria and fungi of ultramafic soils]. In L’Huillier L, T Jaffré, A Wulff, editors. Mines et
369 environnement en Nouvelle-Calédonie: les milieux sur substrats ultramafiques et leur
370 restauration [Mines and environment in New Caledonia : the ultramafic areas and their
371 restoration]. Noumea: IAC Ed.; p. 129–145. French.
- 372 Amir H, A Lagrange, N Hassaine, Y Cavaloc. 2013. Arbuscular mycorrhizal fungi from New
373 Caledonian ultramafic soils improve tolerance to nickel of endemic plant species. *Mycorrhiza.*
374 23:585–595. [doi:10.1007/s00572-013-0499-6](https://doi.org/10.1007/s00572-013-0499-6).
- 375 Amir H, P Jourand, Y Cavaloc, M Ducouso. 2014. Role of mycorrhizal fungi on the alleviation of
376 heavy metal toxicity on plant. In Z Solaiman, L Abbott, A Varma, Editors. *Mycorrhizal Fungi:
377 Use in Sustainable Agriculture and Forestry.* Heidelberg: Springer; p. 241-258.
- 378 Amir H, L L’Huillier, B Fogliani, Y Cavaloc, P Jourand, M Ducouso, C Majorel, L Hannibal, D
379 Saintpierre, P Gunkel, P Pagand, G Echevarria, LC Mouchon, ML Bonis, E Montargues-

- 380 Pelletier, L Maggia, A Wulff. 2015. Caractérisation et fonctionnement du système sol/ plante/
381 microorganismes dans les maquis miniers. Perspectives d'application à la restauration
382 écologique [Characterization and functionality of the system soil/plant/microbes in ultramafic
383 maquis]. Noumea : CNRT Nickel et son environnement ; CNRT Ed. French. Available from :
384 <https://cnrt.nc/ecomine-biotop/>
- 385 Amir H, Y Cavaloc, A Laurent, P Pagand, P Gunkel, M Lemestre, V Medevielle, A Pain, S McCoy.
386 2019. Arbuscular mycorrhizal fungi and sewage sludge enhance growth and adaptation of
387 *Metrosideros laurifolia* on ultramafic soil in New Caledonia: A field experiment. *Sci Tot*
388 *Environ.* 651(1):334-343. doi:[10.1016/j.scitotenv.2018.09.153](https://doi.org/10.1016/j.scitotenv.2018.09.153).
- 389 Amir H, S Gensous, Y Cavaloc, L Wantiez. 2021. Phosphorus Fertilization of an Ultramafic Soil
390 Reduced Effects of Arbuscular Mycorrhizal Fungi but not Mycorrhizal Colonization. *J Soil Sci*
391 *and Plant Nutr.* 21(4):3544-3554.
- 392 Amir H, L Bordez, Y Cavaloc, P Jourand, M Ducouso, F Juillot. 2022. Effects of ultramafic
393 topsoil stockpiling during mine activities on its microbial diversity and other microbiological
394 and physicochemical characteristics. *Ecol Eng.* 177:106563.
- 395 Augé RM, HD Toler, AM Saxton. 2015. Arbuscular mycorrhizal symbiosis alters stomatal
396 conductance of host plants more under drought than under amply watered conditions: a meta-
397 analysis. *Mycorrhiza.* 25:13–24. doi:10.1007/s00572-014-0585.
- 398 Becquer T, E Bourdon, J Pétard. 1995. Disponibilité du nickel le long d'une toposéquence de sol
399 développé sur roches ultramafiques de Nouvelle-Calédonie [Nickel availability along soil
400 toposequence in New Caledonian ultramafic areas]. *CR Acad Sci.* 321:585-592. French.
- 401 Błaszowski J, A Kozłowska, T Crossay, S Symanczik, MN al-Yahya'ei. 2017. A new family,
402 *Pervetustaceae* with a new genus, *Pervetustus*, and *P. simplex* sp. nov. (Paraglomerales), and a
403 new genus, *Innospora* with *I. majewskii* comb. nov. (Paraglomeraceae) in the
404 Glomeromycotina. *Nova Hedwigia.* 105:397–410. doi:10.1127/nova_hedwigia/2017/0419.
- 405 Bordez L, P Jourand, M Ducouso, F Carriconde, Y Cavaloc, S Santini, JM Claverie, L Wantiez, A
406 Leveau, H Amir. 2016. Distribution patterns of microbial communities in ultramafic landscape:
407 a metagenetic approach highlights the strong relationships between diversity and environmental
408 traits. *Mol Ecol.* 25:2258–2272.
- 409 Bordez L, B Fogliani, L L'Huillier, H Amir. 2018. Guide technique de l'utilisation des topsoils en
410 restauration écologique des terrains miniers de Nouvelle-Calédonie [Technical guide on topsoil
411 use for ecological restoration of mining areas in New Caledonia]. Noumea: CNRT ed. French.

- 412 Available from: <https://cnrt.nc/uncategorized/guide-sur-lutilisation-des-topsoils-en->
413 [restauration-ecologique-des-terrains-miniers-lie-au-projet-recosynth-decembre-2018/](https://cnrt.nc/uncategorized/guide-sur-lutilisation-des-topsoils-en-restauration-ecologique-des-terrains-miniers-lie-au-projet-recosynth-decembre-2018/)
- 414 Boulet F, H Lambers. 2005. Characterisation of arbuscular mycorrhizal fungi colonization in cluster
415 roots of *Hakea verrucosa* F. Muell (Proteaceae), and its effect on growth and nutrient
416 acquisition in ultramafic soil. *Plant Soil*. 269:357–367.
- 417 Bourles A. 2019. Les Rhizobactéries des sols ultramafiques de Nouvelle-Calédonie. Caractérisation,
418 rôle dans l’adaptation des plantes à la contrainte édaphique et interaction avec les champignons
419 mycorrhiziens à arbuscules [Rhizobacteria of New Caledonian ultramafic soils.
420 Characterization, role in plant adaptation to edaphic constraints, and interaction with arbuscular
421 mycorrhizal fungi] [dissertation]. Noumea: University of New Caledonia. French.
- 422 Bourles A, L Guentas. S Gensous, E Chalkiadakis, C Majorel, F Juillot, Y Cavaloc, V Burtet-
423 Sarramegna, V Medevielle, P Jourand, H Amir. 2019. New Caledonian ultramafic conditions
424 structure the features of *Curtobacterium citreum* strains that play a role in plant adaptation. *Can*
425 *J Microbiol*. 65: 1-15.
- 426 Bourles A, H Amir, S Gensous, L Guentas, F Gussonneau, V Medevielle, F Juillot, A Basire, M
427 Meyer, V Burtet-Sarramegna, Y Cavaloc, P Jourand, L Guentas. 2019a. Investigating some
428 mechanisms underlying stress metal adaptations of two *Burkholderia* sensus lato species
429 isolated from New Caledonian ultramafic soils. *Eur J Soil Biol*. 97:103166.
- 430 Bourles A, L Guentas, C Charvis, S Gensous, C Majorel, T Crossay, Y Cavaloc, V Burtet-
431 Sarramegna, P Jourand, H Amir. 2020b. Co-inoculation with a bacterium and arbuscular
432 mycorrhizal fungi improves root colonization, plant mineral nutrition, and plant growth of a
433 Cyperaceae plant in an ultramafic soil. *Mycorrhiza*. 30:121–131. [doi:10.1007/s00572-019-
434 00929-8](https://doi.org/10.1007/s00572-019-00929-8).
- 435 Branco S, RH Ree. 2010. Serpentine soils do not limit mycorrhizal fungal diversity. *PLoS ONE*.
436 5(7):e11757. doi:10.1371/journal.pone.0011757.
- 437 Brooks RR. 1987. Serpentine and its vegetation: a multidisciplinary approach. Portland:
438 Dioscorides.
- 439 Brundrett MC. 2009. Mycorrhizal associations and other means of nutrition of vascular plants:
440 understanding the global diversity of host plants by resolving conflicting information and
441 developing reliable means of diagnosis. *Plant Soil*. 320:37–77.
- 442 Cavaloc Y, H Amir. 2019. Fonctionnalité des endomycorhizes. In Carriconde F, N Fernandez
443 Nunez, J Ripoll, A Liopold, B Fogliani, Y Itiaty, M Lelievre, K Letellier, L Maggia, G Bidau,

- 444 L’Huillier, H Amir, Y Cavaloc, M Ducouso, A Galiana, L Hannibal, P Jourand, M Lebrun.
 445 Identification d’indicateurs biologiques du fonctionnement des écosystèmes miniers restaurés.
 446 Acronyme : « Bioindic ». Rapport final de convention CNRT Nickel et son environnement.
 447 Noumea. French. Available from : <https://cnrt.nc/bioindic/>
- 448 Carriconde F, M Gardes, JM Bellanger, K Letellier, S Gigante, V Gourmelon, T Ibanez, S McCoy,
 449 J Goxe, J Read, L Maggia. 2019. Host effects in high ectomycorrhizal diversity tropical
 450 rainforests on ultramafic soils in New Caledonia. *Fungal Ecol.* 39:220-212.
 451 <https://doi.org/10.1016/j.funeco.2019.02.006>
- 452 Crossay T. 2018. Caractérisation taxonomique des champignons mycorrhiziens à arbuscules natifs
 453 des sols ultramafiques de Nouvelle-Calédonie. Analyse de leur synergie permettant l’adaptation
 454 des plantes à ces milieux extrêmes [Taxonomic characterization of arbuscular mycorrhizal
 455 fungi from New Caledonian ultramafic soils. Analysis of their synergy inducing plant
 456 adaptation to extreme conditions] [dissertation]. Noumea: University of New Caledonia.
 457 French.
- 458 Crossay T, A Cilia, Y Cavaloc, H Amir, D Redecker. 2018. Four new species of arbuscular
 459 mycorrhizal fungi (Glomeromycota) associated with endemic plants from ultramafic soils of
 460 New Caledonia. *Mycol Progress.* 17:729–744. doi.org/10.1007/s11557-018-1386-5.
- 461 Crossay T, C Majorel, D Redecker, S Gensous, V Medevielle, G Durrieu, Y Cavaloc, H Amir.
 462 2019. Is a mixture of arbuscular mycorrhizal fungi better for plant growth than single-species
 463 inoculants? *Mycorrhiza.* 29:325–339. [doi:10.1007/s00572-019-00898-y](https://doi.org/10.1007/s00572-019-00898-y).
- 464 Dubus IG, T Becquer. 2001. Phosphorus sorption and desorption in oxide-rich ferrasols of New
 465 Caledonia. *Aust J Soil Res.* 39:403–414. [doi:10.1071/SR00003](https://doi.org/10.1071/SR00003).
- 466 Duhoux E, G Rinaudo, HG Diem, F Auguy, D Fernandez, D Bogusz, C Franche, Y Dommergues,
 467 B Huguenin. 2001. Angiosperm *Gymnostoma* trees produce root nodules colonized by
 468 arbuscular mycorrhizal fungi related to *Glomus*. *New Phytol.* 149:115–125.
- 469 Ferrol N, E Tamayo, P Vargas. 2016. The heavy metal paradox in arbuscular mycorrhizas: from
 470 mechanisms to biotechnological applications. *J Exp Bot.* 67:6253–6265.
 471 [doi:10.1093/jxb/erw403](https://doi.org/10.1093/jxb/erw403).
- 472 Gauthier D, E Navarro, TG Rinaudo, P Jourand, T Jaffré, Y Prin. 1999. Isolation, characterization
 473 and specificity of Abundance of *Frankia* from eight *Gymnostoma* species endemic to New
 474 Caledonia. *Eur J Soil Biol.* 35(4):199-205.

- 475 Gensous S. 2014. Les champignons mycorrhiziens à arbuscules des maquis miniers de Nouvelle-
476 Calédonie : diversité et rôle dans l'adaptation des plantes à la carence en phosphore et à la
477 toxicité métallique [Arbuscular mycorrhizal fungi from New Caledonian ultramafic maquis :
478 diversity and role in plant adaptation to phosphorus deficiency and metal toxicity]
479 [dissertation]. Noumea : University of New Caledonia. French.
- 480 Göhre V, U Paszkowski. 2006. Contribution of the arbuscular mycorrhizal symbiosis to heavy
481 metal phytoremediation. *Planta*. 223:1115–1122.
- 482 Gonin M, S Gensous, A Lagrange, M Ducouso, H Amir, P Jourand. 2013. Rhizosphere bacteria of
483 *Costularia* spp. from ultramafic soils in New Caledonia: diversity, tolerance to extreme edaphic
484 conditions, and role in plant growth and mineral nutrition. *Can J Microbiol*. 59(3):164-74.
- 485 Gourmelon V, L Maggia, JR Powell, S Gigante, S Hortal, C Gueunier, K Letellier, Carriconde F.
486 2016. Environmental and geographical factors structure soil microbial diversity in New
487 Caledonian ultramafic substrates: A metagenomic approach. *PLoS One* 11:e0167405.
- 488 Hart MM, PM Antunes, VB Chaudhary, LK Abbott. 2017. Fungal inoculants in the field: Is the
489 reward greater than the risk? *Funct Ecol*. 00:1–10. doi: 10.1111/1365-2435.12976.
- 490 Herrera A, M Héry, JEM Stach, T Jaffré, P Normand, E Navarro. 2007. Species richness and
491 phylogenetic diversity comparisons of soil microbial communities affected by nickel-mining
492 and revegetation efforts in New Caledonia. *Eur J Soil Biol*. 43:130-139.
- 493 Héry M, S Nazaret, T Jaffré, P Normand, E Navarro. 2003. Adaptation to nickel spiking of bacterial
494 communities in neocaledonian soils. *Environ Microbiol*. 5:3-12. PMID: 12542708.
- 495 Houlès A, K Gotty, F Joussemet, B Vincent, L Hannibal, M Patrois, P Jourand, M Ducouso. 2022.
496 The Nurse Plant *Acacia spirorbis* Enriches Ectomycorrhizal Community Composition of a
497 Target Species: *Tristaniopsis calobuxus*. *Diversity* 14:107. <https://doi.org/10.3390/d14020107>
- 498 Humphreys CP, PJ Franks, M Rees, MI Bidartondo, JR Leake, DJ Beerling. 2010. Mutualistic
499 mycorrhiza-like symbiosis in the most ancient group of land plants. *Nat Commun*. 1(1):1-7.
- 500 Isnard S, L L'huillier, F Rigault, T Jaffré. 2016. How did the ultramafic soils shape the flora of the
501 New Caledonian hotspot? *Plant Soil*. 403:53–76.
- 502 Jaffré T, P Bouchet, JM Veillon. 1998. Threatened plants of New Caledonia: is the system of
503 protected areas adequate? *Biodiv Conserv*. 7:107-135.
- 504 Jaffré T, M Latham, P Quantin. 1971. Les sols des massifs miniers de Nouvelle-Calédonie et leur
505 relation avec la végétation [The soils of New Caledonian ultramafic massifs and their
506 relationships with vegetation]. Noumea : ORSTOM. Report 26. French.

- 507 Jaffré T, M Latham. 1974. Contribution à l'étude des relations sol-végétation sur un massif de
508 roches ultrabasiques de la côte Ouest de la Nouvelle-Calédonie : le Boulinda. [contribution to
509 the study of sol-vegetation relationships in the ultramafic massif of New Caledonian west
510 coast: the Boulinda]. *Adansonia* 14(3):311-336. French.
- 511 Jaffré T, L L'huillier. 2010. Conditions de milieu des terrains miniers [Environmental conditions of
512 the mining areas]. In L L'Huillier, T Jaffré, A Wulff, editors. *Mines et environnement en
513 Nouvelle-Calédonie : les milieux sur substrats ultramafiques et leur restauration* [Mines and
514 environment in New Caledonia : the ultramafic areas and their restoration]. Noumea: IAC Ed.;
515 p. 33–44. French.
- 516 Janoušková M, K Krak, C Wagg, H Storchova, P Caklova, M Vosatka . 2013. Effects of inoculum
517 additions in the presence of a preestablished arbuscular mycorrhizal fungal community. *Appl
518 Environ Microbiol.* 79:6507–6515. doi: 10.1128/AEM.02135-13
- 519 Jourand P, M Ducouso, R Reid, C Majorel, C Richert, J Riss, M Lebrun. 2010a. Nickel-tolerant
520 ectomycorrhizal *Pisolithus albus* ultramafic ecotype isolated from nickel mines in New
521 Caledonia strongly enhance growth of a host plant at toxic nickel concentrations. *Tree Physiol.*
522 30:1311–1319.
- 523 Jourand P, M Ducouso, C Loulergue-Majorel, L Hannibal, S Santoni, Y Prin, M Lebrun. 2010b.
524 Ultramafic soils from New Caledonia structure *Pisolithus albus* in ecotype. *FEMS Microbiol
525 Ecol.* 72:238–249.
- 526 Karatygin, IV, NS Snigirevskaya, KN Demchenko. 2006. Species of the genus *Glomites* as plant
527 symbionts in Early Devonian ecosystems. *Paleont J.* 40:572–579.
- 528 L'Huillier L, S Dupont, I Dubus, T Becquer, E Bourdon, P Laubreaux, B Bonzon. 1998. Carence et
529 fixation du phosphore dans les sols ferrallitiques ferritiques de Nouvelle-Calédonie
530 [Phosphorus deficiency and fixation in New Caledonian lateritic ultramafic soils]. Paper
531 presented at : World Congress of Soil Science 16, august 20-26. Montpellier: ORSTOM.
532 French.
- 533 L'Huillier L, S Edighoffer. 1996. Extractability of nickel and its concentration in cultivated plants
534 in Ni rich ultramafic soils of New Caledonia. *Plant Soil.* 186:255-264.
- 535 L'Huillier L, A Wulff, G Gateblé, B Fogliani, C Zongo, T Jaffré. 2010. La restauration des sites
536 miniers [The restoration of mining areas]. In L L'Huillier, T Jaffré, A Wulff, editors. *Mines et
537 environnement en Nouvelle-Calédonie : les milieux sur substrats ultramafiques et leur*

- 538 restauration [Mines and environment in New Caledonia : the ultramafic areas and their
539 restoration]. Noumea : IAC Ed.; p. 147–230. French.
- 540 Lagrange A, M Ducouso, P Jourand, C Majorel, H Amir. 2011. New insights into the mycorrhizal
541 status of Cyperaceae from ultramafic soils in New Caledonia. *Can J Microbiol.* 57:21–28.
- 542 Lagrange A, L L’Huillier, H Amir. 2013. Mycorrhizal status of Cyperaceae from New Caledonian
543 ultramafic soils: effects of phosphorus availability on arbuscular mycorrhizal colonization of
544 *Costularia comosa* under field conditions. *Mycorrhiza.* 23:655–661. doi:org/10.1007/s00572-
545 013-0503-1.
- 546 Lambers H, C Mougel, B Jaillard, P Hinsinger. 2009. Plant-microbe soil interactions in the
547 rhizosphere: an evolutionary perspective. *Plant Soil.* 321:83–115.
- 548 Latham M. 1975. Les sols d’un massif de roches ultrabasiques de la côte ouest de la Nouvelle-
549 Calédonie : le Boulinda. Les sols à accumulation humifère [The soils of an ultramafic massif of
550 New Caledonian west coast: the Boulinda. *Cah ORSTOM.* 13:27-40. French.
- 551 Latham M, P Quantin, G Aubert. 1978. Etude des sols de Nouvelle-Calédonie. Notice explicative
552 [Study of the New Caledonian soils. Explanatory note]. Report n°78. Paris: ORSTOM ed.
553 French.
- 554 Lebedeva EV, NN Lyalikova, YY Bujelskii. 1978. Participation of nitrifying bacteria in the
555 weathering of serpentized ultrabasic rocks. *Mikrobiologiya.* 47:1101-1107.
- 556 Lee J, RD Reeves, RR Brooks, T Jaffré. 1977. Isolation and identification of citrate-complex of
557 nickel from nickel-accumulating plants. *Phytochem.* 16:1503-1505.
- 558 Leyval C, K Turnau, K Haselwandter. 1997. Effect of heavy metal pollution on mycorrhizal
559 colonization and function: physiological, ecological and applied aspects. *Mycorrhiza.* 7:139–
560 153.
- 561 Maherali H, JN Klironomos. 2007. Influence of Phylogeny on Fungal Community Assembly and
562 Ecosystem Functioning. *Science.* 316:1746–1748. [doi:org/10.1126/science.1143082](https://doi.org/10.1126/science.1143082).
- 563 Majorel C, L Hannibal, M Soupe, F Carriconde, M Ducouso, M Lebrun, P Jourand. 2012.
564 Tracking nickel-adaptive biomarkers in *Pisolithus albus* from New Caledonia using a
565 transcriptomic approach. *Mol Ecol.* 21:2208–2223.
- 566 Marchese C. 2015. Biodiversity hotspots: a shortcut for a more complicated concept. *Global Ecol*
567 *Conserv.* 3:297–309.
- 568 Marschner H. 1995. Mineral Nutrition of Higher Plants. London: Academic Press. 1:869-878

- 569 Mercky S. 1999. Rhizobactéries natives de Nouvelle-Calédonie promotrices de la croissance des
570 plantes [Plant growth promoting rhizobacteria from New Caledonian soils]. *Bois For Trop.*
571 262:80-83. French.
- 572 Myers N, RA Mittermeier, CG Mittermeier, GA da Fonseca, J Kent. 2000. Biodiversity hotspots for
573 conservation priorities. *Nature* 403:853–858.
- 574 Navarro E, T Jaffré, D Gauthier, F Gourbière, G Rinaudo, P Simonet, P Normand. 1999.
575 Distribution of *Gymnostoma* spp. microsymbiotic *Frankia* strains in New Caledonia is related
576 to soil type and to host-plant species. *Mol Ecol.* 8:1781-1788.
- 577 Orłowska E, W Przybyłowicz, D Orłowski, K Turnau, J Mesjasz-Przybyłowicz. 2011. The effect of
578 mycorrhiza on the growth and elemental composition of Ni-hyperaccumulating plant *Berkheya*
579 *coddii* Roessler. *Environ Pollut.* 159:3730–3738.
- 580 Pawłowska TE, RL Chaney, M Chin, I Charvat. 2000. Effects of metal phytoextraction practices on
581 the indigenous community of arbuscular mycorrhizal fungi at a metal contaminated landfill.
582 *Appl Environ Microbiol.* 66:2526–2530.
- 583 Pellegrino E, A Turrini, HA Gamper, G Cafà, E Bonari, JPW Young, M Giovannetti. 2012.
584 Establishment, persistence and effectiveness of arbuscular mycorrhizal fungal inoculants in the
585 field revealed using molecular genetic tracing and measurement of yield components. *New*
586 *Phytol.* 194: 810-822.
- 587 Perrier N. 2005. Bio-Géodiversité fonctionnelle des sols latéritiques miniers : application à la
588 restauration écologique (massif du Koniambo, Nouvelle-Calédonie) [Functional bio-
589 geodiversity of mining lateritic soils : application to ecological resoration (Koniambo massif,
590 New Caledonia)] [dissertation]. Noumea : University of New Caledonia. French.
- 591 Perrier N, H Amir, F Colin. 2006. Occurrence of mycorrhizal symbioses in the metal-rich lateritic
592 soils of the Koniambo massif, New Caledonia. *Mycorrhiza.* 16:449–458. [doi:org/10.1007/
593 s00572-006-0057-6](https://doi.org/10.1007/s00572-006-0057-6).
- 594 Pillon Y, Jaffré T, Birnbaum P, Bruy D, Cluzel D, Ducouso M, Fogliani B, Ibanez T, Jourdan H,
595 Lagarde L, Léopold A, Munzinger J, Pouteau R, Read J & Isnard S. 2021. Infertile landscapes
596 on an old oceanic island: the biodiversity hotspot of New Caledonia. *Biol J Linn Soc.* 133:317–
597 341.
- 598 Proctor J. 2003. Vegetation and soil and plant chemistry on ultramafic rocks in the tropical Far East.
599 *Perspect Plant Ecol.* 6:105–124. [doi :org/10.1078/1433-8319-00045](https://doi.org/10.1078/1433-8319-00045).

- 600 Quantin P, E Bordon, T Becquer. 1997. Minéralogie et contraintes édaphiques des sols
601 « ferritiques » dérivés de roches ultramafiques en Nouvelle-Calédonie : relation entre
602 constituants minéraux et disponibilités en certains éléments (Al, Fe, Si, Mg, Mn, Ni, Co, Cr,
603 Cu, Zn et Mo) facilement solubles [Mineralogy and edaphic constraints of lateritic ultramafic
604 soils in New Caledonia : relationships between mineral constituents and availability of different
605 elements (Al, Fe, Si, Mg, Mn, Ni, Co, Cr, Cu, Zn, Mo) easily soluble]. In T Jaffré, RD Reeves,
606 T Becquer, editors. *Ecologie des milieux sur roches ultramafiques et sur sols métallifères* [The
607 ecology of ultramafic and metalliferous areas]. Noumea : ORSTOM ed.; p. 13-25. French.
- 608 Russell AJ, Bidartondo MI, Butterfield BG. 2002. The root nodules of the Podocarpaceae harbour
609 arbuscular mycorrhizal fungi. *New Phytol.* 156:283–295 doi.org/10.1046/j.1469-
610 8137.2002.00504.x
- 611 Saintpierre D. 2001. Identification de souches originales d'actinomycètes isolés des sols
612 ultramafiques de Nouvelle-Calédonie. Caractérisation chimique de quelques métabolites
613 produits [Identification of original actinomycete strains isolated from New Caledonian
614 ultramafic soils. Chemical characterization of some metabolic products] [dissertation].
615 Toulouse : Institut National Polytechnique. French.
- 616 Smith SE, DJ Read. 2008. *Mycorrhizal symbiosis* (third edition). New York : Academic press.
- 617 Smith SE, FA Smith. 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new
618 paradigms from cellular to ecosystems scales. *Annu Rev Plant Biol.* 62:227–250
- 619 Strullu-Derrien C, P Kenrick, S Pressel, JG Duckett, JP Rioult, DG Strullu. 2014. Fungal
620 associations in *Horneophyton ligneri* from the Rhynie Chert (ca 407 Ma) closely resemble
621 those in extant lower land plants: novel insights into ancestral plant fungus symbioses. *New
622 Phytol.* 203: 964–979.
- 623 Strullu-Derrien C, P Kenrick, MA Selosse. 2017. Structure and phylogeny of mycorrhizal
624 symbioses. In F Martin, editor. *Molecular Mycorrhizal Symbiosis*. New Jersey: Wiley-
625 Blackwell; p. 1-20.
- 626 Tullio M, F Pierandrei, A Salerno, E Rea. 2003. Tolerance to cadmium of vesicular arbuscular
627 mycorrhizae spores isolated from cadmium-polluted and unpolluted soil. *Biol Fertil Soils.*
628 37:211–214.
- 629 Turnau K, J Mesjasz-Przybyłowicz. 2003. Arbuscular mycorrhiza of *Berkheya coddii* and other Ni-
630 hyperaccumulating members of Asteraceae from ultramafic soils in South Africa. *Mycorrhiza.*
631 13:185–190.

- 632 Valliere J, WS Wong, PG Nevill, H Zhong, KW Dixon. 2020. Preparing for the worst: Utilizing
633 stress-tolerant soil microbial communities to aid ecological restoration in the Anthropocene.
634 *Ecol Solut Evid.*1:e12027. [doi:10.1002/2688-8319.12027](https://doi.org/10.1002/2688-8319.12027).
- 635 Wang B, YL Qiu. 2006. Phylogenetic distribution and evolution of mycorrhizas in land plants.
636 *Mycorrhiza.* 16:299–363. [doi:10.1007/s00572-005-0033-6](https://doi.org/10.1007/s00572-005-0033-6).
- 637 Weissenhorn I, A Glashoff, C Leyval, J Berthelin. 1994. Differential tolerance to Cd and Zn of
638 arbuscular mycorrhizal (AM) fungal spores isolated from heavy metal-polluted and unpolluted
639 soils. *Plant Soil.* 167(2):189-196.