



HAL
open science

Genetic parameters of maturing rate index and asymptotic adult weight in French beef cattle

Armance Lepers, S. Aguerre, Julie Promp, Sébastien Taussat, Aurélie Vinet,
Pauline Martin, A. Philibert, A. Laramee, L. Griffon

► To cite this version:

Armance Lepers, S. Aguerre, Julie Promp, Sébastien Taussat, Aurélie Vinet, et al.. Genetic parameters of maturing rate index and asymptotic adult weight in French beef cattle. 74. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2023, Lyon, France. hal-04449124

HAL Id: hal-04449124

<https://hal.science/hal-04449124v1>

Submitted on 9 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Genetic parameters of maturing rate index and asymptotic adult weight in French beef cattle

LEPERS A.¹, AGUERRE S.¹, PROMP J.¹, TAUSSAT S.², VINET A.³, MARTIN P.³, PHILIBERT A.⁴, LARAMEE A.⁴, GRIFFON L.⁴

¹ Institut de l'Élevage, UMR 1313 GABI, F78350 Jouy-en-Josas, ² ELIANCE, UMR 1313 GABI, F78350 Jouy-en-Josas, ³ INRAE, UMR 1313 GABI, F78350 Jouy-en-Josas, ⁴ Institut de l'Élevage, F75595 Paris

INTRODUCTION

Working on the early development of French beef cows is interesting to:

- Reduce fattening time and added concentrates
- Reduce the environmental impact of finished products
- Facilitate 2-year-old calving through early development

**Can early development be estimated from on-farm weighing?
Is genetic selection for early development possible?**

MATERIALS AND METHODS

Phenotypes (5 beef breeds): Use of Brody equation from birth weight declared by breeders, performance control weights and NORMABEV carcass weights of adult females.

Brody equation:


$$Weight(time) = Weight_{adult} - (Weight_{adult} - Weight_{birth})e^{-maturing\ rate\ index \times time}$$

Genetic parameters:


- Estimated adult weight and maturing rate index: data corrected for fixed effects related to birth and slaughter.
- Birth weight and adjusted weights (AW) at 7m, 12m, 18m and 24m: corrected performances with a correction of genetic progress.

RESULTS

Average female performance in each of the five considered breeds:



HERITABILITY ESTIMATES



- Not enough data for the Aubrac breed
- Standard deviation of error between 0.01 and 0.03
- Average genetic variation coefficient equal to 6.16% for adult weight and 9.09% for maturing rate index

Genetic correlation of maturing rate index with bodyweight at different ages (average of the 4 breeds):

	Birth weight	AW at 7m	AW at 12m	AW at 18m	AW at 24m	Adult weight
Maturing rate index	-0.12 (-0.18 to -0.07)	0.32 (0.29 to 0.37)	0.39 (0.32 to 0.45)	0.42 (0.27 to 0.69)	0.40 (-0.03. to 0.70)	-0.64 (-0,65 to -0,62)

DISCUSSION

The maturing rate index seems to be a good phenotype for early development. It would be possible to genetically select the maturing rate index (due to its heritability and genetic variance) without increasing birth weight (average genetic correlation of -0.12). The maturing rate index seems to be positively correlated with adjusted weights between 7 and 24 months. However, the lack of animals with adjusted weights at 18 and 24 months limits the accuracy of the associated correlations and therefore their possible interpretation. The highly negative genetic correlation between maturing rate index and adult weight shows that selecting for heavier adult animals reduces their maturing speed. The disadvantage of the maturing rate index are that the phenotype arrives at the end of the animals' lives and is very dependent on the weaning and post-weaning weights. To obtain breeding values for young animals, a genomic prediction needs to be developed.

CONCLUSION

Maturing rate index is ...

- Measurable from commercial data, although it is highly dependent on weighing at weaning and post-weaning.
- Heritable and genetically variable, so direct selection is possible.
- Genetically correlated with other traits, allowing indirect selection in particular via reduction in adult weight.

Many thanks to the 6 breed societies who participated.