



**HAL**  
open science

## The Severity of Congenital Hypothyroidism With Gland-In-Situ Predicts Molecular Yield by Targeted Next-Generation Sequencing.

Lucie Levailant, Natacha Bouhours-Nouet, Frederic Illouz, Jessica Amsellem Jager, Anne Bachelot, Pascal Barat, Sabine Baron, Candace Bensignor, Aude Brac de la Perriere, Yasmine Braik Djellas, et al.

► **To cite this version:**

Lucie Levailant, Natacha Bouhours-Nouet, Frederic Illouz, Jessica Amsellem Jager, Anne Bachelot, et al.. The Severity of Congenital Hypothyroidism With Gland-In-Situ Predicts Molecular Yield by Targeted Next-Generation Sequencing.. *Journal of Clinical Endocrinology and Metabolism*, 2023, 108 (9), pp.e779-e788. 10.1210/clinem/dgad119 . hal-04452836

**HAL Id: hal-04452836**

**<https://hal.science/hal-04452836v1>**

Submitted on 12 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# The Severity of Congenital Hypothyroidism With Gland-In-Situ Predicts Molecular Yield by Targeted Next-Generation Sequencing

Lucie Levailant,<sup>1,2</sup>  Natacha Bouhours-Nouet,<sup>1,2</sup> Frédéric Illouz,<sup>2,3</sup> Jessica Amsellem Jager,<sup>1,2</sup> Anne Bachelot,<sup>4</sup> Pascal Barat,<sup>5</sup> Sabine Baron,<sup>6</sup> Candace Bensignor,<sup>7</sup> Aude Brac De La Perriere,<sup>8</sup> Yasmine Braik Djellas,<sup>4</sup> Morgane Caillot,<sup>9</sup> Emmanuelle Caldagues,<sup>6</sup> Marie-Neige Campas,<sup>10</sup> Marylène Caquard,<sup>6</sup> Audrey Cartault,<sup>11</sup> Julie Cheignon,<sup>1</sup> Anne Decrequy,<sup>1</sup> Brigitte Delemer,<sup>12</sup> Katherine Dieckmann,<sup>13</sup> Aurélie Donzeau,<sup>1</sup> Emilie Doye,<sup>14</sup> Mélanie Fradin,<sup>15</sup> Mélanie Gaudillière,<sup>8</sup> Frédérique Gatelais,<sup>16</sup> Magali Gorce,<sup>17</sup> Isabelle Hazart,<sup>6</sup> Nada Houcinat,<sup>18</sup> Laure Houdon,<sup>19</sup> Marielle Ister-Salome,<sup>20</sup> Lucie Jozwiak,<sup>21</sup> Patrick Jeannoel,<sup>22</sup> Francois Labarthe,<sup>23</sup> Didier Lacombe,<sup>24</sup> Anne-Sophie Lambert,<sup>25</sup> Christine Lefevre,<sup>20</sup> Bruno Leheup,<sup>26</sup> Clara Leroy,<sup>27</sup> Benedicte Maisonneuve,<sup>28</sup> Isis Marchand,<sup>29</sup> Emeline Marquant,<sup>30</sup> Matthias Muszлак,<sup>31</sup> Letitia Pantalone,<sup>32</sup> Sandra Pochelu,<sup>5</sup> Chloé Quelin,<sup>15</sup> Catherine Radet,<sup>33</sup> Peggy Renoult-Pierre,<sup>34</sup> Rachel Reynaud,<sup>30</sup> Stéphanie Rouleau,<sup>1</sup> Cécile Teinturier,<sup>25</sup> Julien Thevenon,<sup>35</sup> Caroline Turlotte,<sup>36</sup> Aline Valle,<sup>37</sup> Melody Vierge,<sup>30</sup> Carine Villanueva,<sup>8</sup> Alban Ziegler,<sup>17</sup> Xavier Dieu,<sup>2,38</sup> Nathalie Bouzamondo,<sup>2,38</sup> Patrice Rodien,<sup>2,3</sup> Delphine Prunier-Mirebeau,<sup>2,38</sup> and Régis Coutant<sup>1,2</sup>

<sup>1</sup>Department of Pediatric Endocrinology and Diabetology, University Hospital of Angers, 49000 Angers, France

<sup>2</sup>Reference Center for Rare Diseases of Thyroid and Hormone Receptivity, University Hospital of Angers, 49000 Angers, France

<sup>3</sup>Department of Endocrinology, Diabetes and Nutrition, University Hospital of Angers, 49000 Angers, France

<sup>4</sup>Department of Endocrinology and Reproductive Medicine, Hôpital Pitié-Salpêtrière, ICAN, 75651 Paris, France

<sup>5</sup>Pediatric Endocrinology, CHU de Bordeaux, 33000 Bordeaux, France

<sup>6</sup>Pediatrics Department, CHU Nantes, 44000 Nantes, France

<sup>7</sup>Pediatrics Department, CHU de Dijon, 21000 Dijon, France

<sup>8</sup>Hospices Civils de Lyon, Hôpital Femme Mère Enfant, Service d'Endocrinologie Pédiatrique, 69677 Bron, France

<sup>9</sup>Pediatrics Department, CH de Martigues, 13500 Martigues, France

<sup>10</sup>Pediatrician, 64445 Pau, France

<sup>11</sup>Endocrine, Genetics, Bone Diseases, and Paediatric Gynecology Unit, Children's Hospital, CHU Toulouse, 31059 Toulouse, France

<sup>12</sup>Department of Endocrinology, Diabetes and Nutrition, CHU de Reims-Hôpital Robert-Debré, 51100 Reims, France

<sup>13</sup>Pediatrics Department, CH de Blois, 41000 Blois, France

<sup>14</sup>Pediatrician, 69130 Ecully, France

<sup>15</sup>Service de Génétique, CLAD Ouest, CHU Rennes, 35200 Rennes, France

<sup>16</sup>Pediatrician, 49000 Angers, France

<sup>17</sup>Service de Génétique, 49000 Angers Cedex 9, France

<sup>18</sup>CHU Dijon, Centre de référence maladies rares Anomalies du Développement et Syndromes Malformatifs, Centre de Génétique, FHU TRANSLAD, CHU Dijon Bourgogne 21000, France

<sup>19</sup>Pediatric Diabetology, University Hospital, St Pierre de la Reunion 97410, France

<sup>20</sup>Pediatric Endocrinology, Jeanne de Flandre Hospital, 59037 Lille, France

<sup>21</sup>Pediatrics Department, CH de Roubaix, 59100 Roubaix, France

<sup>22</sup>Pediatrics Department, CH de Roanne, 42328 Roanne, France

<sup>23</sup>Reference Center for Inborn Errors of Metabolism, Tours University Hospital, 37044 Tours, France

<sup>24</sup>Department of Medical Genetics, CHU Bordeaux INSERM U1211, Université de Bordeaux, 33076 Bordeaux, France

<sup>25</sup>AP-HP, Bicêtre Paris Saclay Hospital, DMU SEA, Endocrinology and Diabetes for Children, Le Kremlin Bicêtre 94270, France

<sup>26</sup>Service de Génétique clinique, Hôpital Brabois, Centre Hospitalier Universitaire de Nancy, Nancy, Lorraine 54500, France

<sup>27</sup>Service d'Endocrinologie et Maladies Métaboliques, Centre Hospitalier Régional Universitaire de Lille, Hôpital Huriez, 59037 Lille, France

<sup>28</sup>Pediatrics Department, CH de Montluçon, 03100 Montluçon, France

<sup>29</sup>Pediatrics Department, CHI de Créteil, 94010 Créteil, France

Received: 12 April 2022. Editorial Decision: 23 February 2023. Corrected and Typeset: 14 March 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of the Endocrine Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

<sup>30</sup>Assistance-Publique des Hôpitaux de Marseille, Department of Pediatrics, Hôpital de la Timone Enfants, 13005 Marseille, France

<sup>31</sup>Pediatrics Department, CH de Mayotte, 97600 Mayotte, France

<sup>32</sup>Pediatrics Department, CH René Dubos, 95300 Pontoise, France

<sup>33</sup>Pediatrics Department, CH de Cholet, 49300 Cholet, France

<sup>34</sup>Service de Médecine Interne, Unité d'Endocrinologie Diabétologie et Nutrition, Centre Hospitalier Universitaire et Faculté de Médecine, Université de Tours, 37044 Tours, France

<sup>35</sup>Inserm UMR 1231 GAD Team, Genetics of Developmental Anomalies, and FHU-TRANSLAD, CHU/Université de Bourgogne-Franche Comté, 21000 Dijon, France

<sup>36</sup>Pediatrics Department, CH d'Armentières, 59280 Armentières, France

<sup>37</sup>Pediatrics Department, CH de Douai, 59187 Douai, France

<sup>38</sup>Biochemistry and Molecular Biology Laboratory, University Hospital of Angers, 49000 Angers, France

**Correspondence:** Lucie Levailant, MD, Department of Pediatric Endocrinology and Diabetology, University Hospital of Angers, 4 rue Larrey, 49000 Angers, France. Email: [lucie.levailant@chu-angers.fr](mailto:lucie.levailant@chu-angers.fr).

## Abstract

**Introduction:** Congenital hypothyroidism with gland-in-situ (CH-GIS) is usually attributed to mutations in the genes involved in thyroid hormone production. The diagnostic yield of targeted next-generation sequencing (NGS) varied widely between studies. We hypothesized that the molecular yield of targeted NGS would depend on the severity of CH.

**Methods:** Targeted NGS was performed in 103 CH-GIS patients from the French national screening program referred to the Reference Center for Rare Thyroid Diseases of Angers University Hospital. The custom targeted NGS panel contained 48 genes. Cases were classified as solved or probably solved depending on the known inheritance of the gene, the classification of the variants according to the American College of Medical Genetics and Genomics, the familial segregation, and published functional studies. Thyroid-stimulating hormone at CH screening and at diagnosis (TSH<sub>sc</sub> and TSH<sub>dg</sub>) and free T4 at diagnosis (FT4<sub>dg</sub>) were recorded.

**Results:** NGS identified 95 variants in 10 genes in 73 of the 103 patients, resulting in 25 solved cases and 18 probably solved cases. They were mainly due to mutations in the *TG* (n = 20) and *TPO* (n = 15) genes. The molecular yield was, respectively, 73% and 25% if TSH<sub>sc</sub> was ≥ and < 80 mIU/L, 60% and 30% if TSH<sub>dg</sub> was ≥ and < 100 mIU/L, and 69% and 29% if FT4<sub>dg</sub> was ≤ and > 5 pmol/L.

**Conclusion:** NGS in patients with CH-GIS in France found a molecular explanation in 42% of the cases, increasing to 70% when TSH<sub>sc</sub> was ≥ 80 mIU/L or FT4<sub>dg</sub> was ≤ 5 pmol/L.

**Key Words:** congenital hypothyroidism, gland-in-situ, molecular yield, next-generation sequencing, severity

Congenital hypothyroidism (CH) is the most common neonatal endocrine disorder, with an incidence of 1/2500 to 1/3500 newborns (1, 2). Among patients with CH, 15% to 50% have a gland-in-situ (GIS) with or without goiter. In France, the CH screening program relies on thyroid-stimulating hormone (TSH) measurement on dried blood samples collected on filter paper from all newborns at 3 days of life. From the 2000s, an increased incidence of CH has been observed, up to 1/2419 newborns in 2019 (1, 3). This has been driven by an increase in CH-GIS, from 15% of the cases before the 2000s to 50% in 2018 (1). The annual incidence of CH-GIS increased by 5.1% in this period, whereas the incidence of thyroid dysgenesis remained constant (1).

CH-GIS is usually attributed to dysmorphogenesis resulting from mutations in the genes involved in thyroid hormone production or iodide metabolism, typically associated with goiter: thyroglobulin (*TG*), thyroid peroxidase (*TPO*), *SLC26A4* also called pendrin, *SLC5A5* also called Na<sup>+</sup>/I<sup>-</sup> symporter (*NIS*), dual oxidase 2 (*DUOX2*), dual oxidase maturation factor 2 (*DUOX2*), dual oxidase 1 (*DUOX1*), dual oxidase maturation factor 1 (*DUOX1*), iodotyrosine deiodinase (*IYD*) also called iodotyrosine dehalogenase 1 (*DEHAL1*), and the iodide transporter *SLC26A7* (4–10). In addition, mutations in the genes formally involved in thyroid dysgenesis may also be responsible for CH-GIS, typically with thyroid hypoplasia: paired box 8 (*PAX8*), thyroid-stimulating hormone receptor (*TSHR*), Forkhead box E1 (*FOXE1*), and *NKX2-1* genes (6, 11–14).

The proportion of patients with CH-GIS who receive a molecular diagnosis varies widely, from 10% to 93% in studies involving 10 to 290 patients (15–29). Factors contributing to this variability include differences in the clinical characterization of the patients (imaging techniques, perchlorate test),

proportion of familial or sporadic cases, geographic origin, population inbreeding rate, and mainly variant classification. None of these studies has focused on the rate of molecular explanation of CH-GIS according to TSH at screening, TSH, and free T4 (FT4) at diagnosis, all usual markers of the disease severity (30).

Our hypothesis was that the molecular yield would be dependent on the severity of CH. We report 103 patients with CH-GIS referred to our Reference Center for Rare Thyroid Diseases at Angers University Hospital for custom targeted next-generation sequencing (NGS). We assessed the proportion of genetic resolution of patients with CH-GIS according to the TSH at screening, TSH at diagnosis, and FT4 at diagnosis. Cases were considered solved based on known recessive or dominant inheritance of the gene, classification of the variants in agreement with the American College of Medical Genetics and Genomics (ACMG) guidelines (31), the localization in the protein domain, allele frequency in GnomAD genomes (32), in silico prediction based on SIFT and Mutation Taster prediction software (and Polyphen-2 when the 2 previous ones gave discordant results), PhyloP100way conservation score (33), the same variant previously described in the literature in a patient with CH, an in vitro study confirming the pathogenicity of the variant when available, and segregation of variants with CH in multiple affected family members.

## Methods

### Patients

All clinical data and blood samples were collected in our Reference Center for Rare Thyroid Diseases, University Hospital of Angers, France, for targeted NGS testing between January 1, 2016, and May 1, 2020.

Patients were included if they had CH-GIS, with or without goiter, ascertained by thyroid ultrasound (100 cases) and/or thyroid scintigraphy (85 cases). All the families gave written informed consent for genetic testing. In France, the screening is performed in all newborns on day 3. Newborns with a neonatal filter paper TSH > 17 mIU/L with a genetic screening processor (PerkinElmer, Turku, Finland) or > 20 mIU/L with AutoDELFIA (PerkinElmer) underwent a diagnostic procedure consisting of serum TSH, FT4 and free T3 (FT3), and thyroglobulin measurements, thyroid ultrasound, and thyroid scintigraphy, and often a perchlorate discharge test (all performed locally).

CH severity was classified as severe if FT4 was < 5 pmol/L, moderate if FT4 was between 5 and < 10 pmol/L, and mild if FT4 was > 10 pmol/L (34).

For the perchlorate discharge test, partial iodide organification defect and total iodide organification defect were defined as a radioiodine washout of more than 10% and 90%, respectively (35). Reference values for thyroglobulin levels during the first 15 days of life were used (36).

### Genetic Analysis by Next-Generation Sequencing

DNA was extracted from total blood samples with the EZ1 extraction kit (Qiagen, Hilden, Germany). High-throughput sequencing of genes was performed on an Ion Proton sequencer (Thermo Fisher Scientific, Waltham, MA, USA). An amplicon-based enrichment library that focused on 48 genes involved in thyroid dysmorphogenesis and thyroid dysgenesis was prepared using Ion AmpliSeq technology. It included 14 genes known to be involved in primary CH, 3 genes involved in central CH, and 31 in thyroid physiology [Supplementary Table S1 (37)]. The read depth was more than 20, with a high-quality score (Q30). Sanger sequencing was performed for variant confirmation.

### Classification of Variants

We used the 2015 standards and guidelines for the interpretation of sequence variants from the ACMG and the Association for Molecular Pathology (31) and VarSome ACMG implementation (<http://varsome.com>) to classify variants. We also considered the pathogenicity according to the localization in the protein domain (in a mutational hot spot and/or critical and well-established functional domain), allele frequency in GnomAD genomes version 3.1.1 (32) (<https://gnomad.broadinstitute.org/>), in silico prediction of the pathogenicity of the missense variant based on SIFT version 4.1 (<https://sift.bii.a-star.edu.sg/>), and Mutation Taster version 4.2 (<http://www.mutationtaster.org>) prediction software (and Polyphen-2 when the 2 previous ones gave discordant results, (<http://genetics.bwh.harvard.edu/pph2/>), the PhyloP100way conservation score (33) (<http://hgdownload.soe.ucsc.edu/goldenPath/hg38/phyloP100way/>), in vitro study confirming pathogenicity of the variant when available, the same variant previously described in the literature in a patient with CH, and familial segregation. This allowed us to classify the variants as “pathogenic,” “likely pathogenic,” “uncertain significance,” “likely benign,” or “benign.”

### Classification of Cases

From this classification of variants, we then classified the genetic results of each patient into 4 categories according to the

association of variants, potential familial segregation, and the genes involved (according to the recessive or dominant inheritance of the disease): (1) genetically solved cases, (2) probably solved cases, (3) partially unsolved cases, or (4) unsolved cases (Fig. 1).

Genetic variants in the following genes are considered recessive disorders: *TG*, *TPO*, *SLC26A4* (*pendrin*), *SLC5A5* (*NIS*), *FOXE1*, and *IYD* (4, 12–14). The patients had homozygous or compound heterozygous variants in the same gene and were classified according to Supplementary Table S2 (37).

Genetic variants in *PAX8* and *NKX2-1* are considered dominant disorders, as cases have been described in a heterozygous state (4), and patients were classified according to Supplementary Table S3 (37).

Regarding both the *DUOX* gene and its corresponding maturation factor *DUOXA*, cases have been described in heterozygous, compound heterozygous, or homozygous states, in one or several genes of the *DUOX* system (4, 5). This is also the case for the *TSHR* gene.

Thus, for variants in a heterozygous state, we considered cases solved with only 1 variant in 1 gene if this variant had already been described in CH in a heterozygous state and had an in vitro study confirming its pathogenicity. Every other case was considered partially unsolved.

To study the genetic resolution, we only considered solved and probably solved cases. We did not include the partially unsolved cases, because their variants alone could not explain the CH of the patient.

### Statistical Analysis

Continuous variables were expressed as the median (minimum and maximum). Discrete variables were expressed as percent and 95% CI. Significance was defined as  $P < .05$  (SPSS Statistics v25; IBM Corp., Armonk, NY, USA). The receiver operating characteristic curves and the genetic yields according to TSH at screening, TSH at diagnosis, and FT4 at diagnosis were calculated for each hormonal value. Then the curves of the molecular yields were modeled using smoothing splines (GraphPad Prism 9; GraphPad Software Inc.).

## Results


### Clinical and Biological Characteristics of the Whole Group of Patients

One hundred three patients with CH-GIS identified through the French national screening program were referred to our center and underwent genetic testing with NGS: the median TSH at day 3 of the newborn screening program ( $TSH_{sc}$ ) was 60 mIU/L (15–417), the median TSH at diagnosis ( $TSH_{dg}$ ; between day 6 and day 10) was 73 mIU/L (8–840), and the median FT4 ( $FT4_{dg}$ ) was 9.4 pmol/L (1.5–26.7).

Thirty-two patients had a goiter (31%). Twenty-six patients had a familial history of CH in first-degree relatives (25%). Four patients were diagnosed with hearing loss later in the follow-up. Fifteen patients (14.5%) had various abnormalities, such as severe mental retardation, single kidney, or polydactyly.

Fifteen patients had proven transient CH, as their treatment could be stopped: their median  $TSH_{sc}$  was 21.7 mIU/L (15–61), median  $TSH_{dg}$  was 42 mIU/L (11–67), and median  $FT4_{dg}$  was 12.2 pmol/L (8.4–23.3). Fourteen patients failed to stop treatment [median  $TSH_{sc}$  = 36 mIU/L (17–189), median  $TSH_{dg}$  = 18.9 mIU/L (8–558), median  $FT4_{dg}$  = 14.4 pmol/L (1.5–24)],





**Figure 1.** Classification of the cases, according to the genetic testing. See Supplementary Table S5 and Supplementary Table S6 and additional references (37) for a comprehensive classification of each case with variant descriptions. (A) Segregation of variants and CH in multiple affected family members. (B) The patient with *NKX2-1* mutation had pulmonary involvement and neurological delay and a familial segregation of variant and CH. (C) Variants on 2 different genes were considered as partially unsolved because of the lack of evidence in the literature. <sup>†</sup> represents a patient with a transient form. Abbreviations: CH, congenital hypothyroidism.

and the median levothyroxine dose was 2.9 µg/kg/day (0.8-7.5) at a median age of 7.2 years (2.7-21). Thirty-five patients 3 years old or older were considered by their attending physicians as permanent cases as TSH levels were sometimes above the upper limit while on levothyroxine treatment: the median levothyroxine dose was 2.7 µg/kg/day (0.88-9) at a median age of 11.2 years (3.7-37). Last, 15 patients younger than 3 years old were considered by their physician as too young to try stopping treatment: the median levothyroxine dose was 3.95 µg/kg/day (2.4-7.6) at a median age of 1.4 years (0.1-2.7).

### Molecular Explanation of the Cases and Relationships Between Genotype and Phenotype

Genetic testing found 95 variants in 73 of 103 patients: 38 variants were classified as "pathogenic," 14 as "likely pathogenic," 28 of "uncertain significance," and 15 as "likely benign" [Supplementary Table S4 (37)]. Variants classified as "benign" are not reported. After exclusion of "likely benign" variants, the 80 remaining variants were in 10 genes: *TG* (n = 28), *TPO* (n = 26), *DUOX2* (n = 7), *TSHR* (n = 6), *DUOX1* (n = 3), *PAX8* (n = 3), *SLC26A4* (n = 3), *SLC5A5* (n = 2), *DUOX1* (n = 1), and *NKX2-1* (n = 1).

According to the classification of variants, the potential familial segregation of the disease, and the involved genes (associated with either recessive or dominant disease inheritance), 25 cases were solved and 18 were probably solved [Supplementary Table S5; Supplementary Table S6; (37)]. A genetic explanation was found by NGS testing in 43/103 cases (42%; Fig. 1). CH-GIS was due to thyroglobulin mutations in 20 patients and to thyroid peroxidase mutations in 15 patients. Notably, 8 patients had heterozygous variants in genes first involved in thyroid dysgenesis: 5 with variants in the *PAX8* gene (ID 61, 62, 85, 98, 99), 2 in *TSHR* (ID 1 and 12), and 1 in *NKX2-1* (ID 75).

Family history increased the likelihood of genetic resolution, with 15/26 (58%) familial cases with a genetic

explanation (8/15 families), compared with 28/77 (36%) sporadic cases.

The clinical characteristics of the patients with solved or probably solved genetic cases according to the mutated gene are indicated in Table 1.

Some phenotypic characteristics were remarkable:

- Twenty-six of the 32 patients (81%) with goiter were solved or probably solved cases, mainly with *TG* and *TPO* variants (14 and 12 cases, respectively).
- Two of the 4 patients with hearing loss were solved cases due to *TPO* mutations (other cases were not solved).
- Among the 20 patients with probably solved or solved cases with *TG* variants, serum thyroglobulin measurements were available in 14 patients: it was undetectable in 9 patients, low in 4 patients, and in the normal range in 1 patient (ID 64). One other patient, ID 48, with a thyroid scintigraphy ascertaining GIS, had undetectable serum thyroglobulin but no variant found on NGS analysis. A mutation in a noncoding region altering gene expression has yet to be found.

### Genetic Resolution According to Thyroid Function at Screening and at Diagnosis


We studied the proportion of genetic resolution according to the TSH level at neonatal screening and at diagnosis and the FT4 level at diagnosis for all cases and only sporadic cases (Fig. 2). Median TSH<sub>dg</sub> was 100 mIU/L (19-417) in solved cases, 215 mIU/L (17-491) in probably solved cases, 32 mIU/L (8-618) in partially unsolved cases, and 27 mIU/L (17-250) in unsolved cases. Among the partially unsolved or unsolved cases in our study, the median TSH<sub>sc</sub> was 29 mIU/L (17-250), 25 mIU/L (15-275), and 20 mIU/L (15-90) for patients carrying 0 (n = 30), 1 (n = 12), or ≥ 2 (n = 13) nonbenign variants, respectively.

**Table 1. Clinical characteristics of patients with solved or probably solved cases, according to the mutated gene**

Solved or probably solved cases	TG variants	TPO variants	DUOX2 variants	TSHR variants	PAX8 variants	NKX2-1 variants
Number of patients	20	15	2	3	5	1
Median TSH on screening (mIU/L; min-max)	134 (42-210) (n = 14)	113 (56-417) (n = 8)	15	101 and 25	35, 38, and 186	
Median TSH at diagnosis (mIU/L; min-max)	150 (23-792) (n = 17)	213 (100-840) (n = 8)	14 and 100	17 and 25	38 (9-100) (n = 5)	
Median FT4 at diagnosis (pmol/L; min-max)	5.7 (2-9.8) (n = 15)	5.2 (1.8-8.9) (n = 9)	8.9 and 16.3	18.2 et 20.2	12.35 (7-15) (n = 4)	
Goitre (clinical and/or radiological)	70%	80%	50%			
Other clinical signs	30%	13%				
Serum thyroglobulin	n = 14	n = 5	n = 2	n = 2	n = 1	
Undetectable	64%	0				
Low	29%					
Normal	7%	2	1	2	1	
High		3	1			
Thyroid scintigraphy	n = 16	n = 10	n = 2	n = 1	n = 4	n = 1
High level of uptake <sup>d</sup>	94%	100%	2	1	3	1
Low level of uptake or absent <sup>d</sup>	6%				1	
Perchlorate discharge test	n = 6	n = 7		n = 1	n = 1	
Normal	2	3		1	1	
Partial iodide organification defect	4	2				
Total iodide organification defect		2				
Median treatment posology at last follow-up (µg/kg/day; min-max)	3.2 (1.4-5.7) (n = 16)	3 (1.4-5.4) (n = 10)	1.5	3.6, 2.5, and 0.8	2.5, 3.5, and 5.8	1.6
Median age at last follow-up (years; min-max)	8.6 (1.4-17.4) (n = 16)	14.5 (0.3-37) (n = 10)	7.3	23.6, 8.2, and 13.5	5, 5, and 7	4.4
Reevaluation of the thyroid axis	n = 2	n = 2	n = 2	n = 1	n = 2	
Transient CH	0	1	1	1	0	

Abbreviations: CH, congenital hypothyroidism; FT4, free T4.

<sup>d</sup>According to normal laboratory measure.



**Figure 2.** Proportion of genetic resolution according to the TSH level at neonatal screening (A) and at diagnosis (B), and the free T4 level at diagnosis (C) for all cases and only sporadic cases.

Considering the trade-off between sensitivity (population of solved or probably solved cases) and specificity (population of partially unsolved and unsolved cases), the area under the receiver operator characteristic curve (AUC) was 0.84 (CI<sub>95</sub> 0.74-0.94) for TSH at screening. A whole blood TSH at screening (TSH<sub>sc</sub>)  $\geq$  and  $<$  80 mIU/L resulted in 73% (CI<sub>95</sub> 58-92%; cut-off corresponding to the highest positive predictive value) and 25% (CI<sub>95</sub> 15-45%; 1 - negative predictive value) of the molecular yield of the cases, respectively (Fig. 2A).

The AUC was 0.71 (CI<sub>95</sub> 0.60-0.83) for TSH at diagnosis. A serum TSH at diagnosis (TSH<sub>dg</sub>)  $\geq$  and  $<$  100 mIU/L resulted in 60% (CI<sub>95</sub> 44-81%; cut-off corresponding to the highest positive predictive value) and 30% (CI<sub>95</sub> 20-46%) of the molecular yield of the cases, respectively (Fig. 2B).

The AUC was 0.81 (CI<sub>95</sub> 0.70-0.91) for FT4 at diagnosis. A serum FT4 at diagnosis (FT4<sub>dg</sub>)  $\leq$  and  $>$  10 pmol/L resulted in 60% (CI<sub>95</sub> 47-77%) and 15% (CI<sub>95</sub> 7-33%) of the molecular yield of the cases, respectively, with  $\leq$  and  $>$  5 pmol/L resulting in 69% (CI<sub>95</sub> 50-93%) and 29% (CI<sub>95</sub> 19-44%) of the molecular yield, respectively. The highest positive predictive values were observed for the lowest FT4 levels (Fig. 2C). Slightly lower positive predictive values, but with similar trends, were found when studying only sporadic cases (Fig. 2).

### Subjects With Transient Congenital Hypothyroidism

In the 15 patients with transient CH, one was considered probably solved. This patient had compound heterozygous variants in SLC26A4 (1 likely pathogenic and 1 of uncertain significance) and a homozygous TPO deletion c.2722\_2736del, causing the loss of 5 amino acids in the intracellular domain without reading frame modification, of unknown effect (ID 45).

### Discussion

We report 103 cases of CH-GIS referred to our center for genetic testing. In 73 out of 103 patients, genetic testing found 95 variants, including 38 variants classified as “pathogenic,” 14 as “likely pathogenic,” and 28 of “uncertain significance.” Targeted NGS found a genetic explanation with solved or probably solved cases in 42% of the patients (n = 43/103), mainly in the *TG* and *TPO* genes. The molecular yield increased to 70% when TSH at neonatal screening was  $\geq$  80 mIU/L or FT4 at CH diagnosis was  $<$  5 pmol/L.

Numerous studies have described genetic variants in patients with CH-GIS. Some studied only 1 gene (38–41); others reported the frequency of the variants found in patients with CH-GIS but with no individual analysis of genetic resolution (42–48). We identified 15 studies that analyzed the rate of genetic resolution by screening at least 2 genes in CH-GIS (Table 2) (15–29). The resolution varied widely between studies, ranging from 10% to 93%. Only 3 studies included 30 patients or more with CH-GIS and analyzed at least 8 different genes, with a genetic resolution between 41% and 60% (20, 23, 25). The molecular yield in these studies was consistent with the 42% yield in the present study.

The factors that would explain this variability may be the number of genes studied, the proportion of familial cases, the severity of CH, and the classification of solved or unsolved

**Table 2. Literature review of studies that analyzed at least 2 genes in patients having CH-GIS, with the percentage of genetic resolution**

Publication	Year of publication	Country(ies)	Type of CH	Number of patients with GIS	Presence of consanguineous patients	Presence of familial cases	Number of genes analyzed	Percentage of genetic resolution in GIS	More frequently involved gene(s)
Narumi et al.	2011	Japan	GIS	14	NA	Yes	7	93%	<i>DUOX2</i> and <i>TPO</i>
Wang et al.	2014	China	GIS with goiter	67	No	No	3	13%	<i>DUOX2</i>
Jiang et al.	2016	China	GIS	12	No	No	12	92%	<i>DUOX2</i>
Lof et al.	2016	Finland	GIS and TD	21	NA	Yes	13	55% in familial cases, 20% in sporadic cases	<i>TPO</i>
Matsuo et al.	2016	Japan	GIS	48	NA	NA	3	23%	<i>DUOX2</i>
Nicholas et al.	2016	United Kingdom, Oman, Saudi Arabia, United Arab Emirates, Turkey	GIS	49	Yes	Yes	8	59%	<i>TG</i> and <i>TPO</i>
Park et al.	2016	Korea	GIS and TD	NA (among 170)	NA	NA	6	31%	<i>DUOX2</i>
Sun et al.	2018	China	GIS and TD	NA (among 110)	No	Yes	21	52%	<i>DUOX2</i>
Zou et al.	2018	Saudi Arabia	GIS and TD	30	NA	Yes	WES	60%	<i>TPO</i>
Santos-Silva et al.	2019	Portugal	GIS	9	Yes	Yes	28	67%	<i>TG</i> and <i>TPO</i>
Wang et al.	2020	China	GIS and TD	32	No	Yes	29	41%	<i>DUOX2</i>
Zdraveska et al.	2020	Macedonia	GIS and hypoplasia	34	No	Yes	2 to 9	10%	NA
Li et al.	2021	China	GIS and TD	468	NA	Yes	5	4%	<i>PAX8</i>
Shin et al.	2021	South Korea	GIS	20	NA	NA	8	80%	<i>DUOX2</i> and <i>TSHR</i>
Stoupa et al.	2021	France, USA, Turkey	GIS	19	Yes	Yes	78	53%	<i>TG</i>
Present study		France	GIS	103	Yes	Yes	48	42%	<i>TG</i> and <i>TPO</i>

Abbreviations: CH, congenital hypothyroidism; GIS, gland-in-situ; NA, not available; TD, thyroid dysgenesis.



cases between studies. First, the molecular yield was 13% to 23% when 3 genes were studied (16, 19), and this increased to 41% to 93% when at least 8 genes were studied (17, 18, 20, 22–25, 28, 29). In addition, familial cases accounted for 6% to 73% of patients in published studies (15, 18, 20, 23–26, 29), compared with 25% in our study. One study assessed genetic resolution in sporadic (20%) and familial cases (55%) (18). This compares with 36% and 58% in sporadic and familial cases in our study, respectively. None of these studies evaluated the severity of CH as a factor associated with genetic resolution, but 1 reported a median screening TSH of 23.3 mIU/L, with a genetic resolution of 41% (25), while another reported a median TSH at diagnosis of 75 mIU/L with 59% genetic resolution (20). In our study, the median TSH at screening and at diagnosis were 60 and 73 mIU/L, respectively, for a genetic resolution of 42%. These values were consistent with a similar association between CH severity and genetic resolution. Regarding the classification of solved or unsolved cases, the criteria vary between studies. We used strict criteria for the classification of solved or probably solved cases, as we classified as unsolved patients with pathogenic or probably pathogenic variants in silico when they were present only at heterozygous state, without in vitro evidence of their pathogenicity at heterozygous state. Also, variants located in the nonessential splice site were not considered as involved. This may underestimate the genetic resolution of our study.

In most European and Middle Eastern studies, the genes most frequently involved in CH-GIS have been the *TG* and *TPO* genes, in agreement with the present study (18, 20, 23, 24, 26, 29). Conversely, studies from Asia have found cases that were mainly explained by *DUOX2* variants (15–17, 19, 21, 22, 25, 28), whereas *DUOX2* mutations explained none of the solved or probably solved cases in the present study. Our results may not be superimposable to those from other regions of the world, as the mutations found in the CH-GIS depend on the ethnic background. *PAX8* and *NKX2-1* mutations, which were first described in thyroid dysgenesis, have been shown to be more frequently associated with GIS in populations of patients suffering from CH (4, 11, 14, 27), in agreement with the finding in our study, which focused solely on GIS.

Our study has several limitations. The *NNT* (49) and *SLC26A7* (9, 10) genes involved in CH-GIS were not studied, nor were some of the genes involved in CH with thyroid dysgenesis, such as *GLI*-similar zinc finger protein family 3 (*GLIS3*), cell division cycle associated 8 (*CDCA8*) coding boraalin, jagged1 (*JAG1*), netrin 1 (*NTN1*) (4), and *TUBB1* (50). Another limitation of this study is the lack of functional studies for all variants. To assess the responsibility of the variants, we used the referent ACMG classification (31), together with familial segregation, in silico prediction software, location in the protein structure, previous published studies of CH with the same variants, and functional studies when available. Additional new tools for predicting protein structure from computational methods, such as the AlphaFold protein structure database, may prove useful in the future to better characterize the consequences of missense variants (51, 52). In France the median TSH at screening and diagnosis for in situ gland were 34 and 91 mIU/L, respectively, in 2016 (unpublished), whereas median TSH were 60 and 73 mIU/L, respectively, in our study: these higher values are in favor of a bias in referral to the most severe cases. However, we believe that the relationship we have shown between CH severity and

molecular yield would be true over the entire range of TSH at screening and may contribute to explain the variation in the molecular yield between studies.

In conclusion, we report 103 patients with CH-GIS in France, with a genetic resolution for 42% of them. The percentage of genetic resolution increased to 70% when TSH on neonatal screening was  $\geq 80$  mIU/L or FT4 at diagnosis was below 5 pmol/L. Even some transient forms had a molecular explanation, and long-term follow-up will be needed, especially when thyroid hormone requirements are high, such as during puberty and later during pregnancy.

## Acknowledgments

We warmly thank all the pediatricians who sent a blood sample for genetic analysis in our Reference Center for Rare Diseases of Thyroid and Hormone Receptivity.

## Data Availability

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

## Disclosures

The authors have no conflicts of interest relevant to this article to disclose.

## References

- Barry Y, Bonaldi C, Goulet V, *et al*. Increased incidence of congenital hypothyroidism in France from 1982 to 2012: a nationwide multicenter analysis. *Ann Epidemiol*. 2016;26(2):100-105.e4. <https://doi.org/10.1016/j.annepidem.2015.11.005>
- Deladoëy J, Ruel J, Giguère Y, Van Vliet G. Is the incidence of congenital hypothyroidism really increasing? A 20-year retrospective population-based study in Québec. *J Clin Endocrinol Metab*. 2011;96(8):2422-2429. <https://doi.org/10.1210/jc.2011-1073>
- Centre National de Coordination du Dépistage Néonatal. Rapport d'activité du Programmé National du dépistage néonatal, 2019. Available at: <https://depistage-neonatal.fr/espace-pro/depistage/rapport-dactivite-dnn/>
- Peters C, van Trotsenburg ASP, Schoenmakers N. Diagnosis of endocrine disease: congenital hypothyroidism: update and perspectives. *Eur J Endocrinol*. 2018;179(6):R297-R317. <https://doi.org/10.1530/EJE-18-0383>
- Liu S, Han W, Zang Y, *et al*. Identification of two missense mutations in *DUOX1* (p.R1307Q) and *DUOX1* (p.R56W) that can cause congenital hypothyroidism through impairing H<sub>2</sub>O<sub>2</sub> generation. *Front Endocrinol (Lausanne)*. 2019;10:526. <https://doi.org/10.3389/fendo.2019.00526>
- Targovnik HM, Scheps KG, Rivolta CM. Defects in protein folding in congenital hypothyroidism. *Mol Cell Endocrinol*. 2020;501:110638. <https://doi.org/10.1016/j.mce.2019.110638>
- Coscia F, Taler-Verčič A, Chang VT, *et al*. The structure of human thyroglobulin. *Nature*. 2020;578(7796):627-630. <https://doi.org/10.1038/s41586-020-1995-4>
- Le SN, Porebski BT, McCoy J, *et al*. Modelling of thyroid peroxidase reveals insights into its enzyme function and autoantigenicity. *PLoS ONE*. 2015;10(12):e0142615. <https://doi.org/10.1371/journal.pone.0142615>
- Cangul H, Liao X-H, Schoenmakers E, *et al*. Homozygous loss-of-function mutations in *SLC26A7* cause goitrous congenital hypothyroidism. *JCI Insight*. 2018;3(20):e99631. <https://doi.org/10.1172/jci.insight.99631>
- Ishii J, Suzuki A, Kimura T, *et al*. Congenital goitrous hypothyroidism is caused by dysfunction of the iodide transporter *SLC26A7*.

- Commun Biol.* 2019;2(1):270. <https://doi.org/10.1038/s42003-019-0503-6>
11. Ramos HE, Carré A, Chevrier L, *et al.* Extreme phenotypic variability of thyroid dysgenesis in six new cases of congenital hypothyroidism due to PAX8 gene loss-of-function mutations. *European J Endocrinol.* 2014;171(4):499-507. <https://doi.org/10.1530/EJE-13-1006>
  12. Krude H, Biebermann H, Göpel W, Grüters A. The gene for the thyrotropin receptor (TSHR) as a candidate gene for congenital hypothyroidism with thyroid dysgenesis. *Exp Clin Endocrinol Diabetes.* 1996;104(Suppl 4):117-120. <https://doi.org/10.1055/s-0029-1211717>
  13. Tenenbaum-Rakover Y, Almashanu S, Hess O, *et al.* Long-term outcome of loss-of-function mutations in thyrotropin receptor gene. *Thyroid.* 2015;25(3):292-299. <https://doi.org/10.1089/thy.2014.0311>
  14. Carré A, Szinnai G, Castanet M, *et al.* Five new TTF1/NKX2.1 mutations in brain-lung-thyroid syndrome: rescue by PAX8 synergism in one case. *Hum Mol Genet.* 2009;18(12):2266-2276. <https://doi.org/10.1093/hmg/ddp162>
  15. Narumi S, Muroya K, Asakura Y, Aachi M, Hasegawa T. Molecular basis of thyroid dysmorphogenesis: genetic screening in population-based Japanese patients. *J Clin Endocrinol Metab.* 2011;96(11):E1838-E1842. <https://doi.org/10.1210/jc.2011-1573>
  16. Wang F, Lu K, Yang Z, *et al.* Genotypes and phenotypes of congenital goitre and hypothyroidism caused by mutations in dual oxidase 2 genes. *Clin Endocrinol (Oxf).* 2014;81(3):452-457. <https://doi.org/10.1111/cen.12469>
  17. Jiang X, Dias JA, He X. Structural biology of glycoprotein hormones and their receptors: insights to signaling. *Mol Cell Endocrinol.* 2014;382(1):424-451. <https://doi.org/10.1016/j.mce.2013.08.021>
  18. Löf C, Patyra K, Kuulasmaa T, *et al.* Detection of novel gene variants associated with congenital hypothyroidism in a Finnish patient cohort. *Thyroid.* 2016;26(9):1215-1224. <https://doi.org/10.1089/thy.2016.0016>
  19. Matsuo K, Tanahashi Y, Mukai T, *et al.* High prevalence of DUOX2 mutations in Japanese patients with permanent congenital hypothyroidism or transient hypothyroidism. *J Pediatr Endocrinol Metab.* 2016;29(7):807-812. <https://doi.org/10.1515/jpem-2015-0400>
  20. Nicholas AK, Serra EG, Cangul H, *et al.* Comprehensive screening of eight known causative genes in congenital hypothyroidism with gland-in-situ. *J Clin Endocrinol Metab.* 2016;101(12):4521-4531. <https://doi.org/10.1210/jc.2016-1879>
  21. Park K-J, Park H-K, Kim Y-J, *et al.* DUOX2 Mutations are frequently associated with congenital hypothyroidism in the Korean population. *Ann Lab Med.* 2016;36(2):145-153. <https://doi.org/10.3343/alm.2016.36.2.145>
  22. Sun F, Zhang J-X, Yang C-Y, *et al.* The genetic characteristics of congenital hypothyroidism in China by comprehensive screening of 21 candidate genes. *Eur J Endocrinol.* 2018;178(6):623-633. <https://doi.org/10.1530/EJE-17-1017>
  23. Zou M, Alzahrani AS, Al-Odaib A, *et al.* Molecular analysis of congenital hypothyroidism in Saudi Arabia: SLC26A7 mutation is a novel defect in thyroid dysmorphogenesis. *J Clin Endocrinol Metab.* 2018;103(5):1889-1898. <https://doi.org/10.1210/jc.2017-02202>
  24. Santos-Silva R, Rosário M, Grangeia A, *et al.* Genetic analyses in a cohort of Portuguese pediatric patients with congenital hypothyroidism. *J Pediatr Endocrinol Metab.* 2019;32(11):1265-1273. <https://doi.org/10.1515/jpem-2019-0047>
  25. Wang H, Kong X, Pei Y, *et al.* Mutation spectrum analysis of 29 causative genes in 43 Chinese patients with congenital hypothyroidism. *Mol Med Rep.* 2020;22(1):297-309. <https://doi.org/10.3892/mmr.2020.11078>
  26. Zdraveska N, Kocova M, Nicholas AK, Anastasovska V, Schoenmakers N. Genetics of gland-in-situ or hypoplastic congenital hypothyroidism in Macedonia. *Front Endocrinol (Lausanne).* 2020;11:413. <https://doi.org/10.3389/fendo.2020.00413>
  27. Li L, Jia C, Li X, *et al.* Molecular and clinical characteristics of congenital hypothyroidism in a large cohort study based on comprehensive thyroid transcription factor mutation screening in Henan. *Clin Chim Acta.* 2021;518:162-169. <https://doi.org/10.1016/j.cca.2021.03.015>
  28. Shin JH, Kim HY, Kim YM, *et al.* Genetic evaluation of congenital hypothyroidism with gland in situ using targeted exome sequencing. *Ann Clin Lab Sci.* 2021;51(1):73-81.
  29. Stoupa A, Al Hage Chehade G, Chaabane R, *et al.* High diagnostic yield of targeted next-generation sequencing in a cohort of patients with congenital hypothyroidism due to dysmorphogenesis. *Front Endocrinol (Lausanne).* 2020;11:545339. <https://doi.org/10.3389/fendo.2020.545339>
  30. van Trotsenburg P, Stoupa A, Léger J, *et al.* Congenital hypothyroidism: a 2020-2021 consensus guidelines update-an ENDO-European reference network initiative endorsed by the European Society for Pediatric Endocrinology and the European Society for Endocrinology. *Thyroid.* 2021;31(3):387-419. <https://doi.org/10.1089/thy.2020.0333>
  31. Richards S, Aziz N, Bale S, *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med.* 2015;17(5):405-424. <https://doi.org/10.1038/gim.2015.30>
  32. Karczewski KJ, Francioli LC, Tiao G, *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature.* 2020;581(7809):434-443. <https://doi.org/10.1038/s41586-020-2308-7>
  33. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res.* 2010;20(1):110-121. <https://doi.org/10.1101/gr.097857.109>
  34. Léger J, Olivieri A, Donaldson M, *et al.* European Society for Paediatric Endocrinology consensus guidelines on screening, diagnosis, and management of congenital hypothyroidism. *Horm Res Paediatr.* 2014;81(2):80-103. <https://doi.org/10.1159/000358198>
  35. Szinnai G. Clinical genetics of congenital hypothyroidism. *Endocr Dev.* 2014;26:60-78. <https://doi.org/10.1159/000363156>
  36. Sobrero G, Muñoz L, Bazzara L, *et al.* Thyroglobulin reference values in a pediatric infant population. *Thyroid.* 2007;17(11):1049-1054. <https://doi.org/10.1089/thy.2007.0059>
  37. Levallant L. Supplemental data for: Congenital hypothyroidism with gland-in-situ - supplemental material. <https://doi.org/10.7910/DVN/BREA97>. Date of deposit 07 March 2023.
  38. Jin HY, Heo S-H, Kim Y-M, *et al.* High frequency of DUOX2 mutations in transient or permanent congenital hypothyroidism with eutopic thyroid glands. *Horm Res Paediatr.* 2014;82(4):252-260. <https://doi.org/10.1159/000362235>
  39. Muzza M, Rabbiosi S, Vigone MC, *et al.* The clinical and molecular characterization of patients with dysmorphogenic congenital hypothyroidism reveals specific diagnostic clues for DUOX2 defects. *J Clin Endocrinol Metab.* 2014;99(3):E544-E553. <https://doi.org/10.1210/jc.2013-3618>
  40. Hu X, Chen R, Fu C, *et al.* Thyroglobulin gene mutations in Chinese patients with congenital hypothyroidism. *Mol Cell Endocrinol.* 2016;423:60-66. <https://doi.org/10.1016/j.mce.2016.01.007>
  41. Rodrigues C, Jorge P, Soares JP, *et al.* Mutation screening of the thyroid peroxidase gene in a cohort of 55 Portuguese patients with congenital hypothyroidism. *Eur J Endocrinol.* 2005;152(2):193-198. <https://doi.org/10.1530/eje.1.01826>
  42. de Filippis T, Gelmini G, Paraboschi E, *et al.* A frequent oligogenic involvement in congenital hypothyroidism. *Hum Mol Genet.* 2017;26(13):2507-2514. <https://doi.org/10.1093/hmg/ddx145>
  43. Fan X, Fu C, Shen Y, *et al.* Next-generation sequencing analysis of twelve known causative genes in congenital hypothyroidism. *Clin*

- Chim Acta*. 2017;468:76-80. <https://doi.org/10.1016/j.cca.2017.02.009>
44. Makretskaya N, Bezlepina O, Kolodkina A, *et al*. High frequency of mutations in “dysmorphogenesis genes” in severe congenital hypothyroidism. *PLoS ONE*. 2018;13(9):e0204323. <https://doi.org/10.1371/journal.pone.0204323>
45. Bruellman RJ, Watanabe Y, Ebrahim RS, *et al*. Increased prevalence of TG and TPO mutations in Sudanese children with congenital hypothyroidism. *J Clin Endocrinol Metab*. 2020;105(5):1564-1572. <https://doi.org/10.1210/clinem/dgz297>
46. Wang F, Zang Y, Li M, *et al*. DUOX2 And DUOX2 variants confer susceptibility to thyroid dysgenesis and gland-in-situ with congenital hypothyroidism. *Front Endocrinol (Lausanne)*. 2020;11:237. <https://doi.org/10.3389/fendo.2020.00237>
47. Wang F, Xiaole L, Ma R, Zhao D, Liu S. Dual oxidase (DUOX) system genes defects in children with congenital hypothyroidism. *Endocrinology*. 2021;162(8):bqab043. <https://doi.org/10.1210/endo/bqab043>
48. Oliver-Petit I, Edouard T, Jacques V, *et al*. Next-generation sequencing analysis reveals frequent familial origin and oligogenism in congenital hypothyroidism with dysmorphogenesis. *Front Endocrinol (Lausanne)*. 2021;12:657913. <https://doi.org/10.3389/fendo.2021.657913>
49. Roucher-Boulez F, Mallet-Motak D, Samara-Boustani D, *et al*. NNT mutations: a cause of primary adrenal insufficiency, oxidative stress and extra-adrenal defects. *Eur J Endocrinol*. 2016;175(1):73-84. <https://doi.org/10.1530/EJE-16-0056>
50. Stoupa A, Adam F, Kariyawasam D, *et al*. TUBB1 mutations cause thyroid dysgenesis associated with abnormal platelet physiology. *EMBO Mol Med*. 2018;10(12):e9569. <https://doi.org/10.15252/emmm.201809569>
51. Jumper J, Evans R, Pritzel A, *et al*. Highly accurate protein structure prediction with AlphaFold. *Nature*. 2021;596(7873):583-589. <https://doi.org/10.1038/s41586-021-03819-2>
52. Varadi M, Anyango S, Deshpande M, *et al*. AlphaFold protein structure database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res*. 2022;50(D1):D439-D444. <https://doi.org/10.1093/nar/gkab1061>