

## Carotenoids bioproduction by two psychrophilic microalgae: Sphaerocystis sp. and Pleurastrum sp.

Adila GHERABLI<sup>1,2\*</sup>, Madeleine Charbonnier<sup>2</sup>, Fanny Duval<sup>2</sup>, Julien Lemaire<sup>2</sup>, Nabil Grimi<sup>1</sup> <sup>1</sup>Université de Technologie de Compiègne, ESCOM, TIMR (Integrated Transformations of Renewable Matter), Centre de recherche Royallieu - CS 60319 - 60203 Compiègne, Cedex, France. <sup>2</sup>Université Paris-Saclay, CentraleSupélec, Laboratoire de Génie des Procédés et Matériaux, Centre Européen de Biotechnologie et de Bioéconomie (CEBB), 3 rue des Rouges Terres 51110 Pomacle, France.

## Abstract

Among the valuable compounds derived from microalgae, carotenoids are the most studied. Carotenoids in plants, algae, and some photosynthetic bacteria and yeasts are responsible for their red, orange, and yellow pigmentation. Indeed, natural carotenoids receive significant interest because they are potent antioxidants and can protect against certain diseases such as cancer. Therefore, carotenoids derived from microalgae are valued for their diverse applications in food, feed, cosmetic, and pharmaceutical industries.

In recent decades, the carotenoids industry has searched for sources other than *Haematococcus pluvialis* microalga. This mesophilic microalga (temperature growth between 25°- 28° C) represents the most abundant source of natural astaxanthin, a well-known antioxidant and one of the most expensive carotenoids on the market. However, few studies have been conducted on extremophilic microalgae strains that can produce valuable carotenoids at low temperatures and light. The cultivation of microalgae requires artificial lighting to mimic natural sunlight. These lights consume a substantial electricity input. In addition, maintaining the ideal temperature and climate for microalgae growth involves heating or cooling systems, which further contribute to energy consumption. In regions where energy costs are high or renewable sources are limited, the cost of energy consumption can become prohibitive. Hence, studies were performed to find microalgae strains that are more tolerant or adapted. In this context, psychrophilic microalgae represent potent candidates because they are well-adapted to harsh environments in cold-climate regions. Moreover, these photosynthetic organisms cause red, orange, yellow, or brown pigmentations when snow fields are subject to extensive irradiation, high temperature, and low nutrient availability (melting season).

This project is devoted to optimizing carotenoid bioaccumulation by two selected psychrophilic strains *Sphaerocystis sp.* and *Pleurastreum sp.* Both are successfully acclimated to our laboratory conditions for the growth and accumulation of carotenoids. The culture was carried out through two stages (green and red), where parameters such as temperature, lighting, and nutritional medium were carefully monitored. Therefore, the first interesting results of culture pigmentation, cell stages transitioning, and pigment content of Chlorophyll *a*, Chlorophyll *b*, and total carotenoids are presented in this work.

## References

- [1] C. Aflalo, Y. Meshulam, A. Zarka, and S. Boussiba, *Biotechnology and Bioengineering*. 98 (2007) 300–305.
- [2] T. Leya, A. Rahn, C. Lutz, and D. Remias, *FEMS Microbiology Ecology*. 67 (2009) 432–443.
- [3] F. Pagels, V. Vasconcelos, and A. C. Guedes, *Biomolecules*. 11 (2021) 1-21
- [4] A. Gherabli, N. Grimi, J. Lemaire, E. Vorobiev, and N. Lebovka, *Molecules*. 28 (2023) 1-19.



Adila GHERABLI is a third-year Ph.D. student at the University of Technology of Compiègne (TIMR laboratory). She is working on biotransformation and microalgae processing in collaboration with the University of Paris-Saclay, CentraleSupélec (LGPM, CEBB).

- Full Name: Adila GHERABLI
- Email: adila.gherabli@utc.fr
- Mobile Number: +33 699 55 13 37
- Category: Poster
- Postal Address: Centre de recherche Royallieu CS 60319 60203 Compiègne, Cedex, France.