



Increased drought of the mediterranean climate and biotic interactions: functional consequences on plant communities in a shrubland ecosystem

Natalia Rodriguez-Ramirez, Nicolas Montès, Christine Ballini

► To cite this version:

Natalia Rodriguez-Ramirez, Nicolas Montès, Christine Ballini. Increased drought of the mediterranean climate and biotic interactions: functional consequences on plant communities in a shrubland ecosystem. Plant Biology Europe FESPB/EPSO Congress, Jun 2014, Dublin, Ireland. hal-04453930

HAL Id: hal-04453930


<https://hal.science/hal-04453930>

Submitted on 12 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Increased drought of the mediterranean climate and biotic interactions: functional consequences on plant communities in a shrubland ecosystem

Aix-Marseille
UniversitéNatalia RODRIGUEZ-RAMIREZ^{1*}, Nicolas MONTES¹, Christine BALLINI¹


1- Aix-Marseille Université, Institut Méditerranéen de Biodiversité et Ecologie marine et continentale-IMBE



Introduction

Climate Change is predicted to be particularly pronounced on the Mediterranean basin. With up to 40% decrease in summer precipitation and temperature 2° C to 5° C higher by 2080-2099 than in the 1980-1999 (EEA, 2012), important changes on biodiversity and plant-plant interactions are expected on this century (Lloret et al. 2009, Quézel et Médail, 2003).

The aim of the **CLIMED project** (*Climate change effects on Mediterranean biodiversity and consequences for ecosystem functioning*) is to link predicted the decrease of precipitation to changes in biodiversity across multiple trophic levels (plants, litter, soil and their fauna), and quantify their feedback on the ecosystem functioning and environmental services using an experimental approaches



How will the projected diminution of precipitations (-40%) affect the Mediterranean Schrubland vegetation ?

More precisely what effects on:

► Diversity and community structure

- Diversity indices
- Non destructive quantification of biomass production
- Interspecific relationships

► Functional traits and functional diversity:


- Resistance to drought and recovery capacity
- Phenology
- Foliar traits (SLA, DMC C, N and P contents)

► Ecophysiology

- Water Use Efficiency
- Maximum quantum efficiency (Fv/Fm)

► Ecosystem functions: primary production, C and N cycles


Some Results




Mean of shoots with aboveground biomass totally drought, by plot, of the 4 dominant ligneous species in CLIMED for each level of precipitation

- The resistance to total drought of the aboveground biomass depends of the specie and the season
- R. officinalis* seems more resistance to drought than the other dominant species.
- With the decrease precipitation *C. albidus* and *U. parviflorus* are the most vulnerable to total drought of the dominant species

CLIMED Rain Exclusion Experiment



Mediterranean shrubland (2 ha), North of Marseille (280 m)



- 93 plots (4x4m, each):
• 46 with rain exclusion device in « Exclusion »
• 47 with rain exclusion device in « Control »



Over,

« natural » mixtures of 1, 2 ,3 or 4 of the ligneous dominant species:



References:

European Environment Agency (EEA) (2012). Climate change, impacts and vulnerability in Europe
Lloret F et al. (2009) Perspectives in Plant Ecology Evolution and Systematics 11: 53-63.
Médail, F. & Quézel, P. (2003) Bocconea 16(1): 397-422.



Correlation matrices of the cover of the 4 dominant species and diversity by plot, ordered and colored by the Spearman correlation (Rho « p ») magnitude:

CA = *C. albidus*, QC= *Q. coccifera*, RO= *R. officinalis*, UP= *U. parviflorus*

■ significative correlations (p-value <0.05)

- The positively correlation between diversity (H') and *U. parviflorus* cover is almost the same no matter the precipitation level
- The magnitude of the negative correlation between the covers of *Q. coccifera* and *C. albidus* becomes stronger under rain exclusion
- The negative correlation between the covers of *C. albidus* and *R. officinalis* becomes significant under rain exclusion.

Discussion

- Under the precipitation decrease (-40%) projected for the south of France by the end of the century, the diversity and structure of the mediterranean shrubland vegetation could be modified.
- Under higher drought conditions the cover of *C. albidus* and *U. parviflorus* could sharply decrease
- The vulnerability of *U. parviflorus*, a legume, to the decrease of precipitation could have an effect on nitrogen cycle
- The tendency to total drought of *U. parviflorus* with the precipitation decrease, could affect the total diversity of the community as it's positively correlated to it.
- The interspecific relationships in the mediterranean shrubland community seems to be modified with the decrease of water availability. In particular the negative relationships between dominant species could be stronger under a more drought environment.

* natalia.rodriguez@imbe.fr

Aix Marseille Université-Campus Centre, Case 4, 13003 Marseille

Increased drought of the mediterranean climate and biotic interactions: functional consequences on plant communities in a shrubland ecosystem

Natalia RODRIGUEZ-RAMIREZ^{1*}, Nicolas MONTES¹, Christine BALLINI¹


1- Aix-Marseille Université, Institut Méditerranéen de Biodiversité et Ecologie marine et continentale-IMBE



Introduction

Climate Change is predicted to be particularly pronounced on the Mediterranean basin. With up to 40% decrease in summer precipitation and temperature 2° C to 5° C higher by 2080-2099 than in the 1980-1999 (EEA,2012), important changes on biodiversity and plant-plant interactions are expected on this century (Lloret *et al.* 2009, Quézel et Médail, 2003).

The aim of the **CLIMED project** (*Climate change effects on Mediterranean biodiversity and consequences for ecosystem functioning*) is to link predicted the decrease of precipitation to changes in biodiversity across multiple trophic levels (plants, litter, soil and their fauna), and quantify their feedback on the ecosystem functioning and environmental services using an experimental approaches



How will the projected diminution of precipitations (-40%) affect the Mediterranean Schrubland vegetation ?

More precisely what effects on:

► Diversity and community structure

- Diversity indices
- Non destructive quantification of biomass production
- Interspecific relationships

► Functional traits and functional diversity:


- Resistance to drought and recovery capacity
- Phenology
- Foliar traits (SLA, DMC C, N and P contents)

► Ecophysiology

- Water Use Efficiency
- Maximum quantum efficiency (Fv/Fm)


► Ecosystem functions: primary production, C and N cycles

Some Results




- The resistance to total drought of the aboveground biomass depends of the specie and the season
- R. officinalis* seems more resistance to drought than the other dominant species.
- With the decrease precipitation *C. albidus* and *U. parviflorus* are the most vulnerable to total drought of the dominant species

CLIMED Rain Exclusion Experiment




Mediterranean shrubland (2 ha), North of Marseille (280 m)



- 93 plots (4x4m, each):
• 46 with rain exclusion device in « Exclusion »
• 47 with rain exclusion device in « Control »


Over,

« natural » mixtures of 1, 2 ,3 or 4 of the ligneous dominant species:



References:

European Environment Agency (EEA) (2012). Climate change, impacts and vulnerability in Europe
Lloret F *et al.* (2009) Perspectives in Plant Ecology Evolution and Systematics 11: 53-63.
Médail, F. & Quézel, P. (2003) Bocconea 16(1): 397-422.



Correlation matrices of the cover of the 4 dominant species and diversity by plot, ordered and colored by the Spearman correlation (Rho « ρ ») magnitude:
 $\rho > 0 > \rho > \rho$

- The positively correlation between diversity (H') and *U. parviflorus* cover is almost the same no matter the precipitation level
• The magnitude of the negative correlation between the covers of *Q. Coccifera* and *C. albidus* becomes stronger under rain exclusion
• The negative correlation between the covers of *C. albidus* and *R. officinalis* becomes significant under rain exclusion.

Discussion

- Under the precipitation decrease (-40%) projected for the south of France by the end of the century, the diversity and structure of the mediterranean shrubland vegetation could be modified.
- Under higher drought conditions the cover of *C. albidus* and *U. parviflorus* could sharply decrease
- The vulnerability of *U. parviflorus*, a legume, to the decrease of precipitation could have an effect on nitrogen cycle
- The tendency to total drought of *U. parviflorus* with the precipitation decrease, could affect the total diversity of the community as it's positively correlated to it.
- The interspecific relationships in the mediterranean shrubland community seems to be modified with the decrease of water availability. In particular the negative relationships between dominant species could be stronger under a more drought environment.

* natalia.rodriguez@imbe.fr



Aix Marseille Université-Campus Centre, Case 4, 13003 Marseille

En relation avec le suivie de la nécromasse. Anova factoriel avec es données de tous les mois (mois as a factor)					
	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Ttmt	1	37	36.9	10.629	0.00113 **
SP	3	2085	694.9	200.083	< 2e-16 ***
Mois	13	317	24.4	7.016	1.47e-13 ***
Ttmt:SP	3	31	10.5	3.010	0.02908 *
Ttmt:Mois	13	27	2.1	0.602	0.85413
SP:Mois	39	445	11.4	3.285	4.26e-11 ***
Ttmt:SP:Mois	39	49	1.2	0.360	0.99992
Residuals	2702	9384	3.5		

Signif. codes:	0 ***	0.001 **	0.01 *	0.05 .	
	0.1 ' ,	1			

En relation avec le suivie de la nécromasse. Anova factoriel avec es données de tous les mois (mois as a factor)

results QC traits?



What's next:
Bras, isotopes analisis, correaltion with other ecosystems trophic levels

Increased drought of the mediterranean climate and biotic interactions: functional consequences on plant communities in a shrubland ecosystem

Natalia RODRIGUEZ-RAMIREZ^{1*}, Nicolas MONTES¹, Christine BALLINI¹


1- Aix-Marseille Université, Institut Méditerranéen de Biodiversité et Ecologie marine et continentale-IMBE



Introduction

Le bassin méditerranéen est considéré comme une région particulièrement vulnérable face au changement climatique, avec une diminution jusqu'à 40% des précipitations estivales et une augmentation des températures de 2,5° C à 4° C en 2071-2100 par rapport à la période 1961-1990, selon les prévisions (EEA, 2012). On s'attend donc à des changements importants de la biodiversité dans le courant du siècle (Lloret *et al.* 2009, Quézel et Médail, 2003).

Le projet CLIMED (*Climate change effects on Mediterranean biodiversity and consequences for ecosystem functioning, ANR CEP 2009*) a pour objectif d'évaluer l'effet de la sécheresse accrue sur la biodiversité de la garrigue (végétation, litière, sol et faune associée), le fonctionnement de l'écosystème et les services environnementaux




Changements (%) projetés des précipitations annuelles (A) et estivales (B) entre 1961-1990 et 2071-2100. Source: EEA, 2012

Principales questions:

Sur la végétation, quel sera l'effet de la sécheresse accrue sur:


- Relations interspécifiques fonctionnelle:
- Traits fonctionnels et structuraux: S: phénologie, spore, résistance à l'assèchement, la récupération
- Diversité communautaire: morphologiques et chimiques
- Fonctions de l'écosystème: Productivité Primaire, cycles du carbone et de l'azote

Premiers résultats



Nombre moyen de pieds secs (+/- s.d.) par placette et par mois des espèces dominantes - Septembre 2012 à Mars 2014

Site d'étude




Garrigue localisée sur la Chaîne de l'Etoile (280 m) au Nord de Marseille





Rain Exclusion devices

- 93 plots (2x2 m each):
- 46 structures excluant environ 40% des précipitations
 - 47 structures en « Contrôle »



Lignous dominant species : *Cistus albidus*, *Quercus coccifera*, *Rosmarinus officinalis*, *Ulex Parviflorus*



NB: L'encadrement indique que la corrélation est significative ($p < 0.05$). Les matrices sont ordonnées et colorées par la magnitude de la corrélation de Spearman (Rho) obtenues dans chaque cas: les corrélations jaunes sont relativement plus faibles que les vertes et celle-ci que les rouges

- La magnitude des corrélations entre les recouvrements d'une paire d'espèces ou le recouvrement d'une espèce et la diversité de Shannon-Weaver (H') varie dans certains cas avec le niveau de pluie
- Les recouvrements de *Q. coccifera* et *C. albidus* sont dans les deux cas négativement corrélés et la magnitude de cette corrélation est plus élevée sous exclusion de pluie.
- H' est négativement corrélée au recouvrement du *Q. coccifera* quelque soient les conditions de précipitation.
- H' est positivement corrélée avec le recouvrement des autres espèces dominantes mais sous forte sécheresse cette relation est seulement significative avec *U. parviflorus*

Discussion

- La mortalité de la partie aérienne dépend de l'espèce.
- La diminution des précipitations paraît affecter plus fortement la biomasse de *C. albidus* et *U. parviflorus*.
- La partie aérienne de *R. officinalis* paraît la plus résistante face à la diminution des précipitations et semble se récupérer après desséchement total.
- En général, *U. parviflorus* (légumineuse) paraît l'espèce dominante la plus sensible à l'assèchement. Ceci pourrait avoir des effets sur la fixation de N₂.
- Sous exclusion de pluie, la diversité est positivement corrélée au recouvrement d'*U. parviflorus*. Or sous ces conditions, la mortalité de cette espèce augmente. En conséquence, on pourrait donc s'attendre à une diminution de la diversité.
- Le recouvrement de *Q. coccifera* est négativement corrélé à la diversité de la végétation. Or, sous sécheresse accrue sa partie aérienne est moins affectée que d'autres espèces dominantes et semble se récupérer après desséchement total. On pourrait alors s'attendre à un déclin de la diversité de la végétation de garrigue là où *Q. coccifera* est dominant.

Références:

- European Environment Agency (EEA) . 2012. Climate change, impacts and vulnerability in Europe 2012
Lloret F *et al.* (2009). Perspectives in Plant Ecology Evolution and Systematics 11: 53-63.
Médail, F. & Quézel, P. (2003). Bocconeia 16(1): 397-422.

* natalia.rodriguez@imbe.fr

Aix Marseille Université-Campus Centre, Case 4, 13003 Marseille