



**HAL**  
open science

# Ammonia ices under low-energy electron irradiation - Towards quantitative desorption rates and effective cross sections

L Sala, L Amiaud, C Dablemont, A Lafosse

## ► To cite this version:

L Sala, L Amiaud, C Dablemont, A Lafosse. Ammonia ices under low-energy electron irradiation - Towards quantitative desorption rates and effective cross sections. *Journal of Physics: Conference Series*, 2020, 31st International Conference on Photonic, Electronic and Atomic Collisions (ICPEAC XXXI) 23-30 July 2019, Deauville, France, 1412, pp.202003. 10.1088/1742-6596/1412/20/202003 . hal-04455171

**HAL Id: hal-04455171**

**<https://hal.science/hal-04455171v1>**

Submitted on 13 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PAPER • OPEN ACCESS

## Ammonia ices under low-energy electron irradiation - Towards quantitative desorption rates and effective cross sections

To cite this article: L Sala *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 202003

View the [article online](#) for updates and enhancements.

You may also like

- [Rotational Desorption of Ice Mantles from Suprathermally Rotating Grains around Young Stellar Objects](#)  
Thiem Hoang and Le Ngoc Tram
- [Surface Chemical Information from Electron- and Photon-Stimulated Desorption](#)  
M L Knotek
- [Chemistry on Rotating Grain Surfaces: Rotational Desorption of Molecules from Ice Mantles](#)  
Thiem Hoang and Ngo-Duy Tung

**PRIME**  
PACIFIC RIM MEETING  
ON ELECTROCHEMICAL  
AND SOLID STATE SCIENCE

HONOLULU, HI  
Oct 6–11, 2024

Abstract submission deadline:  
**April 12, 2024**

Learn more and submit!

**Joint Meeting of**  
The Electrochemical Society  
•  
The Electrochemical Society of Japan  
•  
Korea Electrochemical Society

## Ammonia ices under low-energy electron irradiation - Towards quantitative desorption rates and effective cross sections

L Sala<sup>1</sup>, L Amiaud<sup>1</sup>, C Dablemont<sup>1</sup> and A Lafosse<sup>1\*</sup>

<sup>1</sup> Institut des Sciences Moléculaires d'Orsay (ISMO), CNRS, Univ Paris Sud, Université Paris-Saclay, Orsay, F-91405, France


**Synopsis** Low-energy (< 20 eV) electron processing of NH<sub>3</sub> multilayer films, systems of interest in astrochemistry, leads to significant NH<sub>3</sub> desorption, fragmentation, and to a complex chemistry arising from recombination and secondary processes induced by reactive products. Quantitative desorption rates of neutral species and effective cross sections for selected processes were obtained by combining three surface-sensitive techniques: High Resolution Electron-Energy Loss Spectroscopy (HREELS), Temperature Programmed Desorption (TPD) and Electron Stimulated Desorption (ESD) of neutral species.

NH<sub>3</sub> is known to be a major component in ice mantles in molecular clouds with up to ~15 % of H<sub>2</sub>O in certain regions of the interstellar medium [1]. Model predictions cannot account for the abundance of NH<sub>3</sub> [2], suggesting that desorption or conversion to other compounds should be considered. Among the various thermal and non-thermal mechanisms responsible [3], this study is dedicated to processes induced by low energy electrons (LEE, energy < 20 eV) [4]. LEE interactions with matter are considered to be the primary driving force in many radiation-induced desorption and chemical reactions in ices [5].

As schematized in figure 1, multilayers of NH<sub>3</sub> isotopologues (<sup>14</sup>NH<sub>3</sub>, <sup>15</sup>NH<sub>3</sub>, and <sup>14</sup>ND<sub>3</sub>) (~10 ML) were deposited on copper at 28 K and crystallized under annealing at 60 K. High-Resolution Electron Energy Loss Spectroscopy (HREELS) and Temperature Programmed Desorption (TPD) were used to control the film morphology and estimate film thickness, respectively. LEE irradiation (1-20 eV) was performed at 28 and 60 K. Electron Stimulated Desorption (ESD) was used to detect neutral fragments/molecules desorbing during irradiation while HREELS and TPD were used to characterize processed residues.

The desorption of intact NH<sub>3</sub> was observed which resulted in the direct erosion of the film proceeding through a mechanism consistent with desorption processes induced by electronic transition (DIET). Different fragmentation and recombination processes were also observed as evidenced by the detection of neutral radicals NH<sub>x</sub> (x = 1,2), and sta-

ble molecules N<sub>2</sub> and H<sub>2</sub>. Aside from desorption, a wealth of chemical processes was also observed at 13 eV. Temporal ESD at this energy allowed for the estimation of the effective cross section of NH<sub>3</sub> desorption and observing the delayed desorption of N<sub>2</sub> and H<sub>2</sub>. TPD analysis of the residues also provided evidence of diazene N<sub>2</sub>H<sub>2</sub> and hydrazine N<sub>2</sub>H<sub>4</sub> synthesis in the film.



**Figure 1.** Scheme of the procedure used to study electron-induced processes in molecular layers of ammonia and its isotopologues.

### References

- [1] Jones A P and Williams D A 1984 *Mon. Not. R. Astron. Soc.* **209** 955
- [2] Brown P D and Charnley S B 1990 *Mon. Not. R. Astron. Soc.* **244** 432
- [3] Bertin M *et al* 2013 *The Astrophysical Journal* **779** 120
- [4] Sala L A 2018 *PhD Thesis*, Université Paris-Sud 11: Orsay, France
- [5] Boamah M D *et al* 2012 *Faraday Discussion* **168** 249

\* E-mail: [anne.lafosse@u-psud.fr](mailto:anne.lafosse@u-psud.fr)