



HAL
open science

Language Kleene Algebra with Complement: A Finite Relational Semantics and (Un)decidability

Yoshiki Nakamura

► **To cite this version:**

Yoshiki Nakamura. Language Kleene Algebra with Complement: A Finite Relational Semantics and (Un)decidability. 2024. hal-04455882v1

HAL Id: hal-04455882

<https://hal.science/hal-04455882v1>

Preprint submitted on 13 Feb 2024 (v1), last revised 13 Jul 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Language Kleene Algebra with Complement: A Finite Relational Semantics and (Un)decidability

Yoshiki Nakamura  

Tokyo Institute of Technology, Japan

Abstract

We study the equational theory of Kleene algebra with complement w.r.t. languages. While the equational theory w.r.t. languages coincides with the language equivalence (under the standard language valuation) for terms of Kleene algebra, this coincidence is broken if we extend the terms with complement. In this paper, we present a *finite relational semantics*, which completely characterizes the equational theory above. As applications, we show that the equational theory is (1) undecidable and Π_1^0 -complete for full terms; (2) decidable and PSPACE-complete if the complement only applies to variables or constants.

2012 ACM Subject Classification Theory of computation \rightarrow Equational logic and rewriting; Theory of computation \rightarrow Logic and verification; Theory of computation \rightarrow Formal languages and automata theory

Keywords and phrases Kleene algebra, Relational model, Complexity

1 Introduction

Kleene algebra (KA) [12, 7] is an algebraic system for regular expressions consisting of union (\cup), composition (\cdot), Kleene-star ($_*$), empty (\perp), and identity ($\mathbb{1}$). In this paper, we consider the [equational theory w.r.t. languages](#) for extended KA terms. We write $\text{LANG} \models t = s$ if $t = s$ holds on all models of [languages](#), i.e., for any sets X , when each [letter](#) maps to any [languages](#) over X , the two expressions, t and s , have the same [language](#). By the completeness of KA, for (non-extended) KA terms, the [equational theory w.r.t. languages](#) coincides¹ with the language equivalence under the standard language valuation (i.e., each x maps to the singleton language $\{x\}$): for all KA terms t, s , we have

$$[t] = [s] \quad \Leftrightarrow \quad \text{LANG} \models t = s \quad (\dagger\text{-KA})$$

where $[u]$ denotes the [language](#) of an expression u under the standard language valuation.

The equivalence above is broken in general if we extend KA terms with some operators. The [equational theory w.r.t. languages](#) for KA terms with some operators was studied, e.g., with reverse [3], with tests [14] (where languages are of guarded strings, not words), with intersection (\cap) [2], with universality (\top) [28, 21], with variable complements (\bar{x}) [19], and combinations of some of them [4, 5]. For example, for (variable) complements, Equation (\dagger -KA) fails, e.g., by the following counter-example [19]:²

$$[\bar{x}] = [\bar{x} \cdot \bar{x}] \quad \text{LANG} \not\models \bar{x} = \bar{x} \cdot \bar{x}.$$

LANG is more compatible with equational reasoning, in that valid [equations](#) in LANG are preserved under substitution (by definition), while those under the language equivalence are not: $[\bar{x}] = \{x\} \neq \{xx\} = [\bar{x} \cdot \bar{x}]$.

Extending with full complement significantly enhances the expressive power. For example, we can define \top and \cap using complement: $\top = \bar{\perp}$ and $t \cap s = \bar{t} \cup \bar{s}$. Moreover, we can encode [quantifier-free formulas](#) (including hypotheses as Horn formulas) by [equations](#) (Sect. 4).

¹ By the completeness theorem of Kleene algebra (e.g., [13]); see also, e.g., [2, 21] and [19, Appendix A].

² $[\bar{x}] = \mathbf{V}^* \setminus \{x\} = [\bar{x} \cdot \bar{x}]$ where \mathbf{V} is the alphabet. For $\text{LANG} \not\models \bar{x} = \bar{x} \cdot \bar{x}$: By the [valuation](#) $x \mapsto \mathbf{V}^* \setminus \{x\}$.

43 In this paper, we present a *finite relational semantics* for KA with complement w.r.t.
 44 languages: [relational subword models](#) RSUB (Sect. 3), which ease us to study the [equational](#)
 45 [theory w.r.t. languages](#), relationally. Our relational semantics can more robustly characterize
 46 the [equational theory](#) than known relational semantics, e.g., for KAs (REL) [22, third page]
 47 and for KAs with \top (GREL) [28, 21] (Remark 3.11). As applications, by transforming the
 48 techniques in [18] from relational models REL to RSUB, we have the following complexity
 49 results: the [equational theory w.r.t. languages](#) is

- 50 1. Π_1^0 -complete for KA terms with full complement (Cor. 4.8 and Thms. 4.7 and 4.10);
 - 51 2. PSPACE-complete for KA terms with variable and constant complements (Thm. 6.10).
- 52 The first result gives a natural example that the [equational theory w.r.t. languages](#) is
 53 undecidable, while the standard language equivalence is decidable (by a standard automata
 54 construction). The second result positively settles the open problem posed in [19].

55 This paper is structured as follows. In Sect. 2, we give basic definitions. In Sect. 3, we
 56 give RSUB and show that the [quantifier-free theory](#) of LANG coincides with that of RSUB.
 57 In Sect. 4, we give a reduction from the [quantifier-free theory](#) into the [equational theory](#)
 58 and show that the [equational theory](#) is Π_1^0 -complete for KA terms with full complement. In
 59 Sect. 5, by using RSUB, we give a graph characterization for KA terms with variable and
 60 constant complements. In Sect. 6, we show that the [equational theory](#) for KA terms with
 61 variable and constant complements is PSPACE-complete. In Sect. 7, we conclude this paper.

62 2 Preliminaries

63 We write \mathbb{N} for the set of non-negative integers. For $l, r \in \mathbb{N}$, we write $[l, r]$ for the set
 64 $\{i \in \mathbb{N} \mid l \leq i \leq r\}$. For a set X , we write $\wp(X)$ for the power set of X .

65 For a set X (of *letters*), we write X^* for the set of *words* over X . We write ϵ for the
 66 empty word. We write wv for the [concatenation](#) of words w and v . A *language* over X is a
 67 subset of X^* . We use w, v to denote [words](#) and use L, K to denote [languages](#), respectively.
 68 For [languages](#) $L, K \subseteq X^*$, the [concatenation](#) $L \cdot K$ and the [Kleene star](#) L^* is defined by:


$$69 \quad L \cdot K \triangleq \{wv \mid w \in L \wedge v \in K\} \quad L^* \triangleq \{w_0 \dots w_{n-1} \mid \exists n \in \mathbb{N}, \forall i < n, w_i \in L\}.$$

71 A [\(2-pointed\) graph](#) G over a set A is a tuple $\langle |G|, \{a^G\}_{a \in A}, 1^G, 2^G \rangle$, where $|G|$ is a
 72 non-empty set (of vertices), each $a^G \subseteq |G|^2$ is a binary relation, and $1^G, 2^G \subseteq |G|$ are
 73 vertices. Let G, H be [graphs](#) over a set A . For a map $f: |G| \rightarrow |H|$, we say that f is a [graph](#)
 74 [homomorphism](#) from G to H , written $f: G \rightarrow H$ if for all x, y , and a , $\langle x, y \rangle \in a^G$ implies
 75 $\langle f(x), f(y) \rangle \in a^H$, $f(1^G) = 1^H$, and $f(2^G) = 2^H$. We say that f is a [graph isomorphism](#) from
 76 G to H if f is bijective and for all x, y , and a , $\langle x, y \rangle \in a^G$ iff $\langle f(x), f(y) \rangle \in a^H$, $f(1^G) = 1^H$,
 77 and $f(2^G) = 2^H$. We say that H is an [edge-extension](#) of G if $|H| = |G|$, $a^H \supseteq a^G$ for all a ,
 78 $1^H = 1^G$, and $2^H = 2^G$. For a set $\{1^G, 2^G\} \subseteq X \subseteq |G|$, the [induced subgraph](#) of G on X is the
 79 [graph](#) $\langle X, \{a^G \cap (X \times X)\}_{a \in A}, 1^G, 2^G \rangle$. For an equivalence relation E , the [quotient graph](#) of G
 80 w.r.t. E is the [graph](#) $G/E \triangleq \langle |G|/E, \{\langle X, Y \rangle \mid \exists x \in X, y \in Y, \langle x, y \rangle \in a^G\}_{a \in A}, [1^G]_E, [2^G]_E \rangle$
 81 where X/E denotes the set of equivalence classes of X by E and $[x]_E$ denotes the equivalence
 82 class of x . Additionally, we use the following operation:

83 ► **Definition 2.1.** For a [graph homomorphism](#) $h: G \rightarrow H$ (G, H over a set A), the [edge-](#)
 84 [saturation](#) of G w.r.t. h is the [graph](#) $\mathcal{S}(h) \triangleq \langle |G|, \{\langle \langle x, y \rangle \mid \langle h(x), h(y) \rangle \in a^H \rangle\}_{a \in A}, 1^G, 2^G \rangle$.

85 ► **Example 2.2.** Let $h: G \rightarrow H$ be the [graph homomorphism](#) indicated by green colored
 86 arrows and also by vertices' colors (graphs are written as unlabelled graphs for simplicity).

87 Then the [edge-saturation](#) $\mathcal{S}(h)$ is the following [graph](#), which is an [edge-extension](#) of G where
88 the extended edges are derived from edges of H :



91 2.1 Syntax: terms of KA with complement

92 We consider [terms](#) over the signature $S \triangleq \{1_{(0)}, \perp_{(0)}, \cdot_{(2)}, \cup_{(2)}, -^*_{(1)}, -^*_{(1)}\}$. Let \mathbf{V} be a set
93 of [variables](#). For a [term](#) t over S , let \bar{t} be s if $t = s^-$ for some s and be t^- otherwise. We use
94 the abbreviations:

$$95 \quad \top \triangleq \perp^- \qquad t \cap s \triangleq (t^- \cup s^-)^-.$$

96

97 For $X \subseteq \{\bar{x}, \bar{1}, \top, \cap, -\}$, let KA_X be the minimal subset A of the set of [terms](#) over S satisfying
98 the following:

$$99 \quad \frac{y \in \mathbf{V}}{y \in A} \quad \frac{}{\bar{1} \in A} \quad \frac{}{\perp \in A} \quad \frac{t \in A \quad s \in A}{t \cdot s \in A} \quad \frac{t \in A \quad s \in A}{t \cup s \in A} \quad \frac{t \in A}{t^* \in A}$$

$$100 \quad \frac{\bar{x} \in X \quad y \in \mathbf{V}}{\bar{y} \in A} \quad \frac{\bar{1} \in X}{\bar{1} \in A} \quad \frac{\top \in X}{\top \in A} \quad \frac{\cap \in X \quad t \in A \quad s \in A}{t \cap s \in A} \quad \frac{- \in X \quad t \in A}{t^- \in A}.$$

101

102 We use parentheses in ambiguous situations. We often abbreviate $t \cdot s$ to ts .

103 An [equation](#) $t = s$ is a pair of [terms](#). An [inequation](#) $t \leq s$ abbreviates the [equation](#)
104 $t \cup s = s$. The set of [quantifier-free formulas](#) of KA_X is defined by the following grammar:

$$105 \quad \varphi, \psi ::= t = s \mid \varphi \wedge \psi \mid \neg \varphi. \qquad (t, s \in \text{KA}_X)$$

106

107 We use the abbreviations: $\varphi \vee \psi \triangleq \neg(\neg\varphi \wedge \neg\psi)$, $\varphi \rightarrow \psi \triangleq \neg\varphi \vee \psi$, $\varphi \leftrightarrow \psi \triangleq (\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$.

108 2.2 Semantics: language models

109 An [S-algebra](#) \mathcal{A} is a tuple $\langle |\mathcal{A}|, \{f^{\mathcal{A}}\}_{f_{(k)} \in S} \rangle$, where $|\mathcal{A}|$ is a non-empty set and $f^{\mathcal{A}}: |\mathcal{A}|^k \rightarrow |\mathcal{A}|$
110 is a k -ary map for each $f_{(k)} \in S$. A [valuation](#) \mathbf{v} of an [S-algebra](#) \mathcal{A} is a map $\mathbf{v}: \mathbf{V} \rightarrow |\mathcal{A}|$.
111 For a [valuation](#) \mathbf{v} , we write $\hat{\mathbf{v}}: \text{KA}_{\{-\}} \rightarrow |\mathcal{A}|$ for the unique homomorphism extending \mathbf{v} .
112 Moreover, for a [quantifier-free formula](#) φ , we define $\hat{\mathbf{v}}(\varphi) \in \{\text{true}, \text{false}\}$ by:

$$113 \quad \hat{\mathbf{v}}(t = s) \triangleq (\hat{\mathbf{v}}(t) = \hat{\mathbf{v}}(s)) \quad \hat{\mathbf{v}}(\varphi \wedge \psi) \triangleq (\hat{\mathbf{v}}(\varphi) \text{ and } \hat{\mathbf{v}}(\psi)) \quad \hat{\mathbf{v}}(\neg\varphi) \triangleq (\text{not } \hat{\mathbf{v}}(\varphi)).$$

114

115 For a [quantifier-free formula](#) φ and a class of [valuations](#) (of [S-algebra](#)) \mathcal{C} ,³ we write

$$116 \quad \mathcal{C} \models \varphi \triangleq \hat{\mathbf{v}}(\varphi) \text{ holds for all } \mathbf{v} \in \mathcal{C}.$$

117

118 We abbreviate $\{\mathbf{v}\} \models \varphi$ to $\mathbf{v} \models \varphi$. The [equational theory of](#) \mathcal{C} is the set of all [equations](#) $t = s$
119 such that $\mathcal{C} \models t = s$. The [quantifier-free theory of](#) \mathcal{C} is the set of all [quantifier-free formulas](#)
120 φ such that $\mathcal{C} \models \varphi$.

121 The [language model](#) \mathcal{A} over a set X , written lang_X , is the [S-algebra](#) defined by $|\mathcal{A}| =$
122 $\wp(X^*)$, $1^{\mathcal{A}} = \{1\}$, $\perp^{\mathcal{A}} = \emptyset$, and for all $L, K \subseteq X^*$,

$$123 \quad L \cdot^{\mathcal{A}} K = L \cdot K \qquad L \cup^{\mathcal{A}} K = L \cup K \qquad L^{*\mathcal{A}} = L^* \qquad L^{-\mathcal{A}} = X^* \setminus L.$$

124

³ This paper considers classes of [valuations](#) rather than classes of [S-algebras](#) (cf. Remark 3.12).

125 We write LANG_X for the class of all valuations of lang_X and write LANG for $\bigcup_X \text{LANG}_X$.
 126 The *equational theory (resp. quantifier-free theory) w.r.t. languages* means that of LANG .

127 The *language* $[t] \subseteq \mathbf{V}^*$ of a *term* t is $\hat{\mathbf{v}}_{\text{st}}(t)$ where \mathbf{v}_{st} is the *valuation* on the *language*
 128 *model* over the set \mathbf{V} defined by $\mathbf{v}_{\text{st}}(x) = \{x\}$ for $x \in \mathbf{V}$. Since $\mathbf{v}_{\text{st}} \in \text{LANG}$, we have

$$129 \quad \text{LANG} \models t = s \quad \Rightarrow \quad [t] = [s] \quad (\dagger\text{-KA}_{\{-\}})$$

131 The converse direction fails, as in the introduction. In the sequel, we consider the *equational*
 132 *theory w.r.t. languages*.

133 2.3 (Generalized) relational models

134 We write Δ_A for the identity relation on a set A : $\Delta_A \triangleq \{\langle x, x \rangle \mid x \in A\}$. For *binary relations*
 135 R, S on a set B , the *composition* $R \cdot S$, the *n -th iteration* R^n (where $n \in \mathbb{N}$), and the *reflexive*
 136 *transitive closure* R^* are defined by:

$$137 \quad R \cdot S \triangleq \{\langle x, z \rangle \mid \exists y, \langle x, y \rangle \in R \wedge \langle y, z \rangle \in S\} \quad R^n \triangleq \begin{cases} R \cdot R^{n-1} & (n \geq 1) \\ \Delta_B & (n = 0) \end{cases} \quad R^* \triangleq \bigcup_{n \in \mathbb{N}} R^n.$$

139 Let U be a *binary relation* on a non-empty set B . A *generalized relational model*⁴ \mathcal{A} on
 140 U is an *S -algebra* such that $|\mathcal{A}| \subseteq \wp(U)$, $\mathbf{1}^{\mathcal{A}} = \Delta_B$, $\perp^{\mathcal{A}} = \emptyset$, and for all $R, S \subseteq U$,

$$141 \quad R \cdot^{\mathcal{A}} S = R \cdot S \quad R \cup^{\mathcal{A}} S = R \cup S \quad R^{*\mathcal{A}} = R^* \quad R^{-\mathcal{A}} = U \setminus R.$$

143 \mathcal{A} is a *relational model* if $U = B \times B$ and $|\mathcal{A}| = \wp(B \times B)$. We write GREL (resp. REL) for
 144 the class of all *valuations of generalized relational models* (resp. *relational models*).⁵

145 3 Finite relational models for language models

146 In this section, we present the finite relational semantics RSUB for LANG . Sect. 3.1 introduces
 147 the class SUB , which is intuitively an intermediate class between LANG and RSUB , and show
 148 that the *quantifier-free theory* of LANG coincides with that of SUB . Sect. 3.2 introduces
 149 RSUB and show that the *quantifier-free theory* of LANG coincides with that of RSUB .

150 3.1 SUB: subword models

151 For the *standard language valuation* (\mathbf{v}_{st}), the membership $w \in \hat{\mathbf{v}}_{\text{st}}(t)$ can be determined from
 152 $w' \in \hat{\mathbf{v}}_{\text{st}}(t')$ where w' ranges over *subwords* of w (and t' ranges over subterms of t). This
 153 situation is the same also for any *valuations* $\mathbf{v} \in \text{LANG}$. To reflect this property to LANG , we
 154 define the following class: *subword models* SUB . These models are *language models* where
 155 the universe is bounded to the set of *subwords* of a *word* $\ell_0 \dots \ell_{n-1}$.

156 **► Definition 3.1.** *Let $n \in \mathbb{N}$ and let $\ell_0, \dots, \ell_{n-1}$ be pairwise distinct letters. Let $U =$
 157 $\{\ell_i \dots \ell_{j-1} \mid 0 \leq i \leq j \leq n\}$. The *subword language model* \mathcal{A} of length n , written SUB_n , is
 158 the *S -algebra* defined by: $|\mathcal{A}| = \wp(U)$, $\mathbf{1}^{\mathcal{A}} = \{1\}$, $\perp^{\mathcal{A}} = \emptyset$, and for all $L, K \subseteq U$,*

$$159 \quad L \cdot^{\mathcal{A}} K = (L \cdot K) \cap U \quad L \cup^{\mathcal{A}} K = L \cup K \quad L^{*\mathcal{A}} = L^* \cap U \quad L^{-\mathcal{A}} = U \setminus L. \quad \lrcorner$$

⁴ By definition, U is a preorder. Reflexivity: By $\Delta_B = \mathbf{1}^{\mathcal{A}} \in |\mathcal{A}| \subseteq \wp(U)$, we have $\Delta_B \subseteq U$. Transitivity:
 By $\emptyset = \perp^{\mathcal{A}} \in |\mathcal{A}|$, $U = \emptyset^{-\mathcal{A}} \in |\mathcal{A}|$, and $U \cdot U = U \cdot^{\mathcal{A}} U \in |\mathcal{A}| \subseteq \wp(U)$, we have $U \cdot U \subseteq U$.

⁵ *Generalized relational models* and *relational models* are variants of proper relation algebras and full
 proper relation algebras (see, e.g., [23]), respectively, where B is non-empty set and the converse operator
 is not introduced (due to this, U is possibly not symmetric, cf. [23, Lem. 3.4]) here.

161 We write SUB_n for the class of all **valuations** of sub_n and write SUB for $\bigcup_{n \in \mathbb{N}} \text{SUB}_n$.

162 We use the following transformation of **valuations** from LANG into SUB .

163 ► **Definition 3.2** (*letters-to-letters valuations*; cf. *words-to-letters valuations* [19]). *Let X be a*
 164 *set and let $\mathbf{v} \in \text{LANG}_X$. Let $n \in \mathbb{N}$ and let a_0, \dots, a_{n-1} be (possibly not pairwise distinct)*
 165 *letters over X . Then we define $\mathbf{v}^{\langle a_0, \dots, a_{n-1} \rangle}$ as the following valuation in SUB_n :*

$$166 \quad \mathbf{v}^{\langle a_0, \dots, a_{n-1} \rangle}(x) \triangleq \{ \ell_i \dots \ell_{j-1} \mid 0 \leq i \leq j \leq n \wedge a_i \dots a_{j-1} \in \mathbf{v}(x) \}. \quad \blacktriangleleft$$

167 ► **Example 3.3.** If $\mathbf{v}(x) = \{aba, a, bb\}$, then $\mathbf{v}^{\langle a, b, a \rangle}(x) = \{\ell_0 \ell_1 \ell_2, \ell_0, \ell_2\}$. Note that $\mathbf{v}^{\langle a, b, a \rangle}$
 168 only reflects w.r.t. **subwords** of aba .

169 Using this, we can embed the membership in LANG into that in SUB .

170 ► **Lemma 3.4.** *Let X be a set and let $\mathbf{v} \in \text{LANG}_X$. Let $n \in \mathbb{N}$ and let a_0, \dots, a_{n-1} be letters*
 171 *over X . For all terms t and $0 \leq i \leq j \leq n$, $\ell_i \dots \ell_{j-1} \in \hat{\mathbf{v}}^{\langle a_0, \dots, a_{n-1} \rangle}(t)$ iff $a_i \dots a_{j-1} \in \hat{\mathbf{v}}(t)$.*

172 **Proof.** By easy induction on t , cf. [19, Lem. 23] (see Appendix D, for detailed proof).⁶ ◀

173 Moreover, this transformation is a surjective map to SUB , as follows:

174 ► **Proposition 3.5.** *Let $n \in \mathbb{N}$ and let $\mathbf{v}' \in \text{SUB}_n$. Let $\mathbf{v} \in \text{LANG}_{\{\ell_0, \dots, \ell_{n-1}\}}$ be the valuation*
 175 *defined by $\mathbf{v}(x) = \mathbf{v}'(x)$. Then $\mathbf{v}^{\langle \ell_0, \dots, \ell_{n-1} \rangle} = \mathbf{v}'$.*

176 **Proof.** By Def. 3.2, $\ell_i \dots \ell_{j-1} \in \mathbf{v}^{\langle \ell_0, \dots, \ell_{n-1} \rangle}(x)$ iff $\ell_i \dots \ell_{j-1} \in \mathbf{v}(x)$ iff $\ell_i \dots \ell_{j-1} \in \mathbf{v}'(x)$. ◀

177 From them, we have that the **quantifier-free theory** of LANG coincides with that of SUB .

178 ► **Lemma 3.6.** *For all quantifier-free formulas φ , we have: $\text{LANG} \models \varphi \Leftrightarrow \text{SUB} \models \varphi$.*

179 **Proof.** By $\mathbf{v} \models t = s \Leftrightarrow (t \leq s \wedge s \leq t)$ and $\mathbf{v} \models t \leq s \Leftrightarrow t \cap s^- \leq \perp$ for $\mathbf{v} \in \text{LANG} \cup \text{SUB}$,
 180 without loss of generality, we can assume that each **equation** in φ is of the form $u \leq \perp$.
 181 By taking the conjunctive normal form, it suffices to show when φ is of the form $(\bigvee_k t_k \leq \perp)$
 182 $\vee (\bigvee_k \neg s_k \leq \perp)$. By $\mathbf{v} \models (t \leq \perp \wedge s \leq \perp) \Leftrightarrow t \cup s \leq \perp$ for $\mathbf{v} \in \text{LANG} \cup \text{SUB}$, it suffices to
 183 show when φ is of the form $(\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp)$. Then, we have:

$$184 \quad \text{SUB} \not\models \left(\bigvee_{k=1}^m t_k \leq \perp \right) \vee (\neg s \leq \perp)$$

$$185 \quad \Rightarrow \exists n, \mathbf{v} \in \text{LANG}_{\{\ell_0, \dots, \ell_{n-1}\}}, \mathbf{v}^{\langle \ell_0, \dots, \ell_{n-1} \rangle} \not\models \left(\bigvee_{k=1}^m t_k \leq \perp \right) \vee (\neg s \leq \perp) \quad (\text{Prop. 3.5})$$

$$186 \quad \Rightarrow \text{LANG} \not\models \left(\bigvee_{k=1}^m t_k \leq \perp \right) \vee (\neg s \leq \perp) \quad (\text{By Lem. 3.4 and } \mathbf{v} \in \text{LANG})$$

$$187 \quad \Rightarrow \exists n \in \mathbb{N}, i_1, j_1, \dots, i_m, j_m \text{ s.t. } 0 \leq i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_m \leq j_m \leq n,$$

$$188 \quad \exists X, \mathbf{v} \in \text{LANG}_X, a_0, \dots, a_{n-1} \in X, \left(\bigwedge_{k=1}^m a_{i_k} \dots a_{j_k-1} \in \hat{\mathbf{v}}(t_k) \right) \wedge \hat{\mathbf{v}}(s) = \emptyset$$

$$189 \quad \Rightarrow \text{SUB} \not\models \left(\bigvee_{k=1}^m t_k \leq \perp \right) \vee (\neg s \leq \perp). \quad (\text{By Lem. 3.4 and } \mathbf{v}^{\langle a_0, \dots, a_{n-1} \rangle} \in \text{SUB}_n) \quad \blacktriangleleft$$

⁶ This equivalence holds thanks to that a_i, \dots, a_{j-1} are **letters** not **words**. For **words-to-letters valuations**, the direction \Rightarrow of Lem. 3.4 only holds [19, Lem. 23].

191 3.2 RSUB: relational subword models

192 Next, we introduce RSUB, the relational version of SUB.


193 ► **Definition 3.7.** Let $n \in \mathbb{N}$. The *relational subword language model* \mathcal{A} of length n , written
 194 rsub_n , is the *generalized relational model* on the set $U = \{\langle i, j \rangle \in [0, n] \times [0, n] \mid i \leq j\}$ s.t.

$$195 \quad |\mathcal{A}| = \{R \in \wp(U) \mid R \supseteq \Delta_{[0, n]} \vee U \setminus R \supseteq \Delta_{[0, n]}\}.$$

196 We write RSUB_n for the class of all *valuations* of rsub_n and write RSUB for $\bigcup_{n \in \mathbb{N}} \text{RSUB}_n$.

197 ► **Proposition 3.8.** For each $n \in \mathbb{N}$, sub_n is *isomorphic* to rsub_n .

Proof. By the bijective⁷ map $f: L \mapsto \{\langle i, j \rangle \mid \ell_i \dots \ell_{j-1} \in L\}$.⁸ Fig. 1 presents illustrative



198 ■ **Figure 1** Illustrative instances of the bijective map from sub_n to rsub_n when $n = 3$ (Prop. 3.8).

199 examples of this map (we display binary relations as graphs in a standard way). ◀

200 ► **Theorem 3.9.** For all *quantifier-free formulas* φ , we have: $\text{LANG} \models \varphi \Leftrightarrow \text{RSUB} \models \varphi$.

201 **Proof.** By Lem. 3.6 and Prop. 3.8. ▶

202 ► **Corollary 3.10.** The *quantifier-free theory w.r.t. languages* is in Π_1^0 for $\text{KA}_{\{-\}}$ terms.

203 **Proof.** By the finite model property of RSUB (the universe $|\text{rsub}_n|$ is finite for each n). ◀

204 ► **Remark 3.11 (RSUB and GREL).** For $\text{KA}_{\{\top\}}$, the *equational theory* of LANG coincides with
 205 that of GREL [21, REL' in Sect. 5][28]. However for $\text{KA}_{\{-\}}$, this coincidence is broken, e.g.,⁹

$$206 \quad \text{LANG} \models ab \cap cd \leq a \top d \cup b \top c \quad \text{GREL} \not\models ab \cap cd \leq a \top d \cup b \top c \quad \begin{array}{c} \circ \xrightarrow{a} \circ \xrightarrow{b} \circ \\ \circ \xrightarrow{c} \circ \xrightarrow{d} \circ \end{array}$$

$$207 \quad \text{LANG} \models a \leq \bar{b} a \bar{b} \cup b a b \quad (\text{G})\text{REL} \not\models a \leq \bar{b} a \bar{b} \cup b a b \quad \begin{array}{c} \circ \xrightarrow{a} \circ \\ \circ \xrightarrow{b} \circ \end{array}$$

209 (Each right figure denotes a *valuation* for $(\text{G})\text{REL} \not\models _$ where some edges are omitted.) ◻

210 ► **Remark 3.12** (the subclass of RSUB for the standard language valuation). Let

$$211 \quad \text{RSUB}_{\text{st}} \triangleq \bigcup_{n \in \mathbb{N}} \left\{ \mathfrak{v} \in \text{RSUB}_n \mid \bigcup_{a \in \mathbf{V}} \mathfrak{v}(a) = \{\langle i-1, i \rangle \mid i \in [1, n]\} \right\}.$$

(where a ranges over \mathbf{V}) are disjoint sets

212 By an analogy of Thm. 3.9, we have that $[t] = [s]$ iff $\text{RSUB}_{\text{st}} \models t = s$ (Appendix A). ◻

⁷ $f(\{1\}) = \Delta_{[0, n]}$ (cf. Def. 3.7) and $f(\{\ell_i \dots \ell_{j-1}\}) = \{\langle i, j \rangle\}$ is singleton if $i < j$.

⁸ This map is the same as Pratt's embedding [22, third page], cf. [28, 21]: $L \mapsto \{\langle w, wv \rangle \mid w \in X^* \wedge v \in L\}$, up to isomorphism, except that the universe is bounded to *subwords* of $\ell_0 \dots \ell_{n-1}$.

⁹ $\text{LANG} \models ab \cap cd \leq a \top d \cup b \top c$: By Levi's inequation [15][5, Example 26]. $\text{LANG} \models a \leq \bar{b} a \bar{b} \cup b a b$: for every *valuation* \mathfrak{v} of LANG, $\mathfrak{v} \models a \leq b a b$ if $1 \in \mathfrak{v}(b)$ and $\mathfrak{v} \models a \leq \bar{b} a \bar{b}$ if $1 \notin \mathfrak{v}(b)$.




Figure 2 Equational theories for $\text{KA}_{\{-\}}$ under GREL.

In Fig. 2, we summarize the equational theories for $\text{KA}_{\{-\}}$ terms above where the inclusions are shown by $\text{REL} \subseteq \text{GREL} \supseteq \text{RSUB} \supseteq \text{RSUB}_{\text{st}}$ and Thm. 3.9 and the non-inclusions are shown by counter-examples.¹⁰

4 From quantifier-free formulas to equations

Using SUB/RSUB, we give a reduction from the quantifier-free theory into the equational theory of LANG. For $n \in \mathbb{N}$, $\mathbf{v} \in \text{SUB}_n$, and i, j s.t. $0 \leq i \leq j \leq n$, we define $\mathbf{v}, i, j \models \varphi$ as

$$\begin{aligned}
\mathbf{v}, i, j \models t = s &\triangleq (\ell_i \dots \ell_{j-1} \in \hat{\mathbf{v}}(t) \text{ iff } \ell_i \dots \ell_{j-1} \in \hat{\mathbf{v}}(s)) \\
\mathbf{v}, i, j \models \varphi \wedge \psi &\triangleq (\mathbf{v}, i, j \models \varphi \text{ and } \mathbf{v}, i, j \models \psi) \quad \mathbf{v}, i, j \models \neg \varphi \triangleq (\text{not } \mathbf{v}, i, j \models \varphi).
\end{aligned}$$

The following shows that to check $\text{SUB} \models t = s$, it suffices to check about the pairs of the left-most and right-most vertices.

► **Lemma 4.1.** For all terms t, s , $\text{SUB} \models t = s \Leftrightarrow \forall n \in \mathbb{N}, \forall \mathbf{v} \in \text{SUB}_n, \mathbf{v}, 0, n \models t = s$.

Proof. Because $t = s \Leftrightarrow (t \leq s \wedge s \leq t)$ and $t \leq s \Leftrightarrow t \cap s^- \leq \perp$ hold for “RSUB \models ” and “ $\mathbf{v}, 0, n \models$ ”, it suffices to show when the equation is of the form $t \leq \perp$. We then have

$$\begin{aligned}
\text{SUB} &\not\models t \leq \perp \\
&\Rightarrow \exists \mathbf{v} \in \text{LANG}, n \in \mathbb{N}, i, j \text{ s.t. } 0 \leq i \leq j \leq n, \ell_i \dots \ell_{j-1} \in \hat{\mathbf{v}}^{\langle \ell_0, \dots, \ell_{n-1} \rangle}(t) \quad (\text{Prop. 3.5}) \\
&\Rightarrow \exists \mathbf{v} \in \text{LANG}, n \in \mathbb{N}, i, j \text{ s.t. } 0 \leq i \leq j \leq n, \ell_i \dots \ell_{j-1} \in \hat{\mathbf{v}}(t) \quad (\text{Lem. 3.4}) \\
&\Rightarrow \exists \mathbf{v} \in \text{LANG}, n \in \mathbb{N}, i, j \text{ s.t. } 0 \leq i \leq j \leq n, \ell_0 \dots \ell_{j-i-1} \in \hat{\mathbf{v}}^{\langle \ell_i, \dots, \ell_{j-1} \rangle}(t) \quad (\text{Lem. 3.4}) \\
&\Rightarrow \exists n \in \mathbb{N}, \mathbf{v} \in \text{SUB}_n, \ell_0 \dots \ell_{n-1} \in \hat{\mathbf{v}}(t) \quad (\text{Let } n \triangleq j - i \text{ as } \mathbf{v}^{\langle \ell_i, \dots, \ell_{j-1} \rangle} \in \text{SUB}_{j-i}) \\
&\Rightarrow \text{SUB} \not\models t \leq \perp. \quad (\text{Trivial})
\end{aligned}$$

(The second line from the bottom is equivalent to $\exists n \in \mathbb{N}, \mathbf{v} \in \text{SUB}_n, \mathbf{v}, 0, n \not\models t \leq \perp$.) ◀

Lem. 4.1 fails for quantifier-free formulas, e.g., $a \leq \perp \vee \bar{a} \leq \perp$ only holds in the right-hand side. We consider replacing each inequation $u \leq \perp$ with $\top u \top \leq \perp$. Then we have that $\mathbf{v} \models u \leq \perp$ iff $\mathbf{v}, 0, n \models \top u \top \leq \perp$, where $n \in \mathbb{N}$ and $\mathbf{v} \in \text{SUB}_n$ (see Lem. 4.2). More generally, for a quantifier-free formula φ , let $\text{Tr}(\varphi)$ be the term defined by:¹¹

$$\text{Tr}(t = s) \triangleq \top((t \cap s^-) \cup (t^- \cap s))\top \quad \text{Tr}(\varphi \wedge \psi) \triangleq \text{Tr}(\varphi) \cup \text{Tr}(\psi) \quad \text{Tr}(\neg \varphi) \triangleq \text{Tr}(\varphi)^-.$$

¹⁰ Note that $\text{EqT}(\text{LANG}) = \text{EqT}(\text{GREL})$ for $\text{KA}_{\{\top\}}$ and $\text{EqT}(\{\mathbf{v}_{\text{st}}\}) = \text{EqT}(\text{GREL})$ for KA [28, 21].

¹¹ $\text{Tr}(t = s)$ can be simplified for specific cases, e.g., $\text{Tr}(t \leq s) = \top(t \cap s^-)\top$ and $\text{Tr}(t \leq \perp) = \top t \top$.

242 ► **Lemma 4.2.** *Let $n \in \mathbb{N}$ and $\mathbf{v} \in \text{SUB}_n$. Let φ be a *quantifier-free formula*. Then,*

$$243 \quad \mathbf{v} \models \varphi \quad \Leftrightarrow \quad \mathbf{v}, 0, n \models \text{Tr}(\varphi) \leq \perp.$$

244 **Proof.** By easy induction on φ . Case $\varphi = (t = s)$: Let $u = (t \cap s^-) \cup (t^- \cap s)$. Then $\mathbf{v} \models t = s$
 245 iff $\hat{\mathbf{v}}(u) = \emptyset$ iff $\ell_0 \dots \ell_{n-1} \notin \hat{\mathbf{v}}(\top u \top)$ iff $\mathbf{v}, 0, n \models \top u \top \leq \perp$ iff $\mathbf{v}, 0, n \models \text{Tr}(t = s)$. For \wedge : By
 246 $\mathbf{v}, 0, n \models ((t \leq \perp) \wedge (s \leq \perp)) \Leftrightarrow (t \cup s \leq \perp)$. For \neg : By $\mathbf{v}, 0, n \models (\neg(t \leq \perp)) \Leftrightarrow (t^- \leq \perp)$. ◀

247 ► **Theorem 4.3** (cf. Schröder-Tarski translation [27]). *There is a polynomial-time reduction*
 248 *from the *quantifier-free theory* of LANG into the *equational theory* of LANG.*

249 **Proof.** By Lems. 4.1 and 4.2, $\text{SUB} \models \varphi$ iff $\text{SUB} \models \text{Tr}(\varphi) \leq \perp$. Thus by Thm. 3.9. ◀

250 ► **Corollary 4.4** (Hoare hypotheses elimination). *For all *terms* t, s, u ,*

$$251 \quad \text{LANG} \models u \leq \perp \rightarrow t \leq s \quad \Leftrightarrow \quad \text{LANG} \models t \leq s \cup \top u \top.$$

252 **Proof.** By set-theoretic equivalences, we have

$$\begin{aligned} 253 \quad \text{SUB} \models u \leq \perp \rightarrow t \leq s &\Leftrightarrow \text{SUB} \models \top(t \cap s^-) \top \leq \top u \top && \text{(By Thm. 4.3)} \\ 254 &\Leftrightarrow \text{SUB} \models t \cap s^- \leq \top u \top && (\Rightarrow: \text{By } \perp \leq \top \Leftarrow: \text{By } \top \top \leq \top) \\ 255 &\Leftrightarrow \text{SUB} \models t \leq s \cup \top u \top && \blacktriangleleft \end{aligned}$$

257 ► **Remark 4.5.** Note that $\text{LANG} \not\models (\neg u \leq \perp) \Leftrightarrow \top \leq \top u \top$, cf. Schröder-Tarski translation
 258 [27, XXXII.] for REL. If $\hat{\mathbf{v}}(u) = \{w\}$ where $\mathbf{v} \in \text{LANG}_X$ and $w \neq \perp$, then $\hat{\mathbf{v}}(\top) = [X^*] \not\subseteq$
 259 $[X^* w X^*] = \hat{\mathbf{v}}(\top u \top)$. This is why we go via “ $\mathbf{v}, 0, n \models _$ ” (Lems. 4.1 and 4.2). Similarly for
 260 Cor. 4.4, $\text{LANG} \not\models (u \leq \perp \rightarrow t \leq s) \Leftrightarrow (t \leq s \cup \top u \top)$, e.g., when $t = \top$ and $s = \perp$. ◻

261 4.1 Undecidability via the Hoare hypotheses elimination

262 Thm. 4.3/Cor. 4.4 implies the undecidability of the *equational theory* of LANG for full *terms*.

263 A *context-free grammar* (CFG) \mathfrak{C} over a finite set A is a tuple $\langle X, \mathcal{R}, s \rangle$, where

- 264 ■ X is a finite set of non-terminal labels s.t. $A \cap X = \emptyset$;
- 265 ■ \mathcal{R} is a finite set of rewriting rules $x \leftarrow w$ of $x \in X$ and $w \in (A \cup X)^*$;
- 266 ■ $s \in X$ is the start label.

267 We write $[\mathfrak{C}]$ for the *language* of \mathfrak{C} . It is well-known that the *universality problem* for CFGs—
 268 given a CFG \mathfrak{C} , does $[\mathfrak{C}] = A^*$ hold?—is Π_1^0 -complete. We can naturally encode this problem
 269 by the *quantifier-free theory* of LANG, as follows (cf. [18, Lem. 47]).

270 ► **Lemma 4.6.** *Let $\mathfrak{C} = \langle X, \mathcal{R}, s \rangle$ be a CFG over a finite set A . Then,*

$$271 \quad [\mathfrak{C}] = A^* \quad \Leftrightarrow \quad \text{LANG} \models \left(\bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x \right) \rightarrow (A^* \leq s).$$

272 *Here, A denotes the term $(a_1 \cup \dots \cup a_n)$ in the right-hand side where $A = \{a_1, \dots, a_n\}$.*

273 **Proof Sketch.** By an analogy of [18, Lem. 47] (for REL). The relational models used in [18]
 274 are of the form of RSUB. See Appendix E for a detail. ◀

275 ► **Theorem 4.7.** *The *equational theory w.r.t. languages* is Π_1^0 -complete for $\text{KA}_{\{\bar{x}, \cap\}}$.*

276 **Proof.** (in Π_1^0): By Cor. 3.10. (Hardness): Let $\mathfrak{C} = \langle X, \{x_i \leftarrow w_i \mid i \in [1, n]\}, s \rangle$ be a
 277 CFG over a finite set A . By Lem. 4.6 and by Cor. 4.4 with $\text{LANG} \models (\bigwedge_{i \in [1, n]} w_i \leq x_i) \Leftrightarrow$
 278 $(\bigcup_{i \in [1, n]} w_i \cap \bar{x}_i) \leq \perp$, we have: $[\mathfrak{C}] = A^*$ iff $\text{LANG} \models A^* \leq s \cup \top(\bigcup_{i \in [1, n]} w_i \cap \bar{x}_i) \top$. Hence,
 279 we can give a reduction from the *universality problem* of CFGs. ◀

280 ▶ **Corollary 4.8.** *The equational theory w.r.t. languages is Π_1^0 -complete for $\text{KA}_{\{-\}}$.*

281 **Proof.** By Cor. 3.10 and Thm. 4.7. ◀

282 Moreover, we also have the following, as $\text{LANG} \models \bar{I} = A\top \rightarrow A^* = \top$ holds (see Appendix E).

283

284 ▶ **Lemma 4.9.** *Let $\mathfrak{C} = \langle X, \mathcal{R}, s \rangle$ be a CFG over a finite set A . Then,*

$$285 \quad [\mathfrak{C}] = A^* \quad \Leftrightarrow \quad \text{LANG} \models (\bar{I} = A\top \wedge \bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x) \rightarrow (\top \leq s).$$

286 Hence, the undecidability still holds even without Kleene star.

287 ▶ **Theorem 4.10.** *The equational theory w.r.t. languages is Π_1^0 -complete for $\text{KA}_{\{-\}}$ without*
288 *Kleene star.*

289 **Proof.** In the same way as Thm. 4.7 (by using Lem. 4.9 instead of Lem. 4.6). ◀

290 5 Graph characterization for $\text{KA}_{\{\bar{x}, \bar{I}, \top, \cap\}}$ terms

291 In Sects. 5 and 6, we consider the decidability of $\text{KA}_{\{\bar{x}, \bar{I}, \top\}}$. We recall Sect. 2 for graphs. In
292 this section, we give a graph characterization of the equational theory of RSUB for $\text{KA}_{\{\bar{x}, \bar{I}, \top, \cap\}}$,
293 by generalizing the graph characterization of REL [18, Thm. 18].

294 5.1 Graph languages for $\text{KA}_{\{\bar{x}, \bar{I}, \top, \cap\}}$

295 Let $\tilde{\mathbf{V}} \triangleq \{x, \bar{x} \mid x \in \mathbf{V}\} \cup \{\bar{I}, \top\}$ and $\tilde{\mathbf{V}}_1 \triangleq \tilde{\mathbf{V}} \cup \{I\}$. For a $\text{KA}_{\{\bar{x}, \bar{I}, \top, \cap\}}$ term t , the graph
296 language $\mathcal{G}(t)$ is a set of graphs over $\tilde{\mathbf{V}}_1$ defined by:¹²

$$297 \quad \mathcal{G}(x) \triangleq \{ \rightarrow \circ - x \rightarrow \circ \rightarrow \} \text{ where } x \in \tilde{\mathbf{V}} \quad \mathcal{G}(\perp) \triangleq \emptyset \quad \mathcal{G}(I) \triangleq \{ \rightarrow \circ \rightarrow \}$$

$$298 \quad \mathcal{G}(t \cap s) \triangleq \{ \rightarrow \circ \xrightarrow[G]{H} \circ \rightarrow \mid G \in \mathcal{G}(t) \wedge H \in \mathcal{G}(s) \} \quad \mathcal{G}(t \cup s) \triangleq \mathcal{G}(t) \cup \mathcal{G}(s)$$

$$299 \quad \mathcal{G}(t \cdot s) \triangleq \{ \rightarrow \circ - G \rightarrow \circ - H \rightarrow \circ \rightarrow \mid G \in \mathcal{G}(t) \wedge H \in \mathcal{G}(s) \} \quad \mathcal{G}(t^*) \triangleq \bigcup_{n \in \mathbb{N}} \mathcal{G}(t^n).$$

300
301 (We use *series-composition* for (\cdot) and *parallel-composition* for (\cap) .)

302 For a valuation $\mathbf{v} \in \text{GREL}$ on a binary relation on a set B and $\langle x, y \rangle \in \hat{\mathbf{v}}(\top)$, let $\mathbf{G}(\mathbf{v}, x, y)$
303 be the graph defined by:

$$304 \quad \mathbf{G}(\mathbf{v}, x, y) \triangleq \langle B, \{\hat{\mathbf{v}}(a)\}_{a \in \tilde{\mathbf{V}}_1}, x, y \rangle.$$

305 For a class $\mathcal{C} \subseteq \text{GREL}$, let $\text{GR}_{\mathcal{C}}$ be the graph language $\{\mathbf{G}(\mathbf{v}, x, y) \mid \mathbf{v} \in \mathcal{C} \text{ and } \langle x, y \rangle \in \hat{\mathbf{v}}(\top)\}$.
306 We say that a graph language \mathcal{G} is *induced subgraph-closed* if every induced subgraph of
307 every $G \in \mathcal{G}$ is isomorphic to a member of \mathcal{G} . We say that a class $\mathcal{C} \subseteq \text{GREL}$ is *induced*
308 *subgraph-closed* if $\text{GR}_{\mathcal{C}}$ is induced subgraph-closed. By the form of rsub_n , RSUB is *induced*
309 *subgraph-closed*.

310 We recall *edge-saturations* $\mathcal{S}(h)$ of Def. 2.1. For a graph G and graph language \mathcal{G} , let

$$311 \quad \mathcal{S}_{\mathcal{C}}(G) \triangleq \{ \mathcal{S}(h) \mid \exists H \in \text{GR}_{\mathcal{C}}, h: G \rightarrow H \} \quad \mathcal{S}_{\mathcal{C}}(\mathcal{G}) \triangleq \bigcup_{H \in \mathcal{G}} \mathcal{S}_{\mathcal{C}}(H).$$

312

¹²We introduce \top -labelled edges, cf. [18, Def. 6], because \top is not fixed to the full relation.

313 ► **Example 5.1.** The following is an instance of $\mathcal{S}_{\text{RSUB}}(G)$ where $\mathbf{V} = \{a\}$:

$$314 \quad \mathcal{S}_{\text{RSUB}}(\rightarrow \circ - a \rightarrow \circ - \bar{a} \rightarrow \circ) = \left\{ \begin{array}{l} \begin{array}{c} \xrightarrow{\bar{1}, \top} \\ \begin{array}{ccccc} \circ & \xrightarrow{a, \bar{1}, \top} & \circ & \xrightarrow{\bar{a}, \bar{1}, \top} & \circ \\ \downarrow \scriptstyle a, \bar{1}, \top & & \downarrow \scriptstyle a, \bar{1}, \top & & \downarrow \scriptstyle a, \bar{1}, \top \\ \circ & & \circ & & \circ \\ \downarrow \scriptstyle \bar{a}, \bar{1}, \top & & \downarrow \scriptstyle \bar{a}, \bar{1}, \top & & \downarrow \scriptstyle \bar{a}, \bar{1}, \top \\ \circ & & \circ & & \circ \end{array} \end{array}, \quad \begin{array}{c} \xrightarrow{\bar{1}, \top} \\ \begin{array}{ccccc} \circ & \xrightarrow{a, \bar{1}, \top} & \circ & \xrightarrow{\bar{a}, \bar{1}, \top} & \circ \\ \downarrow \scriptstyle \bar{a}, \bar{1}, \top & & \downarrow \scriptstyle \bar{a}, \bar{1}, \top & & \downarrow \scriptstyle \bar{a}, \bar{1}, \top \\ \circ & & \circ & & \circ \\ \downarrow \scriptstyle a, \bar{1}, \top & & \downarrow \scriptstyle a, \bar{1}, \top & & \downarrow \scriptstyle a, \bar{1}, \top \\ \circ & & \circ & & \circ \end{array} \end{array}, \right. \\ 315 \end{array}$$

(In the sequel, we often omit \top - or $\bar{1}$ -labelled edges for simplicity.)

By the form of GR_{RSUB} , each H , where $H \in \mathcal{S}_{\text{RSUB}}(H')$ for some H' , satisfies

- 316 ■ \top^H is a total preorder;
- 317 ■ $a^H \supseteq \top^H$ or $\bar{a}^H \supseteq \top^H$ holds for each $a \in \mathbf{V}$. ┘

319 Additionally, let $H^\mathcal{Q} \triangleq H/(\top^H)^\mathcal{Q}$ and $\mathcal{G}^\mathcal{Q} \triangleq \{H^\mathcal{Q} \mid H \in \mathcal{G}\}$ where $R^\mathcal{Q}$ denotes the
 320 equivalence closure of R . We then have the following graph language characterization. This
 321 is an analogy of [18, Theorem 18], but is generalized for including RSUB .

322 ► **Theorem 5.2.** Let $\mathcal{C} \subseteq \text{GREL}$ be *induced subgraph-closed*. For all $\text{KA}_{\{\bar{x}, \bar{1}, \top, \cap\}}$ terms t, s ,

$$323 \quad \mathcal{C} \models t \leq s \quad \Leftrightarrow \quad \forall H \in \mathcal{S}_{\mathcal{C}}(\mathcal{G}(t))^\mathcal{Q}, \exists G \in \mathcal{G}(s), G \rightarrow H.$$

324 **Proof.** As with [18, Theorem 18]. See Sect. 5.3 for a detailed proof. ◀

325 By Thm. 5.2, we can show *inequations* by *graph homomorphisms*, e.g., as follows.

326 ► **Example 5.3.** We recall the *inequations* in Remark 3.11. Gray-colored edges are extended
 327 by *edge-saturations* $\mathcal{S}_{\text{RSUB}}$. We omit unimportant edges.

328 ■ $\text{LANG} \models ab \cap cd \leq a \top d \cup c \top b$: For each *graph* $H \in \mathcal{S}_{\text{RSUB}}(\mathcal{G}(ab \cap cd))^\mathcal{Q}$, we can give a
 329 *graph homomorphism* from some *graph* in $\mathcal{G}(a \top d \cup c \top b)$ as follows:

$$330 \quad \mathcal{G}(a \top d \cup c \top b) = \left\{ \begin{array}{c} \begin{array}{c} \circ \xrightarrow{a} \bullet \xrightarrow{\top} \circ \xrightarrow{d} \circ \\ \downarrow \scriptstyle a, \bullet \\ \circ \xrightarrow{a} \bullet \xrightarrow{b} \circ \\ \downarrow \scriptstyle c, \bullet \\ \circ \xrightarrow{c} \bullet \xrightarrow{d} \circ \end{array}, \quad \begin{array}{c} \circ \xrightarrow{c} \bullet \xrightarrow{\top} \circ \xrightarrow{b} \circ \\ \downarrow \scriptstyle a, \bullet \\ \circ \xrightarrow{a} \bullet \xrightarrow{b} \circ \\ \downarrow \scriptstyle c, \bullet \\ \circ \xrightarrow{c} \bullet \xrightarrow{d} \circ \end{array} \end{array} \right\}$$

(Case $(\bullet, \bullet) \in \top^H$) (Case $(\bullet, \bullet) \in \top^H$)

331 ■ $\text{LANG} \models a \leq \bar{b} \bar{a} \bar{b} \cup b a b$:

$$332 \quad \mathcal{G}(\bar{b} \bar{a} \bar{b} \cup b a b) = \left\{ \begin{array}{c} \begin{array}{c} \bullet \xrightarrow{\bar{b}} \bullet \xrightarrow{a} \bullet \xrightarrow{\bar{b}} \bullet \\ \downarrow \scriptstyle \bar{b} \\ \bullet \xrightarrow{a} \bullet \\ \downarrow \scriptstyle \bar{b} \\ \bullet \xrightarrow{a} \bullet \end{array}, \quad \begin{array}{c} \bullet \xrightarrow{b} \bullet \xrightarrow{a} \bullet \xrightarrow{b} \bullet \\ \downarrow \scriptstyle b \\ \bullet \xrightarrow{a} \bullet \\ \downarrow \scriptstyle b \\ \bullet \xrightarrow{a} \bullet \end{array} \end{array} \right\}$$

(Case $\bar{b}^H \supseteq \top^H$) (Case $b^H \supseteq \top^H$)

333 (Additionally, $_^\mathcal{Q}$ is necessary, e.g., for $\top \leq \top \cup \bar{1}$ [18, Remark 19].) ┘

334 5.2 Word languages for $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$

335 Particularly for $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$, Thm. 5.2 can be rephrased by *word languages*.

336 For a *word* $w = a_1 \dots a_n$ over $\tilde{\mathbf{V}}$, let $\mathcal{G}(w)$ be the following *graph* where $|\mathcal{G}(w)| = [0, n]$:

$$337 \quad \rightarrow \textcircled{0} \xrightarrow{a_0} \textcircled{1} \xrightarrow{a_1} \textcircled{2} \cdots \xrightarrow{a_n} \textcircled{n} \rightarrow .$$

338 $\mathcal{G}(w)$ is the unique *graph* in $\mathcal{G}(w)$ up to *graph isomorphisms*.

339 For a $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ term t , we write $[t]_{\tilde{\mathbf{V}}}$ for the *word language* $[t]$ over $\tilde{\mathbf{V}}$ where $\bar{x}, \bar{1}, \top$
 340 are also viewed as *letters*; e.g., $[\bar{x}]_{\tilde{\mathbf{V}}} = \{\bar{x}\}$ and $[\bar{x}] = \mathbf{V}^* \setminus \{x\}$ for $x \in \mathbf{V}$. Note that
 341 $\mathcal{G}(t) = \{\mathcal{G}(w) \mid w \in [t]_{\tilde{\mathbf{V}}}\}$. Hence, *graph languages* are expressible by *word languages* for
 342 $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ terms.

343 Additionally, we introduce nondeterministic finite word automata with epsilon transitions
 344 (*NFAs*). *NFAs* are (2-pointed) graphs over $\tilde{\mathbf{V}}_1$ where the source and target vertices denote
 345 the initial and (single) accepting states, respectively, and l -labelled edges denote epsilon
 346 transitions. For a graph H and a word $w = a_1 \dots a_n$, we write δ_w^H for the binary relation
 347 $(\mathsf{l}^H)^* \cdot a_1^H \cdot (\mathsf{l}^H)^* \dots a_n^H \cdot (\mathsf{l}^H)^*$. For $q \in |H|$, we let $\delta_w^H(q) \triangleq \{q' \mid \langle q, q' \rangle \in \delta_w^H\}$. For $Q \subseteq |H|$, we
 348 let $\delta_w^H(Q) \triangleq \bigcup_{q \in Q} \delta_w^H(q)$. The word language $[H]_{\tilde{\mathbf{V}}}$ is defined as $\{w \in \tilde{\mathbf{V}}^* \mid \langle 1^H, 2^H \rangle \in \delta_w^H\}$.
 349 Note that $[H]_{\tilde{\mathbf{V}}} = \{w \in \tilde{\mathbf{V}}^* \mid G(w) \longrightarrow H^{\mathcal{Q}}\}$ if l^H is an equivalence relation. Then we have
 350 the following:

351 ► **Corollary 5.4.** *Let $\mathcal{C} \subseteq \text{GREL}$ be induced subgraph-closed. For all $\text{KA}_{\{\bar{x}, \bar{\mathsf{l}}, \top\}}$ terms t, s ,*

$$352 \quad \mathcal{C} \models t \leq s \iff [t]_{\tilde{\mathbf{V}}} \subseteq \{w \in \tilde{\mathbf{V}}^* \mid \forall H \in \mathcal{S}_{\mathcal{C}}(G(w)), [s]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} \neq \emptyset\}.$$

353 **Proof.** By Thm. 5.2. See Sect. 5.3 for a detailed proof. ◀

354 5.3 Proof of Thm. 5.2 and Cor. 5.4

355 Let $\mathbf{v} \in \text{GREL}$. For a graph H and a graph language \mathcal{G} , we write

$$356 \quad \hat{\mathbf{v}}(H) \triangleq \{\langle x, y \rangle \mid H \longrightarrow G(\mathbf{v}, x, y)\} \quad \hat{\mathbf{v}}(\mathcal{G}) \triangleq \bigcup_{H \in \mathcal{G}} \hat{\mathbf{v}}(H).$$

357
 358
 359 ► **Lemma 5.5.** *Let $\mathbf{v} \in \text{GREL}$. For all $\text{KA}_{\{\bar{x}, \bar{\mathsf{l}}, \top, \cap\}}$ terms t , we have $\hat{\mathbf{v}}(t) = \hat{\mathbf{v}}(\mathcal{G}(t))$.*

360 **Proof (cf. [18, Prop. 11]).** By easy induction on t . See Appendix F for a detail. ◀

361 ► **Lemma 5.6.** *Let $\mathcal{C} \subseteq \text{GREL}$, $\mathbf{v} \in \mathcal{C}$, and H be a graph. Then $\hat{\mathbf{v}}(\mathcal{S}_{\mathcal{C}}(H)^{\mathcal{Q}}) = \hat{\mathbf{v}}(H)$.*

362 **Proof.** (\subseteq): Because, for any $J \in \mathcal{S}_{\mathcal{C}}(H)$, we have $J \longrightarrow J^{\mathcal{Q}}$ by $H \longrightarrow J$ and $J \longrightarrow J^{\mathcal{Q}}$. (\supseteq):
 363 Let $h: H \longrightarrow G(\mathbf{v}, x, y)$. Then $\mathcal{S}(h) \longrightarrow G(\mathbf{v}, x, y)$. Because $\mathsf{l}^{G(\mathbf{v}, x, y)}$ is the identity relation,
 364 $\mathcal{S}(h)^{\mathcal{Q}} \longrightarrow G(\mathbf{v}, x, y)$. Hence $\hat{\mathbf{v}}(\mathcal{S}_{\mathcal{C}}(H)^{\mathcal{Q}}) \supseteq \hat{\mathbf{v}}(H)$. ◀

365 ► **Proposition 5.7.** *Let $h: H \longrightarrow G(\mathbf{v}, x, y)$. Then the graph $\mathcal{S}(h)^{\mathcal{Q}}$ is isomorphic to the
 366 induced subgraph of $G(\mathbf{v}, x, y)$ on the range of h .*

367 **Proof.** Easy, by construction. Note that since $\mathsf{l}^{G(\mathbf{v}, x, y)}$ is the identity relation, $\mathsf{l}^{\mathcal{S}(h)}$ is an
 368 equivalence relation (see, e.g., Example 2.2). ◀

369 **Proof of Thm. 5.2.** We have

$$370 \quad \mathcal{C} \models t \leq s \iff \forall \mathbf{v} \in \mathcal{C}, \hat{\mathbf{v}}(t) \subseteq \hat{\mathbf{v}}(s) \iff \forall \mathbf{v} \in \mathcal{C}, \hat{\mathbf{v}}(\mathcal{S}_{\mathcal{C}}(\mathcal{G}(t))^{\mathcal{Q}}) \subseteq \hat{\mathbf{v}}(\mathcal{G}(s)) \quad (\text{Lems. 5.5 and 5.6})$$

$$371 \quad \iff \forall G \in \mathcal{S}_{\mathcal{C}}(\mathcal{G}(t))^{\mathcal{Q}}, \forall J \in \text{GR}_{\mathcal{C}}, (G \longrightarrow J) \text{ implies } (\exists H \in \mathcal{G}(s), H \longrightarrow J) \quad (\text{Def. of } \hat{\mathbf{v}})$$

$$372 \quad \iff \forall G \in \mathcal{S}_{\mathcal{C}}(\mathcal{G}(t))^{\mathcal{Q}}, \exists H \in \mathcal{G}(s), H \longrightarrow G. \quad (\heartsuit)$$

374 Here, for (\heartsuit), \Leftarrow : Let $H \in \mathcal{G}(s)$ be s.t. $H \longrightarrow G$. Then for all J s.t. $G \longrightarrow J$, we have
 375 $H \longrightarrow J$ by transitivity of \longrightarrow . \Rightarrow : By Prop. 5.7 and that $\text{GR}_{\mathcal{C}}$ is induced subgraph-closed,
 376 we have $\mathcal{S}_{\mathcal{C}}(\mathcal{G}(t))^{\mathcal{Q}} \subseteq \text{GR}_{\mathcal{C}}$. Thus by letting $J = G$, this completes the proof. ◀

377 **Proof of Cor. 5.4.** We have

$$378 \quad \mathcal{C} \models t \leq s \iff \forall w \in [t]_{\tilde{\mathbf{V}}}, \forall H \in \mathcal{S}_{\mathcal{C}}(G(w)), \exists v \in [s]_{\tilde{\mathbf{V}}}, G(v) \longrightarrow H^{\mathcal{Q}}$$

$$\quad \quad \quad (\text{Thm. 5.2 and } \mathcal{G}(s) = \{G(v) \mid v \in [s]_{\tilde{\mathbf{V}}}\})$$

$$379 \quad \iff \forall w \in [t]_{\tilde{\mathbf{V}}}, \forall H \in \mathcal{S}_{\mathcal{C}}(G(w)), [s]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} \neq \emptyset \quad ([H]_{\tilde{\mathbf{V}}} = \{v \in \tilde{\mathbf{V}}^* \mid G(v) \longrightarrow H^{\mathcal{Q}}\})$$

$$380 \quad \iff [t]_{\tilde{\mathbf{V}}} \subseteq \{w \in \tilde{\mathbf{V}}^* \mid \forall H \in \mathcal{S}_{\mathcal{C}}(G(w)), [s]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} \neq \emptyset\}. \quad \blacktriangleleft$$

6 PSPACE decidability for $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ terms

In this section, we show that the equational theory w.r.t. languages for $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ is decidable and PSPACE-complete. We consider graphs over $\tilde{\mathbf{V}}_1$ / NFAs, instead of $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ terms / regular expressions over the alphabet $\tilde{\mathbf{V}}$. Relying on the graph characterization (Cor. 5.4), we consider that, given an NFA J (corresponding to the term s in Cor. 5.4), we give an NFA recognizing the following word language:

$$\mathbf{L}_J \triangleq \{w \in \tilde{\mathbf{V}}^* \mid \exists H \in \mathcal{S}_{\text{RSUB}}(\mathbf{G}(w)), [J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset\}.$$

Note that $\text{RSUB} \models t \leq s \Leftrightarrow [t]_{\tilde{\mathbf{V}}} \cap \mathbf{L}_J = \emptyset$ when $[s]_{\tilde{\mathbf{V}}} = [J]_{\tilde{\mathbf{V}}}$. We first give an equivalent notion of “ $w \in \mathbf{L}_J$ ” in Sect. 6.1, and then we give an NFA construction in Sect. 6.2.

6.1 Saturable paths for RSUB

We first give an equivalent notion of $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$, compatible with edge-saturations.

► **Definition 6.1.** Let J and H be graphs. A map $U: |H| \rightarrow \wp(|J|)$ is an *emptiness-witness* for $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$ if the following hold where $U_x \triangleq U(x)$:


- $1^J \in U_{1^H}$ and $\forall a \in \tilde{\mathbf{V}}_1, \forall \langle x, y \rangle \in a^H, \delta_a^J(U_x) \subseteq U_y$;
- $2^J \notin U_{2^H}$. ┘

Intuitively, the first condition denotes that U is a cover of the reachable states from the pair “ $1^J \in U_{1^H}$ ”. If the second condition holds, we can see that the pair “ $2^J \in U_{2^H}$ ” is unreachable. As expected, we have the following (see Appendix B, for a proof).

► **Proposition 6.2.** Let J and H be graphs where 1^H is reflexive. Then

$$[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset \Leftrightarrow \exists U: |H| \rightarrow \wp(|J|), U \text{ is an emptiness-witness for } [J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset.$$

► **Example 6.3.** We consider the following graphs. J denotes the NFA s.t. $[J]_{\tilde{\mathbf{V}}} = \{w \in \{a, \bar{a}\}^* \mid \exists n \in \mathbb{N}, \bar{a} \text{ occurs } 3n + 2 \text{ times in } w\}$. H is a graph in $\mathcal{S}_{\text{RSUB}}(\mathbf{G}(a\bar{a}a))$ where \top - or $\bar{1}$ -labelled edges are omitted. By the form of H , one can see that $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$.



If $U_0 = U_1 = \{\bullet\}$ and $U_2 = U_3 = \{\circ, \bullet\}$, then this U is an *emptiness-witness*; e.g., for $\langle 1, 2 \rangle \in \bar{a}^H$, $\delta_{\bar{a}}^J(U_1) = \delta_{\bar{a}}^J(\{\bullet\}) = \{\bullet\} \subseteq U_2$. Besides this, if $U_0 = U_1 = \{\bullet\}$ and $U_2 = U_3 = \{\circ\}$, this U is also an *emptiness-witness*. These witnesses also show $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$. ┘

Next, we give an equivalent notion of “ $w \in \mathbf{L}_J$ ”. In the following, we abstract saturated edges (gray-colored edges in Example 6.3) using U , as U implies the existence of *edge-saturations*.

► **Definition 6.4.** Let J be a graph and w be a word. A pair $P = \langle H, U \rangle$ is a *saturable path* for $w \in \mathbf{L}_J$ if the following hold:

(P-Ext) H is an *edge-extension* of $\mathbf{G}(w)$ such that¹³

¹³ In this definition, $\top^H, 1^H, \bar{1}^H$ and edges in 1^H are already edge-saturated, for preserving (P-Con) easily.

416 ■ \top^H is a total preorder and $\top^H \supseteq \{\langle i-1, i \rangle \mid i \in [1, n]\}$ where $w = a_1 \dots a_n$;

417 ■ $\bar{1}^H = \top^H \cap \{\langle j, i \rangle \mid \langle i, j \rangle \in \top^H\}$ and $\bar{1}^H = \top^H \setminus \bar{1}^H$;

418 ■ $\forall a \in \mathbf{V}$, $\langle a^H, \bar{a}^H \rangle$ is either $\langle a^{G(w)} \cup \bar{1}^H, \bar{a}^{G(w)} \rangle$ or $\langle a^{G(w)}, \bar{a}^{G(w)} \cup \bar{1}^H \rangle$.

419 **(P-Con)** H is consistent: $\forall a \in \mathbf{V}$, $a^{H^\circ} \cap \bar{a}^{H^\circ} = \emptyset$.

420 **(P-Wit)** $U: |H| \rightarrow \wp(|J|)$ is an emptiness-witness for $[J]_{\bar{\mathbf{V}}} \cap [H]_{\bar{\mathbf{V}}} = \emptyset$.

421 **(P-Sat)** H is saturable: $\forall a \in \mathbf{V}$, $\forall \langle i, j \rangle \in \bar{1}^H$, $\delta_a^J(U_i) \subseteq U_j$ or $\delta_{\bar{a}}^J(U_i) \subseteq U_j$. \lrcorner

422 ► **Lemma 6.5.** Let J be a graph and w be a word. Then,

423 $w \in \mathbf{L}_J \iff$ there is a saturable path for $w \in \mathbf{L}_J$.

424 **Proof.** (\Rightarrow): By Prop. 6.2, let $H' \in \mathcal{S}_{\text{RSUB}}(\mathbf{G}(w))$ and let U be an emptiness-witness for
425 $[J]_{\bar{\mathbf{V}}} \cap [H']_{\bar{\mathbf{V}}} = \emptyset$. We define the graph H as follows:

426 ■ $|H| = |H'|$;

427 ■ $a^H = a^{H'}$ for $a \in \{\top, \bar{1}, \bar{1}\}$;

428 ■ $a^H = a^{G(w)} \cup (a^{H'} \cap \bar{1}^{H'})$ for $a \in \tilde{\mathbf{V}}_1 \setminus \{\top, \bar{1}, \bar{1}\}$.

429 We then have that the pair $P \triangleq \langle H, U \rangle$ is a saturable path for $w \in \mathbf{L}_J$, as follows:

430 ■ (P-Ext): By that H' is an edge-saturation w.r.t. RSUB.

431 ■ (P-Con): Because H' is consistent by $H' \in \mathcal{S}_{\text{RSUB}}(\mathbf{G}(w))$.

432 ■ (P-Wit): Because U is an emptiness-witness for $[J]_{\bar{\mathbf{V}}} \cap [H']_{\bar{\mathbf{V}}} = \emptyset$.

433 ■ (P-Sat): Because $a^{H'} \cup \bar{a}^{H'} = \top^{H'}$ and U is an emptiness-witness for $[J]_{\bar{\mathbf{V}}} \cap [H']_{\bar{\mathbf{V}}} = \emptyset$.

434 (\Leftarrow): Let $P = \langle H, U \rangle$ be the saturable path. By (P-Ext), $\bar{1}^H$ is an equivalence relation.

435 We define the graph H' as follows:

436 ■ $|H'| = |H|$;

437 ■ $a^{H'} = a^H$ for $a \in \{\top, \bar{1}, \bar{1}\}$;

438 ■ for $a \in \mathbf{V}$ and $\langle x, y \rangle \in \top^H$,

439 ■ if $\langle [x]_{\bar{1}^H}, [y]_{\bar{1}^H} \rangle \in a^{H^\circ}$, then $\langle x, y \rangle \in a^{H'} \setminus \bar{a}^{H'}$;

440 ■ else if $\langle [x]_{\bar{1}^H}, [y]_{\bar{1}^H} \rangle \in \bar{a}^{H^\circ}$, then $\langle x, y \rangle \in \bar{a}^{H'} \setminus a^{H'}$;

441 ■ else if $U_y \subseteq \delta_a^J(U_x)$, then $\langle x, y \rangle \in a^{H'} \setminus \bar{a}^{H'}$;

442 ■ else $\langle x, y \rangle \in \bar{a}^{H'} \setminus a^{H'}$.

443 By the construction of H' , we have the following:

444 ■ H' is consistent: If $[x]_{\bar{1}^{H'}} = [y]_{\bar{1}^{H'}}$ then $U_x = U_y$, because $U_x \subseteq \delta_1^J(U_x) \subseteq U_y \subseteq \delta_1^J(U_y) \subseteq U_x$
445 by (P-Wit); thus, if $[x]_{\bar{1}^{H'}} = [x']_{\bar{1}^{H'}}$ and $[y]_{\bar{1}^{H'}} = [y']_{\bar{1}^{H'}}$, then $\langle x, y \rangle \in a^{H'}$ iff $\langle x', y' \rangle \in a^{H'}$.


446 ■ for $a \in \mathbf{V}$, $\bar{a}^{H'} = \top^{H'} \setminus a^{H'}$: Because $a^{H'} \cup \bar{a}^{H'} = \top^{H'}$ and H' is consistent.

447 Combining (P-Ext) and them yields $H' \in \mathcal{S}_{\text{RSUB}}(\mathbf{G}(w))$. Also, U is an emptiness-witness for
448 $[J]_{\bar{\mathbf{V}}} \cap [H']_{\bar{\mathbf{V}}} = \emptyset$ as follows. For edges already in H , it is shown by (P-Wit). For extended
449 edges from H , it is shown by the construction of H' (for the last case of the four cases above,
450 by $U_y \not\subseteq \delta_a^J(U_x)$ and (P-Sat), we have $U_y \subseteq \delta_{\bar{a}}^J(U_x)$). Hence, this completes the proof. \blacktriangleleft

451 ► **Example 6.6.** We recall the graphs J and $H \in \mathcal{S}_{\text{RSUB}}(\mathbf{G}(a\bar{a}a))$ in Example 6.3. The
452 following P is a saturable path for $a\bar{a}a \in \mathbf{L}_J$ where \top - or $\bar{1}$ -labelled edges are omitted:

453
$$P = \left(\begin{array}{cccc} \begin{array}{c} \begin{array}{c} \text{a, } \bar{1} \\ \downarrow \\ \text{0} \end{array} \\ \{\bullet\} \end{array} & \xrightarrow{a} & \begin{array}{c} \begin{array}{c} \text{a, } \bar{1} \\ \downarrow \\ \text{1} \end{array} \\ \{\bullet\} \end{array} & \xrightarrow{\bar{a}} & \begin{array}{c} \begin{array}{c} \text{a, } \bar{1} \\ \downarrow \\ \text{2} \end{array} \\ \{\circ, \bullet\} \end{array} & \xrightarrow{\text{a, } \bar{1}} & \begin{array}{c} \begin{array}{c} \text{a, } \bar{1} \\ \downarrow \\ \text{3} \end{array} \\ \{\circ, \bullet\} \end{array} \end{array} \right).$$

(P is of the form of a *path graph* by taking the *quotient graph* w.r.t. l -labelled edges.) P is an abstraction of edge-saturated *graphs*. From P , we can construct a *graph* $H \in \mathcal{S}_{\text{RSUB}}(\mathcal{G}(a\bar{a}a))$ s.t. $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$. Because both $\delta_a^J(\{\bullet\}) \subseteq \{\circ, \bullet\}$ and $\delta_a^J(\{\circ\}) \subseteq \{\circ, \bullet\}$ hold, in addition to the *graph* H in Example 6.3, for instance, the following are also possible:



Thanks to *saturable paths*, we can replace the existence of such gray-colored edges connecting distant vertices with a locally defined witness U . \lrcorner

Moreover, we can replace (P-Sat)—a condition between distant vertices—with a local condition. Let $\varphi^J(\mathcal{U}, U) \triangleq \forall a \in \mathbf{V}, \forall \langle u, u' \rangle \in \mathcal{U}, \delta_a^J(u) \subseteq U \vee \delta_a^J(u') \subseteq U$.

► **Proposition 6.7.** *Let J and H be graphs. Let $i \in |H|$. Then we have:*

$$(\forall a \in \mathbf{V}, \forall j \text{ s.t. } \langle j, i \rangle \in \bar{l}^H, \delta_a^J(U_j) \subseteq U_i \vee \delta_a^J(U_j) \subseteq U_i) \Leftrightarrow \varphi^J\left(\bigcup_{j; \langle j, i \rangle \in \bar{l}^H} U_j \times U_j, U_i\right).$$

Proof. $(\forall a \in \mathbf{V}, \delta_a^J(U_j) \subseteq U_i \vee \delta_a^J(U_j) \subseteq U_i)$ iff $(\forall a \in \mathbf{V}, (\forall u \in U_j, \delta_a^J(u) \subseteq U_i) \vee (\forall u' \in U_j, \delta_a^J(u') \subseteq U_i))$ iff $\varphi^J(U_j \times U_j, U_i)$ (by taking the prenex normal form). Thus by $(\forall j \text{ s.t. } \langle j, i \rangle \in \bar{l}^H, \varphi^J(U_j \times U_j, U_i))$ iff $\varphi^J(\bigcup_{j; \langle j, i \rangle \in \bar{l}^H} U_j \times U_j, U_i)$, this completes the proof. ◀

6.2 Automata from saturable paths

Let $\mathcal{X} \triangleq \{X \in \wp(\tilde{\mathbf{V}}_1) \mid \bar{l}, \top \in X, \bar{l} \notin X, \text{ and } \forall x \in \mathbf{V}, x \in X \leftrightarrow \bar{x} \notin X\}$. (This set is equivalent to the set $\{\{x \in \tilde{\mathbf{V}}_1 \mid l^H \subseteq x^H\} \mid H \in \text{GR}_{\text{RSUB}}\}$.)

► **Definition 6.8 (NFA construction).** *Let ► and ◀ be two fresh symbols. For a graph J and a set $X \in \mathcal{X}$, let J^{S^X} be the graph G defined as follows:*

- $|G| = \{\blacktriangleright, \blacktriangleleft\} \cup \{\langle \mathcal{U}, U \rangle \in \wp(|J| \times |J|) \times \wp(|J|) \mid \varphi^J(\mathcal{U}, U) \wedge \forall x \in X, \delta_x^J(U) \subseteq U\}$;
- $l^G = \{\blacktriangleright\} \times \{\langle \mathcal{U}, U \rangle \in |G| \mid \mathbf{1}^J \in U \wedge \mathcal{U} = \emptyset\} \cup \{\langle \mathcal{U}, U \rangle \in |G| \mid \mathbf{2}^J \notin U\} \times \{\blacktriangleleft\}$;
- $x^G = \{\langle \langle \mathcal{U}, U \rangle, \langle \mathcal{U}', U' \rangle \rangle \in |G| \times |G| \mid \psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U') \vee \psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U')\}$ for $x \in \tilde{\mathbf{V}}$;
- $\mathbf{1}^G = \blacktriangleright$;
- $\mathbf{2}^G = \blacktriangleleft$.


Here, $\psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U')$ and $\psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U')$ are as follows:

- $\psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U') \triangleq \left(\mathcal{U}' = \mathcal{U} \cup (U \times U) \wedge \bigwedge \left\{ \begin{array}{l} \delta_x^J(U) \subseteq U' \\ \delta_{\bar{l}}^J(\{u \mid \langle u, u \rangle \in \mathcal{U}'\}) \subseteq U' \\ \delta_{\bar{l}}^J(\{u \mid \langle u, u \rangle \in \mathcal{U}'\}) \subseteq U' \end{array} \right\} \right)$;
- $\psi_{x,1}^X(\mathcal{U}, U, \mathcal{U}', U') \triangleq (\mathcal{U}' = \mathcal{U} \wedge U' = U \wedge x \in X)$. \lrcorner

By the form of J^{S^X} , if $a_1 \dots a_n \in [J^{S^X}]_{\tilde{\mathbf{V}}}$, then its run is of the following form:

$$\blacktriangleright \longrightarrow \langle \mathcal{U}_0, U_0 \rangle \xrightarrow{-a_1} \langle \mathcal{U}_1, U_1 \rangle \xrightarrow{-a_2} \langle \mathcal{U}_2, U_2 \rangle \longrightarrow \dots \longrightarrow \langle \mathcal{U}_n, U_n \rangle \longrightarrow \blacktriangleleft \longrightarrow .$$

Intuitively, this run corresponds to the following *saturable path* where some \top -, \bar{l} -, or l -labelled edges are omitted and \bigvee_x^X denotes that an x -labelled edge exists for each $x \in X$:



487 Here, \mathcal{U}_i is used to denote the set $\bigcup_{j:(j,i) \in \bar{H}} U_j \times U_j$ (cf. Prop. 6.7) where H is the [graph](#) of
 488 the [saturable path](#) above.

489 According to the correspondence above, from a [word](#) $w \in \bigcup_{X \in \mathcal{X}} [J^{S^X}]_{\bar{V}}$, we can construct
 490 a [saturable path](#) for $w \in L_J$. Conversely, from a [saturable path](#) for $w \in L_J$, we can show
 491 $w \in \bigcup_{X \in \mathcal{X}} [J^{S^X}]_{\bar{V}}$. Thus, we have the following. See Appendix C for the detailed proof.

492 ► **Lemma 6.9** (Appendix C). *Let J be a [graph](#). We have $L_J = \bigcup_{X \in \mathcal{X}} [J^{S^X}]_{\bar{V}}$.*

493 ► **Theorem 6.10**. *The [equational theory](#) of $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ w.r.t. languages is PSPACE-complete.*

494 **Proof.** (in PSPACE): Let t and s be $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ [terms](#). Let G and J be [graphs](#) s.t. $[G]_{\bar{V}} = [t]_{\bar{V}}$
 495 and $[J]_{\bar{V}} = [s]_{\bar{V}}$. Then we have

$$496 \quad \text{RSUB} \models t \leq s \quad \Leftrightarrow \quad [G]_{\bar{V}} \cap L_J = \emptyset \quad (\text{Cor. 5.4})$$

$$497 \quad \Leftrightarrow \quad [G]_{\bar{V}} \cap \left(\bigcup_{X \in \mathcal{X}} [J^{S^X}]_{\bar{V}} \right) = \emptyset. \quad (\text{Lem. 6.9})$$

499 Hence, we can reduce the [equational theory](#) into the emptiness problem of [NFAs](#) of size
 500 exponential to the size of the input [equation](#). Therefore, by using a standard on-the-fly
 501 algorithm for the non-emptiness problem of NFAs (essentially the graph reachability problem),
 502 we can give a non-deterministic polynomial space algorithm. (Hardness): The [equational](#)
 503 [theory](#) of KA coincides with the language equivalence problem of regular expressions, which
 504 is PSPACE-complete [16]. Hence, the [equational theory](#) of $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ is PSPACE-hard. ◀

505 ► **Remark 6.11.** Cf. [18, Remark 45]: the [equational theory](#) of REL for $\text{KA}_{\{\bar{x}, \bar{1}, \top\}}$ is decidable
 506 in coNEXP, but its PSPACE-decidability is open. In RSUB , each equivalence class induced
 507 from l -labelled edges is always an interval. Thus, the problematic case of [18, Remark 45]
 508 does not occur in RSUB , fortunately. ▽

509 **7 Future directions**

510 We leave open the axiomatizability of the [equational theory](#) of $\text{KA}_{\{\bar{x}\}}/\text{KA}_{\{\bar{1}\}}/\text{KA}_{\{\bar{x}, \bar{1}\}}$ w.r.t.
 511 languages. Note that the [equational theory](#) of $\text{KA}_{\{-\}}$ (without [Kleene star](#)) w.r.t. languages
 512 is not finitely axiomatizable because it is Π_1^0 -complete (Thms. 4.7 and 4.10).

513 A natural interest is to consider variants or fragments of $\text{KA}_{\{-\}}$, e.g., with reverse [3], with
 514 tests [14] (by considering guarded strings) or with (anti-)domain [8]. Also, the combination
 515 of [variables](#) and [letters](#) ([variables](#) under standard [valuations](#)) would be interesting in the
 516 context of regular expressions.

517 Additionally, this paper did not so deal with the expressive power w.r.t. languages. A
 518 weak tool is (\dagger - $\text{KA}_{\{-\}}$): $[t] \neq [s] \Rightarrow \text{LANG} \not\models t = s$; we can separate the expressive powers
 519 w.r.t. languages of given two term sets if their language classes are not equivalent. However,
 520 this cannot show e.g., the expressive power difference between $\text{KA}_{\{\bar{x}\}}$ and KA , while this
 521 difference is an immediate consequence of the counter-example in the introduction and the
 522 completeness theorem of KA . To obtain a more useful tool, it is possibly interesting to modify
 523 existing arguments about the standard relational structures (REL), e.g., [25, 24, 26, 11, 17]
 524 (including finite model theoretical studies such as Ehrenfeucht-Fraïssé games [9, 10]) for
 525 RSUB .

526 **References**

- 527 **1** Hajnal Andréka and D. A. Bredikhin. The equational theory of union-free algebras of relations.
 528 *Algebra Universalis*, 33(4):516–532, 1995. doi:10.1007/BF01225472.

- 529 2 Hajnal Andréka, Szabolcs Mikulás, and István Németi. The equational theory of Kleene lattices.
530 *Theoretical Computer Science*, 412(52):7099–7108, 2011. doi:10.1016/J.TCS.2011.09.024.
- 531 3 S. L. Bloom, Z. Ésik, and Gh. Stefanescu. Notes on equational theories of relations. *algebra*
532 *universalis*, 33(1):98–126, 1995. doi:10.1007/BF01190768.
- 533 4 Paul Brunet. Reversible Kleene lattices. In *MFCS*, volume 83 of *LIPICs*, pages 66:1–66:14.
534 Schloss Dagstuhl, 2017. doi:10.4230/LIPICs.MFCS.2017.66.
- 535 5 Paul Brunet. A complete axiomatisation of a fragment of language algebra. In *CSL*, volume
536 152 of *LIPICs*, page 11:1–11:15. Schloss Dagstuhl, 2020. doi:10.4230/LIPICs.CSL.2020.11.
- 537 6 Paul Brunet and Damien Pous. Petri automata. *Logical Methods in Computer Science*, 13(3),
538 2017. doi:10.23638/LMCS-13(3:33)2017.
- 539 7 John H. Conway. *Regular Algebra and Finite Machines*. Chapman and Hall, 1971.
- 540 8 Jules Desharnais, Bernhard Möller, and Georg Struth. Kleene algebra with domain. *ACM*
541 *Transactions on Computational Logic*, 7(4):798–833, 2006. doi:10.1145/1183278.1183285.
- 542 9 Andrzej Ehrenfeucht. An application of games to the completeness problem for formalized
543 theories. *Fundamenta Mathematicae*, 49(2):129–141, 1961. doi:10.4064/fm-49-2-129-141.
- 544 10 Roland Fraïssé. Sur les classifications des systems de relations. *Publ. Sci. Univ. Alger I*, 1954.
- 545 11 Jelle Hellings. *On Tarski’s Relation Algebra: querying trees and chains and the semi-join*
546 *algebra*. PhD thesis, Hasselt University and Transnational University of Limburg, 2018. URL:
547 <http://hdl.handle.net/1942/25932>.
- 548 12 Stephen C. Kleene. Representation of events in nerve nets and finite automata. In *Au-*
549 *tomata Studies. (AM-34)*, pages 3–42. Princeton University Press, 1956. doi:10.1515/
550 9781400882618-002.
- 551 13 Dexter Kozen. A completeness theorem for Kleene algebras and the algebra of regular events.
552 In *LICS*, pages 214–225. IEEE Comput. Soc. Press, 1991. doi:10.1109/LICS.1991.151646.
- 553 14 Dexter Kozen and Frederick Smith. Kleene algebra with tests: Completeness and decidability. In
554 *CSL*, volume 1258 of *LNCS*, pages 244–259. Springer, 1996. doi:10.1007/3-540-63172-0_43.
- 555 15 F. W. Levi. On semigroups. *Bulletin of the Calcutta Mathematical Society*, (36):141–146, 1944.
- 556 16 A. R. Meyer and L. J. Stockmeyer. The equivalence problem for regular expressions with
557 squaring requires exponential space. In *SWAT*, pages 125–129. IEEE, 1972. doi:10.1109/
558 SWAT.1972.29.
- 559 17 Yoshiki Nakamura. Expressive power and succinctness of the positive calculus of binary
560 relations. *Journal of Logical and Algebraic Methods in Programming*, 127:100760, 2022.
561 doi:10.1016/j.jlamp.2022.100760.
- 562 18 Yoshiki Nakamura. Existential calculi of relations with transitive closure: Complexity and
563 edge saturations. In *LICS*, pages 1–13. IEEE, 2023. doi:10.1109/LICS56636.2023.10175811.
- 564 19 Yoshiki Nakamura and Ryoma Sin’ya. Words-to-letters valuations for language Kleene algebras
565 with variable complements. In *AFL*, volume 386, pages 185–199. EPTCS, 2023. doi:10.4204/
566 EPTCS.386.15.
- 567 20 Damien Pous. On the positive calculus of relations with transitive closure. In *STACS*, volume 96
568 of *LIPICs*, pages 3:1–3:16. Schloss Dagstuhl, 2018. doi:10.4230/LIPICs.STACS.2018.3.
- 569 21 Damien Pous and Jana Wagemaker. Completeness theorems for kleene algebra with top. In
570 *CONCUR*, volume 243 of *LIPICs*, pages 26:1–26:18. Schloss Dagstuhl, 2022. doi:10.4230/
571 LIPICs.CONCUR.2022.26.
- 572 22 V. R. Pratt. Dynamic algebras and the nature of induction. In *STOC*, page 22–28. ACM,
573 1980. doi:10.1145/800141.804649.
- 574 23 Robin Hirsch, Ian Hodkinson. *Relation Algebras by Games*, volume 147 of *Studies in logic and*
575 *the foundations of mathematics*. Elsevier, 1 edition, 2002.
- 576 24 Dimitri Surinx. A framework for comparing query languages in their ability to express boolean
577 queries. 2017. URL: <http://hdl.handle.net/1942/25215>.
- 578 25 Dimitri Surinx, George H. L. Fletcher, Marc Gyssens, Dirk Leinders, Jan Van den Bussche, Dirk
579 Van Gucht, Stijn Vansummeren, and Yuqing Wu. Relative expressive power of navigational

- 580 querying on graphs using transitive closure. *Logic Journal of the IGPL*, 23(5):759–788, 2015.
 581 doi:10.1093/jigpal/jzv028.
- 582 **26** Dimitri Surinx, Jan Van den Bussche, and Dirk Van Gucht. The primitivity of operators
 583 in the algebra of binary relations under conjunctions of containments. In *LICS*, pages 1–10.
 584 IEEE, 2017. doi:10.1109/LICS.2017.8005122.
- 585 **27** Alfred Tarski. On the calculus of relations. *The Journal of Symbolic Logic*, 6(3):73–89, 1941.
 586 doi:10.2307/2268577.
- 587 **28** Cheng Zhang, Arthur Azevedo de Amorim, and Marco Gaboardi. On incorrectness logic
 588 and Kleene algebra with top and tests. *Proceedings of the ACM on Programming Languages*,
 589 6(POPL):29:1–29:30, 2022. doi:10.1145/3498690.

A Proof for Remark 3.12

591 ▶ **Definition A.1.** *Let*

$$592 \text{SUB}_{\text{st}} \triangleq \bigcup_{n \in \mathbb{N}} \left\{ \mathbf{v} \in \text{SUB}_n \mid \bigcup_{a \in \mathbf{V}} \mathbf{v}(a) = \{\ell_i \mid i \in [0, n-1]\} \right\}.$$

593 ▶ **Proposition A.2.** *Let $n \in \mathbb{N}$ and let $\mathbf{v}' \in \text{SUB}_n \cap \text{SUB}_{\text{st}}$. For $i \in [0, n-1]$, let a_i be the
 594 unique one s.t. $\mathbf{v}'(a_i) = \{\ell_i\}$. Then $\mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle} = \mathbf{v}'$.*

595 **Proof.** By Def. 3.2, $\ell_i \in \mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle}(x)$ iff $a_i \in \mathbf{v}_{\text{st}}(x)$ iff $a_i = x$ iff $\ell_i \in \mathbf{v}'(x)$. Also, both
 596 $\mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle}(x)$ and $\mathbf{v}'(x)$ are subsets of $\{\ell_0, \dots, \ell_{n-1}\}$. Thus, $\mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle} = \mathbf{v}'$. ◀

597 ▶ **Lemma A.3** (cf. Lem. 3.6). *For all quantifier-free formulas φ , we have:*

$$598 \mathbf{v}_{\text{st}} \models \varphi \iff \text{SUB}_{\text{st}} \models \varphi.$$

599 **Proof.** As with Lem. 3.6, it suffices to prove when φ is of the form $(\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp)$.

$$600 \text{SUB}_{\text{st}} \not\models (\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp)$$

$$601 \Rightarrow \exists n, a_0, \dots, a_{n-1} \in \mathbf{V}, \mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle} \not\models (\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp) \quad (\text{Prop. A.2})$$

$$602 \Rightarrow \mathbf{v}_{\text{st}} \not\models (\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp) \quad (\text{By Lem. 3.4})$$

$$603 \Rightarrow \exists n \in \mathbb{N}, i_1, j_1, \dots, i_m, j_m \text{ s.t. } 0 \leq i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_m \leq j_m \leq n,$$

$$604 \exists a_0, \dots, a_{n-1} \in \mathbf{V}, (\bigwedge_{k=1}^m a_{i_k} \dots a_{j_k-1} \in \hat{\mathbf{v}}_{\text{st}}(t_k)) \wedge \hat{\mathbf{v}}_{\text{st}}(s) = \emptyset$$

$$605 \Rightarrow \text{SUB}_{\text{st}} \not\models (\bigvee_{k=1}^m t_k \leq \perp) \vee (\neg s \leq \perp).$$

$$606 \quad (\text{By Lem. 3.4 and } \mathbf{v}_{\text{st}}^{\langle a_0, \dots, a_{n-1} \rangle} \in \text{SUB}_n \cap \text{SUB}_{\text{st}}) \quad \blacktriangleleft$$

607 ▶ **Proposition A.4** (Cor. of Prop. 3.8). *For all quantifier-free formulas φ , we have:*

$$608 \text{SUB}_{\text{st}} \models \varphi \iff \text{RSUB}_{\text{st}} \models \varphi.$$

609 **Proof.** By the same bijective map $f: L \mapsto \{\langle i, j \rangle \mid \ell_i \dots \ell_{j-1} \in L\}$ as Prop. 3.8. ◀

610 Combining Lem. A.3 and Prop. A.4 yields

$$611 \quad \mathbf{v}_{\text{st}} \models \varphi \iff \text{RSUB}_{\text{st}} \models \varphi.$$

612 Particularly, we have

$$613 \quad [t] = [s] \iff \text{RSUB}_{\text{st}} \models t = s.$$

614 This matches the following correspondence between [words](#) and binary relations (cf.
615 Sect. 5).

$$616 \quad a_0 a_1 \dots a_{n-1} \quad | \quad \rightarrow \circ \rightarrow a_0 \rightarrow \circ \rightarrow a_1 \rightarrow \circ \rightarrow \dots \rightarrow a_{n-1} \rightarrow \circ \rightarrow .$$

617 **B Proof of Prop. 6.2**

618 Let $R' \subseteq |H| \times |J|$ be the minimal set such that

- 619 ■ $\langle \mathbf{1}^H, \mathbf{1}^J \rangle \in R'$;
- 620 ■ $\forall a \in \tilde{\mathbf{V}}_1, x, y, x', y', \langle x, y \rangle \in R' \wedge \langle x, x' \rangle \in \delta_a^H \wedge \langle y, y' \rangle \in \delta_a^J \Rightarrow \langle x', y' \rangle \in R'$.

621 \triangleright Claim B.1. $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} \neq \emptyset \iff \langle 2^H, 2^J \rangle \in R'$.

622 Proof Sketch. R' is the set of all reachable states of the product [NFA](#) of H and J . Hence
623 this completes the proof. \triangleleft

624 Let $R \subseteq |H| \times |J|$ be the minimal set such that

- 625 ■ $\langle \mathbf{1}^H, \mathbf{1}^J \rangle \in R$;
- 626 ■ $\forall a \in \tilde{\mathbf{V}}_1, x, y, x', y', \langle x, y \rangle \in R \wedge \langle x, x' \rangle \in a^H \wedge \langle y, y' \rangle \in \delta_a^J \Rightarrow \langle x', y' \rangle \in R$.

627 \triangleright Claim B.2. $R = R'$.

628 Proof. (\subseteq): Clear, by $a^H \subseteq \delta_a^H$. (\supseteq): By induction on derivations of R' .

- 629 ■ Case $\langle \mathbf{1}^H, \mathbf{1}^J \rangle \in R'$: Trivial by $\langle \mathbf{1}^H, \mathbf{1}^J \rangle \in R$.
- 630 ■ Case $(\langle x, y \rangle \in R' \wedge \langle x, x' \rangle \in \delta_a^H \wedge \langle y, y' \rangle \in \delta_a^J) \Rightarrow \langle x', y' \rangle \in R'$: By IH, $\langle x, y \rangle \in R$.
 - 631 ■ Sub-Case $a \neq \mathbf{1}$: Let $x_0, \dots, x_{n-1}, x_n, \dots, x_m$ be s.t. $\langle x, x' \rangle = \langle x_0, x_m \rangle$ and
 - 632 * for all $i \in [1, n-1]$, $\langle x_{i-1}, x_i \rangle \in \mathbf{1}^H$;
 - 633 * $\langle x_{n-1}, x_n \rangle \in a^H$;
 - 634 * for all $i \in [n+1, m]$, $\langle x_{i-1}, x_i \rangle \in \mathbf{1}^H$.
 Let $y_0 = \dots = y_{n-1} = y$ and $y_n = \dots = y_m = y'$. Then by applying the second rule
635 multiply, we have $\langle x', y' \rangle \in R$.
 - 636 ■ Sub-Case $a = \mathbf{1}$: By reflexivity of $\mathbf{1}^H$, $\langle x, x' \rangle \in (\mathbf{1}^H)^+$. Let x_0, \dots, x_m ($m > 0$) be s.t.
637 $\langle x, x' \rangle = \langle x_0, x_m \rangle$ and
 - 638 * for all $i \in [1, m]$, $\langle x_{i-1}, x_i \rangle \in \mathbf{1}^H$.
 Let $y_0 = y$ and $y_1 = \dots = y_m = y'$. Then by applying the second rule multiply, we
639 have $\langle x', y' \rangle \in R$. \triangleleft

642 **Proof of Prop. 6.2.** (\Rightarrow): By letting U as the map defined by $U(x) \triangleq \{y \mid \langle x, y \rangle \in R\}$. Here,
643 $2^J \notin U_{2^H}$ is shown by $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$ with Claim B.1 and B.2. (\Leftarrow): Let $R'' \triangleq \{\langle x, y \rangle \mid y \in$
644 $U(x)\}$. By the minimality of R , we have $R \subseteq R''$. By $\langle 2^H, 2^J \rangle \notin R''$, we have $\langle 2^H, 2^J \rangle \notin R$.
645 Hence by Claim B.1 and B.2, we have $[J]_{\tilde{\mathbf{V}}} \cap [H]_{\tilde{\mathbf{V}}} = \emptyset$. \blacktriangleleft

C Proof of Lem. 6.9

646

647 **Proof of Lem. 6.9.** (\subseteq): Let $w = a_1 \dots a_n \in \mathbb{L}_J$. Let $P = \langle H, U \rangle$ be a **saturable path** for $w \in$
 648 \mathbb{L}_J . Let $X \triangleq \{a \in \tilde{\mathbf{V}}_1 \mid a^H \supseteq \mathbb{I}^H\}$ (note that $X \in \mathcal{X}$). For each i , let $\mathcal{U}_i \triangleq \bigcup_{j; \langle j, i \rangle \in \bar{\mathbb{I}}^H} U_j \times U_j$.
 649 Then we have:

650 ■ $\varphi^J(\mathcal{U}_i, U_i)$: By (P-Sat) and Prop. 6.7.

651 ■ $\forall a \in X, \delta_a^J(U_i) \subseteq U_i$: By $a^H \supseteq \mathbb{I}^H \supseteq \Delta_{\mathbb{I}^H}$ and (P-Wit).

652 Thus $\langle \mathcal{U}_i, U_i \rangle \in |J^{\mathcal{S}^X}|$. We consider the following run of the **NFA** $J^{\mathcal{S}^X}$ on w :

653 $\rightarrow \blacktriangleright \longrightarrow \langle \mathcal{U}_0, U_0 \rangle \xrightarrow{-a_1} \langle \mathcal{U}_1, U_1 \rangle \xrightarrow{-a_2} \langle \mathcal{U}_2, U_2 \rangle \longrightarrow \dots \xrightarrow{-a_n} \langle \mathcal{U}_n, U_n \rangle \longrightarrow \blacktriangleleft \rightarrow .$

654 This is indeed a run of the **NFA** $J^{\mathcal{S}^X}$ as follows:

655 ■ $\langle \blacktriangleright, \langle \mathcal{U}_0, U_0 \rangle \rangle \in |J^{\mathcal{S}^X}|$: By $1^J \in U_0$ (P-Wit) and $\mathcal{U}_0 = \emptyset$.

656 ■ $\langle \langle \mathcal{U}_n, U_n \rangle, \blacktriangleleft \rangle \in |J^{\mathcal{S}^X}|$: By $2^J \notin U_n$ (P-Wit).

657 ■ $\forall i \in [1, n], \langle \langle \mathcal{U}_{i-1}, U_{i-1} \rangle, \langle \mathcal{U}_i, U_i \rangle \rangle \in a_i^{J^{\mathcal{S}^X}}$: We distinguish the following cases:

658 ■ Case $\langle i-1, i \rangle \in \mathbb{I}^H$:

659 * $\mathcal{U}_i = \mathcal{U}_{i-1}$: By $\langle j, i \rangle \in \bar{\mathbb{I}}^H$ iff $\langle j, i-1 \rangle \in \bar{\mathbb{I}}^H$, for all j .

660 * $U_i = U_{i-1}$: By (P-Wit), we have $U_{i-1} \subseteq \delta_1^J(U_{i-1}) \subseteq U_i \subseteq \delta_1^J(U_i) \subseteq U_{i-1}$.

661 * $a_i \in X$ ($a_i^H \supseteq \mathbb{I}^H$): By $a_i^H \cap \mathbb{I}^H \neq \emptyset$ and (P-Ext), we have $a_i^H = a_i^{\mathbb{G}(w)} \cup \mathbb{I}^H$ (if not,
 662 this contradicts to (P-Con)).

663 Thus by $\psi_{a_i, 1}^X(\mathcal{U}_{i-1}, U_{i-1}, \mathcal{U}_i, U_i)$, we have $\langle \langle \mathcal{U}_{i-1}, U_{i-1} \rangle, \langle \mathcal{U}_i, U_i \rangle \rangle \in a_i^{J^{\mathcal{S}^X}}$.

664 ■ Case $\langle i-1, i \rangle \in \bar{\mathbb{I}}^H$:

665 * $\mathcal{U}_i = \mathcal{U}_{i-1} \cup (U_{i-1} \times U_{i-1})$: By $\langle j, i \rangle \in \bar{\mathbb{I}}^H$ iff $\langle j, i-1 \rangle \in \bar{\mathbb{I}}^H \vee j = i-1$, for all j .

666 * $\delta_{a_i}^J(U_{i-1}) \subseteq U_i$: By (P-Wit).

667 * $\delta_1^J(\{u \mid \langle u, u \rangle \in \mathcal{U}_i\}) \subseteq U_i$: We have $\delta_1^J(\{u \mid \langle u, u \rangle \in \mathcal{U}_i\}) = \delta_1^J(\bigcup_{j; \langle j, i \rangle \in \bar{\mathbb{I}}^H} U_j) =$
 668 $\bigcup_{j < i} \delta_1^J(U_j) \subseteq U_i$ by (P-Wit).

669 * $\delta_1^J(\{u \mid \langle u, u \rangle \in \mathcal{U}_i\}) \subseteq U_i$: We have $\delta_1^J(\{u \mid \langle u, u \rangle \in \mathcal{U}_i\}) = \delta_1^J(\bigcup_{j; \langle j, i \rangle \in \bar{\mathbb{I}}^H} U_j) =$
 670 $\bigcup_{j < i} \delta_1^J(U_j) \subseteq U_i$ by (P-Wit).

671 Thus by $\psi_{a_i, 1}^X(\mathcal{U}_{i-1}, U_{i-1}, \mathcal{U}_i, U_i)$, we have $\langle \langle \mathcal{U}_{i-1}, U_{i-1} \rangle, \langle \mathcal{U}_i, U_i \rangle \rangle \in a_i^{J^{\mathcal{S}^X}}$.

672 Hence, $w \in [J^{\mathcal{S}^X}]$.

673 (\supseteq): Let $X \subseteq \mathcal{X}$ and $w = a_1 \dots a_n \in [J^{\mathcal{S}^X}]_{\tilde{\mathbf{V}}}$. Let the run of $J^{\mathcal{S}^X}$ on w be as follows:

674 $\rightarrow \blacktriangleright \longrightarrow \langle \mathcal{U}_0, U_0 \rangle \xrightarrow{-a_1} \langle \mathcal{U}_1, U_1 \rangle \xrightarrow{-a_2} \langle \mathcal{U}_2, U_2 \rangle \longrightarrow \dots \xrightarrow{-a_n} \langle \mathcal{U}_n, U_n \rangle \longrightarrow \blacktriangleleft \rightarrow .$

675 Let H be the **edge-extension** of $\mathbb{G}(w)$ defined as follows:

676 ■ $\top^H = \{\langle x, y \rangle \in [0, n] \times [0, n] \mid \forall i \in [y+1, x], \neg \psi_{a_i, 1}^X(\mathcal{U}_{i-1}, U_{i-1}, \mathcal{U}_i, U_i)\}$;

677 ■ $\mathbb{I}^H = \top^H \cap \{\langle x, y \rangle \mid \langle y, x \rangle \in \top^H\}$ and $\bar{\mathbb{I}}^H = \top^H \setminus \mathbb{I}^H$;

678 ■ $\forall a \in \mathbf{V} \cap X, \langle a^H, \bar{a}^H \rangle = \langle a^{\mathbb{G}(w)} \cup \mathbb{I}^H, \bar{a}^{\mathbb{G}(w)} \rangle$.


679 Note that by definition of \top^H , we have

680 ■ $\top^H \supseteq \{\langle x, y \rangle \mid x \leq y\}$;

681 ■ \top^H is transitive by case analysis.

682 Hence, \top^H is a total preorder and each equivalence class w.r.t. \mathbb{I}^H is an interval $[l, r]$.

683 Let $P \triangleq \langle H, U \rangle$ where U is defined as $i \mapsto U_i$ for $i \in [0, n]$. The following depicts P .



685 Then P is a **saturable path** for $w \in \mathbb{L}_J$ as follows:

686 ■ (P-Ext): By the definition of H .

687 ■ (P-Con): Assume that $a^{H^\ominus} \cap \bar{a}^{H^\ominus} \neq \emptyset$. Let x, x', y, y' be s.t. $[x]_{1^H} = [x']_{1^H}$, $[y]_{1^H} = [y']_{1^H}$,
688 $\langle x, y \rangle \in a^H$, and $\langle x', y' \rangle \in \bar{a}^H$. WLOG, we can assume that $a \in X$ and $\bar{a} \notin X$. Then

- 689 ■ $\langle x', y' \rangle \in \bar{a}^{G(w)}$ (so, $x' = y' - 1$ and $a_{y'} = \bar{a}$): By $\bar{a}^H = \bar{a}^{G(w)}$ (since $\bar{a} \notin X$).
- 690 ■ $\langle x, y \rangle \in a^{G(w)}$ (so, $x = y - 1$ and $a_y = a$): If not, then by $a^H = a^G \cup 1^H$, we
691 have $[x]_{1^H} = [y]_{1^H}$. Thus we have $[y' - 1]_{1^H} = [y']_{1^H}$. Then by the definition
692 of \top^H , we have $\neg\psi_{\bar{a},1}^X(\mathcal{U}_{y'-1}, U_{y'-1}, \mathcal{U}_{y'}, U_{y'})$. By the definition of $a^{J^{S^X}}$, we have
693 $\psi_{\bar{a},1}^X(\mathcal{U}_{y'-1}, U_{y'-1}, \mathcal{U}_{y'}, U_{y'})$, so $\bar{a} \in X$. This contradicts $\bar{a} \notin X$.
- 694 ■ $([x, x'] \cup [x', x]) \cap ([y, y'] \cup [y', y]) = \emptyset$ (so, $x = x'$ and $y = y'$): If not, then because the
695 interval between x and x' and that between y and y' have an intersection, we have
696 $[x]_{1^H} = [y]_{1^H}$. Then, in the same manner as above, we have $\bar{a} \in X$. This contradicts
697 $\bar{a} \notin X$.

698 Thus, we reach a contradiction, because $a = a_y = a_{y'} = \bar{a}$. Hence, $a^{H^\ominus} \cap \bar{a}^{H^\ominus} = \emptyset$.

699 ■ (P-Sat): By the the form of J^{S^X} , we have $\mathcal{U}_x = \begin{cases} \mathcal{U}_{x-1} & (\langle x-1, x \rangle \in 1^H) \\ \mathcal{U}_{x-1} \cup (U_{x-1} \times U_{x-1}) & (\langle x-1, x \rangle \in \bar{1}^H) \end{cases}$.

700 Thus, $\mathcal{U}_y = \bigcup_{x; \langle x, y \rangle \in \bar{1}^H} \mathcal{U}_x \times U_x$ (★). Thus by Prop. 6.7, this completes the proof.

701 ■ (P-Wit): For $1^J \in U_0$ and $2^J \notin U_n$, they are shown by the form of J^{S^X} . For $\forall a \in$
702 $\bar{\mathbf{V}}_1, \forall \langle x, y \rangle \in a^H, \delta_a^J(U_x) \subseteq U_y$, we distinguish the following cases:

- 703 ■ Case $a = 1$: Then we have
 - 704 * $U_x = U_y$: By $\langle x, y \rangle \in 1^H$ and the definition of J^{S^X} , we have that $\forall i \in [y +$
705 $1, x]$, $\psi_{a,i,1}^X(\mathcal{U}_{i-1}, U_{i-1}, \mathcal{U}_i, U_i)$. Thus, $U_y = U_{y+1} = \dots = U_x$.
 - 706 * $\delta_1^J(U_x) \subseteq U_x$: By $\langle \mathcal{U}_x, U_x \rangle \in |J^{S^X}|$.
 - 707 Hence, $\delta_1^J(U_x) \subseteq U_y$.
- 708 ■ Case $a = \bar{1}$: Let $z \in [x + 1, y]$ be such that $\psi_{a,z,\bar{1}}^X(\mathcal{U}_{z-1}, U_{z-1}, \mathcal{U}_z, U_z)$ and $\forall z' \in$
709 $[z + 1, y], \neg\psi_{a,z',\bar{1}}^X(\mathcal{U}_{z'-1}, U_{z'-1}, \mathcal{U}_{z'}, U_{z'})$. Then we have

$$\begin{aligned}
 710 \quad \delta_1^J(U_x) &\subseteq \delta_1^J(\{u \mid \langle u, u \rangle \in \mathcal{U}_z\}) && \text{(by (★) and } \langle x, z \rangle \in \bar{1}^H \text{ (by } \langle z-1, z \rangle \in \bar{1}^H)) \\
 711 \quad &\subseteq U_z && \text{(by } \psi_{a,z,\bar{1}}^X(\mathcal{U}_{z-1}, U_{z-1}, \mathcal{U}_z, U_z)) \\
 712 \quad &\subseteq U_{z+1} = \dots = U_y. && \text{(by the form of } J^{S^X}, \psi_{a,z',\bar{1}}^X(\mathcal{U}_{z'-1}, U_{z'-1}, \mathcal{U}_{z'}, U_{z'})) \\
 713
 \end{aligned}$$

714 ■ Case $a = \top$: We distinguish the following two sub-cases:

- 715 * Case $\langle x, y \rangle \in \bar{1}^H$: By the similar argument as Case $a = \bar{1}$.
- 716 * Case $\langle x, y \rangle \in 1^H$: By the similar argument as Case $a = 1$, we have $U_x = U_y$ and
717 $\delta_1^J(U_x) \subseteq U_x$, and thus $\delta_1^J(U_x) \subseteq U_y$.
- 718 ■ Case $a \in \{a, \bar{a} \mid a \in \mathbf{V}\}$: We distinguish the following sub-cases:
 - 719 * Case $\langle x, y \rangle \in \bar{1}^H$: By $\langle x, y \rangle \in a^H \cap \bar{1}^H = a^{G(w)}$, we have $x = y - 1$ and $a_y = a$. Thus
720 by $\psi_{a,y,\bar{1}}^X(\mathcal{U}_{y-1}, U_{y-1}, \mathcal{U}_y, U_y)$, we have $\delta_a^J(U_x) \subseteq U_y$.
 - 721 * Case $a \notin X$: By $a^H = a^{G(w)}$, we have $x = y - 1$ and $a_y = a$. By the form of J^{S^X}
722 with $\neg\psi_{a,y,\bar{1}}^X(\mathcal{U}_{y-1}, U_{y-1}, \mathcal{U}_y, U_y)$ (since $a_y \notin X$), we have $\psi_{a,y,\bar{1}}^X(\mathcal{U}_{y-1}, U_{y-1}, \mathcal{U}_y, U_y)$.
723 Hence, $\delta_a^J(U_x) \subseteq U_y$.
 - 724 * Case $\langle x, y \rangle \in 1^H$ and $a \in X$: By the similar argument as Case $a = 1$, we have
725 $U_x = U_y$ (by $\langle x, y \rangle \in 1^H$) and $\delta_a^J(U_x) \subseteq U_x$ (by $a \in X$). Thus, $\delta_a^J(U_x) \subseteq U_y$. ◀

D

 Explicit proof of Lem. 3.4

(The following proof is almost the same as [19, Lem. 23] except that we consider **letters-to-letters valuations** instead of **words-to-letters valuations**.)

Proof of Lem. 3.4. By easy induction on t .

Case $t = x$: By the definition of $\mathfrak{v}^{(a_0, \dots, a_{n-1})}$, we have $\ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(x) \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(x)$.

Case $t = \perp$: By $\ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(\perp) \Leftrightarrow \text{False} \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(\perp)$.

Case $t = \mathbf{l}$: By $\ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(\mathbf{l}) \Leftrightarrow i = j \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(\mathbf{l})$.

Case $t = s \cup u$: By IH, we have $\ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s \cup u) \Leftrightarrow \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s) \vee \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(u) \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(s) \vee a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(u) \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(s \cup u)$.

Case $t = s^-$: By IH, we have $\ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s^-) \Leftrightarrow \ell_i \dots \ell_{j-1} \notin \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s) \Leftrightarrow a_i \dots a_{j-1} \notin \hat{\mathfrak{v}}(s) \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(s^-)$.

Case $t = s \cdot u$: We have

$$\begin{aligned}
& \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s \cdot u) \\
& \Leftrightarrow \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s) \cdot \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(u) \\
& \quad \text{(By the definition of } \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})} \text{ and that } \ell_i \dots \ell_{j-1} \text{ is a subword of } \ell_0 \dots \ell_{n-1}\text{)} \\
& \Leftrightarrow \bigvee_{i \leq k \leq j} (\ell_i \dots \ell_{k-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s) \wedge \ell_k \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(u)) \quad \text{(By concatenation)} \\
& \Leftrightarrow \bigvee_{i \leq k \leq j} (a_i \dots a_{k-1} \in \hat{\mathfrak{v}}(s) \wedge a_k \dots a_{j-1} \in \hat{\mathfrak{v}}(u)) \quad \text{(IH)} \\
& \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(s \cdot u). \quad \text{(By concatenation)}
\end{aligned}$$

(Note that the last equivalence holds thanks to that a_i, \dots, a_{j-1} are **letters** not **words**, cf. [19, Lem. 23].)

Case $t = s^*$: We have

$$\begin{aligned}
& \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s^*) \\
& \Leftrightarrow \ell_i \dots \ell_{j-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s)^* \\
& \quad \text{(By the definition of } \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})} \text{ and that } \ell_i \dots \ell_{j-1} \text{ is a subword of } \ell_0 \dots \ell_{n-1}\text{)} \\
& \Leftrightarrow \exists m \in \mathbb{N}, \bigvee_{i=k_0 \leq k_1 \leq \dots \leq k_m=j} \bigwedge_{l=1}^m (\ell_{k_{l-1}} \dots \ell_{k_l-1} \in \hat{\mathfrak{v}}^{(a_0, \dots, a_{n-1})}(s)) \quad \text{(By Kleene star)} \\
& \Leftrightarrow \exists m \in \mathbb{N}, \bigvee_{i=k_0 \leq k_1 \leq \dots \leq k_m=j} \bigwedge_{l=1}^m (a_{k_{l-1}} \dots a_{k_l-1} \in \hat{\mathfrak{v}}(s)) \quad \text{(IH)} \\
& \Leftrightarrow a_i \dots a_{j-1} \in \hat{\mathfrak{v}}(s^*). \quad \text{(By Kleene star)} \quad \blacktriangleleft
\end{aligned}$$

E

 Explicit proof for Sect. 4.1

(The following proof is almost the same as [18, Section VI] except that we consider RSUB instead of REL.)

A **context-free grammar** (CFG) \mathfrak{C} over a finite set A is a tuple $\langle X, \mathcal{R}, s \rangle$, where


- X is a finite set of non-terminal labels s.t. $A \cap X = \emptyset$;
- \mathcal{R} is a finite set of rewriting rules $x \leftarrow w$ of $x \in X$ and $w \in (A \cup X)^*$;
- $s \in X$ is the start label.

762 The relation $x \vdash_{\mathfrak{C}} w$, where $x \in X$ and $w \in A^*$, is defined as the minimal relation closed
 763 under the following rule: for all $n \in \mathbb{N}$, $x, x_1, \dots, x_n \in X$ and $w_0, \dots, w_n, v_1, \dots, v_n \in A^*$, if
 764 $x \leftarrow w_0 x_1 w_1 \dots x_n w_n \in \mathcal{R}$, then $\frac{x_1 \vdash_{\mathfrak{C}} v_1 \quad \dots \quad x_n \vdash_{\mathfrak{C}} v_n}{x \vdash_{\mathfrak{C}} w_0 v_1 w_1 \dots v_n w_n}$. The language $[\mathfrak{C}]$ is defined by
 765 $[\mathfrak{C}] \triangleq \{w \in A^* \mid s \vdash_{\mathfrak{C}} w\}$.

766 For a CFG $\mathfrak{C} = \langle X, \mathcal{R}, s \rangle$ and a word $w = a_1 \dots a_n$, we write $\mathbf{v}_{\mathfrak{C}, w}$ for the following
 767 valuation on RSUB_n :

- 768 ■ $\mathbf{v}_{\mathfrak{C}, w}(x) = \{\langle i-1, i \rangle \mid i \in [1, n] \wedge x = a_i\}$ for $x \in A$;
- 769 ■ $\{\mathbf{v}_{\mathfrak{C}, w}(x)\}_{x \in X}$ are the minimal sets such that for all $x \in X$, $v \in (X \cup A)^*$, and $i, j \in [0, n]$,
 770 if $x \leftarrow v \in \mathcal{R}$, then $\frac{\langle i, j \rangle \in \hat{\mathbf{v}}_{\mathfrak{C}, w}(v)}{\langle i, j \rangle \in \mathbf{v}_{\mathfrak{C}, w}(x)}$.

771 For instance, if $A = \{(\cdot, \cdot)\}$, $\mathfrak{C} = \langle \{s\}, \{s \leftarrow (s)s, s \leftarrow ()\}, s \rangle$ (i.e., $[\mathfrak{C}]$ is the Dyck-1 language),
 772 and $w = ((\cdot))$, then $\mathbf{v}_{\mathfrak{C}, w}$ is of the following form:



774 By construction, we have the following:

775 ► **Lemma E.1** (cf. [18, Lem. 47]). Let $\mathfrak{C} = \langle X, \mathcal{R}, s \rangle$ be a CFG. Let \mathcal{C} be s.t. $\{\mathbf{v}_{\mathfrak{C}, w}\} \subseteq \mathcal{C} \subseteq$
 776 GREL. For all $x \in X$ and $w = a_0 \dots a_{n-1} \in A^*$, the following are equivalent:

- 777 1. $x \vdash_{\mathfrak{C}} w$;
- 778 2. $\mathcal{C} \models (\bigwedge_{(x' \leftarrow w') \in \mathcal{R}} w' \leq x') \rightarrow w \leq x$;
- 779 3. $\langle 0, n \rangle \in \mathbf{v}_{\mathfrak{C}, w}(x)$.

780 **Proof.** 1 \Rightarrow 2: By induction on the derivation tree of $\vdash_{\mathfrak{C}}$. Let

$$781 \frac{x_1 \vdash_{\mathfrak{C}} v_1 \quad \dots \quad x_n \vdash_{\mathfrak{C}} v_n}{x \vdash_{\mathfrak{C}} w_0 v_1 w_1 \dots v_n w_m}$$

782 where $x \leftarrow w_0 x_1 w_1 \dots x_n w_m \in \mathcal{R}$. Let $\mathbf{v} \in \mathcal{C}$ be any s.t. $\mathbf{v} \models \bigwedge_{(x' \leftarrow w') \in \mathcal{R}} w' \leq x'$. Particu-
 783 larly, we have $\mathbf{v} \models w_0 x_1 w_1 \dots x_n w_m \leq x$. By IH, we have $\mathbf{v} \models v_k \leq x_k$. Combining them
 784 yields $\mathbf{v} \models w_0 v_1 w_1 \dots v_n w_m \leq x$. (We only need axioms of monoids, so this holds for any
 785 Kleene algebras beyond GREL.)

786 2 \Rightarrow 3: By $\mathbf{v}_{\mathfrak{C}, w} \in \mathcal{C}$, $\mathbf{v}_{\mathfrak{C}, w} \models \bigwedge_{(x' \leftarrow w') \in \mathcal{R}} w' \leq x'$, and $\langle 0, n \rangle \in \hat{\mathbf{v}}_{\mathfrak{C}, w}(w)$, we have $\langle 0, n \rangle \in$
 787 $\mathbf{v}_{\mathfrak{C}, w}(x)$ by 2.

788 3 \Rightarrow 1: By induction on the derivation tree induced from the definition of $\mathbf{v}_{\mathfrak{C}, w}$. Let

$$789 \frac{\langle i, j \rangle \in \hat{\mathbf{v}}_{\mathfrak{C}, w}(w_0 x_1 w_1 \dots x_m w_m)}{\langle i, j \rangle \in \mathbf{v}_{\mathfrak{C}, w}(x)}$$

790 where $x \leftarrow w_0 x_1 w_1 \dots x_m w_m \in \mathcal{R}$. Let $i = i_0 \leq j_0 \leq \dots \leq i_m \leq j_m = j - 1$ be s.t. $\langle i_k, j_k \rangle \in$
 791 $\hat{\mathbf{v}}_{\mathfrak{C}, w}(w_k)$ and $\langle j_{k-1}, i_k \rangle \in \mathbf{v}_{\mathfrak{C}, w}(x_k)$. By construction of $\mathbf{v}_{\mathfrak{C}, w}$, we have $w_k = a_{i_k} \dots a_{j_k}$. By
 792 IH, $x_k \vdash_{\mathfrak{C}} a_{j_{k-1}} \dots a_{i_k}$. Hence by definition of $\vdash_{\mathfrak{C}}$, we have $x \vdash_{\mathfrak{C}} a_{i_0} \dots a_{j-1}$. ◀

793 ► **Lemma E.2** (cf. [18, Lem. 48]). Let $\mathfrak{C} = \langle X, \mathcal{R}, s \rangle$ be a CFG. Let \mathcal{C} be s.t. $\{\mathbf{v}_{\mathfrak{C}, w} \mid w \in$
 794 $A^*\} \subseteq \mathcal{C} \subseteq \text{GREL}$. Then we have

$$795 [\mathfrak{C}] = A^* \Leftrightarrow \mathcal{C} \models \left(\bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x \right) \rightarrow A^* \leq s.$$

796 Here, A^* denotes the term $(a_1 \cup \dots \cup a_n)^*$ in the right-hand side, where $A = \{a_1, \dots, a_n\}$.

797 **Proof.** We have:

$$\begin{aligned}
798 \quad [\mathcal{C}] &= A^* \\
799 \quad &\Leftrightarrow \forall w' \in A^*, s \vdash_{\mathcal{C}} w' \\
800 \quad &\Leftrightarrow \forall w' \in A^*, \mathcal{C} \models \left(\bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x \right) \rightarrow w' \leq s \quad (\text{By Lem. E.1}) \\
801 \quad &\Leftrightarrow \forall i \in \mathbb{N}, \mathcal{C} \models \left(\bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x \right) \rightarrow A^i \leq s \\
802 \quad &\Leftrightarrow \mathcal{C} \models \left(\bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x \right) \rightarrow A^* \leq s. \\
803 \quad &\quad (\Leftarrow: \text{By GREL } \models A^i \leq A^* \Rightarrow: \text{Because GREL is } * \text{-continuous}) \quad \blacktriangleleft
\end{aligned}$$

804 **Proof of Lem. 4.6.** By Lem. E.2. \(\blacktriangleleft\)

805 Furthermore, we give a tool to eliminate [Kleene star](#).

806 **\(\blacktriangleright\) Proposition E.3.** $\text{RSUB} \models \bar{1} = A\top \rightarrow A^* = \top$.

807 **Proof.** Let $n \in \mathbb{N}$ and $\mathbf{v} \in \text{RSUB}_n$. Let $i \in [1, n]$. By $\langle i-1, i-1 \rangle \notin \hat{\mathbf{v}}(\bar{1}) = \hat{\mathbf{v}}(A\top)$, we
808 have $\langle i-1, i-1 \rangle \notin \hat{\mathbf{v}}(A)$. By $\langle i-1, i \rangle \in \hat{\mathbf{v}}(\bar{1}) = \hat{\mathbf{v}}(A\top)$, we have $\langle i-1, i \rangle \in \hat{\mathbf{v}}(A)$ (by
809 $\langle i-1, i-1 \rangle \notin \hat{\mathbf{v}}(A)$). Thus $\hat{\mathbf{v}}(A^*) = \{\langle i, j \rangle \mid 0 \leq i \leq j \leq n\} = \hat{\mathbf{v}}(\top)$. \(\blacktriangleleft\)

810 **Proof of Lem. 4.9.** Because $\mathbf{v}_{\mathcal{C}, w} \models \bar{1} = A\top$, Lem. E.1 holds even if we can replace 2 with

$$811 \quad \mathcal{C} \models (\bar{1} = A\top \wedge \bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x) \rightarrow w' \leq x'.$$

812 Thus for $\mathcal{C} \subseteq \text{RSUB}$, by the same argument as Lem. E.2 with Prop. E.3, we have

$$813 \quad [\mathcal{C}] = A^* \quad \Leftrightarrow \quad \mathcal{C} \models (\bar{1} = A\top \wedge \bigwedge_{(x \leftarrow w) \in \mathcal{R}} w \leq x) \rightarrow \top \leq s. \quad \blacktriangleleft$$

814 **F** Explicit proof of Lem. 5.5

815 (The following proof is almost the same as [18, Prop. 8, 11] except that we consider GREL
816 instead of REL. Similar arguments can also be found, e.g., in [1, 6, 20].)

817 We use the two notations for [graphs](#), [series-composition](#) (\cdot) and [parallel-composition](#) (\cap) :

$$818 \quad G \cdot H \quad \triangleq \quad \circ \text{---} G \text{---} \circ \text{---} H \text{---} \circ \quad \quad \quad G \cap H \quad \triangleq \quad \circ \text{---} \begin{array}{c} G \\ H \end{array} \text{---} \circ \quad .$$

820 **\(\blacktriangleright\) Proposition F.1.** Let $\mathbf{v} \in \text{GREL}$ and G, H be [graphs](#).

$$821 \quad \hat{\mathbf{v}}(G \cap H) = \hat{\mathbf{v}}(G) \cap \hat{\mathbf{v}}(H) \quad (\text{Prop. F.1}\cap)$$

$$822 \quad \hat{\mathbf{v}}(G \cdot H) = \hat{\mathbf{v}}(G) \cdot \hat{\mathbf{v}}(H) \quad (\text{Prop. F.1}\cdot)$$

824 **Proof.** (Prop. F.1\cap): It suffices to prove that for every x, y ,

$$825 \quad \exists f, f: (G \cap H) \longrightarrow \mathbf{G}(\mathbf{v}, x, y) \Leftrightarrow \exists f_G, f_H, f_G: G \longrightarrow \mathbf{G}(\mathbf{v}, x, y) \wedge f_H: H \longrightarrow \mathbf{G}(\mathbf{v}, x, y).$$

826 \Rightarrow : By letting $f_G = \{\langle x', f(x') \rangle \mid x' \in |G|\}$ and $f_H = \{\langle x', f(x') \rangle \mid x' \in |H|\}$. \Leftarrow : By letting

827 $f = f_G \cup f_H$. Note that $f_G(1^G) = x = f_H(1^H)$ and $f_G(2^G) = y = f_H(2^H)$; so f is indeed a

828 map.

829 (Prop. F.1): It suffices to prove that for every x, y ,

$$830 \quad \exists f, f : (G \cdot H) \longrightarrow \mathbf{G}(\mathbf{v}, x, y) \Leftrightarrow \exists z, \exists f_G, f_H, f_G : G \longrightarrow \mathbf{G}(\mathbf{v}, x, z) \wedge f_H : H \longrightarrow \mathbf{G}(\mathbf{v}, z, y).$$

831 \Rightarrow : By letting $z = f(2^G)$, $f_G = \{\langle x', f(x') \rangle \mid x' \in |G|\}$, and $f_H = \{\langle x', f(x') \rangle \mid x' \in |H|\}$. \Leftarrow :

832 By letting $f = f_G \cup f_H$. Note that $f_G(2^G) = z = f_H(\mathbf{1}^H)$; so f is indeed a map. \blacktriangleleft

833 **Proof of Lem. 5.5.** By easy induction on t .

834 Case $t = x$ where $x \in \tilde{\mathbf{V}} = \{a, \bar{a} \mid a \in \mathbf{V}\} \cup \{\bar{\mathbf{1}}, \top\}$: For every $\langle x, y \rangle \in \hat{\mathbf{v}}(\top)$, we have

$$\begin{aligned} 835 \quad \langle x, y \rangle \in \hat{\mathbf{v}}(a) &\Leftrightarrow \rightarrow\circ\text{---}a\rightarrow\circ\rightarrow \longrightarrow \mathbf{G}(\mathbf{v}, x, y) \\ 836 &\Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\rightarrow\circ\text{---}a\rightarrow\circ\rightarrow) && \text{(Def. of } \hat{\mathbf{v}}) \\ 837 &\Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\mathcal{G}(a)). && \text{(Def. of } \mathcal{G}) \end{aligned}$$

839 Case $t = \mathbf{1}$: For every $\langle x, y \rangle \in \hat{\mathbf{v}}(\top)$, we have

$$\begin{aligned} 840 \quad \langle x, y \rangle \in \hat{\mathbf{v}}(\mathbf{1}) &\Leftrightarrow \rightarrow\circ\text{---}\mathbf{1}\rightarrow\circ\rightarrow \longrightarrow \mathbf{G}(\mathbf{v}, x, y) \\ 841 &\Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\rightarrow\circ\text{---}\mathbf{1}\rightarrow\circ\rightarrow) && \text{(Def. of } \hat{\mathbf{v}}) \\ 842 &\Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\rightarrow\circ\rightarrow) && (\mathbf{1}^{\mathbf{G}(\mathbf{v}, x, y)} \text{ is the identity relation)} \\ 843 &\Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\mathcal{G}(\mathbf{1})). && \text{(Def. of } \mathcal{G}) \end{aligned}$$

845 Case $t = \perp$: For every $\langle x, y \rangle \in \hat{\mathbf{v}}(\top)$, we have

$$846 \quad \langle x, y \rangle \in \hat{\mathbf{v}}(\perp) \Leftrightarrow \text{false} \Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\emptyset) \Leftrightarrow \langle x, y \rangle \in \hat{\mathbf{v}}(\mathcal{G}(\perp)). \quad \text{(Def. of } \hat{\mathbf{v}} \text{ and } \mathcal{G})$$

848 Case $t = s \cdot u$:

$$\begin{aligned} 849 \quad \hat{\mathbf{v}}(s \cdot u) &= \hat{\mathbf{v}}(s) \cdot \hat{\mathbf{v}}(u) && \text{(Def. of } \hat{\mathbf{v}}) \\ 850 &= \hat{\mathbf{v}}(\mathcal{G}(s)) \cdot \hat{\mathbf{v}}(\mathcal{G}(u)) && \text{(IH)} \\ 851 &= \bigcup_{G \in \mathcal{G}(s)} \bigcup_{H \in \mathcal{G}(u)} \hat{\mathbf{v}}(G) \cdot \hat{\mathbf{v}}(H) && (\cdot \text{ is distributive w.r.t. } \cup) \\ 852 &= \bigcup_{G \in \mathcal{G}(s)} \bigcup_{H \in \mathcal{G}(u)} \hat{\mathbf{v}}(G \cdot H) && \text{(Equation (Prop. F.1.))} \\ 853 &= \hat{\mathbf{v}}(\mathcal{G}(s \cdot u)). && \text{(Def. of } \mathcal{G}) \end{aligned}$$

855 Case $t = s \cap u$:

$$\begin{aligned} 856 \quad \hat{\mathbf{v}}(s \cap u) &= \hat{\mathbf{v}}(s) \cap \hat{\mathbf{v}}(u) && \text{(Def. of } \hat{\mathbf{v}}) \\ 857 &= \hat{\mathbf{v}}(\mathcal{G}(s)) \cap \hat{\mathbf{v}}(\mathcal{G}(u)) && \text{(IH)} \\ 858 &= \bigcup_{G \in \mathcal{G}(s)} \bigcup_{H \in \mathcal{G}(u)} (\hat{\mathbf{v}}(G) \cap \hat{\mathbf{v}}(H)) && (\cap \text{ is distributive w.r.t. } \cup) \\ 859 &= \bigcup_{G \in \mathcal{G}(s)} \bigcup_{H \in \mathcal{G}(u)} \hat{\mathbf{v}}(G \cap H) && \text{(Equation (Prop. F.1}\cap)) \\ 860 &= \hat{\mathbf{v}}(\mathcal{G}(s \cap u)). && \text{(Def. of } \mathcal{G}) \end{aligned}$$

862 Case $t = s \cup u$:

$$\begin{aligned} 863 \quad \hat{\mathbf{v}}(s \cup u) &= \hat{\mathbf{v}}(s) \cup \hat{\mathbf{v}}(u) && \text{(Def. of } \hat{\mathbf{v}}) \\ 864 &= \hat{\mathbf{v}}(\mathcal{G}(s)) \cup \hat{\mathbf{v}}(\mathcal{G}(u)) && \text{(IH)} \\ 865 &= \hat{\mathbf{v}}(\mathcal{G}(s) \cup \mathcal{G}(u)) = \hat{\mathbf{v}}(\mathcal{G}(s \cup u)). && \text{(Def. of } \mathcal{G}) \end{aligned}$$

867 Case $t = s^*$:

$$868 \quad \hat{\mathbf{v}}(s^*) = \bigcup_{n \in \mathbb{N}} \hat{\mathbf{v}}(s^n) = \bigcup_{n \in \mathbb{N}} \hat{\mathbf{v}}(s)^n \quad (\text{Def. of } \hat{\mathbf{v}})$$

$$869 \quad = \bigcup_{n \in \mathbb{N}} \hat{\mathbf{v}}(\mathcal{G}(s))^n \quad (\text{IH})$$

$$870 \quad = \bigcup_{n \in \mathbb{N}} \bigcup_{G_1, \dots, G_n \in \mathcal{G}(s)} \hat{\mathbf{v}}(G_1) \cdot \dots \cdot \hat{\mathbf{v}}(G_n) \quad (\cdot \text{ is distributive w.r.t. } \cup)$$

$$871 \quad = \bigcup_{n \in \mathbb{N}} \bigcup_{G_1, \dots, G_n \in \mathcal{G}(s)} \hat{\mathbf{v}}(G_1 \cdot \dots \cdot G_n) \quad (\text{Equation (Prop. F.1.)})$$

$$872 \quad = \bigcup_{n \in \mathbb{N}} \hat{\mathbf{v}}(\mathcal{G}(s^n)) = \hat{\mathbf{v}}(\mathcal{G}(s^*)). \quad (\text{Def. of } \mathcal{G})$$

873

874 ◀