

## Two avirulence genes of Leptosphaeria maculans, displaying suppressive interactions, share a common structural pattern and are part of a larger effector family

Noureddine Lazar, Yohann Petit-Houdenot, Nacera Talbi, Inès Li de la Sierra-Gallay, Zelie Emilie, Karine Blondeau, Angela van de Wouw, Bénédicte Ollivier, Carl H. Mesarich, Alexander Idnurm, et al.

## ▶ To cite this version:

Noured dine Lazar, Yohann Petit-Houdenot, Nacera Talbi, Inès Li de la Sierra-Gallay, Zelie Emilie, et al.. Two avirulence genes of Leptosphaeria maculans, displaying suppressive interactions, share a common structural pattern and are part of a larger effector family. Effectome meeting, Oct 2019, Lauret, France. hal-04456391

## HAL Id: hal-04456391 https://hal.science/hal-04456391v1

Submitted on 13 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Two avirulence genes of *Leptosphaeria maculans*, displaying suppressive interactions, share a common structural pattern and are part of a larger effector family

N. Lazar<sup>1</sup>, Y. Petit-Houdenot<sup>2</sup>, N. Talbi<sup>2</sup>, I. Li de la Sierra-Gallay<sup>1</sup>, E. Zélie<sup>1</sup>, K. Blondeau<sup>1</sup>, A. van de Wouw<sup>3</sup>, B. Ollivier<sup>2</sup>, C. Mesarich<sup>4</sup>, A. Idnurm<sup>3</sup>, H. van Tilbeurgh<sup>1</sup> and <u>I. Fudal<sup>2</sup></u>

<sup>1</sup>Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France

<sup>2</sup>UMR1290 BIOGER, INRA / AgroParisTech/ Université Paris-Saclay, Thiverval-Grignon, France <sup>3</sup>School of BioSciences, University of Melbourne, Melbourne, Australia

<sup>4</sup>School of Fundamental Sciences, Massey University, Palmerston North, New Zealand.

Recognition of a pathogen avirulence (AVR) protein triggers a set of immune responses grouped under the term Effector-Triggered Immunity (ETI). Pathogens can escape ETI by different mechanisms including complete deletion or down-regulation of the AVR genes, point mutations allowing recognition to be evaded while maintaining the virulence function of the AVR protein, and acquisition of new effectors able to suppress recognition. In some cases, the effector that suppresses ETI can itself be recognized by an R gene. A few examples of such mechanism were recently described in fungi but the underlying mechanisms have remained unexplained.

The Dothideomycete *Leptosphaeria maculans*, causal agent of oilseed rape stem canker, is one of the fungal pathogens in which suppression of ETI by the presence of an AVR gene has been demonstrated. The AVR gene *AvrLm4-7* was reported to hide *AvrLm3* from *Rlm3*-mediated recognition and *AvrLm5-9* from *Rlm9*-mediated recognition (Plissonneau et al., 2016; Ghanbarnia et al., 2018). The way AvrLm4-7 suppresses recognition of AvrLm3 and AvrLm5-9 by their cognate R genes is not known: the presence of *AvrLm4-7* does not impede *AvrLm3* and *AvrLm5-9* expression and yeast two-hybrid assays suggested the absence of a physical interaction between AvrLm4-7, AvrLm5-9 and AvrLm3. While AvrLm5-9 and AvrLm3 share a low level of identify, no homology was found with AvrLm4-7.

Blondeau et al. (2015) previously determined the crystal structure of AvrLm4-7. We recently determined of the crystal structure of AvrLm5-9. Surprisingly, despite a lack of sequence similarity, AvrLm5-9 shares structural analogies with AvrLm4-7 (a four stranded anti-parallel sheet and two disulfide bridges in common). This suggests that the antagonistic interaction between AvrLm5-9 and AvrLm4-7 could be due to a competition for a same plant target. Structure-informed and pattern-based searches identified a larger number of structural analogues among *L. maculans* effector candidates (including AvrLm3) and a limited number of structural analogues in other phytopathogenic fungi (including ECP11-1 from *Passalora fulva*, which shares homologies with AvrLm3; Mesarich et al., 2018). We determined the crystal structure of ECP11-1 and confirmed it shared structural analogies with AvrLm4-7 and AvrLm5-9. Transformants of *L. maculans* expressing *ECP11-1* triggered *Rlm3*-mediated recognition, and that recognition was suppressed by AvrLm4-7, suggesting either that ECP11-1 has the same function as AvrLm3, or that its structure is sufficiently close to that of AvrLm3 to be recognized by Rlm3.