



Zeta functions interpolating the convolution of the Bernoulli polynomials

Abdelmejid Bayad, Takao Komatsu

► To cite this version:

Abdelmejid Bayad, Takao Komatsu. Zeta functions interpolating the convolution of the Bernoulli polynomials. *Integral Transforms and Special Functions*, 2018, 29 (8), pp.611-622. 10.1080/10652469.2018.1478822 . hal-04463438

HAL Id: hal-04463438

<https://hal.science/hal-04463438>

Submitted on 4 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Zeta functions interpolating the convolution of the Bernoulli polynomials

Abdelmejid Bayad

Département de Mathématiques
 Université d'Evry Val d'Essonne
 23 Bd. de France, 91037 Evry Cedex, France
 abayad@maths.univ-evry.fr

Takao Komatsu

School of Mathematics and Statistics
 Wuhan University, Wuhan, 430072, China
 komatsu@whu.edu.cn

Abstract

We prove nonlinear relation on multiple Hurwitz-Riemann zeta functions. Using analytic continuation of these multiple Hurwitz-Riemann zeta function, we quote at negative integers Euler's nonlinear relation for generalized Bernoulli polynomials and numbers.

1 Introduction and Preliminaries

Let

$$f(x) = \frac{1}{e^x - 1}$$

and

$$b(x) = xf(x) = \frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!} \quad (|x| < 1),$$

where B_n is the n -th Bernoulli number. Then we have the formulae:

Lemma 1 *For $n \geq 0$*

$$f^{(n)}(x) = (-1)^n \sum_{k=1}^{n+1} (k-1)! \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} f(x)^k \quad (1)$$

2010 *Mathematics subject classification.* 11B68,11M35,11F67,

Key words and phrases. Bernoulli numbers, Bernoulli polynomials, Zeta functions, Special values of zeta functions, Convolution identities.

and

$$f(x)^n = \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k+1 \end{bmatrix} f^{(k)}(x), \quad (2)$$

where $\begin{bmatrix} n \\ k \end{bmatrix}$ denotes the (unsigned) Stirling number of the first kind and $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ denotes the Stirling number of the second kind.

Proof. The formulae (1) and (2) are equivalent via the orthogonal relation satisfied by Stirling numbers

$$\sum_{k=0}^{\max\{n,m\}} (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{smallmatrix} k \\ m \end{smallmatrix} \right\} = (-1)^n \delta_{n,m}. \quad (3)$$

It is easy to see that both formulae are valid for $n = 1$. Let $n \geq 2$. If the formula (1) is true, then

$$\begin{aligned} & \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k+1 \end{bmatrix} f^{(k)}(x) \\ &= \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k+1 \end{bmatrix} (-1)^k \sum_{i=1}^{k+1} (i-1)! \left\{ \begin{smallmatrix} n+1 \\ i \end{smallmatrix} \right\} f(x)^i \\ &= \frac{(-1)^{n-1}}{(n-1)!} \sum_{i=1}^n (i-1)! f(x)^i \sum_{k=i-1}^{n-1} (-1)^k \begin{bmatrix} n \\ k+1 \end{bmatrix} \left\{ \begin{smallmatrix} n+1 \\ i \end{smallmatrix} \right\} \\ &= \frac{(-1)^{n-1}}{(n-1)!} \sum_{i=1}^n (i-1)! f(x)^i (-1)^{n-1} \delta_{n,i} \\ &= f(x)^n. \end{aligned}$$

On the other hand, the relation (1) can be proven by induction on n by using the recurrence formula

$$\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\} = m \left\{ \begin{smallmatrix} n-1 \\ m \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n-1 \\ m-1 \end{smallmatrix} \right\} \quad (n, m \geq 1).$$

with

$$\left\{ \begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right\} = 1, \quad \left\{ \begin{smallmatrix} n \\ 0 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} 0 \\ n \end{smallmatrix} \right\} = 0 \quad (n \geq 1).$$

■

Applying the Leibniz derivative formula to $b(x) = xf(x)$, we obtain

$$b^{(n)}(x) = xf^{(n)}(x) + nf^{(n-1)}(x).$$

Then we have

$$b^{(n)}(x) = (-1)^n \sum_{k=1}^{n+1} (k-1)! \left(\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\} x - n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} \right) f(x)^k. \quad (4)$$

Thus, for $n \geq 0$, we have

Proposition 1

$$b^{(n)}(x) = (-1)^n \sum_{k=1}^{n+1} \frac{(k-1)!}{x^k} \left(\begin{Bmatrix} n+1 \\ k \end{Bmatrix} x - n \begin{Bmatrix} n \\ k \end{Bmatrix} \right) b(x)^k. \quad (5)$$

2 Zeta function

Consider the zeta function defined by Mellin integral

$$Z_m(s, x) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} F^{(m)}(t) e^{-xt} dt, \quad (6)$$

where

$$F(t) = -f(-t) = \frac{1}{1 - e^{-t}},$$

which is defined for $\Re(s) > m + 1$ and $\Re(x) > 0$. It is not difficult, by use of the classical Riemann's methods, to prove its analytic continuation to whole complex plane \mathbb{C} except simple poles at $s = 1, 2, \dots, m + 1$. We omit it.

On the other hand, we have the following results.

Theorem 1 *Let m be a nonnegative integer. Then we have*

$$(-1)^m \sum_{k=1}^{m+1} (k-1)! \begin{Bmatrix} m+1 \\ k \end{Bmatrix} \zeta_k(s, x) = \sum_{k=0}^m (-1)^k \binom{m}{k} x^{m-k} \zeta(s-k, x),$$

where

$$\zeta(s, x) = \sum_{n=0}^\infty \frac{1}{(x+n)^s}$$

denotes the Hurwitz zeta function and

$$\zeta_n(s, x) = \sum_{k_1, \dots, k_n=0}^\infty \frac{1}{(x+k_1+\dots+k_n)^s}$$

denotes the multiple Hurwitz zeta function of order n .

Using the orthogonality property (3), from Theorem 1 we obtain

Corollary 1 *For any positive integer n , we have*

$$\begin{aligned} \zeta_n(s, x) &= \frac{(-1)^n}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-k-1}^{(n)}(x) \zeta(s-k, x) \\ &= \frac{(-1)^n}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \sum_{m=0}^{n-k-1} \binom{m+k}{m} \begin{bmatrix} n \\ m+k+1 \end{bmatrix} x^m \zeta(s-k, x), \end{aligned}$$

where $B_j^{(n)}(x)$ is the j -th Bernoulli polynomial of order n .

Proof of Theorem 1. By integration by parts, we have

$$Z_m(s, x) = \frac{1}{\Gamma(s)} \int_0^\infty (t^{s-1} e^{-xt})^{(m)} F(t) dt,$$

where $F(t) = 1/(1 - e^{-t})$. Using the general Leibniz rule, we obtain

$$Z_m(s, x) = \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} x^{m-k} \zeta(s - k, x). \quad (7)$$

On the other hand, from the relation (1), we have

$$\begin{aligned} Z_m(s, x) &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} F^{(m)}(t) e^{-xt} dt \\ &= \sum_{k=1}^{m+1} (k-1)! \left\{ \begin{matrix} m+1 \\ k \end{matrix} \right\} \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \frac{e^{-xt}}{(1-e^{-t})^k} dt. \end{aligned}$$

Therefore,

$$Z_m(s, x) = \sum_{k=1}^{m+1} (k-1)! \left\{ \begin{matrix} m+1 \\ k \end{matrix} \right\} \zeta_k(s, x). \quad (8)$$

The result follows from (7) and (8). ■

Remark. From the analytic continuations of $\zeta_n(s; x)$ and $\zeta(s; x)$ to the whole complex plane, the generalized m -th Bernoulli polynomial $B_m^{(n)}(x)$ of order n and the m -th Bernoulli polynomial $B_m(x)$ are related to $\zeta_n(-m; x)$ and $\zeta(-m; x)$ by the formulas

$$\zeta_n(-m; x) = (-1)^n \frac{m!}{(m+n)!} B_{m+n}^{(n)}(x), \quad (9)$$

$$\zeta(-m; x) = -\frac{B_{m+1}(x)}{m+1}, \quad (10)$$

for any non-negative integer n . In special, if s is a negative integer with $s = -m$ in Corollary 1, then we get the well-known formula [3, Corollary 1.8]:

$$B_{m+n}^{(n)}(x) = (m+n) \binom{m+n-1}{n-1} \sum_{k=0}^{n-1} (-1)^k \binom{m-1}{k} B_{n-k-1}^{(n)}(x) \frac{B_{m+k+1}(x)}{m+k+1}, \quad (11)$$

which recovers once more Euler's identity and Dilcher's results in [5]; it is also reminiscent of the convolution identities in [10]. For more details, on various generalizations on Euler's identity for Bernoulli numbers, see [1, 2, 4, 5, 7, 8, 9] and references therein.

Next, we study the zeta function associated with Mellin integral given by

$$\widehat{Z}_m(s, x) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} G^{(m)}(t) e^{-xt} dt, \quad (12)$$

where

$$G(t) = b(-t) = \frac{t}{1 - e^{-t}}.$$

We have the following.

Theorem 2

$$\begin{aligned} (-1)^m \sum_{k=1}^{m+1} (k-1)! & \left(s \left\{ \begin{matrix} m+1 \\ k \end{matrix} \right\} \zeta_k(s+1, x) - m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} \zeta_k(s, x) \right) \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} (s-k) \zeta(s-k+1, m). \end{aligned}$$

Corollary 2 If s is a negative integer with $s = -n$, we obtain

$$\begin{aligned} (-1)^m \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} B_{n+k}(x) \\ = \sum_{k=1}^{m+1} (-1)^{k-1} \frac{(k-1)! n!}{(n+k)!} \left(\left\{ \begin{matrix} m+1 \\ k \end{matrix} \right\} (n+k) B_{n+k-1}^{(k)}(x) + m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} B_{n+k}^{(k)}(x) \right). \end{aligned}$$

In addition, if $x = 0$, we obtain

$$B_{n+m} = \sum_{k=1}^{m+1} (-1)^{k-1} \frac{(k-1)! n!}{(n+k)!} \left(\left\{ \begin{matrix} m+1 \\ k \end{matrix} \right\} (n+k) B_{n+k-1}^{(k)} + m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} B_{n+k}^{(k)} \right). \quad (13)$$

3 Zeta function of higher order

Let m_1, \dots, m_d be nonnegative integers. We study the zeta function $Z_{m_1, \dots, m_d}(s, x)$ defined by

$$Z_{m_1, \dots, m_d}(s, x) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} G^{(m_1)}(t) \cdots G^{(m_d)}(t) e^{-xt} dt$$

($\Re(s) > m_1 + \cdots + m_d + d$, $\Re(x) > 0$).

Theorem 3 The function $Z_{m_1, \dots, m_d}(s, x)$ can be analytically continued to the whole \mathbb{C} plane except simple poles at $s = 1, 2, \dots, m_1 + \cdots + m_d + d$, and its values at non-positive integers are given as

$$\begin{aligned} Z_{m_1, \dots, m_d}(-n, x) &= (B_{m_1}(x) + \cdots + B_{m_d}(x))^n \\ &:= \sum_{\substack{k_1 + \cdots + k_d = n \\ k_1, \dots, k_d \geq 0}} \binom{n}{k_1, \dots, k_d} B_{m_1+k_1}(x) \cdots B_{m_d+k_d}(x). \end{aligned} \quad (14)$$

Proof. Since

$$\begin{aligned} G^{(m)}(t)e^{-xt} &= (-1)^m b^{(m)}(-t)e^{-xt} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n B_{m+n}(x)}{n!} t^n, \end{aligned}$$

we have

$$\begin{aligned} G^{(m_1)}(t) \cdots G^{(m_d)}(t) e^{-xt} &= \sum_{n=0}^{\infty} \left(\sum_{k_1+\cdots+k_d=n} (-1)^n \frac{B_{m_1+k_1}(x) \cdots B_{m_d+k_d}(x)}{k_1! \cdots k_d!} \right) t^n \\ &= \sum_{n=0}^{\infty} (-1)^n \left(\sum_{k_1+\cdots+k_d=n} \binom{n}{k_1, \dots, k_d} B_{m_1+k_1}(x) \cdots B_{m_d+k_d}(x) \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} (-1)^n (B_{m_1}(x) + \cdots + B_{m_d}(x))^n \frac{t^n}{n!}. \end{aligned}$$

Now,

$$\begin{aligned} &\int_0^\infty t^{s-1} G^{(m_1)}(t) \cdots G^{(m_d)}(t) e^{-xt} dt \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{(B_{m_1}(x) + \cdots + B_{m_d}(x))^k}{k!} \int_0^1 t^{s+k-1} e^{-xt} dt \\ &\quad + \sum_{k=0}^{\infty} (-1)^k \frac{(B_{m_1}(x) + \cdots + B_{m_d}(x))^k}{k!} \int_1^\infty t^{s+k-1} e^{-xt} dt. \end{aligned}$$

The first integral converges if $\Re(s) > m_1 + \cdots + m_d + d$ and the second integral converges absolutely for any $s \in \mathbb{C}$ and represents an entire function of s . Hence, the right-hand side defines holomorphic function for $\Re(s) > m_1 + \cdots + m_d + d$. Since

$$\int_0^\infty t^{s+k-1} e^{-xt} dt = \frac{\Gamma(s+k)}{x^{s+k}}$$

for $\Re(x) > 0$, we obtain

$$\begin{aligned} Z_{m_1, \dots, m_d}(-n, x) &= \lim_{s \rightarrow -n} \left(\frac{1}{\Gamma(s)} \sum_{k=0}^{\infty} (-1)^k \frac{(B_{m_1}(x) + \cdots + B_{m_d}(x))^k}{k!} \frac{\Gamma(s+k)}{x^{s+k}} \right) \\ &= (B_{m_1}(x) + \cdots + B_{m_d}(x))^n \lim_{s \rightarrow -n} \frac{(-1)^n \Gamma(s+n)}{n! \Gamma(s)} \frac{1}{x^{s+n}} \\ &= (B_{m_1}(x) + \cdots + B_{m_d}(x))^n. \end{aligned}$$

■

Theorem 4 For nonnegative integers k_1, \dots, k_d and m_1, \dots, m_d , we have

$$\begin{aligned} \sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} Z_{m_1+k_1, \dots, m_d+k_d}(s, x) \\ = \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} Z_{k_1, \dots, k_d}(s-k, x). \quad (15) \end{aligned}$$

Proof. By the general Leibniz rule, the left-hand side of (15) is equal to

$$\begin{aligned} & \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \left(\sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} G^{(m_1+k_1)}(t) \cdots G^{(m_d+k_d)}(t) \right) e^{-xt} dt \\ &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} (G^{(k_1)}(t) \cdots G^{(k_d)}(t))^{(m)} e^{-xt} dt. \end{aligned}$$

By integration by parts,

$$\int_0^\infty (t^{s-1} e^{-xt})^{(m)} G^{(k_1)}(t) \cdots G^{(k_d)}(t) dt = (-1)^m \int_0^\infty t^{s-1} e^{-xt} (G^{(k_1)}(t) \cdots G^{(k_d)}(t))^{(m)} dt.$$

On the other hand, we have

$$\begin{aligned} & \int_0^\infty (t^{s-1} e^{-xt})^{(m)} G^{(k_1)}(t) \cdots G^{(k_d)}(t) dt \\ &= \sum_{k=0}^m \binom{m}{k} (-x)^{m-k} (s-1)_k \int_0^\infty t^{s-k-1} G^{(k_1)}(t) \cdots G^{(k_d)}(t) e^{-xt} dt. \end{aligned}$$

Hence, we get

$$\begin{aligned} & \sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} Z_{m_1+k_1, \dots, m_d+k_d}(s, x) \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} \frac{(s-1)_k}{\Gamma(s)} \int_0^\infty t^{s-k-1} e^{-xt} dt \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} Z_{k_1, \dots, k_d}(s-k, x), \end{aligned}$$

which is (15). ■

Corollary 3 We have

$$\sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} Z_{m_1, \dots, m_d}(s, x) = \sum_{k=0}^m \binom{m}{k} (-1)^k x^{m-k} \zeta_d(s-k, x). \quad (16)$$

Then by putting $s = -n$, we obtain the following.

Corollary 4

$$\begin{aligned} \sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} (B_{m_1}(x) + \dots + B_{m_d}(x))^n \\ = \sum_{k=0}^m \binom{m}{k} (-1)^{k+d} x^{m-k} \frac{(n+k)!}{(n+k+d)!} B_{n+m+d}^{(d)}(x). \end{aligned}$$

In special, for $x = 0$, we have

$$\sum_{m_1+\dots+m_d=m} \binom{m}{m_1, \dots, m_d} (B_{m_1} + \dots + B_{m_d})^n = \frac{(-1)^{m+d} (n+m)! B_{n+m+d}^{(d)}}{(n+m+d)!}.$$

4 Another method and further formulas

In this section, we study the zeta function

$$Z_{m_1, \dots, m_d}(s, x) := \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} G^{(m_1)}(t) \cdots G^{(m_d)}(t) e^{-xt} dt$$

by another method. First, we consider the function $Z_{m_1, m_2}(s, x)$ when $d = 2$.

Lemma 2

$$Z_{m_1, m_2}(s, x) = (-1)^{m_2} \sum_{0 \leq k_1 \leq k_2 \leq m_2} \binom{m_2}{m_2 - k_2, k, k_2 - k} (-x)^{k_2 - k} Z_{m_1 + m_2 - k_2, 0}(s - k, x). \quad (17)$$

Proof. By integration by parts and the general Leibniz rule, we have

$$\begin{aligned} Z_{m_1, m_2}(s, x) &= \frac{(-1)^m}{\Gamma(s)} \int_0^\infty (t^{s-1} e^{-xt} G^{(m_1)})^{(m_2)} G(t) dt \\ &= (-1)^m \sum_{k_2=0}^{m_2} \binom{m_2}{k_2} \frac{1}{\Gamma(s)} \int_0^\infty (t^{s-1} e^{-xt})^{(k_2)} G^{(m_1+m_2-k_2)}(t) G(t) dt. \end{aligned}$$

Hence,

$$\begin{aligned} Z_{m_1, m_2}(s, x) &= (-1)^{m_2} \sum_{0 \leq k \leq k_2 \leq m_2} \binom{m_2}{k_2} \binom{k_2}{k} (-x)^{k_2 - k} \frac{1}{\Gamma(s - k)} \int_0^\infty t^{s-k-1} e^{-xt} G^{(m_1+m_2-k_2)}(t) G(t) dt \\ &= (-1)^{m_2} \sum_{0 \leq k_1 \leq k_2 \leq m_2} \binom{m_2}{m_2 - k_2, k, k_2 - k} (-x)^{k_2 - k} Z_{m_1 + m_2 - k_2, 0}(s - k, x). \end{aligned}$$

■

By Proposition 1 and Lemma 2, we obtain the following theorem.

Theorem 5 *We have*

$$\begin{aligned} Z_{m_1, m_2}(s, x) &= \sum_{\substack{0 \leq k \leq k_2 \leq m_2 \\ 0 \leq j \leq m_1 + m_2 - k_2}} (-1)^{j+m_2-1} (j-1)! \binom{m_2}{m_2 - k_2, k_2 - k, k} (-x)^{k_2-k} \\ &\quad \times \left(\binom{m_1 + m_2 - k_2 + 1, j}{j} (s-k)(s-k+1)\zeta_{j+1}(s-k+2, x) \right. \\ &\quad \left. + (m_1 + m_2 - k_2) \binom{m_1 + m_2 - k_2}{j} (s-k)\zeta_{j+1}(s-k+1, x) \right). \end{aligned}$$

Proof. By Proposition 1, we have

$$\begin{aligned} G^{(m_1+m_2-k_2)}(t) &= (-1)^{m_1+m_2-k_2} b^{(m_1+m_2-k_2)}(-t) \\ &= \sum_{j=1}^{m_1+m_2-k_2+1} \frac{(j-1)!(-1)^{j-1}}{t^j} \left(\binom{m_1 + m_2 - k_2 + 1}{j} t \right. \\ &\quad \left. + (m_1 + m_2 - k_2) \binom{m_1 + m_2 - k_2}{j} B(-t)^j \right) \\ &= \sum_{j=1}^{m_1+m_2-k_2+1} (j-1)!(-1)^{j-1} \left(\binom{m_1 + m_2 - k_2 + 1}{j} t \right. \\ &\quad \left. + (m_1 + m_2 - k_2) \binom{m_1 + m_2 - k_2}{j} F(t)^j \right). \end{aligned}$$

Note that

$$G(t) = \frac{t}{1 - e^{-t}} = tF(t).$$

Therefore,

$$\begin{aligned} Z_{m_1+m_2-k_2, 0}(s-k, x) &= \frac{1}{\Gamma(s-k)} \int_0^\infty t^{s-k-1} e^{-xt} G^{(m_1+m_2-k_2)}(t) G(t) dt \\ &= \sum_{j=1}^{m_1+m_2-k_2+1} (j-1)!(-1)^{j-1} \left(\binom{m_1 + m_2 - k_2 + 1}{j} \frac{1}{\Gamma(s-k)} \int_0^\infty t^{s-k+1} e^{-xt} F(t)^{j+1} dt \right. \\ &\quad \left. + (m_1 + m_2 - k_2) \binom{m_1 + m_2 - k_2}{j} \frac{1}{\Gamma(s-k)} \int_0^\infty t^{s-k} e^{-xt} F(t)^{j+1} dt \right) \\ &= \sum_{j=1}^{m_1+m_2-k_2+1} (j-1)!(-1)^{j-1} \left(\binom{m_1 + m_2 - k_2 + 1}{j} (s-k+1)(s-k)\zeta_{j+1}(s-k+2, x) \right. \\ &\quad \left. + (m_1 + m_2 - k_2) \binom{m_1 + m_2 - k_2}{j} (s-k)\zeta_{j+1}(s-k+1, x) \right). \end{aligned}$$

Together with Lemma 2, we get the desired result. ■

Putting $s = -n$ in Theorem 5, we obtain the following.

Corollary 5 *We have*

$$\begin{aligned}
 & (B_{m_1} + B_{m_2})^n \\
 &= \sum_{\substack{0 \leq k \leq k_2 \leq m_2 \\ 0 \leq j \leq m_1 + m_2 - k_2}} (-1)^{m_2} \binom{m_2}{k_2} (j-1)! \left(\left\{ \begin{matrix} m_1 + m_2 - k_2 + 1, j \\ j \end{matrix} \right\} \frac{(n+k_2)! B_{n+k_2+j-1}^{(j+1)}}{(n+k_2+j-1)!} \right. \\
 &\quad \left. - (m_1 + m_2 - k_2) \left\{ \begin{matrix} m_1 + m_2 - k_2 \\ j \end{matrix} \right\} \frac{(n+k_2)! B_{n+k_2+j}^{(j+1)}}{(n+k_2+j)!} \right).
 \end{aligned}$$

We find a very intriguing formula in the above corollary. Similarly, we can do the same for any $d \geq 3$ and obtain a generalization of the previous Theorem. However, the formulas obtained for $(B_{m_1} + \dots + B_{m_d})^n$ are not as pretty as in the case $d = 2$. For this reason, we do not wish to state them here.

References

- [1] T. Agoh and K. Dilcher, *Convolution identities and lacunary recurrences for Bernoulli numbers*, J. Number Theory **124** (2007), 105–122.
- [2] T. Agoh and K. Dilcher, *Higher-order recurrences for Bernoulli numbers*, J. Number Theory **129** (2009), 1837–1847.
- [3] A. Bayad and M. Beck, *Relations for Bernoulli Barnes numbers and Barnes zeta functions*, Int. Journal of Number Theory **10** (2014), 1321–1335.
- [4] K.-W. Chen, *Sums of products of generalized Bernoulli polynomials*, Pacific J. Math., **208** (2003), 39–52.
- [5] K. Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996), 23–41.
- [6] M. Eie, *A note on Bernoulli numbers and Shintani generalized Bernoulli polynomials*, Trans. Am. Math. Soc. **348** (1996), 1117–1136.
- [7] K. Kamano, *Sums of products of hypergeometric Bernoulli numbers*, J. Number Theory **130** (2010), 2259–2271.
- [8] M.-S. Kim, *A note on sums of products of Bernoulli numbers*, Appl. Math. Lett. **24** (1) (2011), 55–61.
- [9] A. Petojević, *New sums of products of Bernoulli numbers*, Integral Transforms Spec. Funct. **19** (2008), 105–114.
- [10] Z.-W. Sun and H. Pan, *Identities concerning Bernoulli and Euler polynomials*, Acta Arith. **125** (2006), no. 1, 21–39.