



HAL
open science

Relations for Bernoulli–Barnes numbers and Barnes zeta functions

Abdelmejid Bayad, Matthias Beck

► **To cite this version:**

Abdelmejid Bayad, Matthias Beck. Relations for Bernoulli–Barnes numbers and Barnes zeta functions. International Journal of Number Theory, 2014, 10 (05), pp.1321-1335. 10.1142/S1793042114500298 . hal-04464006

HAL Id: hal-04464006

<https://hal.science/hal-04464006>

Submitted on 4 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RELATIONS FOR BERNOULLI–BARNES NUMBERS AND BARNES ZETA FUNCTIONS

ABDELMEJID BAYAD AND MATTHIAS BECK

Dedicated to the fond memory of Marvin I. Knopp (1933–2011)

ABSTRACT. The *Barnes ζ -function* is

$$\zeta_n(z, x; \mathbf{a}) := \sum_{\mathbf{m} \in \mathbb{Z}_{\geq 0}^n} \frac{1}{(x + m_1 a_1 + \cdots + m_n a_n)^z}$$

defined for $\operatorname{Re}(x) > 0$ and $\operatorname{Re}(z) > n$ and continued meromorphically to \mathbb{C} . Specialized at negative integers $-k$, the Barnes ζ -function gives

$$\zeta_n(-k, x; \mathbf{a}) = \frac{(-1)^n k!}{(k+n)!} B_{k+n}(x; \mathbf{a})$$

where $B_k(x; \mathbf{a})$ is a *Bernoulli–Barnes polynomial*, which can be also defined through a generating function that has a slightly more general form than that for Bernoulli polynomials. Specializing $B_k(0; \mathbf{a})$ gives the *Bernoulli–Barnes numbers*. We exhibit relations among Barnes ζ -functions, Bernoulli–Barnes numbers and polynomials, which generalize various identities of Agoh, Apostol, Dilcher, and Euler.

1. INTRODUCTION

We define, as usual, the *Bernoulli numbers* B_k through the generating function

$$(1) \quad \frac{z}{e^z - 1} = \sum_{k \geq 0} B_k \frac{z^k}{k!}.$$

A fundamental relation of Bernoulli numbers, known at least since Euler’s time, is (for $n \geq 1$)

$$(2) \quad \sum_{j=0}^n \binom{n}{j} B_j B_{n-j} = -n B_{n-1} - (n-1) B_n.$$

Much more recently, multinomial generalizations of (2) were discovered by Agoh and Dilcher [1, 8]. They can be viewed as relations between Bernoulli numbers and *Bernoulli numbers $B_k^{(n)}$ of order n* , defined through

$$(3) \quad \left(\frac{z}{e^z - 1} \right)^n = \sum_{k \geq 0} B_k^{(n)} \frac{z^k}{k!}.$$

Date: 6 September 2013.

2000 Mathematics Subject Classification. Primary 11B68; secondary 11F20, 11M32.

Key words and phrases. Bernoulli–Barnes number, Bernoulli polynomial, Barnes zeta function, Fourier–Dedekind sum.

We thank a referee for helpful comments on a previous version of this paper, in particular, for pointing out Sun’s work on self-dual sequences [22], which gave rise to Corollary 17. This research project was initiated while M. Beck visited the Université d’Evry Val d’Essonne; he thanks their Laboratoire analyse et probabilités for their hospitality. A. Bayad was partially supported by the FDIR of the Université d’Evry Val d’Essonne; M. Beck was partially supported by the US National Science Foundation (DMS-1162638).

Dilcher and others proved generalized formulas relating $B_k^{(n)}$ with B_j . The first few are [8, p. 32]:

(4)

$$B_k^{(2)} = -kB_{k-1} - (k-1)B_k \quad (\text{for any } k \geq 1),$$

$$B_k^{(3)} = k(k-1)B_{k-2} + \frac{3}{2}k(k-2)B_{k-1} + \frac{1}{2}(k-1)(k-2)B_k \quad (\text{for any } k \geq 2),$$

$$B_k^{(4)} = -k(k-1)(k-2)B_{k-3} + \frac{11}{6}k(k-1)(k-3)B_{k-2} - k(k-2)(k-3)B_{k-1} - \frac{1}{6}(k-1)(k-2)(k-3)B_k$$

(for any $k \geq 3$).

Our first goal is to derive relations among *Bernoulli–Barnes numbers* $B_k(\mathbf{a})$, defined for a fixed vector $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \mathbb{R}_{>0}^n$ through

$$(5) \quad \frac{z^n}{(e^{a_1 z} - 1) \cdots (e^{a_n z} - 1)} = \sum_{k \geq 0} B_k(\mathbf{a}) \frac{z^k}{k!}.$$

Note that, with (3) and (5), the Bernoulli–Barnes numbers and Bernoulli numbers are related as

$$B_k(\mathbf{a}) = \sum_{m_1 + \cdots + m_n = k} \binom{k}{m_1, \dots, m_n} a_1^{m_1-1} \cdots a_n^{m_n-1} B_{m_1} \cdots B_{m_n}.$$

Of course, one retrieves the Bernoulli numbers of order n with the special case $a_1 = a_2 = \cdots = a_n = 1$, and the Bernoulli numbers by further specializing $n = 1$. Our first main result is as follows.

Theorem 1. For $n \geq 3$, $m \geq 1$, where m is odd, and $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \mathbb{R}_{>0}^n$,

$$\sum_{j=n-m}^n \binom{n+j-4}{j-2} \frac{1}{(m-n+j)!} \sum_{|I|=j} B_{m-n+j}(\mathbf{a}_I) = \begin{cases} \frac{1}{2} & \text{if } n = m = 3, \\ 0 & \text{otherwise,} \end{cases}$$

where the inner sum is over all subsets $I \subseteq \{1, 2, \dots, n\}$ of cardinality j , and $\mathbf{a}_I := (a_i : i \in I)$.

Here and in what follows below, all binomial coefficients with a negative bottom entry are zero.

Corollary 2. For $n \geq 3$ and odd $m \geq n - 2$,

$$\sum_{j=2}^n \binom{n+j-4}{j-2} \frac{m!}{(m-n+j)!} \binom{n}{j} B_{m-n+j}^{(j)} = \begin{cases} 3 & \text{if } n = m = 3, \\ 0 & \text{otherwise,} \end{cases}$$

For example, for $n = 3, 4$ and odd $m \geq 1$, Theorem 1 gives the relations

$$B_m^{(3)} = \frac{3}{2} \delta_{3,m} - \frac{3}{2} m B_{m-1}^{(2)} \quad (m \geq 3),$$

$$B_m^{(4)} = -2m B_{m-1}^{(3)} - m(m-1) B_{m-2}^{(2)} \quad (m \geq 4).$$

More generally, for any positive integer $n \geq 4$, Corollary 2 gives the following recurrence formula for the numbers $B_m^{(n)}$:

$$(6) \quad \binom{2n-4}{n-2} B_m^{(n)} = - \sum_{j=2}^{n-1} \binom{n+j-4}{j-2} \frac{m!}{(m-n+j)!} \binom{n}{j} B_{m-n+j}^{(j)} \quad \text{for any odd } m \geq n - 2.$$

The novelty of these relations as, e.g., compared with (4) is that they are between Bernoulli numbers of order higher than 1. We suspect that there are relations analogous to (6) for *even* m but leave the search for them as an open problem.

One of the significances of Bernoulli numbers lies in the fact that they are essentially evaluations of the Riemann ζ -function $\zeta(z) := \sum_{m \geq 1} m^{-z}$ (meromorphically continued to \mathbb{C}) at negative integers $-k$:

$$\zeta(-k) = -\frac{B_{k+1}}{k+1}.$$

Bernoulli–Barnes numbers appear in a similar fashion in relation with the Barnes ζ -function [4]

$$\zeta_n(z, x; \mathbf{a}) := \sum_{\mathbf{m} \in \mathbb{Z}_{\geq 0}^n} \frac{1}{(x + m_1 a_1 + \cdots + m_n a_n)^z}$$

defined for $\operatorname{Re}(x) > 0$ and $\operatorname{Re}(z) > n$ and continued meromorphically to \mathbb{C} [9, 10, 11, 16, 19, 21]. Specialized at negative integers $-k$, the Barnes ζ -function gives

$$(7) \quad \zeta_n(-k, x; \mathbf{a}) = \frac{(-1)^n k!}{(m+n)!} B_{k+n}(x; \mathbf{a})$$

where $B_k(x; \mathbf{a})$ is a *Bernoulli–Barnes polynomial*, defined through [4]

$$(8) \quad \frac{z^n e^{xz}}{(e^{a_1 z} - 1) \cdots (e^{a_n z} - 1)} = \sum_{k \geq 0} B_k(x; \mathbf{a}) \frac{z^k}{k!}.$$

Thus the Bernoulli–Barnes numbers are the special evaluations $B_k(\mathbf{a}) = B_k(0; \mathbf{a})$. It is clear that the Barnes zeta function is a multidimensional generalization of various Riemann–Hurwitz zetas functions; e.g., when $n = 1$ and $\mathbf{a} = (a)$, the function $\zeta(s; x, \mathbf{a})$ is the classical *Hurwitz zeta function* $a^{-s} \zeta(s; \frac{x}{a})$. Likewise, the Bernoulli–Barnes numbers and Bernoulli–Barnes polynomials extend the (generalized) Bernoulli numbers and polynomials to higher dimensions. Further generalizations of Bernoulli numbers and polynomials include [13, 20].

Our second main result expresses the Barnes zeta function in terms of Bernoulli–Barnes polynomials, Hurwitz zeta functions, and *Fourier–Dedekind sums* [5], defined as

$$\sigma_r(a_1, \dots, \hat{a}_j, \dots, a_d; a_j) := \frac{1}{a_j} \sum_{\lambda^{a_j} = 1 \neq \lambda} \frac{\lambda^r}{\prod_{1 \leq k \neq j \leq n} (1 - \lambda^{a_k})}.$$

Fourier–Dedekind sums generalize and unify many variants of (generalized) Dedekind sums; see, e.g., [18] or [6, Chapter 8].

Theorem 3. *Let a_1, \dots, a_n be pairwise coprime positive integers. Then*

$$\begin{aligned} \zeta(s; x, \mathbf{a}) &= \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) \zeta(s-k; x) \\ &\quad + \sum_{j=1}^n a_j^{-s} \sum_{r=0}^{a_j-1} \sigma_{-r}(a_1, \dots, \hat{a}_j, \dots, a_n; a_j) \zeta\left(s; \frac{x+r}{a_j}\right). \end{aligned}$$

Theorem 3 has several applications. Specializing s at negative integers gives, with the help of (7):

Corollary 4. *Let a_1, \dots, a_n be pairwise coprime positive integers. Then*

$$\begin{aligned} \sum_{j=1}^n a_j^m \sum_{r=0}^{a_j-1} \sigma_{-r}(a_1, \dots, \hat{a}_j, \dots, a_n; a_j) B_{m+1}\left(\frac{x+r}{a_j}\right) = \\ (-1)^{n-1} \frac{(m+1)!}{(m+n)!} B_{m+n}(x, \mathbf{a}) + \frac{(-1)^{n-1} (m+1)!}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) \frac{B_{m+k+1}(x)}{m+k+1}. \end{aligned}$$

This is reminiscent of a *reciprocity law* for generalized Dedekind sums, due to Apostol [2]; this can be illustrated more easily in the case $n = 2$ and $\mathbf{a} = (a, b)$, for which Theorem 3 specializes to:

Corollary 5. *Let a, b be coprime positive integers. Then*

$$\begin{aligned} \zeta(s; x, (a, b)) &= \frac{1}{ab} \zeta(s-1; x) + \left(1 - \frac{x}{ab}\right) \zeta(s; x) \\ &\quad - a^{-s} \sum_{r=0}^{a-1} \left\{ \frac{b^{-1}r}{a} \right\} \zeta\left(s; \frac{x+r}{a}\right) - b^{-s} \sum_{r=0}^{a-1} \left\{ \frac{a^{-1}r}{b} \right\} \zeta\left(s; \frac{x+r}{b}\right). \end{aligned}$$

Again the specialization of s at negative integers gives, for $n = 2$ and $\mathbf{a} = (a, b)$, using (7):

Corollary 6. *Let a, b be coprime positive integers. Then*

$$\begin{aligned} a^m \sum_{r=0}^{a-1} \left\{ \frac{b^{-1}r}{a} \right\} B_{m+1}\left(\frac{x+r}{a}\right) + b^m \sum_{r=0}^{a-1} \left\{ \frac{a^{-1}r}{b} \right\} B_{m+1}\left(\frac{x+r}{b}\right) = \\ \frac{1}{m+2} B_{m+2}(x, (a, b)) + \frac{1}{ab} \frac{m+1}{m+2} B_{m+2}(x) + \left(\frac{x}{ab} - 1\right) B_{m+1}(x). \end{aligned}$$

This is a ‘‘polynomial generalization’’ of Apostol’s reciprocity law [2]

$$\frac{1}{m} (a^{m-1} s_m(a, b) + b^{m-1} s_m(b, a)) = \frac{B_{m+1}}{(m+1)ab} + \frac{1}{m(m+1)ab} (aB - bB)^{m+1}.$$

Here m is a positive integer, a and b are coprime, we use the umbral notation

$$(aB - bB)^{m+1} = \sum_{i=0}^{m+1} \binom{m+1}{i} (-1)^{m+1-i} a^i b^{m+1-i} B_i B_{m+1-i},$$

and

$$(9) \quad S_m(a, b) = \sum_{r=0}^{a-1} \left\{ \frac{a^{-1}r}{b} \right\} B_m\left(\frac{r}{b}\right)$$

are the *Apostol–Dedekind sums*. The classical Dedekind sums [7, 18] are captured by the special case $m = 1$. Thus in some sense, our study can be viewed as a bridge between Euler-type identities and Dedekind-type reciprocity laws.

Finally, we discuss the special case $\mathbf{a} = (1, \dots, 1)$ of Theorem 3. Denote

$$\zeta_n(s; x) = \zeta(s; x, (1, \dots, 1)),$$

the *Hurwitz zeta function of order n* . Since in this case the sums $\sigma_r(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j)$ vanish, we obtain the following identity.

Corollary 7. *For any positive integer n ,*

$$\zeta_n(s; x) = \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}^{(n)}(x) \zeta(s-k; x).$$

Specializing once more $s = -m$ at a negative integer gives:

Corollary 8. *For any positive integers n, m ,*

$$B_{m+n}^{(n)}(x) = (m+n) \binom{m+n-1}{n-1} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}^{(n)}(x) \frac{B_{m+k+1}(x)}{m+k+1}.$$

Corollary 8 recovers once more Euler's identity (2) and Dilcher's results in [8]; it is also reminiscent of the convolution identities in [23]. We also note that in the above corollaries the coefficients of the polynomials $B_{n-1-k}^{(n)}(x)$ can be explicitly given in terms of Stirling numbers of the first kind $s(n, k)$ as follows:

$$(10) \quad \binom{n-1}{n-1-k} B_{n-1-k}^{(n)}(x) = \sum_{m=0}^{n-1-k} \binom{m+k}{m} s(n, m+k+1) x^m$$

(see, e.g., [12, Equation (52.2.21)]).

2. PROOF OF THEOREM 1

Our proof is based on identities of generating functions. Fix $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \mathbb{R}_{>0}^n$. We define for a subset $I \subseteq \{1, 2, \dots, n\}$

$$f_I(z) := \frac{z^{|I|} e^{z \sum_{i \in I} a_i}}{\prod_{i \in I} (e^{a_i z} - 1)} \quad \text{and} \quad F^{(j)}(z) := \sum_{|I|=j} f_I(z).$$

Proposition 9. For $n \geq 4$,

$$\sum_{j=2}^n \binom{n+j-4}{j-2} (-z)^{n-j} F^{(j)}(z)$$

is an even function in z .

Proof. We need to prove that

$$F(z) := \sum_{j=2}^n \binom{n+j-4}{j-2} (-z)^{n-j} F^{(j)}(z) = \sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} (-z)^{n-|I|} f_I(z)$$

is even, so that it suffices to prove that $F(z)$ equals

$$F(-z) = \sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} z^{n-|I|} e^{-z \sum_{i \in I} a_i} f_I(z),$$

i.e., that

$$F(z) - F(-z) = \sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} z^{n-|I|} f_I(z) \left((-1)^{n-|I|} - e^{-z \sum_{i \in I} a_i} \right)$$

is zero. Written with the denominator $\prod_{i=1}^n (e^{a_i z} - 1)$, the function $F(z) - F(-z)$ has the numerator

$$\sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} z^n e^{z \sum_{i \in I} a_i} \left((-1)^{n-|I|} - e^{-z \sum_{i \in I} a_i} \right) \prod_{i \notin I} (e^{a_i z} - 1)$$

and so we can rephrase our goal to proving that

$$\sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} \left((-1)^{n-|I|} e^{z \sum_{i \in I} a_i} - 1 \right) \prod_{i \notin I} (e^{a_i z} - 1)$$

is zero. With $\prod_{i \notin I} (e^{a_i z} - 1) = \sum_{J \subseteq \bar{I}} (-1)^{n-|I|-|J|} e^{z \sum_{i \in J} a_i}$, we can further rephrase our goal to proving that

$$(11) \quad \sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} \sum_{J \subseteq \bar{I}} (-1)^{|J|} e^{z \sum_{i \in I \cup J} a_i} = \sum_{|I| \geq 2} \binom{n+|I|-4}{|I|-2} \sum_{J \subseteq \bar{I}} (-1)^{n-|I|-|J|} e^{z \sum_{i \in J} a_i}.$$

We will show that the coefficients of $e^{z\sum_{i \in K} a_i}$, for any $K \subseteq \{1, 2, \dots, n\}$, on both sides of (11) are equal. This coefficient is on the left-hand side of (11) equal to

$$\sum_{j=0}^{|K|-2} (-1)^j \binom{|K|}{j} \binom{n+|K|-4-j}{|K|-2-j}.$$

The corresponding coefficient on the right-hand side of (11) is

$$\sum_{I \not\supseteq K} (-1)^{n-|I|-|K|} \binom{n+|I|-4}{|I|-2} = \sum_{j=2}^{n-|K|} (-1)^{n-j-|K|} \binom{n+j-4}{j-2} \binom{n-|K|}{j}.$$

Thus (11) is equivalent to

$$\sum_{j=0}^{k-2} (-1)^j \binom{k}{j} \binom{n+k-4-j}{k-2-j} = \sum_{j=2}^{n-k} (-1)^{n-j-k} \binom{n+j-4}{j-2} \binom{n-k}{j}.$$

But both sides equal $\binom{n-4}{k-2}$, as one can prove, e.g., by putting either side into a generating function for k . \square

Proof of Theorem 1. Recalling that $\mathbf{a}_I = (a_i : i \in I)$, we see that

$$f_I(z) = \frac{z^{|I|} e^{z\sum_{i \in I} a_i}}{\prod_{i \in I} (e^{a_i z} - 1)} = \frac{(-z)^{|I|}}{\prod_{i \in I} (e^{-a_i z} - 1)}$$

is the exponential generating function for $(-1)^k B_k(\mathbf{a}_I)$. Thus, using Proposition 9 and the notation $\text{Odd}(\dots)$ for the odd part of a function, we compute

$$\begin{aligned} 0 &= \text{Odd} \left(\sum_{j=0}^n \binom{n+j-4}{j-2} (-z)^{n-j} F^{(j)}(z) \right) \\ &= \text{Odd} \left(\sum_{j=0}^n \binom{n+j-4}{j-2} (-z)^{n-j} \sum_{|I|=j} \sum_{k \geq 0} (-1)^k B_k(\mathbf{a}_I) \frac{z^k}{k!} \right) \\ &= - \sum_{j=0}^n \binom{n+j-4}{j-2} \sum_{|I|=j} \sum_{\substack{k \geq 0 \\ n-j+k \text{ odd}}} \frac{1}{k!} B_k(\mathbf{a}_I) z^{n-j+k} \\ &= - \sum_{\substack{m \geq 0 \\ m \text{ odd}}} \sum_{j=n-m}^n \binom{n+j-4}{j-2} \frac{1}{(m-n+j)!} \sum_{|I|=j} B_{m-n+j}(\mathbf{a}_I) z^m. \end{aligned}$$

Now read off the coefficients.

In the case $n = 3$, a quick calculation reveals

$$\text{Odd} \left(\sum_{j=0}^3 \binom{j-1}{j-2} (-z)^{3-j} F^{(j)}(z) \right) = -\frac{1}{2} z^3$$

and this explains the special case. \square

We should remark that Theorem 1 was in part motivated by [15] in which Katayama proposed a three-term generalization of the reciprocity theorem for Dedekind–Apostol sums [2]; Apostol’s theorem was a byproduct of another paper of Katayama [14]. Unfortunately, the main theorem of [15] is wrong; to make the central integral of the paper work, one has to use the integrand $\frac{z^{s-1}}{(e^{z/a}-1)(e^{z/b}-1)(e^{z/c}-1)}$ which, unfortunately, does not give rise to Dedekind–Apostol sums. However, using this integrand we discovered Theorem 1.

3. PROOF OF THEOREM 3

The function

$$p_A(t) := \#\{(k_1, \dots, k_n) \in \mathbb{Z}_{\geq 0}^n : k_1 a_1 + \dots + k_n a_n = t\},$$

which counts all partitions of t with parts in the finite set $A := \{a_1, \dots, a_n\}$, is called a *restricted partition function*. For example, basic combinatorics gives

$$p_{\{1, \dots, 1\}}(t) = \binom{n-1+t}{n-1},$$

and a slightly less trivial example was proved by Barlow [3, p. 323–325]: for a and b coprime,

$$p_{\{a, b\}}(t) = \frac{t}{ab} + 1 - \left\{ \frac{b^{-1}t}{a} \right\} - \left\{ \frac{a^{-1}t}{b} \right\},$$

where $\{x\} = x - \lfloor x \rfloor$ denotes the fractional part of x , a^{-1} is computed mod b , and b^{-1} mod a . (The above formulation of Barlow's formula seems to be due to Popoviciu [17]; see also [6, Chapter 1].)

The following theorem was proved in [5]; however, the authors of that paper did not realize the explicit role of Bernoulli–Barnes polynomials.

Theorem 10. *If a_1, \dots, a_n are pairwise coprime positive integers, then*

$$p_A(t) = \frac{(-1)^{n-1}}{(n-1)!} B_{n-1}(-t; (a_1, \dots, a_n)) + \sum_{j=0}^n \sigma_{-t}(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j).$$

Proof. We give an outline of the proof. As in [5], we compute the residues of

$$F_t(z) = \frac{1}{z^{t+1} \prod_{i=1}^n (1 - z^{a_i})}.$$

The residue at $z = 0$ gives $p_A(t)$, whereas the residue at $z = 1$ gives $\frac{(-1)^n}{(n-1)!} B_{n-1}(-t; (a_1, \dots, a_n))$. Finally, if λ is a nontrivial a_i th root of unity,

$$\text{Res}(F_t(z); z = \lambda) = -\frac{1}{a_i \lambda^t \prod_{j \neq i} (1 - \lambda^{a_j})},$$

where the product runs over all $j = 1, \dots, n$ except $j = i$. Thus

$$\sum_{\lambda^{a_i} = 1, \lambda \neq 1} \text{Res}(F_t(z); z = \lambda) = -\sigma_{-t}(a_1, \dots, \widehat{a}_i, \dots, a_n; a_i).$$

The residue theorem completes the proof of Theorem 10. □

Proof of Theorem 3. Set $t = m_1 a_1 + \dots + m_n a_n$. We can rewrite the Barnes zeta function as follows:

$$\zeta(s; x, \mathbf{a}) = \sum_{m_1, \dots, m_n \geq 0} \frac{1}{(x + m_1 a_1 + \dots + m_n a_n)^s} = \sum_{t \geq 0} \frac{p_A(t)}{(x+t)^s}.$$

By applying Taylor's theorem to the function $t \mapsto B_{n-1}(-t; \mathbf{a})$ at $t = -x$,

$$B_{n-1}(-t; \mathbf{a}) = \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) (x+t)^k,$$

and so with Theorem 10 we obtain

$$p_A(t) = \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) (x+t)^k + \sum_{j=0}^n \sigma_{-t}(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j).$$

Hence

$$\zeta(s; x, \mathbf{a}) = \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) \zeta(s-k; x) + \sum_{j=1}^n \sum_{t \geq 0} \frac{\sigma_{-t}(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j)}{(x+t)^s}.$$

Note that $\sigma_{-t}(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j)$ depends only $t \bmod a_j$. Setting $t = ma_j + r$, $0 \leq r \leq a_j - 1$, we obtain

$$\begin{aligned} \zeta(s; x, \mathbf{a}) &= \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) \zeta(s-k; x) \\ &\quad + \sum_{j=1}^n \sum_{r=0}^{a_j-1} \sigma_{-r}(a_1, \dots, \widehat{a}_j, \dots, a_n; a_j) \sum_{m \geq 0} \frac{1}{(x+r+ma_j)^s}. \end{aligned}$$

Writing $(x+r+ma_j)^{-s} = a_j^{-s} \left(\frac{x+r}{a_j} + m \right)^{-s}$ completes the proof of Theorem 3. \square

4. THE SPECIAL CASE $\mathbf{a} = (a, 1, 1, \dots, 1)$

In the special case $A = \{a, 1, 1, \dots, 1\}$ (with n 1's), most of the terms in Theorem 10 disappear and we obtain

$$(12) \quad P_{\{a, 1, 1, \dots, 1\}}(t) = \frac{(-1)^n}{n!} B_n(-t; (a, 1, 1, \dots, 1)) + \sigma_{-t}(1, 1, \dots, 1; a).$$

On the other hand, we can apply [6, Theorem 8.8] to this special case; thus for $t = 1, 2, \dots, a+n-1$,

$$\sigma_t(1, 1, \dots, 1; a) = \frac{(-1)^{n-1}}{n!} B_n(t; (a, 1, 1, \dots, 1)).$$

Since $\sigma_t(1, 1, \dots, 1; a)$ only depends on $t \bmod a$, this range for t is enough to determine $\sigma_t(1, 1, \dots, 1; a)$:

$$(13) \quad \sigma_t(1, 1, \dots, 1; a) = \begin{cases} \frac{(-1)^{n-1}}{n!} B_n(t \bmod a; (a, 1, 1, \dots, 1)) & \text{if } t \not\equiv 0 \pmod{a}, \\ \frac{(-1)^{n-1}}{n!} B_n(a; (a, 1, 1, \dots, 1)) & \text{if } t \equiv 0 \pmod{a}. \end{cases}$$

For the case $t \equiv 0 \pmod{a}$ we can also use [6, Theorem 8.4] which gives

$$\sigma_0(1, 1, \dots, 1; a) = 1 - \frac{(-1)^n}{n!} B_n(0; (a, 1, 1, \dots, 1)).$$

(An easy way to see that our two formulations of $\sigma_0(1, 1, \dots, 1; a)$ are equivalent is through the difference formula

$$B_m(x+a_0; (a_0, a_1, \dots, a_n)) - B_m(x; (a_0, a_1, \dots, a_n)) = m B_{m-1}(x; (a_1, a_2, \dots, a_n))$$

and then specializing this to $m = n$, $x = 0$, $a_0 = a$, and $a_1 = a_2 = \dots = a_n = 1$.) Substituting this back into (12) gives, with $\chi_a(t) := 1$ if $a|t$ and $\chi_a(t) := 0$ otherwise:

Proposition 11. $P_{\{a, 1, 1, \dots, 1\}}(t) = \frac{(-1)^n}{n!} (B_n(-t; (a, 1, 1, \dots, 1)) - B_n(t \bmod a; (a, 1, 1, \dots, 1))) + \chi_a(t)$.

Using (13) and Theorem 3 we obtain:

Proposition 12. *Let $\mathbf{a} = (a, 1, 1, \dots, 1)$, where a is a positive integer. Then*

$$\zeta(s; x, \mathbf{a}) = \frac{(-1)^{n-1}}{(n-1)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} B_{n-1-k}(x; \mathbf{a}) \zeta(s-k; x) + \frac{(-1)^{n-1}}{n!} a^{-s} \sum_{r=1}^a B_n(r; \mathbf{a}) \zeta\left(s; 1 + \frac{x-r}{a}\right).$$

Specializing $s = -m$ at negative integers gives, by Proposition 12 with the help of (7), the following formula.

Corollary 13. Let $\mathbf{a} = (a, 1, 1, \dots, 1)$, where a is a positive integer. Then

$$\frac{m!n!}{(m+n)!} B_{m+n}(x, \mathbf{a}) = \sum_{k=0}^{n-1} (-1)^k (k+1) \binom{n}{k+1} B_{n-1-k}(x; \mathbf{a}) \frac{B_{m+k+1}(x)}{m+k+1} + a^m \sum_{r=1}^a B_n(r; \mathbf{a}) \frac{B_{m+1}\left(1 + \frac{x-r}{a}\right)}{m+1}.$$

5. DIFFERENCE, SYMMETRY AND RECURRENCE FORMULAS FOR $B_m(x; \mathbf{a})$

We conclude by giving various formulas for $B_m(x; \mathbf{a})$, starting with the following difference formula.

Theorem 14. For $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$, we have the difference formula

$$(-1)^m B_m(-x; \mathbf{a}) - B_m(x; \mathbf{a}) = m! \sum_{k=0}^{n-1} \sum_{|I|=k} \frac{B_{m-n+k}(x; \mathbf{a}_I)}{(m-n+k)!},$$

with $B_m(x, \mathbf{a}_I) = x^m$ if $I = \emptyset$. Furthermore,

$$B_m\left(x + \sum_{i=1}^n a_i; \mathbf{a}\right) = (-1)^m B_m(-x; \mathbf{a}).$$

Proof. By use of the identity

$$\sum_{I \subset \{1, \dots, n\}} \prod_{i \in \bar{I}} (e^{a_i t} - 1) = e^{(a_1 + \dots + a_n)t},$$

we obtain the formula

$$(14) \quad \frac{t^n e^{(x + \sum_{i=1}^n a_i)t}}{\prod_{i=1}^n (e^{a_i t} - 1)} = \sum_{I \subset \{1, \dots, n\}} \frac{t^n e^{xt}}{\prod_{i \in I} (e^{a_i t} - 1)},$$

where $\bar{I} = \{1, \dots, n\} \setminus I$. On the other hand, we have the equality

$$(15) \quad \frac{t^n e^{(x + \sum_{i=1}^n a_i)t}}{\prod_{i=1}^n (e^{a_i t} - 1)} = \frac{(-t)^n e^{xt}}{\prod_{i=1}^n (e^{-a_i t} - 1)}.$$

Therefore, by (14) and (15),

$$\sum_{I \subset \{1, \dots, n\}} \frac{t^n e^{xt}}{\prod_{i \in I} (e^{a_i t} - 1)} = \frac{(-t)^n e^{xt}}{\prod_{i=1}^n (e^{-a_i t} - 1)}.$$

Together with (8) this completes the proof. \square

Our next result is a symmetry formula.

Theorem 15. Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$ with $A := a_1 + \dots + a_n > 0$. Then for any integers $l, m \geq 0$,

$$(16) \quad (-1)^m \sum_{k=0}^m \binom{m}{k} A^{m-k} B_{l+k}(x; \mathbf{a}) = (-1)^l \sum_{k=0}^l \binom{l}{k} A^{l-k} B_{m+k}(-x; \mathbf{a}),$$

and

$$(17) \quad \begin{aligned} & \frac{(-1)^m}{m+l+2} \sum_{k=0}^m \binom{m+1}{k} (l+k+1) A^{m+1-k} B_{l+k}(x; \mathbf{a}) \\ & + \frac{(-1)^l}{m+l+2} \sum_{k=0}^n \binom{l+1}{k} (l+k+1) A^{l+1-k} B_{m+k}(-x; \mathbf{a}) \\ & = (-1)^{m+1} B_{l+m+1}(x; \mathbf{a}) + (-1)^{l+1} B_{n+m+1}(-x; \mathbf{a}). \end{aligned}$$

Proof. Observe that from (8) we obtain

$$\frac{d}{dx} B_{l+1}(x; \mathbf{a}) = (l+1) B_l(x; \mathbf{a})$$

and by applying the operator $\frac{d}{dx}$ to (16), this implies (17).

Now we prove (16). Consider the generating function

$$\begin{aligned} & \sum_{m \geq 0} \sum_{l \geq 0} \left((-1)^m \sum_{k=0}^m \binom{m}{k} A^{m-k} B_{l+k}(x; \mathbf{a}) \right) \frac{y^m z^l}{m! l!} \\ &= \sum_{l \geq 0} \sum_{k \geq 0} A^{-k} B_{l+k}(x; \mathbf{a}) \frac{z^l}{l!} \sum_{m=k}^{\infty} (-1)^m \binom{m}{k} \frac{(Ay)^m}{m!} \\ &= \sum_{l \geq 0} \sum_{k \geq 0} B_{l+k}(x; \mathbf{a}) \frac{z^l}{l!} \frac{(-y)^k}{k!} e^{-Ay} = e^{-Ay} \sum_{i \geq 0} \sum_{k=0}^i B_i(x; \mathbf{a}) \frac{z^{i-k}}{(i-k)!} \frac{(-y)^k}{k!} \\ &= \sum_{l \geq 0} \sum_{k \geq 0} B_{l+k}(x; \mathbf{a}) \frac{z^l}{l!} \frac{(-y)^k}{k!} e^{-Ay} = e^{-Ay} \sum_{i \geq 0} \sum_{k=0}^i \frac{B_i(x; \mathbf{a})}{i!} \binom{i}{k} z^{i-k} (-y)^k \\ &= e^{-Ay} \sum_{i \geq 0} B_i(x; \mathbf{a}) \frac{(z-y)^i}{i!} = e^{-Ay} \frac{(z-y)^n e^{x(z-y)}}{\prod_{i=1}^n (e^{a_i(z-y)} - 1)} = \frac{(z-y)^n e^{x(z-y)}}{\prod_{i=1}^n (e^{a_i z} - e^{a_i y})}. \end{aligned}$$

Similarly the generating function is also equal to

$$\sum_{m \geq 0} \sum_{l \geq 0} \left((-1)^l \sum_{k=0}^l \binom{l}{k} A^{l-k} B_{m+k}(-x; \mathbf{a}) \right) \frac{y^m z^l}{m! l!} = \frac{(z-y)^n e^{x(z-y)}}{\prod_{i=1}^n (e^{a_i z} - e^{a_i y})}. \quad \square$$

Specializing Theorem 15 at $l = m$ we obtain a recurrence formula for the polynomials

$$P_m(x) := (m+1) A^{-m} \left(B_m(-x; \mathbf{a}) + B_m(x; \mathbf{a}) \right)$$

as follows.

Corollary 16. For any positive integer $m \geq 1$,

$$P_{2m+1}(x) = -\frac{2m+1}{2(m+1)} \sum_{k=0}^m \binom{m+1}{k} P_{m+k}(x).$$

In the case $n = 1$, $a_1 = 1$, the polynomials $P_m(x)$ are reduced to $(m+1) (B_m(-x) + B_m(x))$.

The form of (16) is reminiscent of identities for self-dual sequences, in particular, [22]. Namely, setting

$$Q_k(x; \mathbf{a}) := (-1)^k A^{-k} B_k(x; \mathbf{a}),$$

the case $l = 0$ in Theorem 15 gives

$$Q_m(-x; \mathbf{a}) = \sum_{k=0}^m \binom{m}{k} (-1)^k Q_k(x; \mathbf{a}) =: Q_k^*(x; \mathbf{a}),$$

the *dual* of $Q_k(x; \mathbf{a})$ (viewed as a sequence with index k). In particular, the sequence $Q_k(0; \mathbf{a})$ is *self dual*:

$$Q_m(0; \mathbf{a}) = Q_k^*(0; \mathbf{a}),$$

Let's record this:

Corollary 17. Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$ with $A := a_1 + \dots + a_n > 0$. Then $((-1)^n A^{-n} B_n(\mathbf{a}))_{n \in \mathbb{Z}_{\geq 0}}$ is a self-dual sequence.

It would be interesting to prove this corollary directly (i.e., without referring to Theorem 15). In fact, if this is possible, one could then apply [22, Theorem 1.1] to give an independent proof of Theorem 15.

As a final result, we obtain the following recurrence formula for the Bernoulli–Barnes numbers.

Theorem 18. For any positive integer $m \geq 1$,

$$B_{2m+1}(\mathbf{a}) = -\frac{1}{2(m+1)} \sum_{k=0}^m \binom{m+1}{k} (m+k+1) A^{m+1-k} B_{m+k}(\mathbf{a})$$

and

$$B_{2m}(\mathbf{a}) = -\frac{1}{(m+1)(2m+1)} \sum_{k=0}^m \binom{m+1}{k} (m+k+1) A^{m-k} B_{m+k}(\mathbf{a}) \\ - (2m)! A^{-1} \sum_{k=0}^{n-1} \sum_{|I|=k} \frac{B_{2m+1-n+k}(\mathbf{a}_I)}{(2m+1-n+k)!}.$$

Note that for $n = 1$, the above results specialize to the well-known difference, symmetry and recurrence concerning the ordinary Bernoulli numbers and polynomials.

REFERENCES

1. Takashi Agoh and Karl Dilcher, *Higher-order recurrences for Bernoulli numbers*, J. Number Theory **129** (2009), no. 8, 1837–1847.
2. Tom M. Apostol, *Generalized Dedekind sums and transformation formulae of certain Lambert series*, Duke Math. J. **17** (1950), 147–157.
3. Peter Barlow, *An Elementary Investigation of the Theory of Numbers*, J. Johnson & Co., London, 1811.
4. Ernest W. Barnes, *The theory of the double gamma function*, Philos. Trans. R. Soc. London, Ser. A **196** (1901), 265–387.
5. Matthias Beck, Ricardo Diaz, and Sinai Robins, *The Frobenius problem, rational polytopes, and Fourier–Dedekind sums*, J. Number Theory **96** (2002), no. 1, 1–21, [arXiv:math.NT/0204035](https://arxiv.org/abs/math.NT/0204035).
6. Matthias Beck and Sinai Robins, *Computing the Continuous Discretely: Integer-point Enumeration in Polyhedra*, Undergraduate Texts in Mathematics, Springer, New York, 2007, Electronically available at <http://math.sfsu.edu/beck/ccd.html>.
7. Richard Dedekind, *Erläuterungen zu den Fragmenten xxviii*, Collected Works of Bernhard Riemann, Dover Publ., New York, 1953, pp. 466–478.
8. Karl Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996), no. 1, 23–41.
9. Emilio Elizalde, *Some analytic continuations of the Barnes zeta function in two and higher dimensions*, Appl. Math. Comput. **187** (2007), no. 1, 141–152.
10. Chelo Ferreira and José L. López, *Asymptotic expansions of the double zeta function*, J. Math. Anal. Appl. **274** (2002), no. 1, 134–158.
11. Eduardo Friedman and Simon N. M. Ruijsenaars, *Shintani–Barnes zeta and gamma functions*, Adv. Math. **187** (2004), no. 2, 362–395.
12. Eldon J. Hansen, *A table of series and products*, Prentice-Hall, Englewood Cliffs, N.J., 1975.
13. Ken Kamano, *Sums of products of hypergeometric Bernoulli numbers*, J. Number Theory **130** (2010), no. 10, 2259–2271.
14. Koji Katayama, *Barnes’ double zeta function, the Dedekind sum and Ramanujan’s formula*, Tokyo J. Math. **27** (2004), no. 1, 41–56.
15. ———, *Barnes’ multiple zeta function and Apostol’s generalized Dedekind sum*, Tokyo J. Math. **27** (2004), no. 1, 57–74.
16. Kohji Matsumoto, *Asymptotic expansions of double zeta-functions of Barnes, of Shintani, and Eisenstein series*, Nagoya Math. J. **172** (2003), 59–102.
17. Tiberiu Popoviciu, *Asupra unei probleme de partiție a numerelor*, Acad. Republicii Populare Romane, Filiala Cluj, Studii și cercetări științifice **4** (1953), 7–58.
18. Hans Rademacher and Emil Grosswald, *Dedekind Sums*, The Mathematical Association of America, Washington, D.C., 1972.
19. Simon N. M. Ruijsenaars, *On Barnes’ multiple zeta and gamma functions*, Adv. Math. **156** (2000), no. 1, 107–132.
20. Junya Satoh, *Sums of products of two q -Bernoulli numbers*, J. Number Theory **74** (1999), no. 2, 173–180.

21. Mauro Spreafico, *On the Barnes double zeta and Gamma functions*, J. Number Theory **129** (2009), no. 9, 2035–2063.
22. Zhi-Wei Sun, *Combinatorial identities in dual sequences*, European J. Combin. **24** (2003), no. 6, 709–718.
23. Zhi-Wei Sun and Hao Pan, *Identities concerning Bernoulli and Euler polynomials*, Acta Arith. **125** (2006), no. 1, 21–39.

LABORATOIRE ANALYSE ET PROBABILITÉS, UNIVERSITÉ D'EVRY VAL D'ESSONNE, 23 BD. DE FRANCE,, 91037 EVRY, FRANCE

E-mail address: abayad@maths.univ-evry.fr

DEPARTMENT OF MATHEMATICS, SAN FRANCISCO STATE UNIVERSITY, 1600 HOLLOWAY AVE, SAN FRANCISCO, CA 94132, U.S.A.

E-mail address: mattbeck@sfsu.edu