



**HAL**  
open science

## Effects of agricultural practices on biodiversity. A review

Corentin Babin, Sandrine Espagnol, Joël Aubin

### ► To cite this version:

Corentin Babin, Sandrine Espagnol, Joël Aubin. Effects of agricultural practices on biodiversity. A review. 2023. hal-04465927

**HAL Id: hal-04465927**

**<https://hal.science/hal-04465927v1>**

Preprint submitted on 19 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open licence - etalab

1 **Effects of agricultural practices on biodiversity. A review**

2 Corentin Babin<sup>1</sup>, Sandrine Espagnol<sup>1</sup>, Joël Aubin<sup>2</sup>

3 ✉ Babin Corentin  
4 babincorentin@live.fr

5 <sup>1</sup> IFIP, 35650 Le Rheu, France

6 <sup>2</sup> SAS, INRAE, Institut Agro, 35000 Rennes, France

7

8 Abstract

9 The loss of biodiversity has become a major concern in the 21<sup>st</sup> century. The current agricultural model  
10 is one of the main drivers of biodiversity loss; however, agriculture can also help maintain and even  
11 promote biodiversity. Therefore, understanding of the positive and negative effects of agricultural  
12 practices on biodiversity needs to be increased. We developed a method to summarize the results of  
13 meta-analyses on this subject. We reviewed 27 meta-analyses from the literature to provide a detailed  
14 overview of their results and quality. These meta-analyses examined effects of several agricultural  
15 practices on terrestrial biodiversity at the local scale: inorganic or organic fertilization, crop rotation,  
16 intercropping, use of cover crops, use of plant-protection products, tillage, landscape management and  
17 grazing. We found that (i) the meta-analyses considered certain taxonomic or functional groups much  
18 more than others, depending on the agricultural practices considered and feasibility criteria; (ii)  
19 compositional biodiversity, especially at the species level, was usually considered through changes in  
20 abundance, richness or diversity; (iii) the taxonomic/functional groups and metrics chosen determined  
21 the biodiversity observed and (iv) all of the agricultural practices studied influence biodiversity  
22 dynamics, and some have a dominant influence, such as the use of plant-protection products and  
23 landscape management. However, the intensity of practices, interactions between practices and the  
24 sensitivity of environments influenced the effects observed and must be better understood to classify  
25 practices objectively according to their effects on biodiversity. Further research and data are required  
26 to qualify in more detail the relations between agricultural practices and biodiversity, as well as to  
27 increase understanding of the relations between ecosystem functioning and the compositional and  
28 structural components of biodiversity.

29

30 Keywords: systematic review, meta-analyses, fertilization, tillage, plant diversification, grazing, plant-  
31 protection products, landscape complexity

32

33 Contents

34 1. Introduction ..... 3

35 2. Scope and methodology ..... 5

36 3. Analyzing the meta-analyses ..... 6

37 4. Effects of the main agricultural practices ..... 7

38 4.1. Effects of fertilization ..... 7

39 4.1.1 Effects of inorganic fertilization ..... 8

40 4.1.2. Effects of organic fertilization ..... 9

41 4.2. Effects of crop diversification at the field scale ..... 10

42 4.2.1. Effects of crop rotation ..... 10

43 4.2.2. Effects of intercropping ..... 11

44 4.2.3. Effects of cover crops ..... 12

45 4.3. Effects of tillage ..... 12

46 4.4. Effects of livestock grazing ..... 13

47 4.5. Effects of landscape management ..... 14

48 4.6. Effects of plant-protection products ..... 16

49 5. Overview of biodiversity in meta-analyses of agricultural practices ..... 18

50 6. Limits of reviewing meta-analyses ..... 19

51 7. Conclusion ..... 19

52 Acknowledgements ..... 20

53 Bibliography ..... 21

54

55

## 56 1. Introduction

57

58 The 1992 United Nations Conference on Environment and Development solidified the concept of  
59 biodiversity by establishing the Convention on Biological Diversity (CBD), an international treaty  
60 initially signed and ratified by nearly 150 governments and 196 countries. The CBD resulted in  
61 recognizing the challenges of protecting biodiversity as a whole at the global scale by strengthening  
62 national and international actions. The CBD's general objectives were conservation of biodiversity,  
63 sustainable use of its components and fair and equitable sharing of the benefits of using genetic  
64 resources. The CBD defined biological diversity as "the variability among living organisms from all  
65 sources including, inter alia, terrestrial, marine and other aquatic ecosystems and the ecological  
66 complexes of which they are part; this includes diversity within species, between species and of  
67 ecosystems" (article 2 of the CBD (1992)). This definition focuses on "variability" at three levels of  
68 organization of living organisms: (1) genes (i.e., intraspecific diversity), (2) species (i.e., interspecific  
69 diversity) and (3) ecosystems (i.e., ecosystem diversity). Diversity can also be understood in terms of  
70 time (e.g., ecological successions, changes in crop rotations) (Peeters et al., 2004).

71 From an operational viewpoint, biodiversity has been defined by three inseparable components in the  
72 literature: composition, structure and functioning (Noss, 1990). In theory, they are positively  
73 correlated, and they operate at all geographic scales and all levels of organization of living organisms  
74 (Figure 1). Composition refers to the identity and variety of the elements of the diversity, such as  
75 allelic richness in a population, species richness in a community or taxonomic diversity in an  
76 ecosystem. Structure refers to the elements that support the composition and the "physical"  
77 organization or arrangement of the system, such as genetic structure at the population level or the size  
78 or distribution of habitats at the landscape level. Functioning involves ecological and evolutionary  
79 processes (e.g., gene flow, nutrient cycles) and refers more generally to all interactions at the levels of  
80 organization of living organisms (e.g., genetic, demographic, interspecific, ecosystem).

81 As climate change, biodiversity loss is currently one of the most critical environmental issues. At the  
82 species level, evidence increasingly reveals that current extinction rates for several taxa are up to 100  
83 times as high as the background rate (Ceballos et al., 2015; IPBES, 2019). According to the latest  
84 version of the IUCN Red List (version 2022.2), 42,108 out of 150,388 species studied are classified as  
85 threatened with extinction, including 41% of amphibians, 37% of sharks and rays, 36% of reef-  
86 building corals, 34% of conifers, 27% of mammals and 13% of birds. Besides a decrease in the  
87 number of species, the abundance of biodiversity has decreased drastically in recent decades. The  
88 Living Planet Index reveals a mean decrease of 69% in wild animal populations monitored since 1970,  
89 and Hallmann et al. (2017) estimated a decrease of more than a 75% in the biomass of flying insects in  
90 German protected areas (representative of those in western Europe) protected areas from 1989 to 2016.

91 The Millennium Ecosystem Assessment (2005) indicated that biodiversity loss was due to five  
92 pressures: land use/land-use change, invasive species, climate change, overexploitation and pollution  
93 (Reid et al., 2005). Among land-use change, conversion of natural habitats was considered the main  
94 driver of biodiversity loss, and agriculture is one of the main reasons that natural habitats are  
95 converted (IPBES, 2019). Agriculture covers ca. 40% of the world's land area (Ellis et al., 2020; *World*  
96 *Bank Open Data*, 2020) and is expanding at the expense of forests (Gibbs et al., 2010). From 2010-  
97 2015, the area of tropical forests decreased by 5.5 million ha per year (Keenan et al., 2015), a trend  
98 that is predicted to continue along with a growing demand for food (Kehoe et al., 2017; Obersteiner et  
99 al., 2016). Nonetheless, although concerns about biodiversity loss focus mainly on the loss of species  
100 due to conversion of natural habitats, human-managed landscapes maintain a certain amount of  
101 biodiversity, and some of them can have a species richness similar to that of natural habitats (Altieri,  
102 1999) and sometimes maintain populations of species in decline (Tucker & Murphy, 1997). It is clear  
103 that consideration of biodiversity must include not only natural areas but also human-managed areas  
104 (Tscharntke et al., 2005).

105 Agricultural production systems were intensified during the 20<sup>th</sup> century under the impetus of  
106 productivist public policies (Hazell & Wood, 2008), which resulted in highly specialized production  
107 systems that exceeded the carrying capacity of the environment. The focus on increasing yields  
108 required reducing crop and livestock diversity and a massively increasing the use of inputs, which  
109 changed soil quality (Foley et al., 2005), the water cycle (Shiklomanov & Rodda, 2004),  
110 biogeochemical cycles (X. Liu et al., 2010; Y. Liu et al., 2008; Smil, 2000), the climate (Wollenberg et  
111 al., 2016) and biodiversity in general (Burel et al., 2008). Agriculture thus plays an active role in  
112 increasing eutrophication, water pollution, greenhouse gas emissions, soil erosion and water scarcity;  
113 decreasing soil organic matter; dependence on non-renewable resources and thus loss of biodiversity  
114 and its associated ecosystem services, such as carbon sequestration and pollination (Campbell et al.,  
115 2017; Emmerson et al., 2016). However, although agriculture is known to influence global  
116 biodiversity, the detailed relations between agricultural practices and biodiversity are not sufficiently  
117 understood, and they need to be understood better to promote practices that combine biodiversity  
118 preservation and productivity.

119 Understanding the environmental impacts of human activities, particularly agriculture, has resulted in  
120 environmental policies influenced by the adoption of objectives at international (e.g., CBD), regional  
121 (e.g., European Environment Agency) and national (e.g., national biodiversity strategy in France)  
122 scales that are based largely on the CBD's Aichi biodiversity targets (*Convention on Biological*  
123 *Diversity*, 2016)). To this end, assessment receives particular attention, which reflects an increased  
124 need for assessment methods to guide policies, consumers and research towards the design and  
125 adoption of more sustainable production systems. However, one major challenge is constructing  
126 reliable and feasible assessment methods, which depends on data gaps, standardization, availability

127 and accessibility. Consequently, technical tools and sustainable digital infrastructure need to be  
128 developed that can support the discovery, analysis, access, dissemination and permanent storage of the  
129 increasingly complex datasets needed to quantify changes in biodiversity at the global scale (Kissling  
130 et al., 2018). A large amount of data is collected and available worldwide, but it remains difficult to  
131 assemble knowledge to assess biodiversity, particularly in an agricultural context. These difficulties  
132 are due to (i) the complexity and scope of the concept of biodiversity (Duelli & Obrist, 2003), (ii) the  
133 variety of agricultural practices and soil and climate conditions (Burel et al., 2008; Le Roux et al.,  
134 2012) and (iii) the intrinsic characteristics of assessment methods, such as scale effects (Feest et al.,  
135 2010), spatial and temporal limits of the systems, the values chosen (e.g., functional unit, reference  
136 value) and sensitivity of the models (Bockstaller et al., 2013). Data on biodiversity can be used to  
137 calibrate predictive models and monitor and assess the effectiveness of initiatives (e.g., public  
138 policies), as well as to develop and prioritize methods and indicators.

139 In this context, the aim of this review was to collect and summarize data from the literature on effects  
140 of agricultural practices on biodiversity. To this end, we reviewed the main knowledge about the  
141 relations between agricultural practices and biodiversity by summarizing the results of meta-analyses  
142 that were selected using several criteria.

143

144 Figure 1.

## 145 2. Scope and methodology

146

147 A robust and operational database was created to assess the effects of the main agricultural practices  
148 on biodiversity at the local scale. Other criteria were also identified, such as the metrics used to assess  
149 biodiversity, the distributions of the components and levels of biodiversity organization and the  
150 anthropogenic factors related to differences in the metrics used. The literature consulted was mainly  
151 meta-analyses that focused on relations between agricultural practices and biodiversity dynamics (in  
152 time and space), because the many studies of the subject analyze only some of these relations due to  
153 its extent and spatiotemporal variability, the variability in soil and climate conditions and the large  
154 number of agricultural practices. Meta-analyses provide several advantages:

- 155 • provide an overview of existing studies and thus cover a larger perimeter
- 156 • can assess the variability in results among many studies and combine a large amount of  
157 information about a given subject, which increases the robustness of the effects observed
- 158 • are based on precise protocols that require a certain degree of homogeneity in the metrics of  
159 the studies analyzed, which reflects the main methodological choices in the literature

160 The literature was searched using the Web of Science and Google Scholar databases from January to  
161 March 2023. Meta-analyses were identified using the following query: “meta-analysis AND agricultu\*  
162 OR farm\* OR crop\* AND biological diversity OR biodiversity OR richness OR evenness OR  
163 composition\* OR abundance OR function\* OR structur\*”. The search identified 279 studies, of which  
164 76 were selected based on the title or abstract. After further reading, only 27 met the selection criteria  
165 (Table 1) and were included in the review.

166

167 Table 1.

168

169 A total of 583 effects were identified in the 27 meta-analyses, and they were summarized and  
170 organized in a spreadsheet (i.e., one row per effect) (Babin et al. 2023). Each row includes (i) the type  
171 and (ii) subtype of agricultural practice, (iii) the direction (i.e., positive, neutral or negative) of the  
172 effect on biodiversity and its statistical significance, (iv) the number of studies for each effect, (v) the  
173 number of observations for each effect, (vi) the baseline situation used for comparison, (vii) the scale  
174 considered, (viii) the meta-analysis reference, (ix) the subject of the study (i.e., taxonomic, functional  
175 and/or other group) and (x) the metrics used for each effect, which were grouped into four categories  
176 (Table 2): abundance, diversity, richness and biological activity, which represented 45%, 31%, 19%  
177 and 5% of the identified effects, respectively.

178

179 Table 2.

180

### 181 3. Analyzing the meta-analyses

182

183 Of the 27 meta-analyses, 24 were published after 2015, illustrating the increasing importance of  
184 biodiversity issues and the development of meta-analyses. For the 20 meta-analyses that specified the  
185 locations of the studies they analyzed, most of the studies had been conducted in western Europe,  
186 China or North America (Figure 2). Overall, 11 types of farming practices were identified in the meta-  
187 analyses (Table 3).

188

189 Figure 2.

190 Table 3.

## 191 4. Effects of the main agricultural practices

### 192 4.1. Effects of fertilization

193

194 Fertilization was one of the practices studied most, and nearly 25% of the meta-analyses addressed  
195 fertilization, particularly nitrogen fertilization. Fertilization impacts biodiversity in two main ways:  
196 directly changing physico-chemical properties of the soil and impacting soil microorganisms related to  
197 resource acquisition, thus changing trophic chains and influencing the nutritional status of plants.  
198 Overall, fertilization has more obvious effects on fields with higher plant diversity, such as grassland,  
199 than on fields with crops (Burel et al., 2008). In general, fertilization has positive effects on the  
200 abundance and growth of soil organisms, and increases soil activity (e.g., rates of litter decomposition  
201 and soil respiration) (de Graaff et al., 2019) and the abundance of genes involved in the nitrogen cycle  
202 (Ouyang et al., 2018). However, several studies found contrasting effects on species richness and  
203 diversity when distinguishing organic and inorganic fertilization, but few of the studies described these  
204 two main types of fertilizers in detail. This is a disadvantage, since more detailed description is  
205 required to study effects of fertilization on biodiversity. For organic fertilization, this includes  
206 distinguishing effects of different types of manure and their "quality" (defined by Köninger et al.,  
207 2021). For inorganic fertilization, this includes distinguishing the elements supplied (e.g., nitrogen  
208 alone, nitrogen and phosphorus), the forms of the fertilizer and the doses applied. In addition, some  
209 taxonomic groups that fertilization can impact directly, such as fungi and arthropods, are less well  
210 documented. The scale of studies should also be considered, since above certain thresholds determined  
211 by soil and climate conditions, fertilizers (particularly nitrogen and phosphorus) can degrade the  
212 environment. Overfertilization inevitably leads to emissions to the environment, which contribute to  
213 impacts on ecosystems (e.g., eutrophication, acidification, ecotoxicity), human health (e.g., water  
214 quality) and climate change (i.e., greenhouse gas emissions) (Burel et al., 2008; Parnaudeau et al.,  
215 2012). Fertilization can influence multiple processes that determine the fate of nutrients, which can in  
216 turn influence biodiversity, nitrogen immobilization, crop uptake and removal, denitrification (and to  
217 some extent, nitrification), volatilization, leaching, run-off and erosion (Hofman & Cleemput, 2004).  
218 The relative magnitudes of these processes depend on environmental variables such as soil pH,  
219 texture, carbon content, bulk density; slope, water supply and temperature, as well as human  
220 interventions such as the type, amount, location and timing of inputs; crop-residue management;  
221 tillage; drainage; irrigation; land-use change and the stocking rate of grasslands (Burel et al., 2008;  
222 Khan et al., 2018; Le Roux et al., 2012). Many factors contributed to the variability in the effects  
223 measured in the studies, which makes it challenging to summarize their results.



#### 224 4.1.1 Effects of inorganic fertilization

225

226 Effects of inorganic fertilization on biodiversity varied (Figure 3). Compared to an unfertilized control,  
227 Dai et al., (2018) found an increase in microbial diversity with the application of nitrogen, phosphorus  
228 and potassium together and, conversely, a decrease in diversity with nitrogen alone. Some fertilizers,  
229 particularly urea and ammonium, can inhibit the development of microorganisms, mainly due to  
230 toxicity and increased ionic strength, or soil acidification (especially ammonium) (Geisseler & Scow,  
231 2014; Hofman & Cleemput, 2004). The results of de Graaff et al. (2019) differed, indicating that  
232 inorganic nitrogen fertilization has a neutral effect (i.e., no effect) on bacterial diversity. Geisseler &  
233 Scow (2014) suggested that inorganic fertilization can change the composition of microbial  
234 communities greatly in the short term, but that environmental factors and crop management influence  
235 longer-term effects. Inorganic fertilization seems to select for more copiotrophs (e.g., proteobacteria)  
236 and fewer oligotrophs (e.g., many acidobacteria) (Dai et al., 2018). In several diversified agricultural  
237 ecosystems such as grasslands, nitrogen inputs can influence the composition of microbial  
238 communities (Geisseler & Scow, 2014). Nonetheless, soil microbial activity increases (de Graaff et al.,  
239 2019) without increasing the abundance of microbial genes involved in soil nitrogen processes  
240 (Ouyang et al., 2018).

241 Some studies found that nitrogen fertilization has an extremely negative effect overall on the  
242 abundance, but not the diversity, of soil fauna (de Graaff et al., 2019). Among the few studies of  
243 arthropods documented in the meta-analyses, Hu et al. (2022) found a neutral effect on the abundance  
244 of springtails and mites. Puissant et al. (2021) found that inorganic fertilization had a neutral effect on  
245 the abundance, as well as a negative effect on the species richness and diversity, of nematode trophic  
246 groups and other nematodes communities. Liu et al. (2016) found somewhat different results for  
247 fertilization with nitrogen, phosphorus and potassium together, which had a neutral effect on the  
248 abundance, species richness and diversity of nematodes. With nitrogen fertilization only, Hu et al.  
249 (2022) found negative effects on the abundance of nematodes in all trophic groups, except for a neutral  
250 effect on bacterivores, while Liu et al. (2016) found neutral effects on the abundance and species  
251 richness of nematodes and, surprisingly, a positive effect on their diversity.

252 Analyzing the effects of inorganic fertilization on biodiversity is complex given the controversy  
253 surrounding most of the effects identified. The results of the meta-analyses may differ for several  
254 reasons. The definition of inorganic fertilization is not always clear: fertilizer can contain only  
255 nitrogen, phosphorus or potassium, or combinations of two or all three of the elements. There are also  
256 several major compounds of nitrogen (e.g., nitrate, ammonium, urea), phosphorous (e.g., phosphorus  
257 pentoxide, calcium phosphate, ammonium phosphate) and potassium (e.g., potassium oxide, potassium  
258 chloride, potassium nitrate, potassium sulfate), which can have distinct direct or indirect effects. For  
259 example, Bouman et al. (1995) found that applying ammonia for nine years resulted in greater

260 acidification than applying urea did. Despite this, little information exists on direct effects of specific  
261 compounds. Indirect effects (i.e., on the ecosystem) are better documented (e.g., Khan et al. (2018))  
262 but are related to the dynamics of biodiversity mainly by theoretical relations or response laws.  
263 Analyzing the effects of inorganic fertilization is complicated, since biodiversity responses depend on  
264 the amount applied, the soil and climate contexts, the duration of application and the previous  
265 agronomic operations (de Graaff et al., 2019; Hu et al., 2022; Humbert et al., 2016; Liu et al., 2016;  
266 Puissant et al., 2021). Future research is needed to provide more precise information.

267 Figure 3.

268

#### 269 4.1.2. Effects of organic fertilization

270

271 The effects of organic fertilization on biodiversity varied (Figure 4). Compared to inorganic  
272 fertilization, organic fertilization has positive effects on the abundance and diversity of microbial  
273 communities and soil fauna (de Graaff et al., 2019; Morugan-Coronado et al., 2022). Compared to  
274 non-fertilization, organic fertilization increases nematode abundance, particularly for certain trophic  
275 groups (i.e., bacterivores, fungivores and omnivores), without increasing the abundance of herbivores.  
276 It also increases the diversity, but not the species richness, of nematodes (Liu et al., 2016; Puissant et  
277 al., 2021). Liu et al. (2016) suggested that applying carbon-rich crop residues (e.g., straw) is the most  
278 effective practice for improving soil biodiversity in intensively managed agroecosystems, which  
279 highlights the importance of doing so. Organic fertilizers are important sources of nutrients that ensure  
280 a succession of communities and development of a complex food web (Burel et al., 2008; de Graaff et  
281 al., 2019; Liu et al., 2016; Puissant et al., 2021). The content of soil organic matter and soil  
282 biodiversity are closely related. Organic fertilizers are a source of organic carbon that feeds soil  
283 organisms, enabling the soil community to function properly (Thiele-Bruhn et al., 2012). de Graaff et  
284 al. (2019) and Liu et al. (2016) found that large inputs of nitrogen fertilizers only simplify the structure  
285 and functioning of communities and that organic fertilizers buffer the stress generated and maintain the  
286 functions of the soil food web. In addition, carbon-rich fertilizers support larger populations, such as  
287 those of free-living nematodes (Liu et al., 2016).

288 Many factors influence the effects of organic fertilization on biodiversity, such as soil and climate  
289 conditions or the type of organic matter (e.g., green manure; composted or uncomposted cattle,  
290 chicken or pig manure; composted plant waste; crop residues; straw) (Burel et al., 2008; de Graaff et  
291 al., 2019; Köninger et al., 2021; Puissant et al., 2021). According to Köninger et al. (2021), the quality  
292 of organic fertilizer (e.g., concentrations of heavy metals, antibiotics, hormones and pathogens)  
293 influences biodiversity more than does the amount applied. Despite the many influencing factors

294 identified, the effects of different types of organic fertilizer cannot be distinguished, especially since  
295 studies generally have considered only a few taxa, mainly bacteria and nematodes.

296 Figure 4.

297

## 298 4.2. Effects of crop diversification at the field scale

299

300 Increasing plant diversity is a major practice for reducing the use of chemical inputs, managing pests  
301 and controlling biogeochemical cycles (Burel et al., 2008). Modifying cropping systems via crop  
302 rotation, intercropping and cover crops is a relevant way to increase the functioning of agrosystems.  
303 Nonetheless, productivist public policies have promoted development of a dominant agricultural  
304 model defined by system “lock-ins” that prevent crop diversification both upstream (e.g., varietal  
305 selection) and downstream (e.g., consumer demand) of agricultural production (Meynard et al., 2013).  
306 Although detailed information about the effects of crop diversification exists, it has not been the  
307 subject of meta-analyses, and a large proportion of studies combine all crop-diversification practices  
308 together, which removes the necessary detail. Several aspects require more research, especially the  
309 effects of incorporating different crops such as legumes, different rotation durations and incorporating  
310 perennial crops. Increased plant diversity is associated with a decrease in pests and chemical inputs,  
311 and the closing of biogeochemical cycles (Altieri, 1999; Isbell et al., 2017; Tilman et al., 2002).  
312 Increasing crop diversification via rotation, intercropping and cover crops is thus a relevant way to  
313 increase agroecosystem functioning (Chen et al., 2019; Kim et al., 2020; Puissant et al., 2021; Venter  
314 et al., 2016).

### 315 4.2.1. Effects of crop rotation

316

317 The effects of crop rotation on biodiversity varied (Figure 5). In theory, increasing the number of plant  
318 species in a crop rotation can increase agroecosystem functioning (Hooper & Vitousek, 1997; Smith et  
319 al., 2008; Venter et al., 2016). Compared to monoculture, crop rotation (i.e., a multi-year sequence of  
320 different annual crops grown in the same field), especially with legumes, is associated with increased  
321 soil fertility, by maintaining soil structure, disrupting pest cycles and reducing weeds (Smith et al.,  
322 2008). However, the processes that underlie these potential positive effects can be due to incorporating  
323 certain species into rotations (e.g., legumes) or to niche facilitation and differentiation (Hooper et al.,  
324 2005). In practice, soil microorganisms influence these processes through their complex biochemical  
325 processes and their interactions with plants. However, little empirical evidence exists; for example,  
326 Venter et al. (2016) provided one of the first meta-analyses of the effects of temporal heterogeneity in  
327 plant diversity on soil biodiversity. Crop rotation can increase the abundance and species richness of  
328 soil organisms (Morugan-Coronado et al., 2022; Venter et al., 2016), particularly fungi (Morugan-

329 Coronado et al., 2022), but has much smaller effects on other organisms. For example, Puissant et al.  
330 (2021) found that crop rotation had no effect on the abundance, diversity or species richness of  
331 nematode trophic groups, except for a decrease in the abundance of herbivores. Nonetheless, certain  
332 biases may remain. For example, methods of microbial analysis may differ greatly in the effects that  
333 they detect (Venter et al., 2016). Most of the rotations studied are short, but long rotations (> 7 years)  
334 have larger positive effects on microbial richness (Venter et al., 2016), and when they include  
335 grassland, they are particularly favorable for increasing the abundance and species richness of most  
336 soil organisms (Burel et al., 2008). Studies usually aggregate data on annual and perennial crops,  
337 making it impossible to differentiate the effects of two different rotations, and little summary  
338 information exists on the inclusion of specific species in rotations, except for legumes. Venter et al.  
339 (2016) found no significant effect of including legumes in rotations on microbial diversity or richness.  
340 Rotations are studied mainly as a way to reduce the incidence of pests (e.g., insects, mites, pathogens,  
341 nematodes, weeds), whose populations can be maintained at acceptable levels (Reddy, 2017) by  
342 introducing non-host plants into rotations to interrupt the pests' development cycles (Burel et al.,  
343 2008) (e.g., reducing the abundance of herbivorous nematodes; Puissant et al. (2021)).

344 Figure 5.

345

#### 346 4.2.2. Effects of intercropping

347

348 Effects of intercropping on biodiversity varied (Figure 6). Intercropping is defined as growing two or  
349 more different crops in the same field at the same time. In practice, different types of intercropping can  
350 be distinguished:

- 351 (i) intercropping of crop varieties, which provide intraspecific (i.e., genetic) diversity to a field and  
352 are studied mainly for their effects on productivity (i.e., pest control and yield) (Montazeaud et  
353 al., 2022; Reiss & Drinkwater, 2018)
- 354 (ii) intercropping of perennial species (e.g., multi-species orchards), which were not included in the  
355 meta-analyses
- 356 (iii) intercropping of annual species, such as meslin (i.e., several cereals) and tremois (i.e., cereals  
357 and legumes), which have different spatial patterns (e.g., mixed, row, strip or relay  
358 intercropping) that can have different effects on biodiversity (Neamatollahi et al., 2013). Nearly  
359 all of the related studies in the meta-analyses referred to this type of intercropping. Like those of  
360 crop rotation, effects of intercropping on the immediate environment are influenced by soil  
361 microorganisms and feed back to increase the abundance and species richness of these  
362 microorganism communities (Chen et al., 2019; Morugan-Coronado et al., 2022). Older and  
363 more diversified intercrops had more influence on microbial abundance (Chen et al., 2019), and

364 these effects concerned only fungi (Morugan-Coronado et al., 2022). Intercropping also had  
365 positive effects on arthropod abundance and species richness (Sanchez et al., 2022) but not on  
366 nematode trophic groups (Puissant et al., 2021).

367 (iv) intercropping of annual and perennial species (i.e., agroforestry), which provides many  
368 ecosystem services (e.g., water purification, soil-quality maintenance, fiber and wood supply)  
369 and food and habitats for many species (Beillouin et al., 2021). Agroforestry has particularly  
370 positive effects on the abundance and species richness of birds (Torralba et al., 2016). Beillouin  
371 et al., (2019) found that agroforestry consistently improved biodiversity and soil quality,  
372 particularly soil organic carbon content. However, few data have been collected, and the  
373 existing data are context-dependent because agroforestry is studied mainly in tropical and  
374 subtropical ecosystems (Beillouin et al., 2019).

375 Figure 6.

376

#### 377 4.2.3. Effects of cover crops

378

379 Effects of cover crops on biodiversity varied (Figure 7). Plant cover provides an organic resource and  
380 helps maintain favorable microclimates for soil organisms and water quality (i.e., catch crops) (Burel  
381 et al., 2008; Le Roux et al., 2012). Cover crops are used mainly to reduce weed populations, nutrient  
382 loss and soil erosion (Puissant et al., 2021). Kim et al. (2020) found that cover crops increased the  
383 abundance, biological activity and species richness of soil microbial communities compared to bare  
384 fallows. This effect is due mainly to the influence of above- and below-ground plant biomass and root  
385 exudates of cover crops (Vukicevich et al., 2016). Thus, cover crops indirectly promote the abundance  
386 of certain groups, such as nematodes, especially herbivores and bacterivores (Puissant et al., 2021).  
387 Moreover, cover crop mixtures increase nitrogen uptake and biomass production (via  
388 complementarity) (Elhakeem et al., 2019) and reduce the abundance of pests without modifying the  
389 composition of their communities (Beaumelle et al., 2021).

390 Figure 7.

391

#### 392 4.3. Effects of tillage

393

394 Effects of tillage on biodiversity varied (Figure 8). Soil tillage prepares fields for sowing, buries the  
395 seeds of annual crops, reduces weed density and, to a lesser extent, can break cycles of pests (e.g.,  
396 slugs, moths) (Burel et al., 2008). Tillage practices vary in the type of plow used, whether the soil is  
397 turned over or not, depth and frequency. Along with other factors, such as soil, climate and related

398 agronomic operations (e.g., crop-residue management, fertilization), this variability makes it difficult  
399 to quantify the effects of tillage on biodiversity (Briones & Schmidt, 2017; Li et al., 2020). The effects  
400 of plowing on soil organisms are controversial and vary greatly. They may be influenced by direct  
401 effects (e.g., exposure to predators, mechanical mortality) or indirect effects via changes in habitats  
402 (e.g., water content, porosity, temperature) or in the spatial distribution of resources (Burel et al., 2008;  
403 de Graaff et al., 2019). More specifically, tillage influences soil communities directly and indirectly by  
404 changing soil structure and decreasing organic matter in the surface layer (de Graaff et al., 2019). The  
405 intensity of tillage influences its effects on soil biodiversity, but since there are many ways to  
406 characterize intensity, the precise effects of different types of tillage are difficult to quantify. In  
407 general, reduced tillage (i.e., decreased frequency and depth) has positive effects on total soil  
408 microbial abundance (Morugan-Coronado et al., 2022) and the abundance and biomass of earthworms  
409 (Moos et al., 2017). Reduced tillage or no-tillage can increase soil bacterial diversity (de Graaff et al.,  
410 2019; Li et al., 2020; Morugan-Coronado et al., 2022) without significantly changing the structure of  
411 fungal communities (de Graaff et al., 2019; Li et al., 2020). This is unexpected, since physical soil  
412 disturbances are thought to impact soil fungi in particular by forcing them to heal their hyphae or  
413 continually increase their mycelial network (Verbruggen & Toby Kiers, 2010). Plowing impacts  
414 macrofauna the most, especially epigeic and anecic earthworms (Briones & Schmidt, 2017; Burel et  
415 al., 2008). Even using reduced tillage for several years selected for earthworm species with a higher  
416 mean individual biomass (Moos et al., 2017). Tillage impacts micro- and mesofauna less than  
417 macrofauna, but decreased abundance of these communities has been observed (Burel et al., 2008).  
418 Puissant et al. (2021) observed an overall negative impact on the abundance and species richness of  
419 nematodes, with strong differences among trophic groups: a strong decrease in the abundance of  
420 omnivores but no impact on bacterivores or fungivores. Physical soil disturbance decreases the  
421 stability of nematode trophic groups, which form a food web of opportunistic taxa that have a strong  
422 ability to colonize and disperse. However, the effects of using reduced tillage or no-tillage must be  
423 compared to those of practices that consequently increase, especially herbicide use (Winkler et al.,  
424 2023).

425 Figure 8.

426

#### 427 4.4. Effects of livestock grazing

428

429 Effects of livestock grazing on biodiversity varied (Figure 9). Many studies have analyzed the  
430 influence of livestock grazing on biodiversity, but most are highly specific, and few reviews or meta-  
431 analyses seem to exist, perhaps due to controversy related to the variability in study results among  
432 sites. The initial “diversity” in grasslands seems to lead to biases in interpretation, especially for plant

433 diversity, whose responses to livestock grazing have been studied the most. Several other factors  
434 complicate the ability to perform reviews or meta-analyses, such as the baseline used for comparison  
435 (usually ungrazed grasslands) differing among studies (especially the duration since last grazing), the  
436 existence (or not) of randomized controlled trials and multi-site observations, differing intensity  
437 measurements (usually the number of animal units/ha) and the fact that the intensity of grassland  
438 management (e.g., fertilization) and frequency of livestock grazing influence the effects of livestock  
439 grazing on biodiversity, but they are rarely considered (Burel et al., 2008; Filazzola et al., 2020; Wang  
440 & Tang, 2019).

441 However, the intensity of livestock grazing seems to be the management factor that influences  
442 biodiversity the most in grazed grasslands. Wang & Tang (2019) found that low- or medium-intensity  
443 livestock grazing can increase the species richness and diversity of microorganisms and decrease those  
444 of arthropods compared to high-intensity grazing. However, they did not explain the baseline used for  
445 comparison, define the intensity levels or describe the methods used. Another meta-analysis (Filazzola  
446 et al. 2020) explored the effects of excluding livestock grazing (followed by abandoning the land) and  
447 found an overall positive effect on the diversity and abundance of many organisms, such as certain  
448 vertebrates and arthropods, but not of detritivores. These positive effects particularly concerned  
449 species that interacted directly with plants (e.g., herbivores, pollinators). However, the effects of  
450 grazing grasslands can persist, while those of excluding livestock decrease after 10-20 years and could  
451 become negative in certain environments (Wang & Tang, 2019). Livestock grazing directly influences  
452 the composition of plant communities and indirectly influences many invertebrate and vertebrate  
453 communities (especially their habitats and food) due to activities that may occur along with or on a  
454 different timescale on the grasslands (e.g., defoliation, fertilization, mowing, health-related treatments,  
455 trampling, excretion, dissemination of seeds, disturbance) (Burel et al., 2008). Livestock grazing thus  
456 exerts direct selection pressure on plants by favoring annual species, plants with a small stature or  
457 prostrate habit, stoloniferous plants or those with a "rosette" architecture (Díaz et al., 2007).

458

459 Figure 9.

460

#### 461 4.5. Effects of landscape management

462

463 Effects of landscape complexity on biodiversity varied (Figure 10). Land-use change is the main cause  
464 of the decrease in the abundance and species richness of communities, which, along with field-based  
465 practices, reduces the complexity of landscapes (i.e., simplification) (IPBES, 2019). Using land for  
466 agricultural production changes the landscape greatly, mainly by increasing habitat fragmentation and

467 decreasing habitat connectivity (Reidsma et al., 2006). Consequently, habitat patches become smaller  
468 and isolated, which decreases the sizes of populations and in turn decreases dispersal and  
469 metapopulation exchanges. To describe the complexity of landscapes and study their effects on  
470 biodiversity, many studies examine landscape heterogeneity, which combines habitat composition (i.e.,  
471 type, size or relative proportion of elements) and configuration (i.e., structure, shape and spatial  
472 arrangement of elements). Several metrics are used to describe the complexity of the composition  
473 (e.g., proportion of habitat types, habitat richness, evenness of distribution) and configuration (e.g.,  
474 connectivity, proximity, patchiness). The variety of metrics used in studies led all meta-analyses to  
475 categorize the metrics, but the elements of complexity were usually poorly defined, which resulted in  
476 comparing systems with blurry boundaries (e.g., "complex" vs. "simple"), which varied among studies.  
477 There are several ways to define landscape complexity, such as a percentage of natural or semi-natural  
478 elements (e.g., 25% representing "complex"), the mean size of fields or habitat connectivity. In  
479 general, the degree of complexity of agricultural landscapes influences how well they can maintain  
480 local populations within themselves (Burel et al., 2008; Estrada-Carmona et al., 2022). In addition,  
481 many invertebrate, vertebrate and plant communities have been observed to have higher species  
482 richness, abundance and diversity in complex landscapes than in simplified landscapes (Chaplin-  
483 Kramer et al., 2011; Estrada-Carmona et al., 2022; Gonthier et al., 2014; Prevedello et al., 2018).  
484 These positive effects influence mainly pollinators and crop auxiliary insects, with much smaller  
485 effects on pests and weeds (Chaplin-Kramer et al., 2011; Estrada-Carmona et al., 2022). Other effects  
486 of landscape simplification are more difficult to quantify, such as the decrease in the abundance of rare  
487 and specialist species in certain habitats and the dominance of the most common generalist species.  
488 Effects can also combine, such as the combined influence of linear elements (e.g., hedges) and surface  
489 elements (e.g., woods), which can be nearly double the size of each effect observed separately  
490 (Estrada-Carmona et al., 2022). The duration of a study can also influence the magnitude and direction  
491 of the results. In addition, Gonthier et al. (2014) highlighted that most studies focus on the influence of  
492 landscape complexity on local management and do not assess the actual influence of the landscape on  
493 biodiversity. They also found that the variability observed in the effects of local landscape  
494 management on organisms could be due to the latter's mobility. For example, local management  
495 practices influence the species richness of plant communities more than that of invertebrate or  
496 vertebrate communities, whose species' greater mobility can protect them from small-scale changes in  
497 local management practices. In addition, some studies suggest that intensively managed agricultural  
498 landscapes may increase the positive effects or decrease the negative effects of certain practices on  
499 species richness by ensuring a pool of biodiversity that is sufficiently large to support recolonization  
500 (e.g., Bengtsson et al. (2005) and Prevedello et al. (2018)). The effects of landscape complexity could  
501 be due to a positive relation between the number and types of habitats (especially the proportion of  
502 semi-natural habitats) and abundance and species richness. The effect of landscape heterogeneity often  
503 refers to the "landscape complementation effect", which refers to the fact that greater habitat



504 variability provides habitats for more (site-specific) species, which require different habitat types  
505 simultaneously (i.e., spatial diversity) or successively (i.e., temporal diversity) to meet their needs  
506 (e.g., feeding, nesting) (Fahrig et al., 2011). The precise relations between landscape heterogeneity and  
507 biodiversity still need to be clarified, both structurally and compositionally, to better identify which  
508 characteristics of the complexity of human-managed landscapes influence biodiversity the most.

509

510 Figure 10.

511

#### 512 4.6. Effects of plant-protection products

513

514 The use of synthetic products is one of the main drivers of biodiversity loss (IPBES, 2019), but the  
515 body of knowledge is heterogeneous. We identified few meta-analyses on the effects of plant-  
516 protection products on biodiversity. A few studies have examined the effects of using certain plant-  
517 protection products (e.g., neonicotinoids, glyphosate) on certain emblematic taxonomic groups,  
518 usually bees, or used a more integrated approach by comparing organic and conventional farming  
519 systems. Some reviews are more specific to the effects of applying plant-protection products on  
520 biodiversity and ecosystem services, in particular the collective scientific expertise of Leenhardt et al.  
521 (2023).

522 Generalizing the effects of plant-protection products on the environment is complicated due to the  
523 large number of products used (particularly toxic ones), the variety of environmental contexts that  
524 influence these effects, the limits of assessing biodiversity and the changes in analytical techniques  
525 and sampling strategies. In 2022, 449 plant-protection substances were registered in the European  
526 Union and marketed in more than 1500 products (Leenhardt et al., 2023). When they are applied, a  
527 certain amount does not reach the target and/or is transferred to the surrounding environment,  
528 especially broad-spectrum products. For example, plants absorb only 1.6-20% of the active ingredients  
529 in insecticides in seed coatings during germination, while the rest is transferred and may persist in the  
530 soil or be transferred to surface water or groundwater (Grout et al., 2020). We distinguished three  
531 types of transfers: horizontal (i.e., runoff, drainage and drift), vertical (i.e., leaching and volatilization)  
532 and those via living organisms (i.e., by plants through absorption or contact and by animals through  
533 bioaccumulation or biomagnification via trophic chains). Plant-protection products can thus  
534 contaminate all environmental compartments (i.e., land, water and air), but the soil is contaminated the  
535 most (Leenhardt et al., 2023). In 2018, 760 substances were detected in groundwater in France, of  
536 which 48% were authorized substances, 37% were banned substances, 9% were metabolites of banned  
537 substances and 5% were metabolites of authorized substances (Beaulaton et al., 2020). Some

538 substances, despite being banned for over 20 years, were still detected, and sometimes were among  
539 those detected the most often, whether in their original form or as metabolites (e.g., atrazine, sodium  
540 chlorate, simazine, alachlor, oxadixyl, chloridazone) (Beaulaton et al., 2020). In another study, Pelosi  
541 et al. (2021) detected at least one pesticide in 100% of 180 soils studied and 92% of 155 earthworms  
542 sampled in both untreated semi-natural areas (i.e., hedges, meadows and organically farmed cereals)  
543 and treated fields in France.

544 Effects of plant-protection products on biodiversity depend on their properties and the degree of  
545 exposure. According to Leenhardt et al. (2023), applying plant-protection products decreases the  
546 abundance and species richness of invertebrates, mainly in agricultural areas. Insects (i.e.,  
547 Lepidoptera, Hymenoptera and Coleoptera) are impacted most by direct effects (i.e., mortality from  
548 insecticides or broad-spectrum products) and indirect effects (e.g., loss of food and habitats due to  
549 herbicide use). At the European scale, ca. 40% of macroinvertebrates are under threat (Leenhardt et al.,  
550 2023). Many studies focus on the decline in bird populations, especially in agricultural areas. In  
551 Europe, agricultural intensification, particularly application of pesticides and fertilizers, is the main  
552 cause of the decline in most bird populations, especially those that feed on invertebrates (Rigal et al.,  
553 2023). The indicator Temporal Monitoring of Common Birds in France shows that the abundance of  
554 75 bird species in agricultural areas decreased by ca. 30% from 1990-2020 (Leenhardt et al., 2023).  
555 The decline is due in part to direct poisoning (e.g., ingestion of treated seeds by granivores or  
556 contaminated bait by birds of prey) and indirect effects (e.g., shortage of food, mainly for  
557 insectivores). High doses of pesticides can cause acute mortality of birds, but longer-term effects of  
558 chronic exposure have been studied much less. For example, Moreau et al. (2021) found that ingesting  
559 low doses of pesticides over a long period had long-term effects on several major physiological  
560 pathways in partridges, without increasing mortality rates, particularly for breeding adults. Among  
561 other effects, individuals can show decreased fitness traits, which can impact populations by reducing  
562 the investment in reproduction and productivity. European Union approval protocols do not consider  
563 the long-term effects of plant-protection products. According to Miles et al. (2018), 79% of herbicides,  
564 75% of fungicides and 92% of insecticides would not have been registered had the risks of chronic  
565 exposure been considered, as proposed in the protocol of the European Food Safety Authority in 2013,  
566 based on a risk assessment of chronic exposure of bees. The meta-analysis of Beaumelle et al. (2021)  
567 found that broad-spectrum or multi-substance products have larger effects on soil fauna, suggesting  
568 that such mixtures, which are frequently detected in agricultural soils or water, would have additive  
569 and/or synergistic effects that are much more harmful to non-target organisms (so called "cocktail  
570 effects"). For example, Pelosi et al. (2021) found that applying neonicotinoids decreased earthworm  
571 abundance significantly more when combined with other pesticides.

572 Effects of organic farming on biodiversity varied (Figure 11). Puissant et al. (2021) found that  
573 transitioning from conventional to organic farming increases the abundance of nematode trophic

574 groups, particularly bacterivores, without altering their composition. Arthropod communities have  
575 much larger abundance and higher species richness under organic farming than conventional farming  
576 (Bengtsson et al., 2005; Lichtenberg et al., 2017; Tuck et al., 2014) but less evenness, due to the  
577 promotion of rare species (Lichtenberg et al., 2017). These effects are stronger at the local scale than at  
578 the regional scale (Lichtenberg et al., 2017). Organic farming influences certain arthropod functional  
579 groups more strongly, such as pollinators and predators (Lichtenberg et al., 2017), and has positive  
580 effects on vertebrate abundance and diversity but not species richness (Bengtsson et al., 2005;  
581 Gonthier et al., 2014; Tuck et al., 2014). Lori et al., (2017) observed significantly higher microbial  
582 biomass and activity under organic farming than conventional farming, which they attributed to more  
583 complex crop rotations and larger organic matter inputs.

584

585 Figure 11.

586

## 587 5. Overview of biodiversity in meta-analyses of agricultural practices

588

589 The components of biodiversity were considered to differing degrees, with species being considered  
590 the most, mainly through changes in abundance, diversity (Shannon index) and species richness. Two  
591 main approaches were used: compositional, which quantifies effects on certain taxonomic groups, and  
592 functional, which quantifies effects on functional groups defined by either their position in the food  
593 web or a functional trait. The studies focused on major taxonomic groups, which were selected  
594 depending on the agricultural practices studied. Nematodes were the group studied the most in the  
595 meta-analyses, usually to study the dynamics and stability of communities (e.g., abundance of trophic  
596 groups) in response to disturbances at the field scale. Nematodes were also used as an indicator of  
597 community stability (e.g., reproduction, food-web complexity) and a bioindicator of soil quality.  
598 Microorganisms were the second-most common group, mainly bacteria and fungi, usually to study  
599 plant-soil interactions at the field scale and more generally as an indicator of soil biological activity.  
600 Arthropods were also frequently studied, mainly to quantify effects of plant diversity (i.e., complexity)  
601 at the landscape scale and, to a lesser extent, at the field scale (especially in intercropping). Like  
602 nematodes, they were usually categorized by their position in the food web, but also by their  
603 “agronomic functions”. Surprisingly, earthworms were studied almost exclusively as an indicator of  
604 “soil health” in studies of tillage effects. Vertebrates were also studied, almost all of them birds at the  
605 landscape scale. While the choice of these groups may suggest their relative importance (e.g., plays an  
606 essential ecological role, conservation status), they may also have been chosen for feasibility and  
607 practicality. For example, nematodes are ubiquitous, diverse (> 25,000 species) and abundant (80% of

608 terrestrial multicellular organisms); occupy nearly all trophic levels and have a strong influence on  
609 ecosystem functioning, particularly biogeochemical cycles, carbon fluxes and the composition of soil  
610 microbial populations (Puissant et al., 2021; van den Hoogen et al., 2019). Thus, they are a particularly  
611 interesting group to use to describe biodiversity dynamics.

## 612 6. Limits of reviewing meta-analyses

613

614 While reviewing meta-analyses can summarize large amounts of information and demonstrate  
615 robustness in the results, it also has limitations. The main limitation is the relatively few combinations  
616 of practices, effects and components of biodiversity studied, especially the absence of certain practices  
617 (e.g., irrigation) and the lack of detail about subtypes of these practices (e.g., adding legumes to  
618 rotations). Even if the data exist, there must be enough of it to perform a meta-analysis, the meta-  
619 analysis must be found, and it must then meet the criteria necessary to be included in a review. The  
620 second main limitation is the inability to assess the magnitude of effects, due to differences in the  
621 methods of the meta-analyses and because such information may lack relevance, particularly given its  
622 high uncertainty. Finally, our approach to the meta-analyses precluded studying interactions between  
623 practices at the system level and their ultimate effects on biodiversity. For example, reduced tillage  
624 decreases impacts on soil biodiversity but can increase herbicide applications. A systems approach is  
625 crucial for assessing crop and livestock systems.

626

## 627 7. Conclusion

628

629 In this review, we compiled the most well-documented and meta-analyzed direct effects of agricultural  
630 practices on biodiversity. At the field scale, the high intensification and simplification of practices  
631 ultimately had negative impacts on multiple biodiversity components, but lower intensification can  
632 promote biodiversity in agroecosystems. However, the variability in results greatly limits the ability to  
633 assess how specific practices influence biodiversity or which practice has the most influence on  
634 biodiversity. Establishing universal response laws seems elusive for most practices and remains a  
635 challenge.

636 We emphasize that the impacts of agricultural land use on biodiversity are related to the intensification  
637 and simplification of systems, mainly more use of plant-protection products and inorganic fertilizers,  
638 lower spatial and temporal diversity of crop species and varieties, larger crop fields and landscape  
639 homogenization, which often decreases landscape complexity. The practices discussed in this review  
640 were those described in the meta-analyses, which focused mainly on large-scale crop and livestock  
641 farming and rarely on horticulture, viticulture, vegetable crops or agroforestry. In addition, many  
642 relatively common practices were not described, such as irrigation, biological pest control and

643 mowing. Intensification of cropping systems is accompanied by intensification of livestock practices,  
644 which relies less on grazing and more on purchased feeds. The effects of practices and systems on  
645 biodiversity must be considered in relation to other characteristics, such as the amount of production  
646 required to meet the demand for food, the simultaneous use of "harmful" and "favorable" practices and  
647 long-term effects, which are more difficult to estimate. These characteristics are currently the subject  
648 of much debate, particularly related to "land sharing" vs "land sparing". Most approaches to  
649 conserving biodiversity rely on monitoring and assessment, which are based largely on the types of  
650 data used in this review. We demonstrate that the taxa and metrics chosen are crucial since the effects  
651 on biodiversity observed depend on them. Given the variety of effects of agricultural practices and  
652 system characteristics on different organisms, agricultural production systems cannot conserve  
653 biodiversity without explicitly clarifying biodiversity objectives.

654 The agricultural practices summarized influence local biodiversity dynamics and should be considered  
655 mechanisms for preserving biodiversity. Nonetheless, it remains unlikely that it will be possible to  
656 accurately and comprehensively quantify the impacts of agricultural practices on biodiversity and thus  
657 classify practices according to these impacts. Many factors contribute to the high variability in the  
658 effects measured and should be explored in future studies, such as the taxa and metrics chosen, soil  
659 and climate conditions, the sensitivity of environments, the degree to which practices are intensified  
660 and interactions between practices. However, landscape complexity has a particularly large influence  
661 because it can provide habitats for more species and enable mobile species to recolonize fields.  
662 Application of plant-protection products also has major direct and indirect negative effects on all  
663 ecosystems and populations of organisms, but their long-term effects and "cocktail effects" remain  
664 poorly understood.

#### 665 [Acknowledgements](#)

666 The authors thank the scientific interest groups Avenir élevage and Grandes cultures for their financial  
667 support, as well as T. Faure for producing the graphics and the members of the working group set up  
668 for this project for their advice: R. Baumont, J-L. Peyraud, C. Bocktaller, B. Dellaporta, F. Angevin, F.  
669 Lescourret, N. Sautereau, E. Andre, M. Fossey, M. Cornelus, H. Gross, G. Fleurance, J. Charef, G.  
670 Chiron, N. Bataille and M. Marcon.

#### 671 [Authors' contributions](#)

672 All authors contributed to the review conceptualization and design. Methodology, data collection,  
673 analysis and writing were performed by C. B. Supervision was performed by J. A. and S. E. The first  
674 draft of the manuscript was written by C. B. all authors commented on previous versions of the  
675 manuscript. All authors read and approved the final version of the manuscript.

#### 676 [Funding](#)

677 This research was financially supported by INRAE.

678 **Data availability**

679 The dataset generated during and analyzed during the current study is available in the data.gouv.fr  
680 repository, <https://doi.org/10.57745/EBRFIP>

681 **Declarations**

682 **Ethics approval** Not applicable.

683 **Consent to participate** Not applicable.

684 **Consent for publication** Not applicable.

685 **Conflict of interest** The authors declare no competing interests

686 **Code availability** Not applicable

687

688 **Bibliography**

689

690 Altieri, M. A. (1999). The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystems &*  
691 *Environment*, 74(1), 19- 31. [https://doi.org/10.1016/S0167-8809\(99\)00028-6](https://doi.org/10.1016/S0167-8809(99)00028-6)

692 Babin, C. , Aubin, J. , Espagnol, S. , 2023, "Effects of agricultural practices on biodiversity",  
693 <https://doi.org/10.57745/EBRFIP>, Recherche Data Gouv, VERSION PROVISOIRE

694 Beaulaton, L., Blard-Zakar, A., Boulenger, C., Bréjoux, E., Cosson, E., Dequesne, J., Kreutzenberger, K.,  
695 Nowak, C., Antoni, V., Cerisier-Auger, A., Dossa-Thauvin, V., Eumont, D., Guilhen, J.-M., &  
696 Guzmova, L. (2020). *Eau et milieux aquatiques—Les chiffres clés*.

697 Beaumelle, L., Auriol, A., Grasset, M., Pavy, A., Thiéry, D., & Rusch, A. (2021). Benefits of increased  
698 cover crop diversity for predators and biological pest control depend on the landscape context.  
699 *Ecological Solutions and Evidence*, 2(3), e12086. <https://doi.org/10.1002/2688-8319.12086>

700 Beillouin, D., Ben-Ari, T., & Makowski, D. (2019). Evidence map of crop diversification strategies at the  
701 global scale. *Environmental Research Letters*, 14(12), 123001. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/ab4449)  
702 [9326/ab4449](https://doi.org/10.1088/1748-9326/ab4449)

703 Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V., & Makowski, D. (2021). Positive but variable effects  
704 of crop diversification on biodiversity and ecosystem services. *Global Change Biology*, 27(19),  
705 4697- 4710. <https://doi.org/10.1111/gcb.15747>

706 Bengtsson, J., Ahnström, J., & Weibull, A.-C. (2005). The effects of organic agriculture on biodiversity  
707 and abundance : A meta-analysis. *Journal of Applied Ecology*, 42(2), 261- 269.  
708 <https://doi.org/10.1111/j.1365-2664.2005.01005.x>

709 Bockstaller, C. C., Cariolle, M., Galan, M.-B., Guichard, L., Leclercq, C., Morin, A., & Surleau-  
710 Chambenoit, C. (2013). Evaluation agri-environnementale et choix des indicateurs : Acquis, enjeux  
711 et pistes. *Innovations Agronomiques*, 31, 1.

712 Bouman, O.T., Curtin, D., Campbell, C.A., Biederbeck, V.O. and Ukrainetz, H. (1995), Soil Acidification  
713 from Long-Term Use of Anhydrous Ammonia and Urea. *Soil Science Society of America Journal*, 59:  
714 1488-1494. <https://doi.org/10.2136/sssaj1995.03615995005900050039x>

715 Briones, M., & Schmidt, O. (2017). Conventional tillage decreases the abundance and biomass of  
716 earthworms and alters their community structure in a global meta-analysis. *Global Change*  
717 *Biology*, 23(10), 4396- 4419. <https://doi.org/10.1111/gcb.13744>

718 Burel, F., Garnier, E., Amiaud, B., Aulagnier, S., Butet, A., Chauvel, B., Carré, G., Cortet, J., Couvet, D.,  
719 Joly, P., Lescourret, F., Plantureux, S., Sarthou, J.-P., Steinberg, C., Tichit, M., Vaissière, B.,

720 van Tuinen, D., & Villenave, C. (2008). Les effets de l'agriculture sur la biodiversité. *Agriculture et*  
721 *biodiversité. Valoriser les synergies.*

722 Campbell, B., Beare, D., Bennett, E., Hall-Spencer, J., Ingram, J., Jaramillo, F., Ortiz, R., Ramankutty, N.,  
723 Sayer, J., & Shindell, D. (2017). Agriculture production as a major driver of the Earth system  
724 exceeding planetary boundaries. *Ecology and Society*, 22(4). [https://doi.org/10.5751/ES-09595-](https://doi.org/10.5751/ES-09595-220408)  
725 220408

726 Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M. (2015).  
727 Accelerated modern human-induced species losses : Entering the sixth mass extinction. *Science*  
728 *Advances*, 1(5), e1400253. <https://doi.org/10.1126/sciadv.1400253>

729 Chaplin-Kramer, R., O'Rourke, M. E., Blitzer, E. J., & Kremen, C. (2011). A meta-analysis of crop pest  
730 and natural enemy response to landscape complexity. *Ecology Letters*, 14(9), 922- 932.  
731 <https://doi.org/10.1111/j.1461-0248.2011.01642.x>

732 Chen, C., Chen, H., Chen, X., & Huang, Z. (2019). Meta-analysis shows positive effects of plant  
733 diversity on microbial biomass and respiration. *Nature Communications*, 10.  
734 <https://doi.org/10.1038/s41467-019-09258-y>

735 *Convention on Biological Diversity*. (2016). [https://www.cbd.int/conferences/2016/cop-](https://www.cbd.int/conferences/2016/cop-13/documents)  
736 13/documents

737 Dai, Z., Su, W., Chen, H., Barberan, A., Zhao, H., Yu, M., Yu, L., Brookes, P., Schadt, C., Chang, S., & Xu,  
738 J. (2018). Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of  
739 Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Global Change Biology*,  
740 24(8), 3452- 3461. <https://doi.org/10.1111/gcb.14163>

741 de Graaff, M.-A., Hornslein, N., Throop, H. L., Kardol, P., & van Diepen, L. T. A. (2019). Effects of  
742 agricultural intensification on soil biodiversity and implications for ecosystem functioning : A meta-  
743 analysis. In *Advances in Agronomy* (Vol. 155, p. 1- 44). Elsevier.  
744 <https://doi.org/10.1016/bs.agron.2019.01.001>

745 Díaz, S., Lavorel, S., de Bello, F., Quétier, F., Grigulis, K., & Robson, T. M. (2007). Incorporating plant  
746 functional diversity effects in ecosystem service assessments. *Proceedings of the National*  
747 *Academy of Sciences*, 104(52), 20684- 20689. <https://doi.org/10.1073/pnas.0704716104>

748 Duelli, P., & Obrist, M. (2003). Biodiversity indicators : The choice of values and measures. *Agriculture,*  
749 *Ecosystems & Environment*, 98, 87- 98. [https://doi.org/10.1016/S0167-8809\(03\)00072-0](https://doi.org/10.1016/S0167-8809(03)00072-0)

750 Elhakeem, A., van der Werf, W., Ajal, J., Lucà, D., Claus, S., Vico, R. A., & Bastiaans, L. (2019). Cover  
751 crop mixtures result in a positive net biodiversity effect irrespective of seeding configuration.  
752 *Agriculture, Ecosystems & Environment*, 285, 106627. <https://doi.org/10.1016/j.agee.2019.106627>

753 Ellis, E., Beusen, A., & Klein Goldewijk, K. (2020). Anthropogenic biomes : 10,000 BCE to 2015 CE.  
754 *Land*, 9, 129. <https://doi.org/10.3390/land9050129>

755 Emmerson, M., Morales, M. B., Oñate, J. J., Batáry, P., Berendse, F., Liira, J., Aavik, T., Guerrero, I.,  
756 Bommarco, R., Eggers, S., Pärt, T., Tsharntke, T., Weisser, W., Clement, L., & Bengtsson, J. (2016).  
757 How Agricultural Intensification Affects Biodiversity and Ecosystem Services. In *Advances in*  
758 *Ecological Research* (Vol. 55, p. 43- 97). Elsevier. <https://doi.org/10.1016/bs.aecr.2016.08.005>

759 Estrada-Carmona, N., Sánchez, A. C., Remans, R., & Jones, S. K. (2022). Complex agricultural  
760 landscapes host more biodiversity than simple ones : A global meta-analysis. *Proceedings of the*  
761 *National Academy of Sciences*, 119(38), e2203385119. <https://doi.org/10.1073/pnas.2203385119>

762 Fahrig, L., Brotons, L., Burel, F., Crist, T., Fuller, R., Sirami, C., Siriwardena, G., & Martin, J.-L. (2011).  
763 Functional Landscape Heterogeneity and Animal Biodiversity in Agricultural Landscapes. *Ecology*  
764 *letters*, 14, 101- 112. <https://doi.org/10.1111/j.1461-0248.2010.01559.x>

765 Feest, A., Aldred, T. D., & Jedamzik, K. (2010). Biodiversity quality : A paradigm for biodiversity.  
766 *Ecological Indicators*, 10(6), 1077- 1082. <https://doi.org/10.1016/j.ecolind.2010.04.002>

767 Filazzola, A., Brown, C., Dettlaff, M. A., Batbaatar, A., Grenke, J., Bao, T., Peetoom Heida, I., & Cahill Jr,  
768 J. F. (2020). The effects of livestock grazing on biodiversity are multi-trophic : A meta-analysis.  
769 *Ecology Letters*, 23(8), 1298- 1309. <https://doi.org/10.1111/ele.13527>

770 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T.,  
771 Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C.,

772 Patz, J. A., Prentice, I. C., Ramankutty, N., & Snyder, P. K. (2005). Global Consequences of Land Use.  
773 *Science*, 309(5734), 570- 574. <https://doi.org/10.1126/science.1111772>

774 Geisseler, D., & Scow, K. M. (2014). Long-term effects of inorganic fertilizers on soil microorganisms –  
775 A review. *Soil Biology and Biochemistry*, 75, 54- 63. <https://doi.org/10.1016/j.soilbio.2014.03.023>

776 Gibbs, H. K., Ruesch, A. S., Achard, F., Clayton, M. K., Holmgren, P., Ramankutty, N., & Foley, J. A.  
777 (2010). Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s.  
778 *Proceedings of the National Academy of Sciences*, 107(38), 16732- 16737.  
779 <https://doi.org/10.1073/pnas.0910275107>

780 Gonthier, D. J., Ennis, K. K., Farinas, S., Hsieh, H.-Y., Iverson, A. L., Batáry, P., Rudolphi, J., Tschardtke, T.,  
781 Cardinale, B. J., & Perfecto, I. (2014). Biodiversity conservation in agriculture requires a multi-scale  
782 approach. *Proceedings of the Royal Society B: Biological Sciences*, 281(1791), 20141358.  
783 <https://doi.org/10.1098/rspb.2014.1358>

784 Grout, T. A., Koenig, P. A., Kapuvári, J. K., & McArt, S. H. (2020). *Neonicotinoid Insecticides in New York*  
785 *State*. Cornell University.

786 Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., Stenmans, W., Müller, A.,  
787 Sumser, H., Hören, T., Goulson, D., & Kroon, H. de. (2017). More than 75 percent decline over 27  
788 years in total flying insect biomass in protected areas. *Plos One*, 12(10), e0185809.  
789 <https://doi.org/10.1371/journal.pone.0185809>

790 Hazell, P., & Wood, S. (2008). Drivers of change in global agriculture. *Philosophical Transactions of the*  
791 *Royal Society of London. Series B, Biological Sciences*, 363(1491), 495- 515.  
792 <https://doi.org/10.1098/rstb.2007.2166>

793 Hofman, G., & Cleemput, O. (2004). *Soil and Plant Nitrogen*.

794 Hooper, D. U., Chapin III, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D.  
795 M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., & Wardle, D. A.  
796 (2005). Effects of Biodiversity on Ecosystem Functioning : A Consensus of Current Knowledge.  
797 *Ecological Monographs*, 75(1), 3- 35. <https://doi.org/10.1890/04-0922>

798 Hooper, D. U., & Vitousek, P. M. (1997). The Effects of Plant Composition and Diversity on Ecosystem  
799 Processes. *Science*, 277(5330), 1302- 1305. <https://doi.org/10.1126/science.277.5330.1302>

800 Hu, J., Zhou, S., Tie, L., Liu, X., Liu, X., Zhao, A., Lai, J., Xiao, L., You, C., & Huang, C. (2022). Effects of  
801 nitrogen addition on soil faunal abundance : A global meta-analysis. *Global ecology and*  
802 *biogeography*, 31(8), 1655- 1666. <https://doi.org/10.1111/geb.13528>

803 Humbert, J., Dwyer, J., Andrey, A., & Arlettaz, R. (2016). Impacts of nitrogen addition on plant  
804 biodiversity in mountain grasslands depend on dose, application duration and climate : A  
805 systematic review. *GLOBAL CHANGE BIOLOGY*, 22(1), 110- 120. <https://doi.org/10.1111/gcb.12986>

806 IPBES. (2019). *Summary for policymakers of the global assessment report on biodiversity and*  
807 *ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem*  
808 *Services*.

809 Isbell, F., Adler, P. R., Eisenhauer, N., Fornara, D., Kimmel, K., Kremen, C., Letourneau, D. K., Liebman,  
810 M., Polley, H. W., Quijas, S., & Scherer-Lorenzen, M. (2017). Benefits of increasing plant diversity in  
811 sustainable agroecosystems. *Journal of Ecology*, 105(4), 871- 879. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2745.12789)  
812 [2745.12789](https://doi.org/10.1111/1365-2745.12789)

813 Keenan, R. J., Reams, G. A., Achard, F., de Freitas, J. V., Grainger, A., & Lindquist, E. (2015). Dynamics  
814 of global forest area : Results from the FAO Global Forest Resources Assessment 2015. *Forest*  
815 *Ecology and Management*, 352, 9- 20. <https://doi.org/10.1016/j.foreco.2015.06.014>

816 Kehoe, L., Romero-Muñoz, A., Polaina, E., Estes, L., Kreft, H., & Kuemmerle, T. (2017). Biodiversity at  
817 risk under future cropland expansion and intensification. *Nature Ecology & Evolution*, 1(8), Article  
818 8. <https://doi.org/10.1038/s41559-017-0234-3>

819 Khan, M. N., Mobin, M., Abbas, Z. K., & Alamri, S. A. (2018). Fertilizers and Their Contaminants in  
820 Soils, Surface and Groundwater. In *Encyclopedia of the Anthropocene* (p. 225- 240). Elsevier.  
821 <https://doi.org/10.1016/B978-0-12-809665-9.09888-8>



822 Kim, N., Zabaloy, M. C., Guan, K., & Villamil, M. B. (2020). Do cover crops benefit soil microbiome? A  
823 meta-analysis of current research. *Soil Biology and Biochemistry*, *142*, 107701.  
824 <https://doi.org/10.1016/j.soilbio.2019.107701>

825 Kissling, W. D., Ahumada, J. A., Bowser, A., Fernandez, M., Fernández, N., García, E. A., Guralnick, R. P.,  
826 Isaac, N. J. B., Kelling, S., Los, W., McRae, L., Mihoub, J.-B., Obst, M., Santamaria, M., Skidmore, A.  
827 K., Williams, K. J., Agosti, D., Amariles, D., Arvanitidis, C., ... Hardisty, A. R. (2018). Building essential  
828 biodiversity variables (EBVs) of species distribution and abundance at a global scale. *Biological*  
829 *Reviews*, *93*(1), 600- 625. <https://doi.org/10.1111/brv.12359>

830 Köninger, J., Lugato, E., Panagos, P., Kochupillai, M., Orgiazzi, A., & Briones, M. J. I. (2021). Manure  
831 management and soil biodiversity : Towards more sustainable food systems in the EU. *Agricultural*  
832 *Systems*, *194*, 103251. <https://doi.org/10.1016/j.agsy.2021.103251>

833 Le Roux, X., Barbault, R., Baudry, J., Burel, F., Doussan, I., Garnier, E., Herzog, F., Lavorel, S., Lifran, R.,  
834 Roger-Estrade, J., Sarthou, J.-P., & Trommetter, M. (2012). *Agriculture et biodiversité*. Editions  
835 Quae (Matières à débattre et décider). <https://hal.archives-ouvertes.fr/hal-01173714>

836 Leenhardt, S., Mamy, L., Pesce, S., & Sanchez, W. (2023). *Impacts des produits phytopharmaceutiques*  
837 *sur la biodiversité et les services écosystémiques*. éditions Quae. [https://doi.org/10.35690/978-2-](https://doi.org/10.35690/978-2-7592-3657-2)  
838 [7592-3657-2](https://doi.org/10.35690/978-2-7592-3657-2)

839 Li, Y., Song, D., Liang, S., Dang, P., Qin, X., Liao, Y., & Siddique, K. (2020). Effect of no-tillage on soil  
840 bacterial and fungal community diversity : A meta-analysis. *Soil & tillage research*, *204*.  
841 <https://doi.org/10.1016/j.still.2020.104721>

842 Lichtenberg, E. M., Kennedy, C. M., Kremen, C., Batáry, P., Berendse, F., Bommarco, R., Bosque-Pérez,  
843 N. A., Carvalheiro, L. G., Snyder, W. E., Williams, N. M., Winfree, R., Klatt, B. K., Åström, S.,  
844 Benjamin, F., Brittain, C., Chaplin-Kramer, R., Clough, Y., Danforth, B., Diekötter, T., ... Crowder, D.  
845 W. (2017). A global synthesis of the effects of diversified farming systems on arthropod diversity  
846 within fields and across agricultural landscapes. *Global Change Biology*, *23*(11), 4946- 4957.  
847 <https://doi.org/10.1111/gcb.13714>

848 Liu, T., Chen, X., Hu, F., Ran, W., Shen, Q., Li, H., & Whalen, J. K. (2016). Carbon-rich organic fertilizers  
849 to increase soil biodiversity : Evidence from a meta-analysis of nematode communities.  
850 *Agriculture, Ecosystems & Environment*, *232*, 199- 207.  
851 <https://doi.org/10.1016/j.agee.2016.07.015>

852 Liu, X., Song, L., HE, C., & ZHANG, F. (2010). Nitrogen deposition as an important nutrient from the  
853 environment and its impact on ecosystems in China : Nitrogen deposition as an important nutrient  
854 from the environment and its impact on ecosystems in China. *Journal of Arid Land*, *2*, 137- 143.  
855 <https://doi.org/10.3724/SP.J.1227.2010.00137>

856 Liu, Y., Villalba, G., Ayres, R. U., & Schroder, H. (2008). Global Phosphorus Flows and Environmental  
857 Impacts from a Consumption Perspective. *Journal of Industrial Ecology*, *12*(2), 229- 247.  
858 <https://doi.org/10.1111/j.1530-9290.2008.00025.x>

859 Lori, M., Symnaczik, S., Mader, P., De Deyn, G., & Gattinger, A. (2017). Organic farming enhances soil  
860 microbial abundance and activity-A meta-analysis and meta-regression. *Plos one*, *12*(7).  
861 <https://doi.org/10.1371/journal.pone.0180442>

862 Meynard, J.-M., Messéan, A., Charlier, A., Charrier, F., Fares, M., Bail, M. L., Magrini, M.-B., & Savini, I.  
863 (2013). Freins et leviers à la diversification des cultures : Étude au niveau des exploitations  
864 agricoles et des filières. *OCL*, *20*(4), Article 4. <https://doi.org/10.1051/ocl/2013007>

865 Miles, M., Alix, A., Becker, R., Coulson, M., Dinter, A., Oger, L., Pilling, E., Sharples, A., & Weyman, G.  
866 (2018, juillet 31). *Improving pesticide regulation by use of impact analyses : A case study for bees*.  
867 <https://doi.org/10.5073/jka.2018.462.021>

868 Montazeaud, G., Flutre, T., Ballini, E., Morel, J.-B., David, J., Girodolle, J., Rocher, A., Ducasse, A.,  
869 Violle, C., Fort, F., & Fréville, H. (2022). From cultivar mixtures to allelic mixtures : Opposite effects  
870 of allelic richness between genotypes and genotype richness in wheat. *New Phytologist*, *233*(6),  
871 2573- 2584. <https://doi.org/10.1111/nph.17915>

872 Moos, J., Schrader, S., & Paulsen, H. (2017). Reduced tillage enhances earthworm abundance and  
873 biomass in organic farming : A meta-analysis. *Landbauforschung - Journal of Sustainable and*  
874 *Organic Agriculture*, 67(3- 4), 123- 128. <https://doi.org/10.3220/LBF1512114926000>

875 Moreau, J., Monceau, K., Crépin, M., Tochon, F. D., Mondet, C., Fraikin, M., Teixeira, M., &  
876 Bretagnolle, V. (2021). Feeding partridges with organic or conventional grain triggers cascading  
877 effects in life-history traits. *Environmental Pollution*, 278, 116851.  
878 <https://doi.org/10.1016/j.envpol.2021.116851>

879 Morugan-Coronado, A., Perez-Rodriguez, P., Insolía, E., Soto-Gomez, D., Fernandez-Calvino, D., &  
880 Zornoza, R. (2022). The impact of crop diversification, tillage and fertilization type on soil total  
881 microbial, fungal and bacterial abundance : A worldwide meta-analysis of agricultural sites.  
882 *Agriculture, Ecosystems & Environment*, 329. <https://doi.org/10.1016/j.agee.2022.107867>

883 Neamatollahi, E., Jahansuz, M. R., Mazaheri, D., & Bannayan, M. (2013). Intercropping. In E.  
884 Lichtfouse (Éd.), *Sustainable Agriculture Reviews : Volume 12* (p. 119- 142). Springer Netherlands.  
885 [https://doi.org/10.1007/978-94-007-5961-9\\_4](https://doi.org/10.1007/978-94-007-5961-9_4)

886 Noss, R. (1990). Indicators for Monitoring Biodiversity : A Hierarchical Approach. *Conservation*  
887 *Biology*, 4, 355- 364. <https://doi.org/10.1111/j.1523-1739.1990.tb00309.x>

888 Obersteiner, M., Walsh, B., Frank, S., Havlík, P., Cantele, M., Liu, J., Palazzo, A., Herrero, M., Mosnier,  
889 A., Valin, H., Riahi, K., Kraxner, F., Fritz, S., & Vuuren, D. (2016). Assessing the land resource-food  
890 price nexus of the Sustainable Development Goals. *Science Advances*, 2, e1501499- e1501499.  
891 <https://doi.org/10.1126/sciadv.1501499>

892 Ouyang, Y., Evans, S., Friesen, M., & Tiemann, L. (2018). Effect of nitrogen fertilization on the  
893 abundance of nitrogen cycling genes in agricultural soils : A meta-analysis of field studies. *Soil*  
894 *Biology and Biochemistry*, 127, 71- 78. <https://doi.org/10.1016/j.soilbio.2018.08.024>

895 Parnaudeau, V. V., Reau, R. R., & Dubrulle, P. P. (2012). Un outil d'évaluation des fuites d'azote vers  
896 l'environnement à l'échelle du système de culture : Le logiciel Syst'N. *Innovations Agronomiques*.

897 Peeters, A., Maljean, J. F., Biala, K., & Brouckaert, V. (2004). Les indicateurs de biodiversité pour les  
898 prairies : Un outil d'évaluation de la durabilité des systèmes d'élevage. *Fourrages*, 17.

899 Pelosi, C., Bertrand, C., Daniele, G., Coeurdassier, M., Benoit, P., Néliu, S., Lafay, F., Bretagnolle, V.,  
900 Gaba, S., Vulliet, E., & Fritsch, C. (2021). Residues of currently used pesticides in soils and  
901 earthworms : A silent threat? *Agriculture, Ecosystems & Environment*, 305, 107167.  
902 <https://doi.org/10.1016/j.agee.2020.107167>

903 Prevedello, J. A., Almeida-Gomes, M., & Lindenmayer, D. B. (2018). The importance of scattered trees  
904 for biodiversity conservation : A global meta-analysis. *Journal of Applied Ecology*, 55(1), 205- 214.  
905 <https://doi.org/10.1111/1365-2664.12943>

906 Puissant, J., Villenave, C., Chauvin, C., Plassard, C., Blanchart, E., & Trap, J. (2021). Quantification of  
907 the global impact of agricultural practices on soil nematodes : A meta-analysis. *Soil Biology and*  
908 *Biochemistry*, 161, 108383. <https://doi.org/10.1016/j.soilbio.2021.108383>

909 Reddy, P. P. (2017). Crop Rotation. In P. P. Reddy (Éd.), *Agro-ecological Approaches to Pest*  
910 *Management for Sustainable Agriculture* (p. 229- 242). Springer. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-981-10-4325-3_15)  
911 [981-10-4325-3\\_15](https://doi.org/10.1007/978-981-10-4325-3_15)

912 Reid, W., Mooney, H., Cropper, A., Capistrano, D., Carpenter, S., Chopra, K., Dasgupta, P., Dietz, T.,  
913 Duraiappah, A., Hassan, R., Kasperson, R., Leemans, R., May, R., Mcmichael, A., Pingali, P., Samper,  
914 C., Scholes, R., Watson, R., Zakri, A. H., & Zurek, M. (2005). *Millenium Ecosystem Assessment*  
915 *Synthesis Report*.

916 Reidsma, P., Tekelenburg, T., van den Berg, M., & Alkemade, R. (2006). Impacts of land-use change on  
917 biodiversity : An assessment of agricultural biodiversity in the European Union. *Agriculture,*  
918 *Ecosystems & Environment*, 114(1), 86- 102. <https://doi.org/10.1016/j.agee.2005.11.026>

919 Reiss, E. R., & Drinkwater, L. E. (2018). Cultivar mixtures : A meta-analysis of the effect of intraspecific  
920 diversity on crop yield. *Ecological Applications: A Publication of the Ecological Society of America*,  
921 28(1), 62- 77. <https://doi.org/10.1002/eap.1629>

922 Rigal, S., Dakos, V., Alonso, H., Auniņš, A., Benkő, Z., Brotons, L., Chodkiewicz, T., Chylarecki, P.,  
923 de Carli, E., del Moral, J. C., Domşa, C., Escandell, V., Fontaine, B., Foppen, R., Gregory, R., Harris,

924 S., Herrando, S., Husby, M., Ieronymidou, C., ... Devictor, V. (2023). Farmland practices are driving  
925 bird population decline across Europe. *Proceedings of the National Academy of Sciences*, 120(21),  
926 e2216573120. <https://doi.org/10.1073/pnas.2216573120>

927 Sanchez, A., Jones, S., Purvis, A., Estrada-Carmona, N., & De Palma, A. (2022). Landscape complexity  
928 and functional groups moderate the effect of diversified farming on biodiversity : A global meta-  
929 analysis. *AGRICULTURE ECOSYSTEMS & ENVIRONMENT*, 332.  
930 <https://doi.org/10.1016/j.agee.2022.107933>

931 Shiklomanov, I., & Rodda, J. (2004). *World Water Resources at the Beginning of the Twenty-First*  
932 *Century*. 13.

933 Smil, V. (2000). Smil, V. Phosphorus in the environment : Natural flows and human interferences.  
934 *Annu. Rev. Energy Environ.* 25, 53-88. *Annual Review of Energy and the Environment*, 25, 53- 88.  
935 <https://doi.org/10.1146/annurev.energy.25.1.53>

936 Smith, R. G., Gross, K. L., & Robertson, G. P. (2008). Effects of Crop Diversity on Agroecosystem  
937 Function : Crop Yield Response. *Ecosystems*, 11(3), 355- 366. <https://doi.org/10.1007/s10021-008-9124-5>

938

939 Thiele-Bruhn, S., Bloem, J., de Vries, F. T., Kalbitz, K., & Wagg, C. (2012). Linking soil biodiversity and  
940 agricultural soil management. *Current Opinion in Environmental Sustainability*, 4(5), 523- 528.  
941 <https://doi.org/10.1016/j.cosust.2012.06.004>

942 Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability  
943 and intensive production practices. *Nature*, 418(6898), Article 6898.  
944 <https://doi.org/10.1038/nature01014>

945 Torralba, M., Fagerholm, N., Burgess, P., Moreno, G., & Plieninger, T. (2016). Do European agroforestry  
946 systems enhance biodiversity and ecosystem services ? A meta-analysis. *Agriculture, Ecosystems &*  
947 *Environment*, 230, 150- 161. <https://doi.org/10.1016/j.agee.2016.06.002>

948 Tschardtke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I., & Thies, C. (2005). Landscape  
949 perspectives on agricultural intensification and biodiversity – ecosystem service management.  
950 *Ecology Letters*, 8(8), 857- 874. <https://doi.org/10.1111/j.1461-0248.2005.00782.x>

951 Tuck, S. L., Winqvist, C., Mota, F., Ahnström, J., Turnbull, L. A., & Bengtsson, J. (2014). Land-use  
952 intensity and the effects of organic farming on biodiversity : A hierarchical meta-analysis. *Journal*  
953 *of Applied Ecology*, 51(3), 746- 755. <https://doi.org/10.1111/1365-2664.12219>

954 Tucker, N. I. J., & Murphy, T. M. (1997). The effects of ecological rehabilitation on vegetation  
955 recruitment : Some observations from the Wet Tropics of North Queensland. *Forest Ecology and*  
956 *Management*, 99(1), 133- 152. [https://doi.org/10.1016/S0378-1127\(97\)00200-4](https://doi.org/10.1016/S0378-1127(97)00200-4)

957 van den Hoogen, J., Geisen, S., Routh, D., Ferris, H., Traunspurger, W., Wardle, D. A., de Goede, R. G.  
958 M., Adams, B. J., Ahmad, W., Andriuzzi, W. S., Bardgett, R. D., Bonkowski, M., Campos-Herrera, R.,  
959 Cares, J. E., Caruso, T., de Brito Caixeta, L., Chen, X., Costa, S. R., Creamer, R., ... Crowther, T. W.  
960 (2019). Soil nematode abundance and functional group composition at a global scale. *Nature*,  
961 572(7768), Article 7768. <https://doi.org/10.1038/s41586-019-1418-6>

962 Venter, Z. S., Jacobs, K., & Hawkins, H.-J. (2016). The impact of crop rotation on soil microbial  
963 diversity : A meta-analysis. *Pedobiologia*, 59(4), 215- 223.  
964 <https://doi.org/10.1016/j.pedobi.2016.04.001>

965 Verbruggen, E., & Toby Kiers, E. (2010). Evolutionary ecology of mycorrhizal functional diversity in  
966 agricultural systems. *Evolutionary Applications*, 3(5- 6), 547- 560. <https://doi.org/10.1111/j.1752-4571.2010.00145.x>

967

968 Vukicevich, E., Lowery, T., Bowen, P., Úrbez-Torres, J. R., & Hart, M. (2016). Cover crops to increase  
969 soil microbial diversity and mitigate decline in perennial agriculture. A review. *Agronomy for*  
970 *Sustainable Development*, 36(3), 48. <https://doi.org/10.1007/s13593-016-0385-7>

971 Wang, C., & Tang, Y. (2019). A global meta-analyses of the response of multi-taxa diversity to grazing  
972 intensity in grasslands. *Environmental Research Letters*, 14(11), 114003.  
973 <https://doi.org/10.1088/1748-9326/ab4932>

974 Winkler, J., Dvořák, J., Hosa, J., Martínez Barroso, P., & Vavřková, M. D. (2023). Impact of  
975 Conservation Tillage Technologies on the Biological Relevance of Weeds. *Land*, 12(1), Article 1.  
976 <https://doi.org/10.3390/land12010121>  
977 Wollenberg, E., Richards, M., Smith, P., Havlík, P., Obersteiner, M., Tubiello, F. N., Herold, M., Gerber,  
978 P., Carter, S., Reisinger, A., van Vuuren, D. P., Dickie, A., Neufeldt, H., Sander, B. O., Wassmann, R.,  
979 Sommer, R., Amonette, J. E., Falcucci, A., Herrero, M., ... Campbell, B. M. (2016). Reducing  
980 emissions from agriculture to meet the 2 °C target. *Global Change Biology*, 22(12), 3859- 3864.  
981 <https://doi.org/10.1111/gcb.13340>  
982 *World Bank Open Data*. (2020). World Bank Open Data. <https://data.worldbank.org>  
983