



HAL
open science

Gambierol, a marine dinoflagellate toxin, potently increases evoked quantal transmitter release and reverses pre- and post-synaptic neuromuscular block at vertebrate junctions

Jordi Molgó, Sébastien Schlumberger, Haruhiko Fuwa, Evelyne Benoit, Denis Servent

► To cite this version:

Jordi Molgó, Sébastien Schlumberger, Haruhiko Fuwa, Evelyne Benoit, Denis Servent. Gambierol, a marine dinoflagellate toxin, potently increases evoked quantal transmitter release and reverses pre- and post-synaptic neuromuscular block at vertebrate junctions. 20th World Congress of the International Society on Toxinology - Toxinology in the 21st century: Public health impact from basic, translational and clinical sciences, International Society on Toxinology, Sep 2019, Buenos Aires, Argentina. hal-04465977

HAL Id: hal-04465977

<https://hal.science/hal-04465977v1>

Submitted on 19 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

5. Gambierol, a marine dinoflagellate toxin, potently increases evoked quantal transmitter release and reverses pre- and post-synaptic neuromuscular block at vertebrate junctions

Jordi Molgó, Sébastien Schlumberger, Haruhiko Fuwa, Evelyne Benoit, Denis Servent

CEA, Institut des sciences du vivant Frédéric Joliot, Service d'Ingénierie Moléculaire des Protéines, Université Paris-Saclay, bâtiment 152, 91191 Gif sur Yvette, France

Institut des Neurosciences Paris-Saclay, UMR 9197 CNRS / Université Paris-Sud, CNRS, Gif sur Yvette, France

Department of Applied Chemistry, Faculty of Science and Engineering, Chuo University 1-13-27 Kasuga, Bunkyo, Tokyo 112-8551, Japan

CEA, Institut des sciences du vivant Frédéric Joliot, Service d'Ingénierie Moléculaire des Protéines, Université Paris-Saclay, bâtiment 152, 91191 Gif sur Yvette, France

Gambierol is a marine polycyclic ether toxin that was first isolated together with ciguatoxins from cultured *Gambierdiscus toxicus* dinoflagellates collected in French Polynesia. The chemical synthesis of gambierol permitted the analyses of its mode of action, which includes the selective inhibition of voltage-gated K⁺ (KV) channels in various cells and tissues expressing such channels. In the present study, we investigated the action of synthetic gambierol at vertebrate skeletal neuromuscular junctions using conventional techniques. Nanomolar concentrations of gambierol inhibited the fast K⁺ current and prolonged the duration of the presynaptic action potential in motor nerve terminals, as revealed by presynaptic focal current recordings, and increased stimulus-evoked quantal transmitter release in neuromuscular junctions blocked either by a high Mg²⁺-low Ca²⁺ medium, or by botulinum neurotoxin type-A. Also, gambierol reversed the postsynaptic block produced by d-tubocurarine. In motor nerve terminals loaded with fluo-3/AM, gambierol increased the transient Ca²⁺-signals in response to nerve-stimulation at 1-10 Hz. The results suggest that gambierol, which on equimolar basis is more potent than 3,4-diaminopyridine, can have potential application in pathologies in which it is necessary to antagonize pre- or post-synaptic neuromuscular block, or both.