



HAL
open science

The modulation of vesicular volume and its effects on neurotransmitter secretion through the fusion pore in PC12 Cells

Leslie A Sombers, H. Jacob Hanchar, Tom L. Colliver, Nathan J Wittenberg, Ann-Sofie Cans, Stéphane Arbault, Christian Amatore, Andrew G Ewing

► **To cite this version:**

Leslie A Sombers, H. Jacob Hanchar, Tom L. Colliver, Nathan J Wittenberg, Ann-Sofie Cans, et al.. The modulation of vesicular volume and its effects on neurotransmitter secretion through the fusion pore in PC12 Cells. 12th International Symposium on Chromaffin Cell Biology (ISCCB-12), Sep 2003, La Palma, Spain. hal-04467837

HAL Id: hal-04467837

<https://hal.science/hal-04467837>

Submitted on 21 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The modulation of vesicular volume and its effects on neurotransmitter secretion through the fusion pore in PC12 Cells.

Leslie A. Sombers^a, H. Jacob Hanchar^a, Tom L. Colliver^{a,b}, Nathan J. Wittenberg^a, Ann-Sofie Cans^c, Stéphane Arbault^d, Christian Amatore^d and Andrew G. Ewing^{a,b}

a. Department of Chemistry, The Pennsylvania State University, USA.

b. Department of Neuroscience and Anatomy, The Pennsylvania State University, USA.

c. Department of Chemistry, Göteborg University, Sweden.

d. Ecole Normale Supérieure. Département de Chimie. UMR CNRS-ENSUPMC 8640. Paris. France.

Understanding vesicular exocytosis is fundamental to developing insight into the biology, chemistry and physics of neurotransmission. Through the direct 'presynaptic' observation of quantal release using amperometry it becomes strikingly evident that the exocytotic release event is regulated by multiple mechanisms¹. For instance, many transients in amperometric records of exocytosis events exhibit a pre-spike feature, or foot, which represents a steady-state flux of neurotransmitter through a stable fusion pore connecting the vesicle lumen to the extracellular fluid². This flux is thought to be driven by diffusion along a gradient of chemical potential. Recent work in our laboratory indicates that vesicular volume prior to secretion is strongly correlated with the characteristics of amperometric foot events, and furthermore, suggests that the protein dense core in many vesicles acts to pressurize the inner solution and drive catecholamine through the fusion pore³.

RESULTS AND DISCUSSION

In order to alter vesicular volume, we have used pharmacological manipulations that directly affect the VMAT-mediated transport of catecholamines into PC12 cell vesicles. Cells were incubated in 100 μ M L-3,4-dihydroxyphenylalanine or 100 nM reserpine for 90 minutes in order to increase and decrease, respectively, the volume of single pheochromocytoma (PC12) cell vesicles⁴. Amperometry experiments utilizing a same-cell paradigm were used in conjunction with transmission electron microscopy to determine that as vesicle size is decreased the frequency with which foot events are observed increases. Consistent with this, fewer events with feet are observed as vesicle size is increased. Additionally, both the amount and duration of neurotransmitter released in the foot portion of the event decrease as vesicle size decreases, whereas both of these characteristics increase as vesicle size increases. Finally, smaller vesicles release a greater percentage of their total contents in the foot portion of the event. These trends provide evidence for a nanotube-like intermediate structure for the exocytotic fusion pore. This is shown in Figure 1B, where a model is presented to describe the effect of vesicle volume on membrane tension and fusion pore dynamics. A brief description of the mechanisms regulating neurotransmitter release via the fusion pore is presented in the figure legend.




Figure 1: A) Pharmacological manipulation of vesicle volume alters the dense core to vesicular halo volume ratio. Treatment with reserpine diminishes the vesicular halo; however the volume of the dense core changes relatively little. In contrast, treatment with L-DOPA swells the vesicular halo, and again, the dense core volume remains relatively constant. **B)** Upon opening of the fusion pore, the dense core expands resulting in enhanced pressure-driven mass transport through the fusion pore and fluidic pressure on the interior vesicle walls. As the dense core to vesicular halo volume ratio is greatest for reserpine treated vesicles, it appears that the intravesicular pressure is greatest for these cells. An increase in intravesicular pressure increases vesicular membrane tension. If the fusion pore is of a nanotube geometry, this will lead to an immediate constriction of the fusion pore, as shown³. Subsequently, the increase in the difference in membrane tension from the cell to the vesicle should result in increased lipid flow through the fusion pore, decreasing its overall stability and resulting in an abrupt opening of the pore.

These studies are noteworthy because the dynamics of the exocytotic fusion pore have recently been proposed to play a role in synaptic plasticity⁵. Additionally, physiological⁶⁻⁸, genetic⁸ and pharmacological^{3,10} conditions can affect vesicle size, and in several instances vesicle size has been found to be inversely related to levels of synaptic activity⁹. This work provides new insight into the biological consequences of such vesicular volume modification. It 46 *Cell Biology of the Chromaffin Cell* provides evidence that cells may modulate the rate of neurotransmitter release, or the duration of that release, in the synaptic cleft by manipulating vesicle volume. This appears to be true in the case of release via the fusion pore that is formed in the early stages of exocytosis from PC12 cells.

REFERENCES

1. Sulzer, D. and E.N. Pothos, *Regulation of quantal size by presynaptic mechanisms*. Rev Neurosci, 2000. **11**:159-212.
2. Chow, R.H., et al., *Delay in vesicle fusion revealed by electrochemical monitoring of single secretory events in adrenal chromaffin cells*. Nature, 1992. **356**:60-63.
3. Sombers, L.A., et al., *The effects of vesicular volume on secretion through the fusion pore in exocytotic release from PC12 cells*. J Neurosci, 2004. **24**:303-309.
4. Colliver, T.L., et al., *VMAT-mediated changes in quantal size and vesicular*

volume. J Neurosci, 2000. **20**:5276-5282.

5. Choi, S., et al., *Postfusional regulation of cleft glutamate concentration during LTP at 'silent synapses'*. Nat Neurosci, 2000. **3**:330-336.

6. Zimmermann, H. and V.P. Whittaker, *Effect of electrical stimulation on the yield and composition of synaptic vesicles from the cholinergic synapses of the electric organ of Torpedo: a combined biochemical, electrophysiological and morphological study*. J Neurochem, 1974. **22**:435-450.

7. Maler, L. and W.B. Mathieson, *The effect of nerve activity on the distribution of synaptic vesicles*. Cell Mol Neurobiol, 1985. **5**:373-387.

8. Wickelgren, W.O., et al., *Ultrastructural correlates of transmitter release in presynaptic areas of lamprey reticulospinal axons*. J Neurosci, 1985. **5**:118-201.

9. Karunanithi, S., et al., *Quantal size and variation determined by vesicle size in normal and mutant Drosophila glutamatergic synapses*. J Neurosci, 2002. **22**:10267-10276.

10. Sulzer, D. and R. Edwards, *Vesicles: equal in neurotransmitter concentration but not in volume*. Neuron, 2000. **28**:5-7.