



HAL
open science

Gambierol blocks a K(+) current component affecting action potential duration in fetal adrenomedullary chromaffin cells. Is this action sufficient to modulate catecholamine release?

Jordi Molgó, Roland Bournaud, Sébastien Schlumberger, Makoto Sasaki, Haruhiko Fuwa, Denis Servent, Evelyne Benoit

► To cite this version:

Jordi Molgó, Roland Bournaud, Sébastien Schlumberger, Makoto Sasaki, Haruhiko Fuwa, et al.. Gambierol blocks a K(+) current component affecting action potential duration in fetal adrenomedullary chromaffin cells. Is this action sufficient to modulate catecholamine release?. ICHA 2021 - 19th International conference on Harmful algae online, Oct 2021, La Paz, Mexico. Proceedings of the 19th international conference on harmful algae. hal-04471073v1

HAL Id: hal-04471073

<https://hal.science/hal-04471073v1>

Submitted on 21 Feb 2024 (v1), last revised 21 Feb 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Gambierol effects on k⁺ currents and catecholamine release in single rat fetal adrenomedullary cultured chromaffin cells

Evelyne Benoit **1**, Sébastien Schlumberger **2**, Jordi Molgó **1***, Makoto Sasaki **3**, Haruhiko Fuwa **4**, Roland Bournaud **2**

1 : Service d'Ingénierie Moléculaire pour la Santé (SIMoS), Université Paris-Saclay, CEA, Département Médicaments et Technologies pour la Santé (DMTS), Institut des Sciences du Vivant Frédéric Joliot, ERL CNRS 9004 , F-91191 Gif-sur-Yvette - France

2 : Laboratoire de neurobiologie cellulaire et moléculaire, CNRS UPR9040, F-91198 Gif-sur-Yvette - France

3 : Graduate School of Life Sciences, Tohoku University, 980-8577 Sendai - Japan

4 : Department of Applied Chemistry, Faculty of Science and Engineering, Chuo University, 112-8551 Tokyo - Japan

* : Auteur correspondant

The purpose of this work was to study the action of the polycyclic ether gambierol on K⁺ currents and catecholamine secretion in single rat fetal adrenomedullary cultured chromaffin (AMC) cells using perforated whole-cell voltage-clamp recordings, and current-clamp and electrochemical recordings. From the several types of voltage-gated K⁺ channels (KV) contributing to the total outward current of rat fetal AMC cells, gambierol only partly inhibited the total K⁺ current, when added after or before K_{Ca} and K_{ATP} blockers, and affected neither K_{Ca} nor K_{ATP} channels. After blocking of Nav and K_{ATP} channels, and preventing the activation of K_{Ca} channels, gambierol blocked 50% of the maximal K⁺ current fraction with an inhibitory concentration (IC₅₀) of 5.8 ± 1.56 nM (n = 9). In marked contrast to ciguatoxins, gambierol slowed the kinetics of K⁺-current activation by $75.4 \pm 10.1\%$ (n = 4) with respect to controls (p = 0.031). Hence, before and after gambierol the activation time constants of K⁺ current were 3.82 ± 0.39 ms (n = 4) and 6.80 ± 1.02 ms (n = 4), respectively. Simultaneous current-clamp and single-cell amperometry recordings revealed that gambierol did not modify the membrane potential following 11-seconds depolarizing current-steps, in both quiescent and active cells displaying repetitive firing of action potentials, and it did not increase the number of exocytotic catecholamine release events, with respect to controls. The subsequent block of K_{Ca} channels, both depolarized the membrane and enhanced by 2.7 and 3.5-fold the exocytotic event frequency in quiescent and active cells respectively, highlighting the key modulatory role played by K_{Ca} channels in the control of exocytosis from rat fetal AMC cells

Mots-Clés : gambierol ; potassium currents ; calcium ; activated K⁺ channels ; ATP ; sensitive K⁺ channels ; catecholamine release
