



## Multi-scale evaluation of spider toxins as potential anti-nociceptive agents: example of cyriotoxin-1a

Evelyne Benoit, Michel de Waard, Rémy Béroud, Michel Partiseti, Denis Servent

### ► To cite this version:

Evelyne Benoit, Michel de Waard, Rémy Béroud, Michel Partiseti, Denis Servent. Multi-scale evaluation of spider toxins as potential anti-nociceptive agents: example of cyriotoxin-1a. RT27 – 27th Meeting of the French Society of Toxinology – Toxins: Mr Hyde or Dr Jekyll?, Dec 2021, Paris (Virtual), France. hal-04471737

HAL Id: hal-04471737

<https://hal.science/hal-04471737>

Submitted on 21 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Multi-scale evaluation of spider toxins as potential anti-nociceptive agents: Example of Cyriotoxin-1a

Evelyne Benoit<sup>1</sup>, Michel De Waard<sup>2</sup>, Rémy Béroud<sup>3</sup>, Michel Partiset<sup>4</sup>, and Denis Servent<sup>1</sup>



<sup>1</sup> Service d'Ingénierie Moléculaire pour la Santé, ERL CNRS/CEA 9004, Gif-sur-Yvette, France

<sup>2</sup> Institut du Thorax, Inserm UMR 1087 / CNRS UMR 6291, Nantes, France

<sup>3</sup> Smartox Biotechnology, Saint-Egrève, France

<sup>4</sup> Sanofi R & D, Integrated Drug Discovery – High Content Biology, Vitry-sur-Seine, France



*Na<sub>v</sub>1.7 channel subtype is highly expressed in dorsal root ganglia (DRG) and plays a key role in pain signaling. We performed high-throughput screening of an original animal venom library, using automated patch-clamp experiments on human Na<sub>v</sub> channel subtypes, to find new selective inhibitors of this target ①, and then in vitro and in vivo electrophysiological experiments to characterize the active peptides that have been purified, sequenced and chemically synthesized ②. We identified a novel peptide, Cyriotoxin-1a (CyrTx-1a), isolated from Cyriopagopus schoedtei spider venom ①. This 33 amino acids toxin inhibits hNa<sub>v</sub>1.1-1.3 and 1.6-1.7 in the low nanomolar range, compared to the micromolar range for hNa<sub>v</sub>1.4-1.5 and 1.8. CyrTx-1a is 920 times more efficient at inhibiting tetrodotoxin (TTX)-sensitive than TTX-resistant Na currents of adult mouse DRG neurons in vitro, and approximately 170 times less efficient than huwentoxin-IV at altering mouse skeletal neuromuscular excitability in vivo. Finally, CyrTx-1a exhibits analgesic effects in mice by significantly increasing reaction time in the hot-plate assay ②. These results pave the way for further engineering studies aimed to optimize the potential anti-nociceptive properties of CyrTx-1a.*

