

Pedo-sedimentary dynamics of the Grenzbank zone in the Sangiran dome (Java central, Indonesia): the witness of a complex transitional period in the hominid bearing series

Boris Brasseur, François Semah, Anne-Marie Sémah, Tony Djubiantono

▶ To cite this version:

Boris Brasseur, François Semah, Anne-Marie Sémah, Tony Djubiantono. Pedo-sedimentary dynamics of the Grenzbank zone in the Sangiran dome (Java central, Indonesia): the witness of a complex transitional period in the hominid bearing series. 16th Annual Meeting of EAA, European Association of Archaeologists, 2010, The Hague, Netherlands. hal-04473282

HAL Id: hal-04473282

https://hal.science/hal-04473282

Submitted on 22 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pedo-sedimentary dynamics of the *Grenzbank* zone in the Sangiran dome (Java central, Indonesia):

the witness of a complex transitional period in the hominid bearing series

Boris BRASSEUR¹, François SEMAH², Anne-Marie SEMAH³, Tony DJUBIANTONO⁴

- **1-** Dépt. de Préhistoire du Muséum national d'Histoire naturelle CNRS UMR 7194, Institut de Paléontologie Humaine, 1 rue René Panhard, 75013 PARIS ; brasseur@mnhn.fr
- **2-** Dépt. de Préhistoire du Muséum national d'Histoire naturelle CNRS UMR 7194, Institut de Paléontologie Humaine, 1 rue René Panhard, 75013 PARIS ; semahf@mnhn.fr
- **3-** .I.R.D. LOCEAN, Paleoproxus UMR 7159, 32 avenue Henri Varagnat, 93143 Bondy Cedex, France; seemah@bondy.ird.fr.
- **4-** Pusat Penelitian Arkeologi Nasional, Jl. Raya Condet Pejaten 4, Pasar Minggu, JAKARTA Selatan, Indonesia ; tonydjubiantono@yahoo.com

Abstract

The thick sedimentary series of the Sangiran dome are known to reflect the geological evolution of Central Java since the beginning of the Quaternary until the Lower to Middle Pleistocene boundary. The site is furthermore famous for numerous discoveries of *Homo erectus*, the oldest ones dated at c. 1.5 M.a.

An orogenic phase took place during the late Lower Pleistocene, resulting into the shaping of hill ranges in the Northern and Southern borders of the Solo basin, from which emerge the Sangiran boutonnière. It resulted into synorogenic deposits which abruptly interrupt the clayey sedimentation in the depression. These often conglomeratic deposits are cemented by a calcareous matrix (either contemporaneous of the sedimentation or post-depositional in a phreatic context). They are likely to have formed during a quite long time, but have often been used to mark a stratigraphic boundary (*Grenzbank* in German).

This study combines field survey and micromorphological approach in order to precise the pedosedimentary dynamics of the Grenzbank unit. In a palaeogeographical perspective, we sample 25 sections, which yielded 95 micromorphological samples.

We define three lithostratigraphic microfacies with various geographical distributions. Facies 1 constitutes an independent unit present in the NW part of the dome. Important quantities of mud aggregates and carbonate rhizolithes are also described. It reflects the development of alluvial fans coming from the north during a tectonic episode, associated with strong erosion of pedological covers. Two other facies are defined as the basal part of the overlying Kabuh volcano-alluvial series, and reflect complex changes of river pathways over Central Java. They are more developed in the southern part, and indicate the transport of both siliceous raw material (quartzite, gneiss, chalcedony, jasper, silicified limestone) and small sized artifacts (whose type matches the so called 'Sangiran flakes') that are the oldest discovered in Sangiran.