

## Effect of Rootstock on Some Aspects of Pistachio (Pistacia vera L.): A Review

Aram Akram Mohammed, Fakhraddin Mustafa Hama Salih

### ▶ To cite this version:

Aram Akram Mohammed, Fakhraddin Mustafa Hama Salih. Effect of Rootstock on Some Aspects of Pistachio (Pistacia vera L.): A Review. European Journal of Theoretical and Applied Sciences, 2024, 2 (1), pp.648-659. 10.59324/ejtas.2024.2(1).56 . hal-04473836

### HAL Id: hal-04473836 https://hal.science/hal-04473836

Submitted on 22 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



### Effect of Rootstock on Some Aspects of Pistachio (*Pistacia vera* L.): A Review

Aram Akram Mohammed <sup>™</sup> <sup>™</sup> Horticulture Department, College of Agricultural Engineering Sciences, University of Sulaimani, Sulaimani, Kurdistan Region, Iraq

Fakhraddin Mustafa Hama Salih <sup>10</sup> Horticulture Department, College of Agricultural Engineering Sciences, University of Sulaimani, Sulaimani, Kurdistan Region, Iraq

#### **Suggested Citation**

Mohammed, A. A. & Hama Salih, F. M. (2024). Effect of Rootstock on Some Aspects of Pistachio (Pistacia vera L.): A Review. *European Journal of Theoretical and Applied Sciences, 2*(1), 648-659. DOI: <u>10.59324/ejtas.2024.2(1).56</u>

#### Abstract:

Budding and grafting are the strategies employed to combat unfavorable environmental conditions and improve some physiological defects in the *Pistacia vera* tree. Drought and salinity stresses are the most prominent adverse conditions encountered in pistachio production. It has been observed in different studies that various pistachio rootstocks can be used to ameliorate the effect of those two stresses. Besides, rootstock has a role in some physiological performances of pistachios such as nutrient uptake and photosynthesis. Furthermore, nut blank, unsplit nut, and alternate

are three physiological disorders found in pistachio. Relationships have been found between the degree of these physiological disorders and the rootstock effect. The impact of rootstock on drought and salinity stresses, physiological performances, and physiological disorders in *P. vera* will be discussed in this review.

Keywords: drought, salinity, nutrient uptake, blank nut, alternate bearing, nut splitting.

#### Introduction

Pistachio (*Pistacia vera*) is essentially produced via budding and grafting onto a convenient rootstock, on the one hand, to obtain true-totype seedlings with a determined sex type at the early stages of growth. This is because *P. vera* is a dioecious species, has a long vegetative stage, and there is no reliable marker to distinguish the sex type of the seedlings at the initial growth stages (Bai et al., 2019). Also, rooting of the *P. vera* cuttings is difficult and not vailed on a large scale (Noori et al., 2019). Thus, to generate trueto-type trees, budding and grafting are applied (Moriana et al., 2018). On the other hand, to combat unfavorable environmental conditions that may obstacle the production of P. vera, budding and grafting is used onto a resistible rootstock. However, P. vera is highly cultivated in arid and semi-arid areas, but the resistance of the different cultivars and genotypes to adverse conditions is variable (Hakimnejad et al., 2019). In contrast, there are certain cultivars of P. vera can withstand some of the tough conditions and can be used as a rootstock for the susceptible cultivars (Sheikhi et al., 2019). Besides, some other species in the Pistacia genus compatible with P. vera can be used as rootstock which have different degrees of resistance to adverse circumstances (Rahemi and Tavallali, 2007). Selection of a certain rootstock is a key component of P. vera cultivation. Various

This work is licensed under a Creative Commons Attribution 4.0 International License. The license permits unrestricted use, distribution, and reproduction in any medium, on the condition that users give exact credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if they made any changes.



features of a tree's growth, productivity, and adaptability to different conditions are affected by the choosing of rootstock (Mir et al., 2023). This selection inevitably relies on the nature of the problem P. vera cultivation has, or the site and the country production. Several P. vera cultivars are employed as rootstock in Iran, such as 'Badami', 'Sarakhs', and 'Qazvini', and in Turkey P. vera is also used in most cases (Acar et al., 2017). The reasons that force the growers to use P. vera as a rootstock are the survival rate after transplanting is high due to lateral roots that are fully grown, and the seedlings of this species reach budding and grafting size in a shorter period. Whereas, in the USA, more specifically in California, P. vera has been completely replaced by other rootstocks in the last two decades due to susceptibility to Verticillium, thus P. integerrima and 'UCB-I' rootstocks have frequently been used for their tolerance to Verticillium (Kaska et al., 2002).

Growth and productivity of pistachio and other fruit trees are influenced by rootstock (Font et al., 2020). The relationship between scion and rootstock may reflect on variable responses originating from physiological and genetic factors (Warschefsky et al., 2016). To improve adaptation and productivity, the best tool is the diversification of cultivars and rootstocks. The main focus of recent studies is on the evaluation of pistachio rootstocks and scion-rootstock behavior from physiological and agronomic perspectives and their consequences on agrounder food characteristics, distinct environments (Noguera-Artiaga et al., 2020a). According to the interaction of scion-rootstock revealed in experiments, the responses were not consistent and occasionally contradicting (Ouni et al., 2022). This review aims to show the effect of rootstocks on some physiological aspects and adaptation for adverse conditions of P. vera that were obtained in numerous researches.

# Response of *P. vera* Rootstocks to Drought Stress

The growth and productivity of plants are negatively affected by drought stress (Anjum et al., 2011). Comparing with other environmental

factors, drought stress resulted in the maximum obstruction (Shao et al., 2009). P. vera tolerates drought at a high level and can be cultivated under rainfed in several regions (Tadros et al., 2021). However, irrigation imparts better nut quality, a high shell splitting rate, and minimizes alternate bearing (Goldhamer and Beede, 2004). Managing good production of *P. vera* in drylands at all stages is an important issue (Fereres et al., 2003). Using efficient rootstock is one of the strategies to reduce the undesirable consequences of water deficit (Opazo et al., 2020). There are expectations that tolerable rootstocks contribute to the growth and yield of scion during water scarcity (Serra et al., 2014). Grafted P. vera onto Pistacia atlantica, Pistacia integerrima, and Pistacia terebinthus rootstocks were three irrigation subjected to treatments (regulated deficit irrigation (RDI) at stage two (shell hardening), i) to avoid water stress the grafted trees were regularly irrigated. ii) suppression irrigation until stem water potential (SWP) reached below -1.3 MPa, after that irrigated to maintain SWP below but near this threshold. iii) Irrigation to keep SWP at -1.5 MPa. It was observed that P. atlantica resulted in better yield and quality in comparison with the other rootstocks (Carbonell-Barrachina et al., 2015). Moreover, budded P. vera cv. Kerman onto P. terebinthus, P. atlantica, and P. integerrima were evaluated for their resistance to RDI in shallow soils at stage two by midday stem water potential  $(\psi_{stem})$ . The recommended stress was determined around -1.5 MPa, but at -2.0 MPa will be detrimental to pistachio production in the long term. P. atlantica followed by P. terebinthus were better than P. integerrima, and the worst physiological status was found with the latter one (Memmi et al., 2016). In a somewhat similar to previous results, Noguera-Artiaga et al. (2020b) exposed budded P. vera onto P. terebinthus, P. atlantica and P. integerrima rootstocks to RDI at SWP -1.5 MPa and <-2.0 MPa at stage two. In conclusion, they recommended that P. terebinthus at SWP -1.5 MPa was the best in terms of nut composition, functionality, and sensory quality. Apart from the RDI, Gijón et al. (2010) withdrew irrigation from 1-year-old 'Kerman' seedlings grafted on P. terebinthus L., P. atlantica, and a hybrid of P. atlantica×P. vera on



"day of the year" (DOY) 204 till 218 in pots. The rootstocks had an apparent role in the response of the Kerman cultivar to the water stress. A higher degree of stomatal control and low leaf senescence were observed in the seedlings grafted onto the hybrid. P. terebinthus encouraged powerful stomatal control and rapid leaf senescence. P. atlantica had similar levels of water stress as the two other rootstocks. In another pot experiment, 'Kerman' scions were budded onto 'UCB-I', P. atlantica, and P. terebinthus rootstocks and subjected to water stress for 28 days outdoor, in the second year after budding. The data showed that the three rootstocks gave rise to osmotic adjustment. Furthermore, the parameters confirmed that P. atlantica was more tolerant to drought, which could be due to greater activity of the roots. Whereas, the vegetative growth data implied 'UCB-I' was resistant more than P. atlantica. This may be because one of the parents of 'UCB-I' is P. atlantica, thus having an absorption capacity greater than P. atlantica (Moriana et al., 2018). Among the rootstocks belonging to P. vera species, 'Qazvini' had a better tolerance to drought than 'Badami' (Bagheri et al., 2012).

### Adaptation of *P. vera* to Salinity Through the Application of Rootstocks

Salinity is another issue that will face pistachio production worldwide. The majority of P. vera all over the world is cultivated in saline soils, or low quality and saline water is used for irrigation (Dehghani et al., 2023). Salinity causes deleterious effects on plants by inducing osmotic and specific-ion toxicity (Abbaspour et al., 2012). The worst outcomes that could occur in plants because of salinity are growth reduction, nutritional deficiencies or imbalances, toxicity symptoms, and reduction in the quality and quantity of fruit (Zörb et al., 2019). In the response of plants to soil salinity, rootstock has a crucial role because the first plant organ facing salinity is the root. Therefore, having precise knowledge about scion/rootstock interaction is a key strategy to avoid the injurious effects of salinity (Ahmad and Anjum, 2020). In many fruit

tree species, sensitive scions to salinity are grafted onto the salt-tolerant rootstocks to combat salinity (Yin et al., 2010). In many different studies, the ability of *P. vera* rootstocks has been investigated to reveal the ability to withstand salinity stress. In this context, salinity stress of combined SO42-, Cl-, and boron (B) at 3.5, 8.7,12, or 16 dS $\cdot$ m<sup>-1</sup> electrical conductivity (ECiw) was evaluated in P. vera cv. Kerman grafted onto P. atlantica, P. integerrima, and 'UCB-I' rootstocks. The salinity treatments were added The performance through irrigation. of 'Kerman' pistachio was significantly better onto P. atlantica and 'UCB-I' rootstocks than on P. *integerrima* at 16 dS·m<sup>-1</sup> (Ferguson et al., 2002). Besides, 'Qazvini', 'Sarakhs', and 'Badamin-ezarand' rootstocks which belong to P. vera species were subjected to salinity stress after treatment with 0, 75, 150, or 225 mM NaCl in 8-L polyethylene pots via irrigation every 3 days. The data were calculated at 0, 30, and 60 day intervals. The most salt-tolerant rootstock was 'Qazvini' in comparison to 'Sarakhs' and 'Badami-e-zarand' rootstocks (Hokmabadi et al., 2005). In another similar study, Karimi et al. (2009) added 0, 800, 1600, and 3200 mg NaCl kg <sup>1</sup> soil to the pots of 30-day-old 'Qazvini' and 'Badami' seedling rootstocks. The seedlings were irrigated with distilled water at 7-day interval after adding the NaCl amounts each time. The analysed data affirmed that 'Qazvini' was more resistant to salinity than 'Badami'. Panahi (2009) applied iso-osmotic agents (150 mM NaCl and 20% PEG-6000 solutions) by means of irrigation to P. atlantica subsp. mutica, P. vera 'Sarakhs', and P. vera 'Badami' rootstocks at 3-month-old. Isoosmotic treatments caused 2-fold ABA in P. atlantica subsp. mutica, and P. vera 'Badami', but it was 1.5-fold in P. vera 'Sarakhs'. In addition, 'Badami-e-Zarand A', 'Badami-e-Zarand B', 'Qazvini', and 'Sarakhs' rootstocks were irrigated with saline water at 0.75, 5, 10, and 15 ds.m<sup>-1</sup>. The assessment of salinity tolerance established that 'Badami-e-Zarand B' could be used as a salttolerant rootstock, and 'Sarakhs' was a sensitive rootstock (Karimi et al., 2011). Among the subspecies of P. atlantica, P. atlantica (standard) as resistant, and P. atlantica subsp. Kurdica as sensitive to salinity was determined under 0.75,



5, 10, and 15 dS.m<sup>-1</sup> (Karimi, et al., 2012). In their study, Karimi et al. (2014) selected the hybrid rootstock ('Badami-Riz-e-Zarand'-Female×P.atlantica-male) as the best rootstock for salinity over 'Qazvini' and 'Badami-Riz-e-Zarand' rootstocks under 0, 60, and 120 mM  $NaCl+CaCl_2+MgCl_2$  (3:2:1) for 45 days. Furthermore, P. vera and P. atlantica were employed to adapt Mateur cultivar to salinity. The grafted trees were irrigated with three levels of saline water (tap water (EC<sub>w</sub>:1.95 dSm<sup>-1</sup>), moderately saline water (EC<sub>w</sub>: 5 dSm<sup>-1</sup>), and highly saline water (EC<sub>w</sub>:  $12 \text{ dSm}^{-1}$ ), three times a month for three years. The findings explained that P. atlantica rootstock was an outstanding option to avoid P. vera cultivars from saline stress. This may be because P. atlantica capacity for nutrient uptake, control, and accumulation of salt ions (Na<sup>+</sup> and Cl<sup>-</sup>), selectivity for K<sup>+</sup>-Na<sup>+</sup> and Ca<sup>2+</sup>-Na<sup>+</sup>, and resuming normal growth and fruiting after relieving from saline stress are high (Mehdi-Tounsi et al., 2017). Six rootstocks for P. vera ('Badami', 'Akbari', 'Italyayi', 'Ahmad-Aghaee', 'Qazvini', and 'UCB-I') were tested for their resistance to salinity stress. The salinity treatment was NaCl at 0.5, 12, and 18 dSm<sup>-1</sup>. 'UCB-I' and 'Akbari' followed by 'Ahmad-Aghaee' had better performance in the saline conditions (Raoufi et al., 2019). Also, Goharriz et al. (2020) recognized 'UCB-I' and 'Badami' as most salinity- and drought-resistant the rootstocks (35% of field capacity + 250 mM NaCl) condition, 'Qazvini' as moderate, and 'Kalleh-ghuchi' as sensitive. Additionally, the type of salt determines the severity of the stress. Adish et al. (2010) studied the effect of 0, 100, and 200 mM NaCl and CaCl<sub>2</sub> on pistachio rootstock 'Badami' at the seedling stage. A negative salinity effect was observed owing to both NaCl and CaCl<sub>2</sub>, however impact of NaCl was more extreme than CaCl<sub>2</sub>. In spite of these studies, the salinity tolerance of rootstocks for P. vera has been examined in vitro. Chelli-Chaabouni et al. (2010) culture seeds of *P. vera* and *P. atlantica* in vitro in media supplemented with low NaCl at 0 to 80mM for 45 days, or high NaCl at 0, 131, and 158.5mM for *P. vera* and 0, 131, and 240mM for P.atlantica. The lethal concentrations were 158.5 mM for P. vera and 240 mM for P.atlantica.

*P.atlantica* controlled sodium and chloride uptake more efficiently than *P. vera*. In an additional *in vitro* investigation, the multiplied shoots of 'Akbari', 'Ahmad-Aghaee', 'Italyayi', 'Badami', 'Qazvini', and 'UCB-I' pistachios were grown in media containing 0, 60, and 120 mM NaCl. 'Akbari' had the best performance at 120 mM NaCl (Raoufi et al., 2021).

# Nutrient Uptake is Affected by Rootstock in *P. vera*

Plant nutrition is regulated by soil, environment and plant factors. Among them, plant factors associated with variety, age, growth stage, and root structure (Erdal and Nazli, 2019). The researches have been explained that rootstock is another plant factor that plays a role in the uptake of nutrients (Valverdi and Kalcsits, 2021). Mineral nutrition of the grafted plant can be enhanced due to rootstock. For that reason, employing efficient rootstock may ameliorate mild nutrient deficiency and minimize the application of nutrients in some cases (Ibacache and Sierra, 2009). The level of nutrients in scion is variable depending on the type of rootstock, and this variability likely originates from interspecific variation among rootstocks with regard to nutrient absorption and translocation to scion (Vahdati et al., 2021). Caruso et al. (2003) demonstrated that P. atlantica rootstock delivered more N, P, K, Ca, Mg, Fe, Mn, Zn, and Cu to P. vera cv. Bianca scion than P. integerrima and P. terebinthus rootstocks. In P. vera, the type of rootstock governs the nutrient content and the type of the mineral in leaf and kernel of the nuts. Tavallali and Rahemi (2007) analyzed the nutrient amounts in the leaf and kernel of 'Ohadi', 'Kalleh-ghuchi', and Ahamd-aghaii grafted onto P. vera cv. Badami and Sarakhs, and P. atlantica subsp. mutica rootstocks. The highest P, Zn, K, and lower Na and Mg were measured in the leaves of the cultivars grafted onto Beneh rootstock. 'Badami' resulted in higher leaf Ca and lower Zn. Higher leaf Fe and Cu were detected in the grafted trees onto 'Sarakhs'. Besides, kernel nutrient contents (P, K, Mg, Fe, Zn, and Cu) of the P. vera cultivars were higher onto 'Sarakhs' rootstock. Sherafati et al. (2009)



found that the acquisition of potassium and zinc by 'Akbari' scion was the best onto 'Badami' rootstock, but the worst onto Daneshmandi rootstock. The highest amount of iron into 'Barg-seyah' scion was onto 'Kalleh-ghuchi' rootstock. Moreover, the P. vera rootstocks are different in response to symbiotic fungi to improve nutrient uptake. Bagheri et al. (2012) observed that 'Qazvini' rootstock was better at absorbing P and Mn than 'Badami' under Glomus mosseae inoculation. Meanwhile, the same rootstock does not accumulate the same nutrient in different scions at similar concentration. At the beginning of fruit maturation, Surucu et al. (2020) collected leaf samples from many P. vera cultivars grafted onto Pistacia khinjuk rootstock and analyzed for Mg, Ca, N, P, K, Fe, Cu, Mn, and Zn contents. The highest N, P, and K in Haciserifi, Mg, Ca, and Cu in 'Mumtaz', Fe and Zn in 'Vahidi', and Mn in 'Sel-15' were recorded.

### Relationship Between Rootstock and Photosynthesis Performance in *P. vera*

The photosynthetic rate of scion varieties is distinctly influenced by rootstock. Rootstock can control canopy size, leaf morphology, and gas exchange which in turn affect photosynthesis competence (Losciale et al., 2008). Thus, as a crucial criterion for selection, Jover et al. (2012) reported that the effect of rootstock on photosynthesis should be taken into account. Also, Fullana-Pericàs et al. (2020) referred to grafting onto a convenient rootstock can be a beneficial technique to make better photosynthetic performance in plants. In the case of P. vera rootstocks, the photosynthesis rate was assessed in 'Ohadi', 'Kalleh-ghuchi', and 'Ahmad-Aghaee' scions grafted on P. vera cv. Sarakhs and Badami riz, P. atlantica subsp. atlantica and P. atlantica subsp. mutica rootstocks. The trees grown onto P. vera cv. Sarakhs and P. atlantica subsp. atlantica had the maximum photosynthetic rates, in contrast, the rate of photosynthesis was the least in the trees their rootstocks were P. atlantica subsp. mutica and P. vera cv. Badami (Fotouhi et al., 2006). Scionrootstock combinations of P. vera cv. Mateur and

Kerman scions onto *P. vera* and *P. atlantica* rootstocks were evaluated by Ghrab et al. (2014). It was obtained that *P. atlantica* rootstocks provide the maximum photosynthesis and chlorophyll content for the two scions.

### Rootstock to Increase Nut Splitting Rate

One of the desirable characteristics of pistachio nuts is the splitting of the shell (longitudinal dehiscence). It is a much more attractive trait that encourages a variety choosing in the markets (Ouni et al., 2022). During the harvesting weeks, the amount of split nuts increased because of the enlargement of the inner kernel which forces the outer shell to open. However, after harvesting, a large quantity of closed nuts remains due to irrigation scheduling, imbalanced nutrition, fertilizer application, cultivar, salinity, and temperature upon blooming stage (Sukunza et al., 2023). Hence, special treatment should be conducted artificially on unsplit nuts aiming splitting them, such as longitudinal force on the shell suture and thermal processing (Foroutanaliabad and Foroutanaliabad, 2006). Nevertheless, these methods are expensive, require special equipment, and nut quality may be reduced. Regarding nut splitting factors, the rootstock is among the factors that have been reported to impact pistachio nut splitting (Ouni et al., 2018). Early nut splitting was examined by Tajabadipour et al. (2006) in 'Ohadi', 'Kallehghuchi', and 'Ahmad-Aghaee' tree cultivars grafted onto P. vera cv. Badami-e-Riz, P. vera var 'Sarakhs', P. atlantica subsp. mutica, and P. atlantica subsp. atlantica rootstocks. 'Kalleh-ghuchi' onto P. atlantica subsp. atlantica gave the highest percentage of earlier splitting, but the same cultivar delayed nut splitting onto P. vera cv Badami-e-Riz rootstock. The rootstocks did not induce earlier nut splitting in 'Ohadi' and 'Ahmad-Aghaee'. In addition, three P. vera cultivars ('Ohadi', 'Kalleh-ghuchi', and 'Ahmad-Aghaee') were compared to their differences in nut quality including splitting rate onto *P. vera* cv. Sarakhs and Badami, and P. atlantica subsp. mutica rootstocks. The highest splitting percentage was recorded in the P. vera cultivars whose rootstock



was P. vera cv. Badami (Rahemi and Tavallali, 2007). Effects of pistachio rootstocks (P. vera, P. atlantica, and P. khinjuk) on nut splitting of 'Siirt' and 'Ohadi' cultivars were examined. Shell splitting rate was outstanding in both cultivars as produced nuts onto P. atlantica rootstock, but the worst onto P. khinjuk rootstock. At the same time, the 'Siirt' cultivar was better than 'Ohadi' regarding nut splitting (Turker and Ak, 2010). Whereas, there is reports that nut splitting is characteristic of cultivar rather than the effect of rootstock. Parfitt et al. (2005) stated that cultivar and shell splitting are closely linked. Some P. vera cultivars were studied to determine the amount of nut splitting that grew onto P. khinjuk rootstock. It was obtained that Sel-2, Sel-5, and 'Siirt' cultivars had the highest percentage of split nuts related to 'Haciserifi', 'Kerman', Mumtaz, 'Ohadi', 'Sel-1', 'Sel-10', 'Sel-11', 'Sel-14', 'Sel-15', 'Uzun', and 'Vahidi' cultivars (Surucu et al., 2020). Additionally, Caruso et al. (2003) declared that rootstock did not significantly affect nut splitting, when they studied P. vera cv. Bianca onto P. atlantica, P. integerrima, P. terebinthus rootstocks.

## Role of Rootstock in Blank Nut in *P. vera*

A pistachio nut without a kernel is called a blank (empty) nut. This occurs as a result of failure of the embryo to develop, development of ovary tissues without successful fertilization, embryo abortion, tree inability to provide assimilates to complete kernel growth to a sufficient extent, and inadequate irrigation and boron leaf levels (Kashaninejad and Tabil, 2011). The severity of the blank is associated with cultivar, rootstock, and year to year (Hormaza and Wünsch, 2007). There are reports that rootstock lessens blank nut in P. vera. In this respect, P. vera cv. Ohadi, 'Kalleh-ghuchi', and 'Ahmad-Aghaee Ahmad-Aghaee' trees were investigated for blankness rate onto P. vera cv. Sarakhs and Badami, and P. atlantica subsp. mutica. P. vera cv. Badami rootstock produced the minimum of nuts with blankness (Rahemi and Tavallali, 2007). In another experiment, Turker and Ak (2010) showed that 'Siirt' and 'Ohadi' pistachio cultivars

onto *P. vera*, *P. atlantica*, and *P. khinjuk* had various nut filling percentage. The lowest blank nuts were calculated from the two cultivars onto *P. atlantica* rootstock, on the contrary the two cultivars onto *P. khinjuk* exhibited the peak blank nuts. Whereas, not all studies found that rootstock determines the amount of blank nuts in pistachio. Caruso et al. (2003) announced the ineffectiveness of rootstocks (*P. atlantica*, *P. integerrima*, *P. terebinthus*) to change the ratio of blank nut significantly in *P. vera* cv. Bianca.

# Rootstock and Alternate Bearing in *P. vera*

The prominent obstacle in the production of pistachio is alternate bearing. It is a physiological disorder observed in pistachio and other fruit species in which a high yield quantity is obtained in one year (on year), but in the following year (off year) little or no yield may be harvested. Despite nut quantity, this disorder relates to nut quality from blank and non-split nut perspectives. In 'on' year, non-split nuts are profound and blanking is much more common in 'off' years (Kashaninejad and Tabil, 2011). Alternate bearing has been attributed to certain reasons. Competition between the developed inflorescence buds and heavy crop in the summer of 'on' year on carbohydrate may be one of the causes in which the produced nuts deprives the inflorescence buds from reaching sufficient carbohydrate and eventually abscised (Baninasab and Rahemi, 2006). On the other hand, the nutrient status of the tree is an additional factor that may influence the degree of alternate bearing. Kumar et al. (2016) found that the application of NPK and B decreased the flowering and fruit set along with reduction in bud abscission during 'on' and 'off' years. Warm climate and rain-fed conditions accentuate this phenomenon; also it is variable according to cultivar (Ghrab et al., 2014; Carbonell-Barrachina et al., 2015). The rootstocks may impact the intensity of alternating bearing in pistachio (Kaska et al., 2002). Pistachio rootstocks have different ability to uptake nutrient from the soil (Surucu et al., 2020), and as mentioned above nutrient is one of the factors



that influences alternate bearing. Also, it was hypothesized that the vigor of pistachio rootstock mitigated alternate bearing because vigorous rootstock with large leaf surface area can manufacture more carbohydrates (Khezri et al., 2020). Beede et al. (2017) showed that Kerman P. vera cultivar onto 'PGII' and 'UCB-I' rootstocks had lower alternate bearing than onto P. atlantica and 'PGI' rootstocks. Lower bud abscission was detected in 'Kalleh-ghuchi' pistachio cultivars onto P. vera cv. Badami-Riz rootstock in comparison to P. atlantica, P. atlantica subsp. mutica, and P. khinjuk (Khezri, 2010). P. vera and P. atlantica rootstocks were tested for their effect on the functionality of P. vera cv. Mateur and Achouri in 2014 and 2015. Yield quantity indicated 2014 as 'on' year and 2015 as 'Off' year. In 2015, P. atlantica rootstock provided Mateur cultivar with a better yield than Achouri onto P. vera rootstock. In contrast, the impact of the rootstocks was insignificant in 2014 on both cultivars (Ouni et al., 2018). However, Ouni et al. (2022) referred that alternate bearing in P. vera cv. Mateur and Achouri onto P. vera and P. atlantica rootstocks varied depending on the year to year and cultivar but not rootstock, 'Mateur' was better than 'Achouri'.

### Conclusion

P. vera is a species that can tolerate different adverse conditions, but the degree of tolerance is different according to genotypes and cultivars. Rootstock has been observed that effective for adapting this species to unfavorable conditions. Through this review, it was revealed that P. vera 'Badami', 'Qazvini', 'Sarakhs', 'Akbari', and some other cultivars along with other species from the Pistacia genus such as P. atlantica, P. terebinthus, P. integerrima, P. khinjuk, and a hybrid of P. atlantica× P. integerrima ('UCB-I') can be used for different purposes. The studies on drought stress recommended P. atlantica and 'Qazvini' as drought-resistant rootstock. Also, P. atlantica and 'UCB-I' along with 'Qazvini' followed by 'Akbari' had the best performance in saline conditions. Regarding nutrient uptake, each rootstock had the best absorption capacity for specific nutrient elements, but, generally, P.

atlantica, 'Badami', 'Sarakhs' rootstocks absorb the majority of the nutrient elements at a high level. The investigation indicated that rootstocks affected some physiological characteristics and physiological disorders of *P. vera*. In this context, *P. atlantica* and 'Sarakhs' caused the best photosynthesis. Besides. *P. atlantica* and 'Badami' resulted in the lowest blank and alternate bearing but the maximum nut splitting rate. However, some studies do not support the influence of rootstocks on nut splitting, blank nut, and alternate bearing in pistachio. Thus, rigorous investigations are needed to confirm the role of rootstocks in these physiological disorders.

### **Conflict of Interests**

No conflict of interest.

### References

Abbaspour, H., Afshari, H., & ABDEL-WAHHAB, M.A. (2012). Influence of salt stress on growth, pigments, soluble sugars and ion accumulation in three pistachio cultivars. *Journal of Medicinal Plants Research, 6*, 2468-2473. https://doi.org/10.5897/JMPR11.1710

Acar, I., Kafkas, S., Kapchina-Toteva, V. & Ercişli, S. (2017). Effect of rootstock on fat content and fatty acid composition of immature pistachio kernels. *Comptes rendus de l'Acad'emie bulgare des Sciences, 70*(7), 1049-1056.

Adish, A., Fekri, M. & Hokmabadi, H. (2010). Response of 'Badami'-Zarand pistachio rootstock to salinity stress. *International Journal of Nuts Related Sciences*, 1(1), 1-11. <u>https://doi.org/10.22034/JON.2010.515761</u>

Ahmad, R. & Anjum, M. A. (2020). Physiological and molecular basis of salinity tolerance in fruit crops. In *Fruit Crops* (pp. 445-464). Elsevier.

Anjum, S. A., Xie, X., Wang, L. C., Saleem, M. F., Man, C. & Lei, W. (2011). Morphological, physiological and biochemical responses of plants to drought stress. *African journal of agricultural research*, 6(9), 2026-2032. http://dx.doi.org/10.5897/AJAR10.027



Bagheri, V., Shamshiri, M. H., Shirani, H. & Roosta, H. R. (2012). Nutrient uptake and distribution in mycorrhizal pistachio seedlings under drought stress. *Journal of Agricultural Science and Technology*, 14, 1591-1604.

Bai, Q., Ma, Z., Zhang, Y., Su, S. & Leng, P. (2019). The sex expression and sex determining mechanism in *Pistacia* species. *Breeding science*, *69*(2), 205-214. https://doi.org/10.1270%2Fjsbbs.18167

Baninasab, B. & Rahemi, M. (2006). Possible role of non-structural carbohydrates in alternate bearing of pistachio. *European journal of horticultural science*, 71(6), 277–282. https://doi.org/10.3390/su152115300

Beede, R.H., Ferguson, L., Lampinen, B. & Metcaf, S. (2017). Effect of hedging and topping on pistachio alternate bearing: 2012–2017. Retrieved from https://americanpistachios.org/sites/default/fi les/inlinefiles/3.%20Effect%20of%20Hedging%20and%

files/3.%20Effect%20of%20Hedging%20and%20Topping%20on%20Pistachio%20Alternate%20BearingEdited%2012-29-2017.pdf

Carbonell-Barrachina, Á. A., Memmi, H., Noguera-Artiaga, L., Gijón-López, M. D. C., Ciapa, R. & Pérez-López, D. (2015). Quality attributes of pistachio nuts as affected by rootstock and deficit irrigation. *Journal of the Science of Food and Agriculture*, 95(14), 2866-2873. https://doi.org/10.1002/jsfa.7027

Caruso, T., Barone, E., Marra, F. P., Sottile, F., La Mantia, M. & De Pasquale, C. (2003). Effect of rootstock on growth, yield and fruit characteristics in cv. Bianca pistachio (*Pistacia vera* L.) trees. *Options Méditerranéennes, 63*, 117-122.

Chelli-Chaabouni, A., Mosbah, A. B., Maalej, M., Gargouri, K., Gargouri-Bouzid, R. & Drira, N. (2010). In vitro salinity tolerance of two pistachio rootstocks: *Pistacia vera* L. and *P. atlantica* Desf. *Environmental and Experimental Botany*, 69(3), 302-312. <u>http://dx.doi.org/10.1016/j.envexpbot.2010.05</u> .010 Dehghani, F., Rahnemaie, R., Dalir, N., Saadat, S. & Zarebanadkouki, M. (2023). Interactive effect of salinity and Ca to Mg ratio of irrigation water on pistachio growth parameters and its ionic composition in a calcareous soil. *New Zealand Journal of Crop and Horticultural Science*, *51*(3), 432-450. <u>https://doi.org/10.1080/01140671.2021.20079</u> 58

Erdal, İ. & Nazli, İ. (2019). Influence of rootstock and variety on leaf nutrient concentration of pear grown on a nutrientsufficient soil. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi, 22*, 141-147. https://doi.org/10.18016/ksutarimdoga.vi.563 740

Fereres, E., Goldhamer, D. A. & Parsons, L. R. (2003). Irrigation water management of horticultural crops. *HortScience*, *38*(5), 1036-1042.

http://dx.doi.org/10.21273/HORTSCI.38.5.10 36

Ferguson, L. O. U. I. S. E., Poss, J. A., Grattan, S. R., Grieve, C. M., Wang, D. O. N. G., Wilson, C., ..., & Chao, C. T. (2002). Pistachio rootstocks influence scion growth and ion relations under salinity and boron stress. *Journal of the American Society for Horticultural Science*, 127(2), 194-199. http://dx.doi.org/10.21273/JASHS.127.2.194

Font i Forcada, C., Reig, G., Mestre, L., Mignard, P., Betrán, J. Á. & Moreno, M. Á. (2020). Scion× rootstock response on production, mineral composition and fruit quality under heavycalcareous soil and hot climate. *Agronomy*, *10*(8), 1159.

https://doi.org/10.3390/agronomy10081159

Foroutanaliabad, A. & Foroutanaliabad, A. (2006). Methods for splitting pistachio nuts. US patent 7695750B2. Retrieved from https://patents.google.com/patent/US7695750 B2/en

Fotouhi, R., Sajadian, H. & Hokmabadi, H. (2006). Effects of some pistachio rootstocks on leaf photosynthesis rate of three Iranian commercial pistachio cultivars. In *XXVII International Horticultural Congress-IHC2006:* 

655

International Symposium on Enhancing Economic and Environmental 772, 241-244.

Fullana-Pericàs, M., Conesa, M. À., Pérez-Alfocea, F. & Galmés, J. (2020). The influence of grafting on crops' photosynthetic performance. *Plant Science*, 295, 110250. https://doi.org/10.1016/j.plantsci.2019.110250

Ghrab, M., Chelli-Chaabouni, A. & Ben Mimoun, M. (2014). Behaviour of scionrootstock combinations of pistachio under Mediterranean climate. In XXIX International Horticultural Congress on Horticulture: Sustaining Lives, Livelihoods and Landscapes (IHC2014): 1109, 155-160.

Goldhamer, D. & Beede, R. (2004). Regulated deficit irrigation effects on yield, nut quality and water-use efficiency of mature pistachio trees. *The Journal of Horticultural Science and Biotechnology*, 79(4), 538-545. https://doi.org/10.1080/14620316.2004.11511 802

Hakimnejad, S., Karimi, H. R., Sahhafi, S. R. & Esmaeilizadeh, M. (2019). Morphological, ecophysiological and photosynthetic diversity of some *Pistacia* species for use in breeding programs. *Genetic Resources and Crop Evolution*, *66*, 1399-1419. <u>https://doi.org/10.1007/s10722-019-00804-7</u>

Hokmabadi, H., Arzani, K. & Grierson, P. F. (2005). Growth, chemical composition, and carbon isotope discrimination of pistachio (*Pistacia vera* L.) rootstock seedlings in response to salinity. *Australian Journal of Agricultural Research*, 56(2), 135-144.

Hormaza, J. I. & Wünsch, A. (2007). Pistachio. In *Fruits and nuts* (pp. 243-251). Berlin, Heidelberg: Springer Berlin Heidelberg.

Ibacache, A. G. & Sierra, C. B. (2009). Influence of rootstocks on nitrogen, phosphorus and potassium content in petioles of four table grape varieties. *Chilean Journal of Agricultural Research*, 69(4), 503-508. <u>http://dx.doi.org/10.4067/S0718-</u> <u>58392009000400004</u>

Jover, S., Martínez-Alcántara, B., Rodríguez-Gamir, J., Legaz, F., Primo-Millo, E., Forner, J.

& Forner-Giner, M. A. (2012). Influence of rootstocks on photosynthesis in Navel orange leaves: effects on growth, yield, and carbohydrate distribution. *Crop science*, 52(2), 836-848.

https://doi.org/10.2135/cropsci2011.02.0100

Karimi, H. R., Ebadi, A., Zamani, Z. & Fatahi, R. (2011). Effects of water salinity on growth indices and physiological parameters in some pistachio rootstocks. *Acta Horticulrurae*, *912*, 171-178.

https://doi.org/10.17660/ActaHortic.2011.912 .24

Karimi, H. R., Maleki-Kuhbanani, A. & Roosta, H. R. (2014). Evaluation of inter-specific hybrid of *P. atlantica* and *P. vera* L. cv. 'Badami'-Riz-e-Zarand' as Pistachio rootstock to salinity stress according to some growth indices and ecophysiological and biochemical parameters. *Journal of Stress Physiology & Biochemistry*, 10(3), 5-17.

Karimi, H. R., Zamani, Z., Ebadi, A. & Fatahi, R. (2012). Effects of water salinity on growth indices and physiological parameters in some wild pistachio. *International Journal of Nuts and Related Sciences*, 3(1), 41-48. <u>http://dx.doi.org/10.1080/01904167.2011.555</u> 577

Karimi, S., Rahemi, M., Maftoun, M. & Tavallali, V. (2009). Effects of long-term salinity on growth and performance of two pistachio (*Pistacia* L.) rootstocks. *Australian Journal of Basic and Applied Sciences*, 3(3), 1630-1639.

Kashaninejad, M. & Tabil, L. G. (2011). Pistachio (*Pistacia vera* L.). In *Postharvest biology and technology of tropical and subtropical fruits* (pp. 218-247). Woodhead Publishing.

Kaska, N., Nikpeyma, Y. & Kafkas, S. A. L. İ. H. (2002). Interactions between pistachio rootstock and cultivar in K. Maras/Turkey-Preliminary results. *Acta Horticulrurae*, 591,67-71. https://doi.org/10.17660/ActaHortic.2002.591 .6

Khezri, M. (2010). Effect of rootstocks and spray application of nitrogen, boron, zinc and free polyamines on some physiological problems and yield of pistachio (*Pistacia vera* cv. Kaleh-Ghoochi). Ph.D dissertation. University of Tehran, Iran.

Khezri, M., Heerema, R., Brar, G. & Ferguson, L. (2020). Alternate bearing in pistachio (*Pistacia vera* L.): A review. *Trees*, *34*, 855-868. <u>https://doi.org/10.1007/s00468-020-01967-y</u>

Kumar, P., Sharma, S. K., Chandel, R. S., Singh, J. & Kumar, A. (2016). Nutrient dynamics in pistachios (*Pistacia vera* L.): The effect of mode of nutrient supply on agronomic performance and alternate-bearing in dry temperate ecosystem. *Scientia Horticulturae*, 210, 108-121. http://dx.doi.org/10.1016/j.scienta.2016.07.01 8

Losciale, P., Zibordi, M., Manfrini, L. & Grappadelli, L. C. (2008). Effects of rootstock on pear photosynthetic efficiency. *Acta Horticulturae*, *800*, 241-248. https://doi.org/10.17660/ActaHortic.2008.800 .28

Mehdi-Tounsi, H., Chelli-Chaabouni, A., Mahjoub-Boujnah, D. & Boukhris, M. (2017). Long-term field response of pistachio to irrigation water salinity. *Agricultural Water Management*, 185, 1-12. https://doi.org/10.1016/j.agwat.2017.02.003

Memmi, H., Gijón, M. C., Couceiro, J. F. & Pérez-López, D. (2016). Water stress thresholds for regulated deficit irrigation in pistachio trees: Rootstock influence and effects on yield quality. *Agricultural Water Management*, 164, 58-72.

https://doi.org/10.1016/j.agwat.2015.08.006

Mir, M. M., Parveze, M. U., Iqbal, U., Rehman, M. U., Kumar, A., Simnani, S. A., ... & Bhat, M. A. (2023). Development and Selection of Rootstocks. In *Temperate Nuts* (pp. 45-78). Singapore: Springer Nature Singapore.

Moriana, A., Memmi, H., Centeno, A., Martín-Palomo, M. J., Corell, M., Torrecillas, A. & Pérez-López, D. (2018). Influence of rootstock on pistachio (*Pistacia vera* L. cv Kerman) water relations. *Agricultural water management*, 202, 263-270.

https://doi.org/10.1016/j.agwat.2017.12.026

Noguera-Artiaga, L., Sánchez-Bravo, P., Hernández, F., Burgos-Hernández, A., Pérez-López, D. & Carbonell-Barrachina, Á. A. (2020b). Influence of regulated deficit irrigation and rootstock on the functional, nutritional and sensory quality of pistachio nuts. *Scientia Horticulturae*, 261, 108994. https://doi.org/10.1016/j.scienta.2019.108994

Noguera-Artiaga, L., Sánchez-Bravo, P., Pérez-López, D., Szumny, A., Calin-Sánchez, Á., Burgos-Hernández, A. & Carbonell-Barrachina, Á. A. (2020a). Volatile, sensory and functional properties of hydrosos pistachios. *Foods*, 9(2),

Noori, I., Ahmad, F., Aziz, R., & Mohammed, A. (2019). Propagation of pistachio Pistacia vera L. by air layering under the effects of IBA and GA 3 treatments. *Eurasia J Biosci, 13*, 2001-2004.

158. https://doi.org/10.3390/foods9020158

Opazo, I., Toro, G., Salvatierra, A., Pastenes, C. & Pimentel, P. (2020). Rootstocks modulate the physiology and growth responses to water deficit and long-term recovery in grafted stone fruit trees. *Agricultural Water Management*, *228*, 105897. https://doi.org/10.1016/j.agwat.2019.105897

Ouni, S., Chaabouni, A. C., Noguera-Artiaga, L., Hernández, F., Hamda, H. B., Ouerghui, I., ..., & Rhouma, A. (2018). Effects of two rootstocks (*Pistacia vera* L. and *Pistacia atlantica* Desf.) on the yield, morphology, chemical composition, and functional properties of two pistachio varieties ("Mateur" and "Achoury"). *The Journal of Microbiology, Biotechnology and Food Sciences*, 8(2), 853.

https://doi.org/10.15414/JMBFS.2018.8.2.853 -856

Ouni, S., Noguera-Artiaga, L., Carbonell-Barrachina, A., Ouerghui, I., Jendoubi, F., Rhouma, A. & Chelli-Chaabouni, A. (2022). Cultivar and Rootstock Effects on Growth, Yield and Nut Quality of Pistachio under Semi-Arid Conditions of South Mediterranean. *Horticulturae*, 8(7), 606. https://doi.org/10.3390/horticulturae8070606

Panahi, B. (2009). Effects of osmotic and salt stresses on water relation parameters of pistachio seedlings. *Plant Ecophysiology*, *1*, 1–8.

657

Parfitt, D., Kallsen, C., & Maranto, J. (2005). The orchard. Pistachio cultivars. In L. Ferguson, R.H. Beede, & M.W. Freeman, Haviland, D.R., Holtz, B.A., Kallsen, C.E. (Eds.), *Pistachio Production Manual* (4th ed., pp.62–66). University of California Fruit & Nut Research & Information Center: Davis, CA, USA.

Rahemi, M. & Tavallali, V. (2007). Effects of rootstock on Iranian pistachio scion cultivars. *Fruits*, 62(5), 317-323. http://dx.doi.org/10.1051/fruits:2007027

Raoufi, A., Rahemi, M., Salehi, H. & Javanshah, A. (2019). Selecting high performance rootstocks for pistachio cultivars under salinity stress based on their morpho-physiological characteristics. *International Journal of Fruit Science*, 20(2), 29-47. https://doi.org/10.1080/15538362.2019.17016

<u>16</u>

Raoufi, A., Salehi, H., Rahemi, M., Shekafandeh, A. & Khalili, S. (2021). In vitro screening: The best method for salt tolerance selection among pistachio rootstocks. *Journal of the Saudi Society of Agricultural Sciences*, 20(3), 46-154. https://doi.org/10.1016/j.jssas.2020.12.010

Serra, I., Strever, A., Myburgh, P. A. & Deloire, A. (2014). The interaction between rootstocks and cultivars (*Vitis vinifera* L.) to enhance drought tolerance in grapevine. *Australian Journal* of grape and wine Research, 20(1),1-14. https://doi.org/10.1111/ajgw.12054

Shao, H.B., Chu, L.Y., Jaleel, C.A., Manivannan, P., Panneerselvam, R. & Shao, M.A. (2009). Understanding water deficit stress-induced changes in the basic metabolism of higher plants-biotechnologically and sustainably improving agriculture and the ecoenvironment in arid regions of the globe. *Critical reviews in biotechnology*, 29(2), 31-151. https://doi.org/10.1080/07388550902869792

Sheikhi, A., Arab, M. M., Brown, P. J., Ferguson, L. & Akbari, M. (2019). Pistachio (*Pistacia* spp.) breeding. *Advances in Plant Breeding Strategies: Nut* and Beverage Crops, 4, 353-400. http://dx.doi.org/10.1007/978-3-030-23112-5\_10 Sherafati, A., Hokmabadi, H. & Taheri, M. (2009). Effects of some rootstocks on mineral nutrient uptake in two pistachio cultivars (*Pistacia vera* L.). In *V International Symposium on Pistachios and Almonds 912*, 197-201.

Sukunza, X., Bolaños, M., Tellabide, M., Estiati, I., Aguado, R. & Olazar, M. (2023). A novel alternative method for pistachio nuts splitting using conical spouted beds. *Chemical Engineering Science*, 282, 119348.

https://doi.org/10.1016/j.ces.2023.119348

Surucu, A., Acar, I., Demirkiran, A. R., Farooq, S. & Gokmen, V. (2020). Variations in nutrient uptake, yield and nut quality of different pistachio cultivars grafted on *Pistacia khinjuk* rootstock. *Scientia Horticulturae*, *260*, 108913. https://doi.org/10.1016/j.scienta.2019.108913

Tadros, M. J., Al-Mefleh, N. K., Othman, Y. A. & Al-Assaf, A. (2021). Water harvesting techniques for improving soil water content, and morpho-physiology of pistachio trees under rainfed conditions. *Agricultural Water Management, 243*, 106464. https://doi.org/10.1016/j.agwat.2020.106464

Tajabadipour, A., Panahi, B. & Zadehparizi, R. (2006). The effects of rootstock and scion on early splitting and cracked hull of pistachio. *Acta Horticulturae*, 726, 193-198. https://doi.org/10.17660/ActaHortic.2006.726 .29

Tavallali, V. & Rahemi, M. (2007). Effects of rootstock on nutrient acquisition by leaf, kernel and quality of pistachio (*Pistacia vera* L.). *American-Eurasian Journal of Agricultural and Environmental Science*, 2(3), 240-246.

Turker, S. & Ak, B. E. (2010). Effects of different rootstocks on physical traits of 'Siirt' and 'Ohadi' pistachio cultivars. *Options Méditerranéennes*, 94, 245-250.

Vahdati, K., Sarikhani, S., Arab, M. M., Leslie, C. A., Dandekar, A. M., Aletà, N., ... & Mehlenbacher, S. A. (2021). Advances in rootstock breeding of nut trees: objectives and strategies. *Plants*, *10*(11), 2234. https://doi.org/10.3390/plants10112234



Valverdi, N. A. & Kalcsits, L. (2021). Rootstock affects scion nutrition and fruit quality during establishment and early production of 'Honeycrisp' apple. *HortScience*, *56*(2), 261-269. https://doi.org/10.21273/HORTSCI15488-20

Warschefsky, E. J., Klein, L. L., Frank, M. H., Chitwood, D. H., Londo, J. P., von Wettberg, E. J. & Miller, A. J. (2016). Rootstocks: diversity, domestication, and impacts on shoot phenotypes. *Trends in plant science*, *21*(5), 418-437. https://doi.org/10.1016/j.tplants.2015.11.008 Yin, R., Bai, T., Ma, F., Wang, X., Li, Y. & Yue, Z. (2010). Physiological responses and relative tolerance by Chinese apple rootstocks to NaCl stress. *Scientia Horticulturae*, *126*(2), 247-252. https://doi.org/10.1016/J.SCIENTA.2010.07. 027

Zörb, C., Geilfus, C. M. & Dietz, K. J. (2019). Salinity and crop yield. *Plant biology*, *21*, 31-38. <u>https://doi.org/10.1111/plb.12884</u>

