

The voltage-dependent sodium channel: A well-known molecular target of saxitoxins (cyanotoxins), ciguatoxins and brevetoxins (phycotoxins)

Evelyne Benoit, Denis Servent, Jordi Molgó

▶ To cite this version:

Evelyne Benoit, Denis Servent, Jordi Molgó. The voltage-dependent sodium channel: A well-known molecular target of saxitoxins (cyanotoxins), ciguatoxins and brevetoxins (phycotoxins). PHYCOTOX 2023 - GDR Annual Conference on Toxic Microalgae, May 2023, Nantes, France. hal-04473937

HAL Id: hal-04473937 https://hal.science/hal-04473937v1

Submitted on 22 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The voltage-dependent sodium channel: A well-known molecular target of saxitoxins (cyanotoxins), ciguatoxins and brevetoxins (phycotoxins)

Evelyne Benoit*†1, Denis Servent¹, and Jordi Molgo¹

¹Université Paris-Saclay – CEA – Institut des sciences du vivant Frédéric Joliot, Département Médicaments et Technologies pour la Santé (DMTS), Service d'Ingénierie Moléculaire pour la Santé (SIMoS), EMR CNRS/CEA 9004, 91191 Gif-sur-Yvette, France

Résumé

Because of its highly selectivity to Na ions, and activation and inactivation gating properties, the voltage-dependent sodium channel plays a fundamental role in generating and conducting action potentials. The channel alpha-subunit is a transmembrane protein composed of 4 domains (DI-DIV), each of them consisting in 6 segments (S1-S6). The segment S4 is responsible for the channel activation and the P loop, connecting S5 and S6, is involved in the channel selectivity. Receptor-sites for toxins have been identified on this alpha-subunit. Indeed, aquatic neurotoxins interacting specifically with this channel are widely distributed among the various organisms responsible for intoxications or envenomations. In particular, saxitoxins, gonyautoxins and derivates, a family of divalent and tricyclic cations produced by some dinoflagellates and cyanobacteria and responsible for Paralytic Shellfish Poisoning, interact with the receptor-site 1 of sodium channels, leading to pore blockade and thus to current and action potential inhibition. This is exemplified by an electrophysiological study of oyster cerebro-visceral nerve sensitivity to saxitoxin which decreases during recent exposure of bivalve mollusks to contaminated dinoflagellates in natural and, in a more subtle way, experimental environments, providing better understanding of the mechanisms associated with the contamination of bivalve mollusks and, therefore, of the means of detecting and counterbalancing this contamination (1). Ciguatoxins and brevetoxins, two families of cyclic polyethers produced by some dinoflagellates and responsible for Ciguatera Fish Poisoning and Neurologic Shellfish Poisoning, respectively, interact with the receptor-site 5 of sodium channels, leading to partly blockade of inactivation and thus to spontaneous and/or repetitive action potentials. Consequently, toxin-induced cell swelling occurs. This is exemplified by electrophysiological and confocal-imaging studies of toxin effects on vertebrate myelinated nerve fibers (2,3). (1) Boullot et al., Mar. Drugs, 2021, 19(7):380. (2) Mattei et al., J. Neurosci. Res., 1999, 55(6):666-673. (3) Mattei et al., Neuropharmacology, 2014, 85:417-426.

Mots-Clés: Sodium channels, Biophysic properties, Action potential, Saxitoxins, Ciguatoxins, Brevetoxins, Cell swelling

^{*}Intervenant

[†]Auteur correspondant: evelyne.benoit@cea.fr