



New Genotype of *Coxiella burnetii* Causing Epizootic Q Fever Outbreak in Rodents, Northern Senegal

Joa Mangombi-Pambou, Laurent Granjon, Clément Labarrere, Mamadou Kane, Youssoupha Niang, Pierre-Edouard Fournier, Jérémy Delerce, Florence Fenollar, Oleg Mediannikov

► To cite this version:

Joa Mangombi-Pambou, Laurent Granjon, Clément Labarrere, Mamadou Kane, Youssoupha Niang, et al.. New Genotype of *Coxiella burnetii* Causing Epizootic Q Fever Outbreak in Rodents, Northern Senegal. *Emerging Infectious Diseases*, 2023, 29 (5), pp.1078-1081. 10.3201/eid2905.221034 . hal-04475772

HAL Id: hal-04475772

<https://hal.science/hal-04475772>

Submitted on 23 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Acknowledgments

We thank Pieter Monsieurs for help with generating whole-genome sequence data and Isabel Micalessi and Lara Balcaen for molecular diagnosis of the case.

Genomic sequence reads of MHOM/CR/2020/StPierre are available on the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) under accession no. PRJNA881292. The *hsp70* sequence is available from GenBank under accession no. OQ200658.

F.V.D.B. and S.H. acknowledge support from the Research Foundation Flanders (grant nos. 1226120N and G092921N).

About the Author

Dr. Van den Broeck is a researcher specializing in genetics and population genomics of eukaryote parasites, particularly parasites that cause neglected tropical diseases. He has contributed multiple research papers investigating the complex hybrid ancestry of protozoan parasites, such as *Trypanosoma cruzi* (agent of Chagas disease) and *Leishmania braziliensis* (agent of cutaneous leishmaniasis).

References

1. Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R. Leishmaniasis: a review. *F1000 Res.* 2017;6:750. <https://doi.org/10.12688/f1000research.11120.1>
2. Pan American Health Organization. Guideline for the treatment of leishmaniasis in the Americas. 2nd ed. Washington: The Organization; 2022.
3. van Griensven J, van Henten S, Mengesha B, Kassa M, Adem E, Endris Seid M, et al. Longitudinal evaluation of asymptomatic *Leishmania* infection in HIV-infected individuals in North-West Ethiopia: a pilot study. *PLoS Negl Trop Dis.* 2019;13:e0007765. <https://doi.org/10.1371/journal.pntd.0007765>
4. Van der Auwera G, Maes I, De Doncker S, Ravel C, Cnops L, Van Esbroeck M, et al. Heat-shock protein 70 gene sequencing for *Leishmania* species typing in European tropical infectious disease clinics. *Euro Surveill.* 2013;18:20543. <https://doi.org/10.2807/1560-7917.ES2013.18.30.20543>
5. Van den Broeck F, Savill NJ, Imamura H, Sanders M, Maes I, Cooper S, et al. Ecological divergence and hybridization of Neotropical *Leishmania* parasites. *Proc Natl Acad Sci U S A.* 2020;117:25159–68. <https://doi.org/10.1073/pnas.1920136117>
6. Miranda ADC, González KA, Samudio F, Pineda VJ, Calzada JE, Capitan-Barrios Z, et al. Molecular identification of parasites causing cutaneous leishmaniasis in Panama. *Am J Trop Med Hyg.* 2021;104:1326–34. <https://doi.org/10.4269/ajtmh.20-1336>
7. Tihon E, Imamura H, Dujardin JC, Van Den Abbeele J, Van den Broeck F. Discovery and genomic analyses of hybridization between divergent lineages of *Trypanosoma congolense*, causative agent of Animal African Trypanosomiasis. *Mol Ecol.* 2017;26:6524–38. <https://doi.org/10.1111/mec.14271>
8. Belli AA, Miles MA, Kelly JM. A putative *Leishmania panamensis/Leishmania braziliensis* hybrid is a causative agent of human cutaneous leishmaniasis in Nicaragua. *Parasitology.* 1994;109:435–42. <https://doi.org/10.1017/S0031182000080689>
9. Brito MEF, Andrade MS, Mendonça MG, Silva CJ, Almeida EL, Lima BS, et al. Species diversity of *Leishmania* (*Viannia*) parasites circulating in an endemic area for cutaneous leishmaniasis located in the Atlantic rainforest region of northeastern Brazil. *Trop Med Int Health.* 2009;14:1278–86. <https://doi.org/10.1111/j.1365-3156.2009.02361.x>
10. Tabbabi A, Cáceres AG, Bustamante Chauca TP, Seki C, Choochartpong Y, Mizushima D, et al. Nuclear and kinetoplast DNA analyses reveal genetically complex *Leishmania* strains with hybrid and mito-nuclear discordance in Peru. *PLoS Negl Trop Dis.* 2020;14:e0008797. <https://doi.org/10.1371/journal.pntd.0008797>

Address for correspondence: Frederik Van den Broeck or Gert Van der Auwera, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium; email: fvandenbroeck@gmail.com or gvdauwera@itg.be

New Genotype of *Coxiella burnetii* Causing Epizootic Q Fever Outbreak in Rodents, Northern Senegal

Joa Mangombi-Pambou, Laurent Granjon, Clément Labarrere, Mamadou Kane, Youssoupha Niang, Pierre-Edouard Fournier, Jérémy Delerce, Florence Fenollar, Oleg Mediannikov

Author affiliations: Centre Interdisciplinaire de Recherches Médicales de Franceville, Franceville, Gabon (J. Mangombi-Pambou); Aix-Marseille University, Marseille, France (J. Mangombi-Pambou, C. Labarrere, P.-E. Fournier, J. Delerce, F. Fenollar, O. Mediannikov); University Hospital Institute Méditerranée Infection, Marseille (J. Mangombi-Pambou, C. Labarrere, P.-E. Fournier, F. Fenollar, O. Mediannikov); Centre de Biologie pour la Gestion des Populations, Montpellier, France (L. Granjon); Biologie des Populations Animales Sahelo-Soudaniennes, Dakar, Senegal (M. Kane, Y. Niang)

DOI: <https://doi.org/10.3201/eid2905.221034>

In Senegal, *Coxiella burnetii*, which causes Q fever, has often been identified in ticks and humans near livestock, which are considered to be reservoirs and main sources of infection. We describe the emergence of *C. burnetii* in rodents, not previously known to carry this pathogen, and describe 2 new genotypes.

Coxiella burnetii is a causative agent of Q fever, a worldwide zoonosis. The disease may be acute (relatively benign) or chronic (with a wide range of clinical manifestations that can lead to high human mortality) (1). Humans are infected by inhaling contaminated environmental dust and aerosol particles from the birth products of infected animals, as well as through direct contact with milk, urine, or feces containing *C. burnetii* (2,3). Humans are not considered natural hosts of *C. burnetii* (4). A wide spectrum of animals can serve as hosts (4), but the reservoirs are livestock, mainly sheep, cattle, and goats (3), which are also the main sources of human infection (1).

In Senegal, *C. burnetii* has been reported in humans (4–6) and ticks (4). It has been isolated from rodent-associated soft ticks (*Ornithodoros sonrai*) and detected in several species of hard ticks collected from ruminants (4). Our previous study of zoonotic pathogens in rodents collected in 2017 revealed no presence of *C. burnetii* in rodent populations from the Ferlo region in northern Senegal (7). However, in this study, we tested rodent samples collected during 2019–2020 from the same region and found high prevalence of a new *C. burnetii* genotype, which might indicate an ongoing epizootic outbreak.

We screened 125 rodent samples for *C. burnetii*; the rodents were collected in the Ferlo region in northern Senegal near Widou Thiengoly (15.99°N, 15.32°W) under framework agreements between the French National Research Institute for Development and Senegal (7). None of the rodent species investigated were listed as

protected with the International Union for Conservation of Nature or the Convention on International Trade in Endangered Species of Wild Fauna and Flora. Handling procedures were performed under Centre de Biologie Pour la Gestion des Populations agreement no. D-34-169-1 for experiments on wild animals and followed the guidelines of the American Society of Mammologists (8).

Rodents sampled belonged to the species *Arvicanthis niloticus* (n = 29), *Desmodilliscus braueri* (n = 3), *Gerbillus nancillus* (n = 9), *G. nigeriae* (n = 71), *Jaculus jaculus* (n = 4), *Taterillus* spp. (probably *T. pygargus*) (n = 8), and *Xerus erythropus* (n = 1). We extracted DNA from the spleen as described elsewhere (7) and stored it at –20°C. We detected bacterial DNA using *C. burnetii*-specific quantitative real-time PCR with primers and probes targeting IS1111 and IS30A spacers (4). For positive samples with a cycle threshold value <38, we first amplified 3 pairs of intergenic spacer primers, Cox2F/R, Cox5F/R, and Cox18F/R (5). Multispacer sequence typing (MST) genotyping of *C. burnetii* strains using sequences from the amplification of these 3 primer pairs revealed a potential new genotype. We amplified the other 7 primer pairs, Cox20F/R, Cox22F/R, Cox37F/R, Cox51F/R, Cox56F/R, Cox57F/R, and Cox61F/R, to describe this genotype.

Overall, 22.4% (28/125 for IS1111) and 19.2% (24/125 for IS30A) of rodents screened were positive for *C. burnetii*-specific quantitative PCR: *Desmodilliscus braueri* (33.3%; 1/3), *G. nancillus* (33.3%; 3/9), *G. nigeriae* (28.2%; 20/71), *Jaculus jaculus* (25%; 1/4), and *Taterillus* spp. (37.5%, 3/8). We found no *Arvicanthis niloticus* or *Xerus erythropus* rodents positive for *C. burnetii*. We performed complete MST genotyping of positive *C. burnetii* strain samples, and the sequences obtained from the primer pairs showed 100% identity for all positive samples. Nevertheless, 1 sample showed an insertion of 5 nucleotide bases in the amplified sequence of Cox56 spacer (Table), indicating

Table. Characterization of new *Coxiella burnetii* MST75 and MST76 genotypes described in study of new genotype of *C. burnetii* causing epizootic Q fever outbreak in rodents, northern Senegal*

Species	No.	Is1111 positive, no. (%)	Profile of spacers										Genotype
			Cox2	Cox5	Cox18	Cox20	Cox22	Cox37	Cox51	Cox56	Cox57	Cox61	
<i>Arvicanthis niloticus</i>	29	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
<i>Desmodilliscus braueri</i>	3	1 (33.3)	3	14	6	6	5	4	10	10	13	5	MST75
<i>Gerbillus nancillus</i>	9	3 (33.3)	3	14	6	6	5	4	10	10	13	5	MST75
<i>G. nigeriae</i> †	71	20 (28.2)	3	14	6	6	5	4	10	20/21‡	13	5	MST75/ MST76‡
<i>Jaculus jaculus</i>	4	1 (25)	3	14	6	6	5	4	10	10	13	5	MST75
<i>Taterillus</i> spp.	8	3 (37.5)	3	14	6	6	5	4	10	10	13	5	MST75
<i>Xerus erythropus</i>	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total	125	28	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

*MST, multispacer sequence typing; NA, not applicable.

†Invasive (expanding) species.

‡Genotype MST76 has the same profile as MST75 except for Cox 56.




Figure. Neighbor-joining circular unrooted tree showing the relationship between *Coxiella burnetii* genotypes described in study of new genotype of *C. burnetii* causing epizootic Q fever outbreak in rodents, northern Senegal. MST75 and MST76 (red) were compared with genotypes already found in Senegal, MST19 and MST6 (blue), and other genotypes. The analysis involved 64 nt sequences. All positions containing gaps and missing data were eliminated. There were a total of 4,692 positions in the final dataset. Evolutionary analyses were conducted in MEGA7 (<https://www.megasoftware.net>). MST, multispacer sequence typing.

the presence of a probable variant. All of these sequences demonstrated the presence of ≥ 1 new genotypes of *C. burnetii* according to the BLAST database (<https://ifr48.timone.univ-mrs.fr/mst/coxiellaburnetii/blast.html>). Phylogenetic analysis based on concatenated sequences of spacers revealed that the new genotypes, MST75 (major) and MST76 (with an insertion), are close together and cluster with other genotypes, including those already found in Senegal, such as MST19, and those infecting animals and humans, such as MST16, MST17, MST20, and MST61 (Figure).

Our finding of *C. burnetii* MST75 and MST76 genotypes in the Ferlo rodent community suggests the emergence of a Q fever epizootic outbreak. Previously identified *C. burnetii* strains in Senegal were related to the proximity of livestock near the villages of Dielmo and Ndiop (9). In Ferlo, a previous study conducted on rodents sampled in 2017 did not find *C. burnetii* (7), indicating a relatively recent, possibly still ongoing epizootic outbreak. High *C. burnetii*

prevalences (28%–38%) were observed in different species of gerbilline rodents, including *G. nigeriae*, which has recently colonized northern Senegal and is now the dominant species in outdoor rodent communities of Ferlo (10). The possibility of animal transmission from farms located near the rodent sampling area should also be explored. Our study shows the emergence in Senegal of new *C. burnetii* genotypes in susceptible animals, such as rodents (1), which might be a source of human infections. Although the pathogenicity of these new genotypes for humans is yet unknown, our findings signal the urgent need for epidemiologic surveillance for *C. burnetii* infection in humans in Senegal and neighboring countries.

Acknowledgments

We thank Philippe Gauthier for precise taxonomic identification of some rodent specimens using barcoding (based on *cytb* sequences), and Fabien Fliriden for his technical support on molecular biology.

This study was supported by the Institut Hospitalo-Universitaire Méditerranée Infection, the French National Research Agency under the Investissements d'avenir programme, (reference ANR-10-IAHU-03), and the fourth Make Our Planet Great Again (MOPGA 4) program. This work was cofunded by the Labex Dispositif de Recherche Interdisciplinaire sur les Interactions Hommes-Milieus, and the Investissements d'Avenir program (ANR-11-LABX-0010), which is managed by the Agence nationale de la recherche.

About the Author

Dr. Mangombi-Pambou is a postdoctoral researcher at the Institut Hospitalo Universitaire Méditerranée Infection in Marseille, France. His research interests include infectious diseases of zoonotic origin and the ecology and role of rodents in the dynamics of zoonotic diseases.

References

1. Angelakis E, Raoult D. Q fever. *Vet Microbiol*. 2010;140:297–309. <https://doi.org/10.1016/j.vetmic.2009.07.016>
2. Tissot-Dupont H, Raoult D. Q fever. *Infect Dis Clin North Am*. 2008;22:505–14, ix. <https://doi.org/10.1016/j.idc.2008.03.002>
3. Vanderburg S, Rubach MP, Halliday JEB, Cleaveland S, Reddy EA, Crump JA. Epidemiology of *Coxiella burnetii* infection in Africa: a OneHealth systematic review. *PLoS Negl Trop Dis*. 2014;8:e2787. <https://doi.org/10.1371/journal.pntd.0002787>
4. Mediannikov O, Fenollar F, Socolovschi C, Diatta G, Bassene H, Molez J-F, et al. *Coxiella burnetii* in humans and ticks in rural Senegal. *PLoS Negl Trop Dis*. 2010;4:e654. <https://doi.org/10.1371/journal.pntd.0000654>
5. Angelakis E, Mediannikov O, Socolovschi C, Mouffok N, Bassene H, Tall A, et al. *Coxiella burnetii*-positive PCR in febrile patients in rural and urban Africa. *Int J Infect Dis*. 2014;28:107–10. <https://doi.org/10.1016/j.ijid.2014.05.029>
6. Sokhna C, Mediannikov O, Fenollar F, Bassene H, Diatta G, Tall A, et al. Point-of-care laboratory of pathogen diagnosis in rural Senegal. *PLoS Negl Trop Dis*. 2013;7:e1999. <https://doi.org/10.1371/journal.pntd.0001999>
7. Dahmana H, Granjon L, Diagne C, Davoust B, Fenollar F, Mediannikov O. Rodents as hosts of pathogens and related zoonotic disease risk. *Pathogens*. 2020;9:202. <https://doi.org/10.3390/pathogens9030202>
8. Sikes RS, Gannon WL; Animal Care and Use Committee of the American Society of Mammalogists. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *J Mammal*. 2011;92:235–53. <https://doi.org/10.1644/10-MAMM-F-355.1>
9. Diouf FS, Ndiaye EHI, Hammoud A, Diamanka A, Bassene H, Ndiaye M, et al. Detection of *Coxiella burnetii* and *Borrelia* spp. DNA in cutaneous samples and in household dust in rural areas, Senegal. *Vector Borne Zoonotic Dis*. 2021;21:659–66. <https://doi.org/10.1089/vbz.2020.2723>
10. Granjon L, Ba K, Diagne C, Ndiaye A, Piry S, Thiam M. The Ferlo small rodent community: historical trends and characteristics of the current population [in French]. In: Boëtisch G, Duboz P, Guissé A, Sarr P, editors. *The Great*

Green Wall, an African response to climate change [in French]. Paris: Centre national de la recherche scientifique; 2019. p. 1–19.

Address for correspondence: Oleg Mediannikov, IHU Méditerranée Infection, 19-21 Boulevard Jean Moulin, 13005 Marseille, France; email: olegusss1@gmail.com

Therapeutic Failure and Acquired Bedaquiline and Delamanid Resistance in Treatment of Drug-Resistant TB

James Millard, Stephanie Rimmer, Camus Nimmo, Max O'Donnell

Author affiliations: Guys and St Thomas' NHS Foundation Trust, London, UK (J. Millard); Wellcome Trust Liverpool Glasgow Centre for Global Health Research, Liverpool, UK (J. Millard); University of Liverpool, Liverpool (J. Millard); Africa Health Research Institute, Durban, South Africa (J. Millard, C. Nimmo); Imperial College Healthcare, London (S. Rimmer); Francis Crick Institute, London (C. Nimmo); University College London, London (C. Nimmo); Columbia University Medical Center, New York, New York, USA (M. O'Donnell); CAPRISA MRC-HIV-TB Pathogenesis and Treatment Research Unit, Durban (M. O'Donnell)

DOI: <http://doi.org/10.3201/eid2905.221716>

New classes of antitubercular drugs, diarylquinolines and nitroimidazoles, have been associated with improved outcomes in the treatment of drug-resistant tuberculosis, but that success is threatened by emerging drug resistance. We report a case of bedaquiline and delamanid resistance in a 55-year-old woman in South Africa with extensively drug-resistant tuberculosis and known HIV.

Major improvements have been achieved in drug-resistant tuberculosis (TB) treatment in recent years; 2 new drug classes, diarylquinolines (bedaquiline) and nitroimidazoles (pretomanid and delamanid), have been central to this success (1). The Ze-NiX trial demonstrated cure rates of >90% in complex