



HAL
open science

Deviations for generalized tiling billiards in cyclic polygons

Magali Jay

► **To cite this version:**

| Magali Jay. Deviations for generalized tiling billiards in cyclic polygons. 2024. ⟨hal-04476038⟩

HAL Id: hal-04476038

<https://hal.science/hal-04476038v1>

Preprint submitted on 24 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-SA 4.0 - Attribution - Non-commercial use - ShareAlike - International License

DEVIATIONS FOR GENERALIZED TILING BILLIARDS IN CYCLIC POLYGONS

MAGALI JAY

Abstract. This work continues the study of tiling billiards, a class of dynamical system introduced by Davis et al. in 2018. We develop the study of generalized tiling billiards in a cyclic polygon. This work shows that the behavior of generalized tiling billiards in cyclic N -gons with $N \geq 5$ is considerably different from that of triangular and quadrilateral tiling billiards studied before. Indeed, we exhibit an open set of generalized tiling billiard trajectories deviating sublinearly from their asymptotic direction, whereas for $N = 3$ or 4 almost every trajectory stays at a bounded distance from a line. Moreover, we establish the rate of deviations both in the generic case and in some non generic cases.

1. INTRODUCTION

1.1. A quick bibliographical overview of tiling billiards. Davis, DiPietro, Rustad and St-Laurent [DDRS18] defined and studied a class of dynamical systems that they called *tiling billiards*. In a tiling billiard, the trajectory is refracted across the boundary of a plane tiling (the angle of incidence and the angle of reflection are opposite), see Figure 2. Nogueira had already suggested to study such billiards at the end of [Nog89], in relation with interval exchange transformations with flips (IETFs).




FIGURE 1. Trajectories of regular tiling billiards




FIGURE 2. Example of two trajectories of a polygonal tiling billiard
Only a part of the tiling is drawn, hence only a part of each trajectory.

For a fixed tiling of the plane, we are interested in the following questions. What sort of trajectories exist? Periodic, unbounded, dense in open sets of the plane? Are they stable under perturbations of initial conditions of the trajectory (starting point and direction) or of the form of the tiling?

There exist examples of tiling billiards with *periodic*, *drift-periodic* or *linearly escaping* trajectories. We call **drift-periodic** a trajectory with a translation symmetry. We say that a trajectory is **linearly escaping** if it is not drift-periodic and the main part of the asymptotic expansion of its coordinates (by arclength parametrization) is linear and the remaining term is bounded by a constant. Another behavior is known: in the trihexagonal tiling billiard, the generic trajectories are dense in open sets of the plane [DH18].

Triangle and cyclic quadrilateral tilings. Let us consider a triangle tiling where each tile is a centrally symmetric copy of each of its neighboring tiles. The qualitative behavior of triangle tiling billiards has been fully classified in [Par19], building upon [BDFI24] and [HP22]. We say that a trajectory is **stable** if a small perturbation of it leads to another trajectory that crosses the same sides (in the same order).


Theorem 1.1 ([BDFI24], [Par19], [HP22]). *Let δ be a triangle. Let γ be a trajectory in the δ -tiling billiard. Then exactly one of the following four cases holds:*

- (1) *The trajectory γ is periodic and stable under perturbation of the tiling or of the initial conditions.*
- (2) *The trajectory γ is drift-periodic and the angles of δ are rationally dependent.*
- (3) *The trajectory γ is linearly escaping.*
- (4) *The trajectory γ is not bounded, has no asymptotic direction and passes through circumcenter of triangles. Moreover δ belongs to the Rauzy gasket, which has a zero Lebesgue measure.*

See Figure 3 for an illustration of the first three cases. The exceptional case of unbounded trajectories with no asymptotic direction is linked to minimal IETFs. Hubert and Paris-Romaskevich with Dynnikov, Mercat and Skripchenko [DHMPS] obtained the analogous result for classification of trajectories on cyclic quadrilateral tilings, and related the dynamics of such billiards to the Novikov's problem of the study of foliations on subsurfaces of genus 3 of the 3-torus given by 1-forms, see also [Par21].

For triangle and cyclic quadrilateral tiling, the dynamical systems have an integral of motion: the oriented distance of a trajectory to the circumcenters of the crossed polygons is preserved. Baird-Smith, Davis, Fromm, Iyer explain this for triangles in [BDFI24] (Lemma 3.2), with a beautiful argument of folding. It was then generalized in [Par19]. This allows us to study such tiling billiards via IETFs. See Section 2.2 for another explanation of the link between tiling billiards and IETFs.

The behaviors of trajectories in tiling billiards tend to differ from the ones in classical inner billiards, as dynamics of IETFs differs from those of interval exchange transformations without flips (IETs). No general study of tiling billiards exists yet. In this article, we will handle open sets of parameters in generalized tiling billiards, made with any cyclic N -gons with $N \geq 5$, although they will not tile the plane. The results are true for polygons with more than five sides but the reader can think of pentagons in a first reading. All polygons will be drawn as pentagons.




(A) Some periodic trajectories and two (stable) perturbations

The green and grey tiling is a perturbation of the green and orange one.

(B) A part of a drift-periodic trajectory and two of its (stable) perturbations

The light and deep blue tiling is a perturbation of the green and orange one.



(c) A part of a linearly escaping trajectory

FIGURE 3. Examples of trajectories in triangle tiling billiards

1.2. The studied system.

The generalized tiling. We first define how we build the generalized tiling with the N -gons ($N \geq 5$). Each one of its tiles is congruent to a fixed cyclic polygon. Moreover if T and T'

are two neighboring tiles then T' is the image of T via a central symmetry with respect to the middle of their shared side. This does not define a proper tiling: polygons will either overlap each other or not fill the entire plane. Indeed when turning around a vertex, all the angles of the N -gon appear and sum up to $(N-2)\pi$, which is not a divisor of 2π , see Figure 4. But we can still define a billiard trajectory in the plane, generalizing tiling billiards for any polygon.




FIGURE 4. The way we put polygons together to play tiling billiards will not tile the plane with N -gons, $N \geq 5$.



Fix a N -gon on the plane, a point inside it and a direction (see Figure 5a). Follow the direction until reaching a side. Then draw a second centrally symmetric polygon, so that the two polygons share the crossed side (see Figure 5b). The trajectory is refracted through the side and continues in a straight line in the second polygon (see Figure 5c). Each time the trajectory reaches a side, we draw the following polygon via central symmetry. The new polygon may overlap a previous one. The trajectory takes into account only the last drawn polygon (see Figure 5e and 5f). A trajectory that hits a corner stops.

Assumptions. Let P be a cyclic polygon. We will denote by $\mathbf{a}_1, \dots, \mathbf{a}_N$ the sides of P , clockwise labelled and \mathbf{a}_N being its longest side. We denote by a_1, \dots, a_N the lengths of the arcs subtended by each side (see Figure 6a). Conversely, if $a_1, \dots, a_N > 0$, with a_N being maximal, we denote by $P(a_1, \dots, a_N)$ the cyclic polygon whose arcs subtended by its sides have lengths a_1, \dots, a_N . We will restrict ourselves to the study of billiards satisfying the following conditions:

$$(C) \begin{cases} P \text{ is inscribed in a circle} & (C0) \\ a_1 + \dots + a_{N-1} < \tau < a_N & (C1) \end{cases}$$

Condition (C0) ensures that a certain parameter τ (see Figure 6b and Definition 2.3) is preserved all along the trajectory and hence allows us to use IETFs to study the system (see Sections 2.2 and 2.3), as in [BDFI24], [HP22], [Par19], [Par21] and [DHMPS]. This is a crucial fact for the now existing methods of study — we know nothing for tiling billiards made with non cyclic polygons. Condition (C1) means both that P is inscribed in a half-circle and that the trajectory is close to the circumcenter. It is crucial for the following since it allows to reduce the system to a standard IET without flips with N intervals of continuity, which permits to use ergodic theory tools (see paragraphs 2.2 and 2.3).

1.3. Results. Let us identify the real plane with \mathbb{C} . Let $p_n \in \mathbb{C}$ be the point of the trajectory on the n -th side crossing.



(G) A part of a generalized pentagon tiling billiard trajectory satisfying condition (C)

FIGURE 5. Examples of trajectories in a generalized pentagonal tiling billiard



FIGURE 6. Notations for cyclic polygons

Lemma 1.2. For every a_1, \dots, a_{N-1} , almost every $a_N > \sum_{i=1}^{N-1} a_i$, and every $\tau \in \left(\sum_{i=1}^{N-1} a_i, a_N \right)$, the cyclic N -gon $P(a_1, \dots, a_N)$ defines a generalized tiling billiard whose trajectories having parameter τ all admit a mean displacement $m = \lim_{n \rightarrow \infty} \frac{p_n}{n}$ that does not vanish.

Theorem A. *Let \mathcal{L} be a line in the asymptotic direction (almost always defined) of the trajectory. For almost every tiling billiard satisfying the conditions (C), the trajectory does not stay at a bounded distance from \mathcal{L} .*

We quantify the order of magnitude of deviations from the asymptotic direction with the following two theorems. They are linked to the behavior of the Teichmüller flow on the moduli space of translation surfaces.

Theorem B. *Let $N \in \mathbb{N}, N \geq 5$. There exist $\theta_1 > \theta_2 > \theta_3 > 0$ such that for almost every N -gon \mathbf{P} and every τ , both satisfying condition (C), for any initial direction having parameter τ in a \mathbf{P} -tiling billiard, the sequence of points $(p_n)_{n \in \mathbb{N}}$ satisfies:*

$$\limsup_{n \rightarrow \infty} \frac{\log |p_n - nm|}{\log n} = \begin{cases} \frac{\theta_2}{\theta_1} > 0 & \text{if } N \text{ is odd} \\ \frac{\theta_2}{\theta_1} \text{ or } \frac{\theta_3}{\theta_1} > 0 & \text{if } N \text{ is even} \end{cases},$$

where $m = \lim_{n \rightarrow \infty} \frac{p_n}{n}$.

Remark 1.3. *The numbers θ_1, θ_2 and θ_3 depend only on the number of sides of the polygon! The numbers $1 + \frac{\theta_i}{\theta_1}$ ($1 \leq i \leq 3$) are the first three Lyapounov exponents of Teichmüller flow in the hyperelliptic component of the stratum $\mathcal{H}(N-3)$ if N is odd, and $\mathcal{H}(\frac{N}{2}-2, \frac{N}{2}-2)$ if N is even.*

When $N = 5$, a result of Eskin, Kontsevich and Zorich [EKZ11] states that $\frac{\theta_2}{\theta_1} = \frac{1}{3}$. Forni [For02] proves that θ_1, θ_2 and θ_3 are distinct.

We prove these results via IETs and their generic properties. The Rauzy induction \mathfrak{R} (see paragraph 3.1) is a powerful tool to study orbits of an IET. If T is an IET on I , $\mathfrak{R}(T)$ is the first return map of T on a well chosen subinterval $J \subset I$, which gives once again an IET with the same number of intervals of continuity. To track the orbit of T through Rauzy induction, we need to know both the sequence of combinatorial data of the induced IETs $(\mathfrak{R}^n(T))_{n \in \mathbb{N}}$ and the lengths of their intervals of continuity. This information can be encoded in an infinite path in a finite graph, called the Rauzy diagram. The lengths of intervals of continuity of $\mathfrak{R}^n(T)$ linearly depend on those of T . A corresponding matrix cocycle (see paragraph 3.1) is important for the study of our dynamical system. The Rauzy induction for IETs corresponds to the Teichmüller flow for their suspensions (that are translation surfaces). See [Via06], [MMY05],[Yoc10].

When the path in the Rauzy diagram corresponding to some IET is a loop, which is the same as to say that the orbit of the IET through the Rauzy induction is periodic, we say that this IET is **self-similar**. The matrix associated to this loop allows us to study the symbolic coding of a point by the IET T . When the loop is complex enough, the matrix is primitive. This case is obviously not generic but important. Indeed it corresponds to the case when the vertical flow on a suspension of the IET is a pseudo-Anosov map. In this case, one can give a more precise estimate of the deviations.

Theorem C. *Let $N \in \mathbb{N}, N \geq 5$ and $a = (a_1, a_2, \dots, a_{N-1})$ be the (normalized) Perron eigenvector (with eigenvalue λ_1) of a primitive matrix $M \in \mathcal{M}_{N-1}(\mathbb{R})$ that corresponds to a loop in the Rauzy diagram. Assume that the second eigenvalue λ_2 of M is such that $|\lambda_2| > 1$ and that no other eigenvalue has the same modulus as λ_2 . For almost every $a_N > \sum_{i=1}^{N-1} a_i$, there exist C_1 and C_2 and an infinite set $\mathcal{S} \subset \mathbb{N}$ such that for every $\tau \in (\sum_{i=1}^{N-1} a_i, a_N)$ and any*

trajectory having parameter τ in a $P(a_1, \dots, a_N)$ -tiling billiard, the sequence of points $(p_n)_{n \in \mathbb{N}}$ satisfies:

$$\begin{cases} \forall n \in \mathbb{N} & |p_n - nm| < C_1 n^{\frac{\log \lambda_2}{\log \lambda_1}} \\ \forall n \in \mathcal{S} & |p_n - nm| > C_2 n^{\frac{\log \lambda_2}{\log \lambda_1}} \end{cases} .$$

Figure 7 illustrates this phenomenon of deviation and the growth rate. This theorem quantifies the second term of the asymptotic expansion of some Birkhoff's sums for a fixed point. Another related problem is to study the uniform speed of convergence of the same Birkhoff's sums, i.e. to study the *discrepancy*. This was done by Adamczewski in [Ada04].




FIGURE 7. The modulus of the deviations, depending on the number of steps of the trajectory, has a sublinear growth

Deviations, depicted in blue, are always smaller than a (strictly) sublinear function, depicted in green, with a growth rate ρ , and infinitely often greater than another function, also in green, with the same growth rate ρ .

To summarize, this article shows that for almost every generalized tiling billiard satisfying condition (C):

- (1) like for triangle or cyclic quadrilateral tiling billiards, the trajectory has an asymptotic direction (Lemma 1.2),
- (2) unlike in triangles and cyclic quadrilaterals case, it deviates sublinearly from it (Theorems A, B and C).

1.4. **Outline of the proof.** The proofs of our theorems are organised as follows:

- (1) Define the IET that corresponds to our generalized tiling billiard.
- (2) Define a piecewise constant function f (that depends on P) and approximates the trajectory. We show that f has almost always a non zero mean, which means that the trajectory of the studied tiling billiard has an asymptotic direction.
- (3) Study the function $h = \text{Im}(\frac{1}{m}f - 1)$ linked to the deviations from the asymptotic direction. We show that for almost every a_N , h is not contained in a codimension 3 subspace (resp. 2 if N is odd).
- (4) Zorich's results on deviations for IETs give the rate given in Theorem B

IETs are generically uniquely ergodic [Mas82] [Vee82]. Hence we know that the orbit of a point equidistributes in the different intervals of continuity. Zorich establishes [Zor99] how much an orbit can deviate from this equidistribution. We can think of such results as central

limit theorems. To prove this, Zorich studies Teichmüller flow of suspension surfaces (that are translation surfaces) of IETs and how it behaves with respect to Oseledets' flag decomposition.

The main contribution of this article is to prove that one can apply Zorich's results to the case of tiling billiards we study. This is not a straightforward verification. We have to show where some specific vector lies in the flag given by the Oseledets' theorem (see Section 3.2). The difficult point is that we know the expression of the vector but, in general, not of the subspaces of the flag. We can think of them as random subspaces. We can still conclude for almost every parameters because the vector has one more degree of freedom than the flag. Moreover it depends analytically on this parameter independent of the flag. This corresponds to step (3) in the following decomposition of the proof:

The most technical parts of the proofs of Theorems B are done first for pentagons and then generalized. We prove Theorem C by direct computation.

1.5. Outline of the paper. We explain step (1) in Section 2. Then we recall briefly some background in Section 3. We introduce functions f and g in Section 4. Section 5 studies the function g and show the properties that allow us to apply Zorich's results. Section 6 handles with the direct computations of deviations in case of a self-similar IET, using techniques of symbolic coding, Rauzy induction and Rokhlin towers, in order to prove Theorem C.

Acknowledgements

I thank my PhD advisors Pascal Hubert and Olga Paris-Romaskevich for having introduced this subject to me.

2. THE STUDIED SYSTEM

2.1. Definition of a generalized tiling billiard.

Definition 2.1 (Generalized tiling billiard). *Let P be a polygon and v a vertex of P . We call a generalized tiling billiard trajectory the triplet $(\mathbf{T}, (P_i)_{i \in \mathbb{N}}, (v_i)_{i \in \mathbb{N}})$ where \mathbf{T} is a piecewise linear curve in the real plane such that for every integer i :*

- (P_i, v_i) is an isometric copy of (P, v) in the plane,
- \mathbf{T} crosses P_i in a segment,
- P_i and P_{i+1} share a side, crossed by \mathbf{T} ,
- P_{i+1} is the image of P_i through a central symmetry,
- the common side of P_i and P_{i+1} is the bissector of the angle made by $\mathbf{T} \cap P_i$ and $\mathbf{T} \cap P_{i+1}$.

This defines a P -generalized tiling billiard as a dynamical system. See Figure 8 for an illustration.

Remark 2.2. *When we think of a generalized tiling billiard trajectory, we mostly think of \mathbf{T} although defining it as a triplet with crossed polygons and vertices v_i is much more convenient for the study of the trajectory. The definition requires no property of P although we will restrict in the following to the case where P is cyclic. We denote by \mathcal{C}_i the circumcircle of P_i .*

2.2. A system of coordinates. Here we describe a convenient frame of reference to study a tiling billiard trajectory. It is the one of an ant that would follow this trajectory. At each step, the ant can see the polygon it is crossing and its circumcircle. We need the following




FIGURE 8. Notations for generalized tiling billiards

data to determine the position of P_i with respect to the ant (in other words, the position of the ant in P):

- (1) On which chord of the circumcircle \mathcal{C}_i of P_i the ant is travelling,
- (2) Where v_i is on \mathcal{C}_i (this determines how P_i is placed in \mathcal{C}_i).

Lemma 3.2 in [BDFI24] applies also for generalized tiling billiards made with any cyclic polygon and shows that the chord of \mathcal{C}_i on which the ant is travelling is preserved (See Section 2.3). The data (1) is hence a parameter of the trajectory. For example if the ant is travelling on a diameter of \mathcal{C}_i , then it will also travel on a diameter of any \mathcal{C}_j . Therefore we will need only the position of v_i on \mathcal{C}_i to determine the position of P with respect to the ant.

Definition 2.3. *We say that the trajectory has parameter τ if the arc between the two endpoints of the chord supporting a step of the trajectory \mathbf{T} has length τ (see Figure 6b).*

Let $(\mathbf{T}, (P_i)_{i \in \mathbb{N}}, (v_i)_{i \in \mathbb{N}})$ be a tiling billiard. We normalize the situation to $a_1 + \dots + a_{N-1} = 1$. We identify the circumcircle \mathcal{C}_i of P_i with the interval $[0, \sum_{k=1}^N a_k) = [0, 1 + a_N)$ by identifying 0 to the point $o_i \in \mathcal{C}_i$ where the line containing $T_i = \mathbf{T} \cap P_i$ would meet \mathcal{C}_i if going on in straight line after the side crossing. Then we locate the vertex v_i of P_i in the frame of \mathcal{C}_i . Let $x_i \in [0, 1 + a_N)$ be the point with which v_i is identified. See Figure 9 for an illustration of notations.

2.3. The obtained Interval Exchange Transformation. We can prove the following by direct computation (see Figure 11) or with an argument of folding, which was the proof of [BDFI24] for triangles. It also applies for generalized tiling billiards made with any cyclic polygon, see Lemma 1 of [Par19].

Theorem 2.4 (Theorem 3.3 of [BDFI24]). *There is an IETF Φ such that $x_{i+1} = \Phi(x_i)$ for every i (see Figure 10a).*

If we add the hypothesis (C1), the IETF Φ of Theorem 2.4 has permutation

$$\pi(\Phi) = \left(\begin{array}{cccccc} \overline{I_1} & \overline{I_2} & \dots & \overline{I_{N-1}} & \overline{I_N^+} & \overline{I_N^-} \\ \overline{I_N^+} & \overline{I_1} & \overline{I_2} & \dots & \overline{I_{N-1}} & \overline{I_N^-} \end{array} \right)$$

where we overlined flipped intervals (here, the IET flips every interval). The intervals of continuity of the IET have lengths

$$\lambda(\Phi) = (a_1 \quad a_2 \quad \dots \quad a_{N-1} \quad a_N - \tau \quad \tau).$$




FIGURE 9. Moving frame of reference

Remark 2.5. Here we consider Φ as an IET on an interval, it has $N + 1$ intervals of continuity. If we define it on a circle it has only N intervals of continuity.

The choice of v ensures that the trajectory crosses the side \mathbf{a}_k if and only if $x \in I_k$.


The condition (C1) implies that the trajectory crosses the longest side every other crossing. It is hence enough to study the squared map Φ^2 , which is an IET without flip, see Figure 10. We consider $T = \Phi^2|_I$ the square of Φ , restricted on $I = I_1 \cup \dots \cup I_{N-1}$. Recall that we have normalized the situation so that $a_1 + \dots + a_{N-1} = 1$ so T is defined on $(0, 1)$. It is an IET (without flip) with permutation

$$\pi(T) = \begin{pmatrix} I_1 & I_2 & \dots & I_{N-1} \\ I_{N-1} & \dots & I_2 & I_1 \end{pmatrix}$$

and lengths

$$\lambda(T) = (a_1 \ a_2 \ \dots \ a_{N-1}).$$

We give here a straightforward way to compute the coordinate x_{i+1} depending on x_i , and on the parameters a_1, \dots, a_N and τ . Because of the refraction law, the lengths α and β on



Orbits of points inside grey parts do not correspond to a trajectory of a tiling billiard.

FIGURE 10. Example of the IETs Φ and Φ^2 , corresponding to a tiling billiard.




FIGURE 11. Coordinates x_i and x_{i+1} are related via a linear equation

the Figure 11 are equal. Since $\alpha = x_i - \sum_{k=1}^{j-1} a_k$, one gets that $\gamma = \tau - \beta = \tau - x_i + \sum_{k=1}^j a_k$. We conclude the expression of x_{i+1} depending on x_i, a_1, \dots, a_N and τ :

$$\begin{aligned} x_{i+1} &= \gamma + \sum_{k=1}^j a_k \\ &= \tau - (x_i - a_j) \end{aligned}$$

which corresponds to the expression of a fully flipped IET.

3. BACKGROUND

3.1. Rauzy induction, Rokhlin towers, and the associated cocycle. The Rauzy induction $\mathfrak{R}(T)$ of an IET T defined on I is the first return map of T to a well chosen subinterval

$J \subset I$. It can be convenient to consider the renormalized Rauzy induction \mathcal{R} that consists to remormalise J to the size of I . For more details, see [Via06]. Zorich [Zor96] showed that one can speed up the Rauzy induction so that the induction is ergodic with respect to an absolutely continuous invariant probability measure μ_Z . The Rauzy induction acts also on the underlying permutation of the IET. We call the **Rauzy class of the permutation** σ the permutations in the orbit of σ by the Rauzy induction.

When we cut off $I' = I \setminus J$ from I , we can think of it being stacked over its preimage by T . The choice of J ensures that the subinterval on which I' is being stacked corresponds exactly to an interval of continuity of $\mathfrak{R}(T)$. We can do that at each iteration of Rauzy induction. We obtain a stack of subintervals of I , that we call a **Rokhlin tower**. Let $J^{(k)}$ be the subinterval on which $\mathfrak{R}^k(T)$ is defined and $x \in J^{(k)}$. The subintervals stacked over x allow to keep track of the orbit of x under T . Their number corresponds to the time of first return of x to $J^{(k)}$. See Figure 12. We call **towers of order** l the ones obtained when applying l times the induction¹.

We denote by $A_{i,j}^{(l)}$ the number of subintervals stacked over the interval of continuity $I_j^{(l)}$ of $\mathfrak{R}^l(T)$ that were initially in the interval of continuity I_i of T . It corresponds to the number of times that the orbit of $x \in I_j^{(l)}$ lies in the *old* interval I_i before first return to $I_j^{(l)}$. This number does not depend on $x \in I_j^{(l)}$ thanks to the choice of subintervals on which we induce. We denote $A_T = \left(A_{i,j}^{(1)} \right)_{1 \leq i,j \leq d}$ the matrix whose entries are these numbers. It is a transvection. This defines a cocycle $T \in \text{IET}(I) \mapsto A_T \in \text{SL}_d(\mathbb{R})$.

Analogously we denote by B the cocycle associated to Zorich induction. We underline the fact that the time related to cocycle B is not the same as the one related to cycle A , there is a speeding up factor. Moreover there is an exponential factor between the time of cocycle B and the time of the iteration of T , see Lemma 4 in [Zor99].

3.2. Oseledets' theorem. We recall Oseledets' theorem on which rely Zorich's results that we will need later in this article.

Theorem 3.1 (Oseledets). *Let X be a topological space, $\varphi : X \rightarrow X$ be a measurable function, and μ be an ergodic φ -invariant probability measure on X . Let $A : \left\{ \begin{array}{l} X \rightarrow \mathcal{M}_d(\mathbb{R}) \\ x \mapsto A_x \end{array} \right\}$ be a cocycle. Assume that the cocycle is log-integrable: $\int_X \log \|A_x\| d\mu < \infty$.*

There exist $\theta_1 > \theta_2 > \dots > \theta_d$, called Lyapunov exponents, such that for almost every $x \in X$, there exists a flag $\mathbb{R}^d = \mathcal{H}_k^{(x)} \supset \mathcal{H}_{k-1}^{(x)} \supset \dots \supset \mathcal{H}_1^{(x)}$ such that:


$$A_x \mathcal{H}_j^{(x)} = \mathcal{H}_j^{(\varphi(x))}$$

and

$$\forall v \in \mathcal{H}_j \setminus \mathcal{H}_{j-1}, \frac{1}{n} \log \|A_x^{(n)} v\| \xrightarrow{n \rightarrow \infty} \theta_j$$

where $A_x^{(n)} = A_{\varphi^{n-1}(x)} \dots A_{\varphi(x)} A_x$.

¹Note that this notion depends on the induction we apply, Rauzy or Zorich one.



The colored subinterval will be cut in the following step of Rauzy induction (and stacked over the interval to form a Rokhling tower).




FIGURE 12. Rokhlin towers of orders 1 to 5 for the Rauzy induction of an IET

In our context we apply Oseledets' theorem in the set

$$X = \bigsqcup_{\sigma \in [\pi(T)]} \Delta_{N-1} \times \{\sigma\}$$

of all IETs whose underlying permutation is in the Rauzy class $[\pi(T)]$ of

$$\pi(T) = \begin{pmatrix} I_1 & I_2 & \dots & I_{N-1} \\ I_{N-1} & \dots & I_2 & I_1 \end{pmatrix}.$$

The space X is equipped with μ_Z , which is an ergodic \mathfrak{R} -invariant probability measure [Zor96]. The cocycle B introduced in last section is invertible, B and B^{-1} are log-integrable. We consider here B^{-1} . This is used in [Zor99] to prove Proposition 5.1.

In this article we call **Oseledets generic** both an IET for which the flag decomposition exists and the corresponding uplet of its lengths (a_1, \dots, a_{N-1}) .

3.3. Decomposition of the symbolic coding. At each step of renormalized Rauzy induction, there is an associated substitution. For example, the substitutions associated to the example of Figure 12 are:

$$\sigma_1 : \begin{cases} a & ad \\ b & b \\ c & c \\ d & d \end{cases} \mapsto \begin{cases} a & a \\ b & bd \\ c & c \\ d & d \end{cases}, \quad \sigma_2 : \begin{cases} a & a \\ b & bd \\ c & c \\ d & d \end{cases} \mapsto \begin{cases} a & a \\ b & b \\ c & c \\ d & cd \end{cases}, \quad \sigma_3 = \sigma_4 : \begin{cases} a & a \\ b & b \\ c & c \\ d & cd \end{cases} \mapsto \begin{cases} a & a \\ b & b \\ c & cd \\ d & d \end{cases}.$$

The composition of the substitutions gives the substitution associated to \mathfrak{R}^5 :

$$\sigma = \sigma_5 \circ \sigma_4 \circ \sigma_3 \circ \sigma_2 \circ \sigma_1 : \begin{cases} a & ad \\ b & bd \\ c & cccd \\ d & ccd \end{cases}.$$

Its associated matrix is

$$M_\sigma = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

See [Del16] for more details about the substitution associated to the Rauzy induction. We mention also that Definition 8 of [AI01] proposes a more general framework. For more details about the matrix of a substitution we refer to [Que10].

Proposition 3.2. *Let T be a self-similar IET on an interval I . Let q be the minimal integer such that $\mathcal{R}^q(T) = T$. Let σ be the substitution associated to \mathfrak{R}^q applied to T . We define $K = \max_{a \in \mathcal{A}} |\sigma(a)|$. Let $x \in I$. Let $\bar{x} = x_0 x_1 x_2 \dots$ be the symbolic coding of x under T . Let $n \in \mathbb{N}$.*

We can decompose the prefix of \bar{x} of length n as

$$\bar{x}_n = s_0 \sigma(s_1) \dots \sigma^{l-1}(s_{l-1}) \sigma^l(m_l) \sigma^{l-1}(p_{l-1}) \dots \sigma(p_1) p_0$$

where each word s_i or p_i is of length at most $K - 1$ and the length of m_l is between 1 and $2K - 2$.

Remark 3.3. *In this decomposition, l is as big as possible, the words s_i (resp. p_i) are suffixes (resp. prefixes) of words of $\sigma(\mathcal{A})$.*

For a proof of this proposition, see Lemma 5.5 in [Que10].

In terms of Rokhlin towers, this means that we consider towers of order l (apply the substitution once corresponds to apply the Rauzy induction once, and consider one order further on the Rokhlin tower). If the point x (whose symbolic coding begins with \bar{x}_n) belongs to the interval on which we induce, then all s_i are empty words. The orbit of x under T can be read on the Rokhlin tower, starting at the bottom of one tower. If not, it starts in the

middle of some tower, goes up until the top of this tower and then returns at the bottom of another (or maybe the same) tower (of order l). The length of the word m_l corresponds to the number of towers (with multiplicity) totally wandered by the first n steps of the orbit of x . If the orbit stops in the top of a tower, all p_i are empty words. Otherwise, the orbit continues to wander a part of a tower of rank l , and this part is the concatenation of $|p_{l-1}|$ towers of rank $l-1$, $|p_{l-2}|$ towers of rank $l-2$, ..., $|p_1|$ towers of rank 1 and finally $|p_0|$ towers of rank 0 (i.e. non-towers).

Bounding the number l of induction steps in terms of the size n of the orbit. We recall some classical facts.

Lemma 3.4. *Let σ be a primitive substitution on an alphabet \mathcal{A} and λ the Perron-Frobenius eigenvalue for the matrix M_σ of the substitution. There exist constants $A, B > 0$ such that for every letter x in \mathcal{A} and every $n \in \mathbb{N}$, $A\lambda^n < |\sigma^n(x)| < B\lambda^n$.*

This lemma results from basic properties of primitive substitutions that are explained in [Que10].

Corollary 3.5. *Let T be an IET on an interval I and $x \in I$. Let \bar{x} be the symbolic coding of x under T . Assume that $\mathcal{R}^q(T) = T$. Denote by λ the Perron-Frobenius eigenvalue of M_σ where σ is the substitution associated to \mathfrak{R}^q applied to T . There exists a constant $\kappa > 0$ such that the decomposition*

$$\bar{x}_n = s_0\sigma(s_1)\dots\sigma^{l-1}(s_{l-1})\sigma^l(m_l)\sigma^{l-1}(p_{l-1})\dots\sigma(p_1)p_0$$

verifies

$$\forall l \in \mathbb{N}, \left| l - \frac{\log(n)}{\log(\lambda)} \right| \leq \kappa.$$

Proof. Let A, B be the constants of Lemma 3.4, and $K = \max_{a \in \mathcal{A}} |\sigma(a)|$. We set $C = \max\left(A, \frac{(2K-2)B\lambda}{\lambda-1}\right)$ and $\kappa = \frac{\log C}{\log \lambda}$. Lemma 3.4 gives upper and lower bounds for $n = \sum_{k=0}^{l-1} (|\sigma^k(s_k)| + |\sigma^k(p_k)|) + |\sigma^l(m_l)|$, which permits to show the corollary. \square

4. ESTIMATION OF THE TRAJECTORY

4.1. Lengths of shifts. The assumption (C1) implies that one out of two crossed sides is the side a_N (the biggest one). Let us remark that every other polygon along the trajectory is a translation of the first one. Therefore we can approach two steps (i and $i+1$) of the trajectory by a translation: the one that sends the polygon P_i to the polygon P_{i+2} .

We identify the space of translation with \mathbb{C} . Let us denote by f_i the translation induced by crossing the side \mathbf{a}_i and then the side \mathbf{a}_N (see Figure 13). We recall that x_0 denotes the position of the vertex v_0 of P_0 in \mathcal{C}_0 . One has:

$$\forall k \in \{1, \dots, N-1\}, f_k = r(e^{X_k} - e^{X_N} + e^{X_{k-1}} - e^{X_0}),$$

where

$$X_k = -\frac{i}{r} \left(x_0 + \sum_{j=1}^k a_j \right) \quad \text{and} \quad r = \frac{1 + a_N}{2\pi} \text{ is the radius of the circumcircle of } P.$$




FIGURE 13. Translations f_i when crossing sides a_i followed by side a_5

4.2. Estimate the trajectory. Let us define the piecewise constant complex function f on the interval $[0, 1)$ by:

$$f(x) = f_i \quad \text{if} \quad \sum_{k=1}^{i-1} a_k \leq x < \sum_{k=1}^i a_k.$$

After $2n$ steps in the plane, the trajectory is in the polygon P_{2n} , polygon which is equal to the first one (P_0) translated by

$$S_n f(x_0) = \sum_{k=0}^{n-1} f(T^k(x_0)).$$

This sum is the *Birkhoff sum* of f with respect to T and x_0 . We approach the trajectory with this value. We highlight that we only need to know how many times the orbit $(T^k x)_{k \leq n}$ goes into each interval $I_i \subset (0, 1)$ to compute Birkhoff sum. The sequence $(S_n f(x_0))_{n \in \mathbb{N}}$ of Birkhoff sums gives us (with uniformly bounded error) an estimate of the position p_{2n} (of the $2n$ -th side crossing of the trajectory) and hence also of p_{2n+1} .

Note that f depends on x_0 . However, a change of x_0 will only multiply f by a complex number. Therefore all the translations f_i depend on x_0 (i.e. on the position of P_0) only up to the same rotation. Whether the trajectory has a limit or not hence does not depend on x_0 . Nor the fact that the trajectory stays at bounded distance of a line. Only the asymptotic direction (if it exists) depends on x_0 .

4.3. Average shift. We first show that the trajectory has almost always an asymptotic direction.

Lemma 1.2. *For every a_1, \dots, a_{N-1} , almost every $a_N > \sum_{i=1}^{N-1} a_i$, and every $\tau \in \left(\sum_{i=1}^{N-1} a_i, a_N \right)$, the cyclic N -gon $P(a_1, \dots, a_N)$ defines a generalized tiling billiard whose trajectories having parameter τ all admit a mean displacement $m = \lim_{n \rightarrow \infty} \frac{p_n}{n}$ that does not vanish.*

Proof. As soon as T is uniquely ergodic (which is the generic case [Mas82], [Vee82]), Birkhoff's theorem tells us that the main part of this sum will be n times the mean value $m = \sum_{k=1}^{N-1} a_k f_k$ of f :

$$\forall x_0 \quad S_n f(x_0) \underset{n \rightarrow \infty}{=} n \sum_{k=1}^{N-1} a_k f_k + o(n).$$

Hence, in terms of the trajectory, m corresponds to the asymptotic direction of the trajectory, as soon as it does not vanish. Let us show that this is almost always the case.

We have:

$$\begin{aligned} m &= \sum_{k=1}^{N-1} a_k f_k \\ &= r e^{\frac{i}{r} x_0} \left[\sum_{k=1}^{N-2} (a_k + a_{k+1}) e^{\frac{i}{r} \sum_{j=1}^k a_j} - \left(\sum_{k=1}^{N-2} a_k \right) e^{\frac{i}{r} \sum_{j=1}^{N-1} a_j} - \left(\sum_{k=2}^{N-1} a_k \right) \right]. \end{aligned}$$

Let us denote $s = \frac{2\pi}{1+a_N} = \frac{1}{r}$ the inverse of the radius of the circumcircles \mathcal{C}_i . If a_1, \dots, a_{N-1} are fixed, the map $s \in [0, \pi] \mapsto sm$ is analytic. As a non trivial sum of linearly independent functions ($s \mapsto e^{is \sum_{j=1}^k a_j}$ for $1 \leq k \leq N$), it cannot be everywhere equal to zero. Therefore the map has a finite number of zeros in the compact set $[0, \pi]$. We conclude that only a finite number of values of $a_N \in (1, +\infty)$ could make m vanish. \square

4.4. Normalized deviations. Our goal is to study the deviations from the mean displacement. We want to estimate their size and their direction, in particular we want to show that they are happening in the direction orthogonal to m . To do so, we renormalize the situation so that the trajectory goes asymptotically in the horizontal direction at unit speed, namely we rotate and contract (or expand) the plane by the complex coefficient $\frac{1}{m}$. Now we have to show that the deviations have an unbounded imaginary part.

Lemma 4.1. *We set $G = \frac{1}{m} \begin{pmatrix} f_1 \\ \vdots \\ f_{N-1} \end{pmatrix} - \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$ and $H = \text{Im}(G)$. Let $h : (0, 1) \rightarrow \mathbb{R}$*

be the map that is constant and equal to H_i on the subinterval $(\sum_{j=1}^i a_j, \sum_{j=1}^{i+1} a_j)$ for every $0 \leq i \leq N-1$. The following statements are equivalent:

- (1) *The trajectory in the plane admits unbounded deviations in the direction orthogonal to the asymptotic one.*
- (2) *The trajectory does not stay at a bounded distance from a line.*
- (3) *The sequence $(S_n(h))_{n \in \mathbb{N}}$ is not bounded.*

Moreover, if T is Osseledets-generic they are also equivalent to:

- (4) *The vector H is not in the contracted plane of the flag decomposition of \mathbb{R}^{N-1} given by the Oseledets' theorem.*

Proof. The first two statements are clearly equivalent.

Quantifying deviations consists in computing the second term of the asymptotic expansion of $S_n f(x_0)$, namely studying $S_n f(x_0) - nm$. We renormalize the situation by dividing by m : $\frac{1}{m} S_n f(x_0) - n$. This corresponds to contract the plane by a factor of $\frac{1}{m}$, the renormalized trajectory admits the real line as an asymptotic direction. Since we are interested in its

deviations from this asymptotic line, we want to compute the imaginary part of the second term in the asymptotic expansion of $S_n f(x_0)$:

$$\operatorname{Im} \left(\frac{1}{m} S_n f(x_0) - n \right) = S_n h(x_0).$$

The function h corresponds to the deviations from the renormalized mean displacement (equal to $(1, 0) \in \mathbb{R}^2$).

If H is in the contracted plane of the Oseledets' flag, then the sum converges (as a geometric sum), i.e. $S_n h(x_0)$ is bounded, and the trajectory stays at a bounded distance from the asymptotic line. On the contrary, if H is not in the contracted plane of the Oseledets' flag, then the sum diverges (see Proposition 5.1), i.e. $S_n h(x_0)$ is not bounded, and the trajectory does not stay at a bounded distance from the asymptotic line. \square

The following property of vector G will be useful.

Lemma 4.2. *The vector G is normal to the vector $V = \begin{pmatrix} a_1 \\ \vdots \\ a_{N-1} \end{pmatrix}$.*

Proof. This is a direct computation:

$${}^t V G = \frac{1}{m} \underbrace{\sum_{k=1}^{N-1} a_k f_k}_{=m} - \underbrace{\sum_{k=1}^{N-1} a_k}_{=1} = 0.$$

\square

We show in the next section that in the generic case we can apply Zorich's result on deviations for IETs. This gives Theorem B. We handle the non generic case of a self-similar IET in Section 6: one has $B_T^{(nq)} = \left(B_T^{(q)} \right)^n$. Direct computations on H and $M = B_T^{(q)}$ allow to prove Theorem C.

5. THE GENERIC CASE

We assume here that T is Oseledets-generic, which means (see Subsection 3.2) that there exists a flag depending on T

$$\mathcal{H}_1 \subset \mathcal{H}_2 \subset \dots \subset \mathcal{H}_{\mathbf{g}(T)} \subset \mathbb{R}^{N-1}$$

with $2\mathbf{g}(T) = \begin{cases} N-1 & \text{if } N \text{ is odd} \\ N-2 & \text{if } N \text{ is even} \end{cases}$, such that

$$\forall v \in \mathcal{H}_k \setminus \mathcal{H}_{k-1}, \frac{1}{n} \log \|B_T^{(n)}(v)\| \xrightarrow{n \rightarrow \infty} \theta_k$$

where the numbers $\theta_1 > \dots > \theta_{\mathbf{g}(T)}$ do not depend on T . The $1 + \frac{\theta_i}{\theta_1}$ are the Lyapunov exponents of the Teichmüller flow on the hyperelliptic stratum ($\mathcal{H}(N-3)$ if N is odd, (resp. $\mathcal{H}(\frac{N}{2}-2, \frac{N}{2}-2)$ if N is even) of the moduli space of abelian differentials. We will apply Proposition 3 of [Zor99]:

Proposition 5.1 (Zorich). *For a generic IET on $(0, 1)$, any point $x \in (0, 1)$, and any function $\varphi \in \text{Ann}(\mathcal{H}_p) \setminus \text{Ann}(\mathcal{H}_{p+1})$, $1 \leq p \leq \mathbf{g}(T) - 1$, the following holds:*

$$\limsup_{n \rightarrow +\infty} \frac{\log |S_n(\varphi(x))|}{\log n} = \frac{\theta_{p+1}}{\theta_1}.$$

We already know that the function h is in $\text{Ann}(\mathcal{H}_1)$ (because it has 0 mean). In this section we will show that it is not in $\text{Ann}(\mathcal{H}_3)$ for almost every choice of parameters (see Proposition 5.5). This will establish Theorems A and B.

To do that, we give an expression of H . It involves a matrix Q whose size increases with N : $Q \in \mathcal{M}_{\frac{(N-1)N}{2}, N-1}(\mathbb{R})$. This is why we do the proof first for $N = 5$ (in Sections 5.1 and 5.2) and then, with the very same proof but a bigger matrix, for $N > 5$ (in Section 5.3).

5.1. Expression of H , when $N = 5$. Let us compute

$$H = \text{Im} \left(\frac{1}{m} \begin{pmatrix} f_1 \\ f_2 \\ f_3 \\ f_4 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \right) = \frac{1}{|m|^2} \text{Im} \left(\bar{m} \begin{pmatrix} f_1 \\ f_2 \\ f_3 \\ f_4 \end{pmatrix} \right)$$

where \bar{m} denotes the complex conjugate of m . One can check that, with $s = \frac{2\pi}{1+a_5}$,

$$H = \frac{1}{|m|^2} Q \begin{pmatrix} \sin(a_1 s) \\ \sin(a_2 s) \\ \sin(a_3 s) \\ \sin(a_4 s) \\ \sin((a_1 + a_2) s) \\ \sin((a_2 + a_3) s) \\ \sin((a_3 + a_4) s) \\ \sin((a_1 + a_2 + a_3) s) \\ \sin((a_2 + a_3 + a_4) s) \\ \sin((a_1 + a_2 + a_3 + a_4) s) \end{pmatrix} = H(a_1, a_2, a_3, a_4, s)$$

where

$${}^t Q = \begin{pmatrix} a_2 + a_3 + a_4 & -a_1 + a_3 + a_4 & -(a_1 + a_2) & -(a_1 + a_2) \\ a_2 + a_3 & -a_1 + a_3 & -(a_1 + a_2) & 0 \\ 0 & a_3 + a_4 & -a_2 + a_4 & -(a_2 + a_3) \\ a_3 + a_4 & a_3 + a_4 & -(a_1 + a_2) + a_4 & -(a_1 + a_2 + a_3) \\ 0 & a_4 & a_4 & -(a_2 + a_3) \\ a_3 + a_4 & a_3 + a_4 & -(a_1 + a_2) & -(a_1 + a_2) \\ a_2 + a_3 & -a_1 & -a_1 & 0 \\ 0 & -(a_3 + a_4) & a_2 & a_2 \\ -a_3 & -a_3 & a_1 + a_2 & 0 \\ -(a_2 + a_3 + a_4) & a_1 - a_4 & a_1 - a_4 & a_1 + a_2 + a_3 \end{pmatrix}.$$

5.2. Vector H is almost never in a fixed plane, when $N = 5$. We show that H does not stay in a fixed plane of \mathbb{R}^4 when s varies.

Proposition 5.2. *Let \mathcal{P} be a plane in \mathbb{R}^4 . Let $a_1, a_2, a_3, a_4 \in \mathbb{R}^+$ summing up to 1. Then the set $\{s \in (0, \pi) \mid H(a_1, a_2, a_3, a_4, s) \in \mathcal{P}\}$ is finite.*

We first show:

Lemma 5.3. *The matrix Q has rank 3.*

Proof. Since the vector $V = {}^t(a_1 \ a_2 \ a_3 \ a_4)$ is normal to G , and $H = \text{Im}(G)$, the relation $\sum_{i=1}^4 a_i R_i(Q) = 0$ holds, where $R_i(Q)$ denotes the i -th row of Q (i.e. the i -th column of tQ). It means that Q has rank at most 3.

Moreover the minor of tQ formed by the first three columns and the fourth, fifth and eighth rows of tQ is:

$$\begin{vmatrix} a_3 + a_4 & a_3 + a_4 & -(a_1 + a_2) + a_4 \\ 0 & a_4 & a_4 \\ 0 & -(a_3 + a_4) & a_2 \end{vmatrix} = (a_3 + a_4)a_4(a_2 + a_3 + a_4) > 0.$$

The matrix Q has hence rank at least 3. \square

Proof of Proposition 5.2. The map

$$\mathbf{H} : \begin{array}{ccc} [0, \pi] & \longrightarrow & \mathbb{R}^4 \\ s & \longmapsto & H(a_1, a_2, a_3, a_4, s) \end{array}$$

is analytic. If $\mathbf{H}(s)$ were lying in \mathcal{P} for all s in a sub-interval J of $[0, \pi]$, then we would have two linearly independent relations

$$\begin{cases} \sum_i \mu_i \sin(\theta_i s) = 0 \\ \sum_i \lambda_i \sin(\theta_i s) = 0 \end{cases}$$

where the θ_i 's are sums of a_j with consecutive indices. Since G is normal to v , the first relation holds if $\mu = v$. But Q has rank 3 and $s \mapsto \sin((a_1 + a_2 + a_2 + a_3 + a_4)s) = \sin(s)$ is linearly independent of all other maps $s \mapsto \sin(\theta s)$ for $\theta \neq \pm 1$. Therefore the second relation cannot hold for all $s \in J$.

Moreover, the zeroes of the analytic map $s \in [0, \pi] \mapsto \sum_i \lambda_i \sin(\theta_i s)$ are discrete, and in finite number in the compact $[0, \pi]$. This proves Proposition 5.2 \square

5.3. For $N > 5$. As in the case $N = 5$, we consider H the imaginary part of G :

$$H = \text{Im} \left(\frac{1}{m} \begin{pmatrix} f_1 \\ \vdots \\ f_{N-1} \end{pmatrix} - \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right) = \frac{1}{|m|^2} \text{Im} \left(\bar{m} \begin{pmatrix} f_1 \\ \vdots \\ f_{N-1} \end{pmatrix} \right).$$

All coordinates of $|m|^2 H$ are of the form $\sum \lambda_i \sin(\theta_i s)$ where $s = \frac{1}{r}$, θ_i are sums of a_j with consecutive indices, and λ_i are sums of $\pm a_j$. We can write $|m|^2 H$ as $|m|^2 H = Q\Theta$ where the matrix $Q \in \mathcal{M}_{\frac{(N-1)N}{2}, N-1}(\mathbb{R})$ has all of its coefficients in the set $\{0, \lambda_i\}$ and the vector $\Theta \in \mathcal{M}_{\frac{(N-1)N}{2}, 1}$ has all of its coordinates of the form $\sin(\theta_i s)$. We order this coordinates by lexical order on (number of terms a_j within the sum $\theta_i = \sum_j a_j$; number of the first index

j), so that:

$$\Theta = \begin{pmatrix} \sin(a_1 s) \\ \vdots \\ \sin(a_{N-1} s) \\ \sin((a_1 + a_2)s) \\ \vdots \\ \sin((a_{N-2} + a_{N-1})s) \\ \sin((a_1 + a_2 + a_3)s) \\ \vdots \\ \sin((a_{N-3} + a_{N-2} + a_{N-1})s) \\ \vdots \\ \vdots \\ \sin((a_1 + a_2 + \dots + a_{N-2})s) \\ \sin((a_2 + a_3 + \dots + a_{N-1})s) \\ \sin((a_1 + a_2 + \dots + a_{N-1})s) \end{pmatrix}.$$

One can check that lots of coordinates of Q are zero. In the following we only need to consider a submatrix of Q .

Lemma 5.4. *The matrix $Q \in \mathcal{M}_{\frac{(N-1)N}{2}, N-1}(\mathbb{R})$ has rank at least $N - 3$, at most $N - 2$, and exactly $N - 2$ when N is odd.*

Proof. As in the case $N = 5$, G is normal to the vector $V = {}^t(a_1 \dots a_{N-1})$, which implies that Q has rank at most $N - 2$.

The submatrix of tQ made of the first $N - 1$ rows and $N - 2$ columns is:

$$\begin{pmatrix} \sum_{k=2}^{N-1} a_k & -a_1 + \sum_{k=3}^{N-1} a_k & -(a_1 + a_2) & \dots & -(a_1 + a_2) \\ \hline a_2 + a_3 & -a_1 + a_3 & -(a_1 + a_2) & 0 & \dots & 0 \\ 0 & a_3 + a_4 & -a_2 + a_4 & \dots & \dots & 0 \\ \vdots & \vdots & a_4 + a_5 & \dots & \dots & \vdots \\ 0 & \dots & \dots & -a_{N-4} + a_{N-2} & \dots & -a_{N-4} - a_{N-3} \\ \vdots & \vdots & \vdots & 0 & a_{N-2} + a_{N-1} & -a_{N-3} + a_{N-1} \\ \hline a_{N-2} + a_{N-1} & \dots & \dots & \dots & a_{N-2} + a_{N-1} & -(\sum_{k=2}^{N-3} a_k) + a_{N-1} \end{pmatrix}$$

Note that the framed submatrix formed by the rows $2, \dots, N - 2$ and first $N - 3$ columns of tQ is an invertible diagonal matrix, so Q has rank at least $N - 3$.

We show now that if N is odd, Q has rank at least $N - 2$. We replace the j -th column C_j of tQ by $C_j + \sum_{k=1}^{j-1} (-1)^{j-k} C_k$. The rows between 2 and $N - 1$ and the first $N - 2$ columns of this new matrix are equal to (changes are written in blue):

$$\left(\begin{array}{cccccccc} a_2 + a_3 & -a_1 - a_2 & 0 & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & a_3 + a_4 & -a_2 - a_3 & 0 & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & a_4 + a_5 & -a_3 - a_4 & \cdots & \cdots & \cdots & \vdots \\ 0 & \vdots & \vdots & -a_{N-4} - a_{N-3} & \cdots & \cdots & \cdots & 0 \\ a_{N-2} + a_{N-1} & 0 & a_{N-2} + a_{N-1} & 0 & a_{N-2} + a_{N-1} & -a_{N-3} - a_{N-2} & \cdots & -(\sum_{k=2}^{N-3} a_k) + a_{N-1} \end{array} \right)$$

We develop the determinant with respect to the last row of this submatrix and get:

$$\begin{aligned} & (a_{N-2} + a_{N-1}) [(-a_1 - a_2)(-a_2 - a_3) \cdots (-a_{N-3} - a_{N-2}) \\ & \quad + (a_2 + a_3)(a_3 + a_4)(-a_3 - a_4) \cdots (-a_{N-3} - a_{N-2}) \\ & \quad + (a_2 + a_3)(a_3 + a_4)(a_4 + a_5)(a_5 + a_6)(-a_5 - a_6) \cdots (-a_{N-3} - a_{N-2}) \\ & \quad + (a_2 + a_3)(a_3 + a_4) \cdots (a_7 + a_8)(-a_7 - a_8) \cdots (-a_{N-3} - a_{N-2}) \\ & \quad \vdots \\ & \quad + (a_2 + a_3)(a_3 + a_4) \cdots (a_{N-4} + a_{N-3})(-a_{N-3} - a_{N-2})] \\ & + \left(-\sum_{k=1}^{N-3} a_k + a_{N-1} \right) (a_2 + a_3)(a_3 + a_4) \cdots (a_{N-3} + a_{N-2})(a_{N-2} + a_{N-1}) \\ = & (a_{N-2} + a_{N-1}) \left[(-1)^{N-3} \prod_{k=1}^{N-3} (a_k + a_{k+1}) + \sum_{\substack{l=3 \\ l \text{ odd}}}^{N-3} \left((a_l + a_{l+1})(-1)^{N-3-l+1} \prod_{k=2}^{N-3} (a_k + a_{k+1}) \right) \right] \\ & + \left(-\sum_{k=1}^{N-3} a_k + a_{N-1} \right) \prod_{k=2}^{N-2} (a_k + a_{k+1}) \\ = & \prod_{k=2}^{N-2} (a_k + a_{k+1}) \left[\sum_{k=1}^{N-3} a_k - \sum_{k=1}^{N-3} a_k + a_{N-1} \right] \\ = & a_{N-1} \prod_{k=2}^{N-2} (a_k + a_{k+1}) \neq 0. \end{aligned}$$

We conclude that if N is odd, then Q has rank exactly $N - 2$. □

As for the case $N = 5$, the map

$$\mathbf{H}: \begin{array}{ccc} [0, \pi] & \longrightarrow & \mathbb{C}^{N-1} \\ s & \longmapsto & H(a_1, \dots, a_{N-1}, s) = Q\Theta \end{array}$$

is analytic. With Lemma 5.4, it allows us to prove the following proposition.

Proposition 5.5. *Let \mathcal{P} be a subspace of dimension $N - 3$ in \mathbb{C}^{N-1} . Let $a = (a_1, \dots, a_{N-1}) \in (\mathbb{R}_+^*)^{N-1}$ summing up to 1 and Oseledets generic. The set $\{s \in (0, \pi) \mid H(a, s) \in \mathcal{P}\}$ is finite. Moreover, if N is odd, it holds also for \mathcal{P} a subspace of dimension $N - 2$ in \mathbb{C}^{N-1} .*

5.4. **Conclusion.** As a corollary we get Theorems A and B.

Theorem A. *Let \mathcal{L} be a line in the asymptotic direction (almost always defined) of the trajectory. For almost every tiling billiard satisfying the conditions (C), the trajectory does not stay at a bounded distance from \mathcal{L} .*

Theorem B. *Let $N \in \mathbb{N}, N \geq 5$. There exist $\theta_1 > \theta_2 > \theta_3 > 0$ such that for almost every N -gon \mathbf{P} and every τ , both satisfying condition (C), for any initial direction having parameter τ in a \mathbf{P} -tiling billiard, the sequence of points $(p_n)_{n \in \mathbb{N}}$ satisfies:*

$$\limsup_{n \rightarrow \infty} \frac{\log |p_n - nm|}{\log n} = \begin{cases} \frac{\theta_2}{\theta_1} > 0 & \text{if } N \text{ is odd} \\ \frac{\theta_2}{\theta_1} \text{ or } \frac{\theta_3}{\theta_1} > 0 & \text{if } N \text{ is even} \end{cases},$$

where $m = \lim_{n \rightarrow \infty} \frac{p_n}{n}$.

Theorem A. *Let \mathcal{L} be a line in the asymptotic direction (almost always defined) of the trajectory. For almost every tiling billiard satisfying the conditions (C), the trajectory does not stay at a bounded distance from \mathcal{L} .*

Proof. Recall that the function g is in $\text{Ann}(\mathcal{H}_1)$ (because it has 0 mean). Let $\mathcal{P} = \text{Ann}(\mathcal{H}_3)$ (or $\text{Ann}(\mathcal{H}_2)$ if Q has rank $N - 2$). Proposition 5.5 tells that for almost every choice of parameters, h is not in \mathcal{P} . The proposition 5.1 applied to h hence establishes Theorem B:

$$\begin{aligned} |p_{2n} - 2nm| &= |S_n h(x_0)| + R_n \quad \text{where } R_n \text{ is uniformly bounded} \\ \limsup_{n \rightarrow +\infty} \frac{\log |p_{2n} - 2nm|}{\log(2n)} &= \limsup_{n \rightarrow +\infty} \frac{\log |S_n h(x_0)|}{\log(n)} \\ &= \frac{\theta}{\theta_2} \end{aligned}$$

where $\theta = \theta_2$ if N is odd, and $\theta \in \{\theta_2, \theta_3\}$ if N is even.

Moreover we have renormalized our problem to the case when the asymptotic direction is horizontal and we have studied the imaginary part of the deviations, this hence proves also Theorem A. \square

6. A SPECIAL CASE

The previous theorems tells nothing when our IET T is not Oseledets generic. The goal of this section is to study a non generic but important case for which we manage to do explicit computations, and prove Theorem C.

Let M be the matrix associated to a loop in the Rauzy diagram. Then M is positive. We assume that the loop is complex enough so that M is primitive. We assume that the second eigenvalue λ_2 of M is simple, that M has no other eigenvalue of the same modulus than λ_2 , and $|\lambda_2| > 1$. The article [Ham23] implies that this is the generic behavior. It follows that $\lambda \in \mathbb{R}$. We denote by V the Perron Frobenius vector for M , normalized so that its coordinates sum up to 1 and set $(a_1, \dots, a_{N-1}) = V$. The IET with lengths (a_1, \dots, a_{N-1}) is self similar and uniquely ergodic. Section 4.3 applies: the mean m of f (corresponding to the asymptotic

direction) does not vanish for almost every choice of $a_N > \sum_{k=1}^{N-1} a_k$. We will now compute the order of magnitude of deviations.

We denote by W_1 the Perron Frobenius eigenvector for tM (normalized so that its coordinates sum up to 1) and by W_2 its (normalized) eigenvector associated to the second eigenvalue λ_2 . Note that W_2 is a real vector since M and λ are real. We decompose $\mathbb{R}^{N-1} = \text{Vect}(W_1) \oplus \text{Vect}(W_2) \oplus E_3$ where E_3 is the direct sum of characteristic spaces of tM (associated to eigenvalues $\lambda_3, \dots, \lambda_d$). By assumption:

$$\forall v \in E_3, \frac{\|{}^tMv\|}{\|v\|} \leq |\lambda_3| < |\lambda_2|.$$

We consider cases where $a_N > \sum_{k=1}^{N-1} a_k$ is such that $H \notin E_3$. According to Proposition 5.5, this is all cases but finitely many ones.

6.1. Upper bound. If $w = w_0w_1\dots w_{n-1}$ is a finite word over the alphabet $\{1, \dots, N-1\}$, we set $e_w = \sum_{k=0}^{n-1} e_{w_k}$.

Let $x \in \mathbb{S}^1$ and let $\bar{x}_n = x_0x_1\dots x_{n-1}$ be the prefix of length n of its symbolic coding \bar{x} under T , decomposed as (see Proposition 3.2)

$$\bar{x}_n = s_0\sigma(s_1)\dots\sigma^{l-1}(s_{l-1})\sigma^l(m_l)\sigma^{l-1}(p_{l-1})\dots\sigma(p_1)p_0.$$

One has:

$$e_{\bar{x}_n} = \sum_{k=0}^{l-1} M^k e_{s_k} + M^l e_{m_l} + \sum_{k=0}^{l-1} M^k e_{p_k}.$$

The deviations from the mean displacement (normalized to be horizontal at unit-speed) are approximated by:

$$\begin{aligned} S_n h(x) &= \sum_{k=0}^{n-1} h(T^k(x)) \\ &= \langle H, e_{\bar{x}_n} \rangle \\ &= \sum_{k=0}^{l-1} \langle {}^tM^k H, e_{s_k} + e_{p_k} \rangle + \langle {}^tM^l H, e_{m_l} \rangle \end{aligned}$$

The vector $H = \text{Im}(G)$ is in the vector space $\text{Vect}(W_2) \oplus E_3$ because G is normal to the Perron-Frobenius vector V of M , and $H \notin E_3$. It can therefore be decomposed as $G_I = \alpha W_2 + U$ with $U \in E_3$ and $\alpha \in \mathbb{R}^*$. Then one has:

$$\begin{aligned} S_n h(x) &= \alpha \left(\sum_{k=0}^{l-1} \langle {}^tM^k W_2, e_{s_k} + e_{p_k} \rangle + \langle {}^tM^l W_2, e_{m_l} \rangle \right) \\ &\quad + \sum_{k=0}^{l-1} \langle {}^tM^k U, e_{s_k} + e_{p_k} \rangle + \langle {}^tM^l U, e_{m_l} \rangle. \end{aligned}$$

We write $P_1 = \alpha \left(\sum_{k=0}^{l-1} \langle {}^tM^k W_2, e_{s_k} + e_{p_k} \rangle + \langle {}^tM^l W_2, e_{m_l} \rangle \right)$ and $P_2 = \sum_{k=0}^{l-1} \langle {}^tM^k U, e_{s_k} + e_{p_k} \rangle + \langle {}^tM^l U, e_{m_l} \rangle$.

The principal part of the deviations is given by:

$$P_1 = \alpha \left(\sum_{k=0}^{l-1} \lambda_2^k \langle W_2, e_{s_k} + e_{p_k} \rangle + z \lambda_2^l \langle W_2, e_{m_l} \rangle \right)$$

whose norm is dominated by:

$$|\alpha| |\lambda_2|^l \frac{|\lambda_2|}{|\lambda_2| - 1} 2(K-1) \|W_2\|_\infty$$

where K is such that all s_k and p_k are at length at most $K-1$ and m_l at most $2K-2$ (see Proposition 3.2). The other part of the deviations, P_2 , whose norm is dominated by $|\lambda_3|^l \frac{|\lambda_3|}{|\lambda_3| - 1} 2(K-1) \|U\|_\infty$, has a smaller order. To summarize, we have, for l big enough:

$$|S_n g(x)| \leq D \lambda_2^l$$

with $D = \alpha \frac{\lambda_2}{\lambda_2 - 1} 2(K-1) \|W_2\|_\infty + 1$.

Finally, using Corollary 3.5, one gets, for n big enough:

$$|S_n g(x)| \leq D' n^{\frac{\log(\lambda_2)}{\log(\lambda_1)}}$$

with $D' = D \exp\left(\frac{\log(\lambda_2) \log(A)}{\log(\lambda_1)}\right)$. One can replace D' by a larger constant C_1 so that the inequality

$$|S_n g(x)| \leq C_1 n^{\frac{\log(\lambda_2)}{\log(\lambda_1)}}$$

is true for all $n \in \mathbb{N}$.

This shows the first inequality of Theorem C. We prove the second one in the following paragraph.

6.2. Lower bound. We want to build an increasing sequence $(n_l)_{l \in \mathbb{N}}$ of indices of "significant deviations", in other words such that

$$(\star) \quad \exists C_2, \forall l \geq l_0, |S_{n_l} h(x)| > C_2 n_l^{\frac{\log \lambda_2}{\log \lambda_1}}.$$

Let l be an integer. We decompose the symbolic coding of x as

$$\bar{x} = s_0 \sigma(s_1) \dots \sigma^{l-1}(s_{l-1}) \sigma^l(w)$$

where the s_i are finite words of length at most $K-1$ (see Proposition 3.2) and w is an infinite word. This corresponds to decompose the orbit of x with Rokhlin towers of order l .

Let us denote by π_j (for $1 \leq j \leq 2$) the projection onto $\text{Vect}(W_j)$ and π_3 the one onto E_3 with respect to the decomposition $\mathbb{R}^{N-1} = \text{Vect}(W_1) \oplus \text{Vect}(W_2) \oplus E_3$.

Let us denote $m_l = |s_0 \sigma(s_1) \dots \sigma^{l-1}(s_{l-1})|$, corresponding to the height of the partial Rokhlin tower before the orbit of x_0 arrives at the bottom of a tower of order l .

If $|S_{m_l} h(x)| > |\lambda_2|^l$, then we define $n_l = m_l$ and we have:

$$|S_{m_l} h(x)| > |\lambda_2|^l \geq D |\lambda_2|^{\frac{\log(n_l)}{\log(\lambda_1)}} = D n_l^{\frac{\log(|\lambda_2|)}{\log(\lambda_1)}}$$

with $D = \exp\left(\frac{-C}{\log(\lambda_1)}\right)$ (where C is the constant given by Corollary 3.5).

Else ($|S_{m_l} h(x)| \leq |\lambda_2|^l$). At least one vector e_k of the canonical basis of \mathbb{R}^{N-1} is such that $\pi_2(e_k) \neq 0$. Let i be such that $\pi_2(e_{w_i}) \neq 0$ and for every index $j < i$, $\pi_2(e_{w_j}) = 0$. The matrix M is primitive, so is the substitution σ . Hence there exists L (which depends only on M , not

on w) such that $i \leq L$. Let us denote $m'_i = |\sigma^l(w_1 \dots w_{i-1})|$ and $m''_i = |\sigma^l(w_i)|$. We define n_l as the length $|s_0 \sigma(s_1) \dots \sigma^{l-1}(s_{l-1}) \sigma^l(w_1 \dots w_i)| = m_l + m'_i + m''_i$. Then we have:

$$\begin{aligned} |S_{n_l} h(x)| &= |S_{m_l} h(x) + S_{m'_i} h(T^{m_l}(x)) + S_{m''_i} h(T^{m_l+m'_i}(x))| \\ &\geq |S_{m''_i} h(T^{m_l+m'_i}(x))| - |S_{m_l} h(x)| - |S_{m'_i} h(T^{m_l}(x))| \end{aligned}$$

By hypothese, one has

$$|S_{m_l} h(x)| \leq |\lambda_2|^l.$$

By definition of i , if we set $c = \max_{1 \leq k \leq N-1} |\pi_3(e_k)|$, one has

$$|S_{m'_i} h(T^{m_l}(x))| \leq c |\lambda_3|^i \leq c |\lambda_3|^L.$$

Finally

$$\begin{aligned} |S_{m''_i} h(T^{m_l+m'_i}(x))| &= \left| \langle {}^t M^l H, e_{w_i} \rangle \right| \\ &\geq |\alpha| \left| \lambda_2^l \langle W_2, e_{w_i} \rangle \right| - \left| \langle {}^t M^l U, e_{w_i} \rangle \right| \\ &\geq |\alpha| |\lambda_2|^l - c |\lambda_3|^l \|U\|. \end{aligned}$$

Putting these inequalities together we conclude that, for l big enough:

$$|S_{n_l} h(x)| \geq \frac{|\alpha|}{2} |\lambda_2|^l \geq \frac{|\alpha|}{2} D |\lambda_2|^{\frac{\log(n_l)}{\log(\lambda_1)}} = \frac{|\alpha|}{2} D n_l^{\frac{\log(|\lambda_2|)}{\log(\lambda_1)}}.$$

In the first case $|S_{m_l} h(x)| > |\lambda|^l$ so m_l goes to infinity as l increases and in the second case $n_l \geq m''_i \geq A \lambda_1^l$ (Lemma 3.4). Therefore the sequence $(n_l)_{l \in \mathbb{N}}$ goes to infinity. If we set

$$C_2 = \frac{|\alpha|}{2} D = \frac{|\alpha|}{4} \exp\left(\frac{-C}{\log(\lambda_1)}\right),$$

the inequality (\star) holds for $l \in \mathbb{N}$ big enough, which gives the second inequality of Theorem C.

7. CONCLUDING REMARKS

In this article we assumed that the N -gons are inscribed in a half-circle (that means a proportion of $\frac{N}{2^N}$ of the cyclic N -gons) and we studied trajectories passing near the circumcenter of the polygons. In this section, we discuss briefly the obstacles to remove such assumptions. We recall the key elements to prove Theorems A, B and C.

- (1) We can study the trajectory via an IETF Φ because the N -gon is cyclic.
- (2) The square Φ^2 of this map stabilizes a subinterval J , and Φ^2_J is an IET with $N - 1$ intervals of continuity. We do not need assumption to ensure that Φ^2 does not flip any interval, but we do need the assumption on the parameter τ of the trajectory to ensure that Φ^2_J has only $N - 1$ intervals of continuity.
- (3) We can define a function f of displacement, that is analytic.
- (4) We can show that for any fixed codimension 3 (or 2 if N is odd) subspace E of \mathbb{R}^{N-1} , the vector H , whose coordinates are the values of the function $h = \text{Im}(\frac{1}{m} f - 1)$, is not in E . This relies strongly on the fact that we have an explicit expression of H .

We could overcome point 2 by studying the Rauzy-Nogueira diagram of the underlying permutation of Φ . We would get many open sets of parameters, each one associated to a new map, that is a renormalization of Φ . So far, we can again define a function f of displacement and even ensure that it is analytic. But if we want to conclude, we need to have an explicit expression of the function f for each open set of parameters. These open sets depend on paths in the Rauzy-Nogueira diagram in question. The trouble is that this diagram, for $N = 5$, has more than 1 million of vertices and more than 3 millions of edges! Hence establishing the appropriate open sets seems out of reach without another idea, either about how we cut the set of parameters, or about how we show that H is not in E .

REFERENCES

- [Ada04] B. Adamczewski, “Symbolic discrepancy and self-similar dynamics,” *Ann. Inst. Fourier (Grenoble)*, vol. 54, no. 7, pp. 2201–2234, 2004.
- [AI01] P. Arnoux and S. Ito, “Pisot substitutions and Rauzy fractals,” in 2, vol. 8, Journées Montoises d’Informatique Théorique (Marne-la-Vallée, 2000), 2001, pp. 181–207.
- [BDFI24] P. Baird-Smith, D. Davis, E. Fromm, and S. Iyer, *Tiling billiards on triangle tilings, and interval exchange transformations*, 2024.
- [DDRS18] D. Davis, K. DiPietro, J. Rustad, and A. St Laurent, “Negative refraction and tiling billiards,” *Adv. Geom.*, vol. 18, no. 2, pp. 133–159, 2018.
- [DH18] D. Davis and W. P. Hooper, “Periodicity and ergodicity in the trihexagonal tiling,” *Comment. Math. Helv.*, vol. 93, no. 4, pp. 661–707, 2018.
- [Del16] V. Delecroix, *Interval exchange transformations*, 2016.
- [DHMPS] I. Dynnikov, P. Hubert, P. Mercat, O. Paris-Romaskevich, and A. Skripchenko, “Novikov gasket has zero lebesgue measure,” work in progress.
- [EKZ11] A. V. Eskin, M. Kontsevich, and A. Zorich, “Sum of lyapunov exponents of the hodge bundle with respect to the teichmüller geodesic flow,” *Publ. Math. Inst. Hautes Études Sci.*, vol. 120, pp. 207–333, 2011.
- [For02] G. Forni, “Deviation of ergodic averages for area-preserving flows on surfaces of higher genus,” *Ann. of Math. (2)*, vol. 155, no. 1, pp. 1–103, 2002.
- [Ham23] U. Hamenstädt, “Typical properties of periodic Teichmüller geodesics: Lyapunov exponents,” *Ergodic Theory Dynam. Systems*, vol. 43, no. 2, pp. 556–584, 2023.
- [HP22] P. Hubert and O. Paris-Romaskevich, “Triangle tiling billiards and the exceptional family of their escaping trajectories: Circumcenters and Rauzy gasket,” *Exp. Math.*, vol. 31, no. 1, pp. 58–87, 2022.
- [MMY05] S. Marmi, P. Moussa, and J.-C. Yoccoz, “The cohomological equation for roth-type interval exchange maps,” *J. Amer. Math. Soc.*, vol. 18, no. 4, pp. 823–872, 2005.
- [Mas82] H. Masur, “Interval exchange transformations and measured foliations,” *Ann. of Math. (2)*, vol. 115, no. 1, pp. 169–200, 1982.
- [Nog89] A. Nogueira, “Almost all interval exchange transformations with flips are nonergodic,” *Ergodic Theory Dyn. Syst.*, vol. 9, no. 3, pp. 515–525, 1989.
- [Par19] O. Paris-Romaskevich, “Trees and flowers on a billiard table,” preprint, Oct. 2019.
- [Par21] O. Paris-Romaskevich, “Tiling billiards and Dynnikov’s helicoid,” *Trans. Mosc. Math. Soc.*, vol. 2021, pp. 133–147, 2021.

- [Que10] M. Queffélec, “Dynamical systems arising from substitutions,” in *Substitution Dynamical Systems - Spectral Analysis*, 2nd ed. Berlin, Heidelberg: Springer Berlin Heidelberg, 2010, pp. 125–160.
- [Vee82] W. A. Veech, “Gauss measures for transformations on the space of interval exchange maps,” *Annals of Mathematics*, vol. 115, no. 2, pp. 201–242, 1982.
- [Via06] M. Viana, “Ergodic theory of interval exchange maps.,” *Rev. Mat. Univ. Complut. Madrid*, vol. 19, no. 1, pp. 7–100, 2006.
- [Yoc10] J.-C. Yoccoz, “Interval exchange maps and translation surfaces,” in *Homogeneous flows, moduli spaces and arithmetic. Proceedings of the Clay Mathematics Institute summer school, Centro di Ricerca Matematica Ennio De Giorgi, Pisa, Italy, June 11–July 6, 2007*, Providence, RI: American Mathematical Society (AMS); Cambridge, MA: Clay Mathematics Institute, 2010, pp. 1–69.
- [Zor96] A. Zorich, “Finite Gauss measure on the space of interval exchange transformations. Lyapunov exponents,” *Ann. Inst. Fourier (Grenoble)*, vol. 46, no. 2, pp. 325–370, 1996.
- [Zor99] A. Zorich, “How do the leaves of a closed 1-form wind around a surface?” In *Pseudoperiodic topology*, ser. Amer. Math. Soc. Transl. Ser. 2, vol. 197, Amer. Math. Soc., Providence, RI, 1999, pp. 135–178.

Magali Jay, AIX-MARSEILLE UNIVERSITÉ, I2M, MARSEILLE, FRANCE

E-mail address: `magali.jay@univ-amu.fr`