

**Insight into the wheat residues-derived adsorbents for the remediation of organic and  
inorganic aquatic contaminants: A review**

Muthanna J. Ahmed<sup>a\*</sup>, Ioannis Anastopoulos<sup>b</sup>, Dimitrios Kalderis<sup>c</sup>, Muhammad Haris<sup>d</sup>,

Muhammad Usman<sup>e</sup>

<sup>a</sup>Department of Chemical Engineering, College of Engineering, University of Baghdad, 10071  
Baghdad, Iraq

<sup>b</sup>Department of Agriculture, University of Ioannina, UoI Kostakii Campus, 47040 Arta, Greece

<sup>c</sup>Laboratory of Environmental Technologies and Applications, Department of Electronic  
Engineering, Hellenic Mediterranean University, Chania 73100, Greece

<sup>d</sup>School of Environmental Science and Engineering, Shaanxi University of Science &  
Technology, Xi'an 710021, P.R. China.

<sup>e</sup>Université de Rennes, École Nationale Supérieure de Chimie de Rennes, CNRS, UMR 6226, F-  
35000, Rennes, France.

\* Corresponding author. Tel.: +9647801642643

E-mail address: [muthanna.ja@gmail.com](mailto:muthanna.ja@gmail.com) (Muthanna J. Ahmed)

## **Supplementary Data**

**Table S1.** The names and forms of common adsorption models and error functions.

**Table S1**

Relation	Formula	Parameters definition	Ref.
Isotherm model			
Langmuir	$q_e = \frac{q_L K_L C_e}{1 + K_L C_e}$ $R_L = \frac{1}{1 + K_L C_o}$	$q_e$ (mg/g): Equilibrium adsorbed amount $C_e$ (mg/L) : Equilibrium residual amount $q_L$ (mg/g): Langmuir predicted adsorbed amount $K_L$ (L/mg): Langmuir affinity parameter $R_L$ : Favorability parameter $C_o$ (mg/L): Maximum inlet amount	(Langmuir, 1916)
Freundlich	$q_e = K_F C_e^{1/n}$	$K_F$ ((mg/g) (L/mg) <sup>1/n</sup> ): Adsorbed amount parameter 1/n: Adsorption intensity parameter	(Freundlich, 1906)
Temkin	$q_e = B_T \ln K_T C_e$	$B_T$ : Temkin constant related to adsorption heat $K_T$ (L/mg): Equilibrium binding constant	(Temkin & Pyzhev, 1940)
Dubinin– Radushkevich	$q_e = q_D e^{\left\{-K_D(RT \ln(1 + \frac{1}{C_e}))^2\right\}}$	$q_D$ (mg/g): D–R predicted adsorbed amount $K_D$ : D–R constant.	(Dubinin & Radushkevich, 1947)
Sips	$q_e = \frac{q_s K_s C_e^{1/m}}{1 + K_L C_e^{1/m}}$	$q_s$ (mg/g) : Sips predicted adsorbed amount $K_s$ (L/mg) <sup>1/m</sup> : Sips affinity parameter 1/m: Heterogeneity index	(Sips, 1948)
Kinetic model			
Pseudo-first Order (PFO)	$q_t = q_e (1 - e^{-k_1 t})$	$q_t$ (mg/g): Instant adsorbed amount $k_1$ (1/min): Apparent PFO rate constant	(Lagergren, 1898)
Pseudo-second Order (PSO)	$q_t = \frac{k_2 q_e^2 t}{1 + k_2 q_e t}$	$k_2$ (g/mg.min): Apparent PSO rate constant	(Ho & McKay, 1999)

Intraparticle diffusion	$q_t = k_{di} t^{1/2} + C_i$	$k_{di}$ (mg/g.min <sup>1/2</sup> ): Rate constant $C_i$ (mg/g): Adsorption degree parameter	(Weber & Morris, 1963)
Elovich	$q_t = \frac{1}{\beta} \ln(1 + \alpha \beta t)$	$\alpha$ (mg/g min): Primary rate of adsorption $\beta$ : Desorption parameter	(Roginsky & Zeldovich, 1934)
Thermodynamic models			
Van't Hoff	$\ln(K_d) = \frac{\Delta S^\circ}{R} - \frac{\Delta H^\circ}{RT}$	$K_d$ : Distribution coefficient for the adsorption. $\Delta S^\circ$ (J/mol.K): Change in entropy $\Delta H^\circ$ (J/mol): Change in enthalpy	(Sari & Tuzen, 2008)
Gibbs	$\Delta G^\circ = -RT \ln(K_d)$	$R$ (J/mol.K): Universal gas constant (8.314) $T$ (K): temperature $\Delta G^\circ$ (J/mol): Change in free energy	(Sari et al., 2008)
Error function			
Coefficient of Determination ( $R^2$ )	$R^2 = 1 - \frac{\sum_{i=1}^n (q_{e,exp} - q_{e,cal})^2}{\sum_{i=1}^n (q_{e,exp})^2 - [(\sum_{i=1}^n q_{e,exp})^2 / n]}$	$n$ : Number of applied data $q_{exp}$ and $q_{cal}$ (mg/g): Observed and predicted adsorbed amounts	(Boulinguez et al., 2008)
Chi-square statics ( $\chi^2$ )	$\chi^2 = \sum_{i=1}^n \frac{(q_{e,exp} - q_{e,cal})^2}{q_{e,cal}}$	$n$ : Number of applied data $q_{exp}$ and $q_{cal}$ (mg/g): Observed and predicted adsorbed amounts	(Boulinguez et al., 2008)

## References:

- Boulinguez B., Le Cloirec P., Wolbert D., 2008. Revisiting the determination of Langmuir parameters application to tetrahydrothiophene adsorption onto activated carbon. *Langmuir* 24, 6420–6424.
- Dubinin M.M. & Radushkevich L.V., 1947. The equation of the characteristic curve of the activated charcoal. *Proceedings of the Union of Soviet Socialist Republics Academy of Sciences* 55, 331–337.
- Freundlich H.M.F., 1906. Over the adsorption in solution. *The Journal of Physical Chemistry* 57, 385–471.
- Ho Y.S. & McKay G., 1999. Pseudo-second order model for sorption processes. *Process Biochem.* 34, 451–65.
- Langmuir I., 1916, The constitution and fundamental properties of solids and liquids. *J. Am. Chem. Soc.* 38, 2221–2295.
- Lagergren S., 1898. About the theory of so-called adsorption of soluble substances, *Kungliga Svenska Vetenskapsakademiens* 4, 1–39.
- Roginsky S. & Zeldovich Y.B., 1934. The catalytic oxidation of carbon monoxide on manganese dioxide. *Acta Physicochem. USSR* 1, 554–595.
- Sari A, Tuzen M, 2008. Biosorption of total chromium from aqueous solution by red algae (*Ceramium virgatum*): equilibrium, kinetic and thermodynamic studies. *Journal of Hazardous Materials* 160, 349–355.
- Sari A, Mendil D., Tuzen M., Soylak M., 2008. Biosorption of Cd(II) and Cr(III) from aqueous solution by moss (*Hylocomium splendens*) biomass: equilibrium, kinetic and thermodynamic studies. *Chemical Engineering Journal* 144, 1–9.

- Sips R., 1948. Combined form of Langmuir and Freundlich equations. *The Journal of Chemical Physics* 16, 490–495.
- Temkin M.I. & Pyzhev V., 1940. Kinetics of ammonia synthesis on promoted iron catalyst. *Acta Physicochimica USSR* 12, 327–356.
- Weber W.J. & Morris J.C., 1963. Kinetics of adsorption on carbon from solution. *Journal of the Sanitary Engineering Division* 89, 31–60.