



**HAL**  
open science

# Molecular mechanisms of cancer cachexia-related loss of skeletal muscle mass: data analysis from preclinical and clinical studies

Agnès Martin, Yann S Gallot, Damien Freyssenet

► **To cite this version:**

Agnès Martin, Yann S Gallot, Damien Freyssenet. Molecular mechanisms of cancer cachexia-related loss of skeletal muscle mass: data analysis from preclinical and clinical studies. *Journal of Cachexia, Sarcopenia and Muscle*, 2023, 14 (3), pp.1150-1167. 10.1002/jcsm.13073 . hal-04480183

**HAL Id: hal-04480183**

**<https://hal.science/hal-04480183>**

Submitted on 27 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Molecular mechanisms of cancer cachexia-related loss of skeletal muscle mass: data analysis from preclinical and clinical studies

Agnès Martin<sup>1,3</sup>, Yann S. Gallot<sup>2,3</sup> & Damien Freyssenet<sup>1\*</sup> 

<sup>1</sup>Laboratoire Interuniversitaire de Biologie de la Motricité EA 7424, Univ Lyon, Université Jean Monnet Saint-Etienne, Saint-Priest-en-Jarez, France; <sup>2</sup>LBEPS, Univ Evry, IRBA, Université Paris Saclay, Evry, France; <sup>3</sup>Share first authorship.

## Abstract

Cancer cachexia is a systemic hypoanabolic and catabolic syndrome that diminishes the quality of life of cancer patients, decreases the efficiency of therapeutic strategies and ultimately contributes to decrease their lifespan. The depletion of skeletal muscle compartment, which represents the primary site of protein loss during cancer cachexia, is of very poor prognostic in cancer patients. In this review, we provide an extensive and comparative analysis of the molecular mechanisms involved in the regulation of skeletal muscle mass in human cachectic cancer patients and in animal models of cancer cachexia. We summarize data from preclinical and clinical studies investigating how the protein turnover is regulated in cachectic skeletal muscle and question to what extent the transcriptional and translational capacities, as well as the proteolytic capacity (ubiquitin–proteasome system, autophagy–lysosome system and calpains) of skeletal muscle are involved in the cachectic syndrome in human and animals. We also wonder how regulatory mechanisms such as insulin/IGF1–AKT–mTOR pathway, endoplasmic reticulum stress and unfolded protein response, oxidative stress, inflammation (cytokines and downstream IL1 $\beta$ /TNF $\alpha$ –NF- $\kappa$ B and IL6–JAK–STAT3 pathways), TGF- $\beta$  signalling pathways (myostatin/activin A–SMAD2/3 and BMP–SMAD1/5/8 pathways), as well as glucocorticoid signalling, modulate skeletal muscle proteostasis in cachectic cancer patients and animals. Finally, a brief description of the effects of various therapeutic strategies in preclinical models is also provided. Differences in the molecular and biochemical responses of skeletal muscle to cancer cachexia between human and animals (protein turnover rates, regulation of ubiquitin–proteasome system and myostatin/activin A–SMAD2/3 signalling pathways) are highlighted and discussed. Identifying the various and intertwined mechanisms that are deregulated during cancer cachexia and understanding why they are decontrolled will provide therapeutic targets for the treatment of skeletal muscle wasting in cancer patients.

**Keywords** Autophagy–lysosome; Cancer cachexia; Glucocorticoids; Inflammation; Myostatin; Oxidative stress; Proteostasis; Skeletal muscle; Ubiquitin–proteasome

Received: 28 October 2021; Revised: 15 June 2022; Accepted: 14 August 2022

\*Correspondence to: Damien Freyssenet, Laboratoire Interuniversitaire de Biologie de la Motricité, Faculté de Médecine, 10 rue de la Marandière, 42 270 Saint Priest en Jarez, France. Email: [damien.freyssenet@univ-st-etienne.fr](mailto:damien.freyssenet@univ-st-etienne.fr)

## Introduction

Cancer cachexia is a systemic hypoanabolic and catabolic syndrome characterized by a progressive unintentional loss of body mass that cannot be reversed by nutritional support.<sup>1</sup> The prevalence of cachexia is quite variable, reaching up to 70% in pancreatic cancer patients.<sup>2</sup> Skeletal muscle represents the primary site of protein loss during cancer cachexia.<sup>3</sup> Image analysis indicates that muscle depletion varies from 7 to 30%,<sup>4,5,S1–S6</sup> (references S1 to S215 are listed as a supplementary reference list) the depletion worsening with the severity of the disease.<sup>5</sup> Another noteworthy feature is muscle weakness, which often may precede muscle loss<sup>S7,S8</sup> and which aggravates with cachexia.<sup>S7</sup> Besides muscle, the syndrome can also affect other tissues including bone, heart, liver and/or adipose tissue.<sup>6</sup> Overall, cancer cachexia increases the risk of surgical complications<sup>7,S9</sup> and chemotherapy toxicity<sup>S10–S13</sup> and lead to functional impairments, respiratory complications and fatigue that markedly reduce patients' quality of life and ultimately patients' survival. It has been estimated that in 2013, 15.8 subjects per 10 000 of the total population in European Union suffered from cancer cachexia.<sup>8</sup>


Over the past 15 years, considerable progress has been made in elucidating the molecular pathways involved in muscle mass loss, thus providing outstanding information about the pathophysiological mechanisms involved in cancer cachexia. However, our current knowledge comes mainly from animal models. Even though the number of human studies

exploring the biological mechanisms of muscle wasting is increasing, human studies remain scarce, thus raising the question of the translatability of animal findings to human clinical research. This is a critical point as therapeutic targets may differ between human and animals. Surprisingly, a detailed comparative analysis between cachectic human patients and cachectic cancer animals is missing.

This review provides a comprehensive analysis of the molecular mechanisms involved in cancer cachexia-associated muscle mass loss in preclinical and clinical studies. Article database of the US National Library of Medicine (PubMed) were searched using 'cancer cachexia' AND (muscle or 'skeletal muscle') AND specific terms (corresponding to the studied scientific area of each section). Any additional relevant literature was obtained from the reference lists of the published papers. This methodology has been applied for each section of the current review. Preclinical and clinical studies dealing with cancer, but for which cachexia was not characterized or demonstrated, were not included.

## Protein synthesis and degradation rates

Little information is presently available on muscle protein turnover in cachectic cancer patients (*Figure 1*). A decline in muscle protein synthesis rate has been reported in cachectic cancer patients compared with healthy subjects.<sup>9,10</sup>



**Figure 1** Relative variation in protein synthesis and degradation rates in clinical and preclinical studies. Variations have been calculated from data reported in quoted references. All data are expressed relative to controls or non-cachectic cancer patients (indicated by an asterisk).

Colorectal cancer patients with lower leg muscle mass than control subjects display unchanged muscle protein synthesis and a trend towards increased muscle protein breakdown, but reduced postprandial muscle protein synthesis,<sup>11</sup> which overall may confer a net catabolic status. Surprisingly, a lower protein degradation rate has been reported in weight-losing cancer patients than in healthy controls.<sup>12</sup> This was also accompanied by a blunted anabolic response to feeding.<sup>12</sup> Higher protein synthesis rates have even been documented in skeletal muscle of cachectic cancer patients,<sup>13,14</sup> suggesting the existence of a compensatory mechanism that may bridle the magnitude of muscle mass loss.

Animal studies clearly show a reduction in muscle protein synthesis<sup>15–21,S14–S17</sup> and an increase in protein breakdown,<sup>15,17–20,22–29,S14,S17–S21</sup> (Figure 1), the extent of the variation being associated with the severity of cachexia.<sup>17,18,S16</sup> Of note, unchanged muscle protein synthesis rates have also been frequently reported in animal experiments.<sup>22–26,S18,S19,S22</sup>

Therefore, currently available human data on muscle protein turnover in cachectic cancer patients are very limited and sometimes contradictory, even if a reduction in protein synthesis emerges as a mechanism that contributes to a net catabolic status. In animal models, a decrease in protein synthesis and an increase in protein degradation lower muscle mass. It is important to remind that cancer cachexia develops much more slowly in patients compared with animal models of cancer cachexia. Thus, it would be expected that protein degradation rate would be less easily identified as increased in human, compared with animals. More sensitive methods applied to the kinetic analysis of protein turnover during disease progression would be therefore necessary to provide a clear picture of skeletal muscle protein turnover in cachectic cancer patients.

## Gene expression capacity during cancer cachexia

Transcriptional and translational capacities are essential for skeletal muscle homeostasis. The transcriptional capacity is determined by the efficiency of the transcriptional machinery and the number of myonuclei, whereas the translational capacity depends on the ribosomal content and the global cellular RNA pool that is available to sustain the synthesis of myofibrillar protein.

### Myonuclear death

Although not systematically observed,<sup>523</sup> hallmarks of nuclear death by apoptosis have been reported in skeletal muscle of cachectic cancer patients (PARP and DNA fragmentation,<sup>524</sup>

increased BAX pro-apoptotic factor mRNA level,<sup>525</sup> increased p53 phosphorylation,<sup>30</sup> activation of caspase-8 and caspase-9<sup>30</sup>). Animal studies also show the presence of TUNEL-positive nuclei,<sup>27,31,S20,S26–S28</sup> an increase in PARP cleavage<sup>529</sup> and DNA fragmentation<sup>32,S30–S32</sup> in cachectic skeletal muscle. Increases in BAX pro-apoptotic-to-BCL2 anti-apoptotic protein and mRNA ratios<sup>S30,S33–S35</sup> and BAX mRNA level<sup>S36</sup> as well as increased expression<sup>S33,S34</sup> and cleavage<sup>S35</sup> of caspase-3, along with increased caspase-1, caspase-3, caspase-6, caspase-7, caspase-8 and caspase-9 activities,<sup>32,S29,S37</sup> have also been described. Accordingly, transcriptomic studies highlight heightened expression of apoptosis genes in muscle of cachectic cancer mice.<sup>33,34</sup>

As skeletal muscle is a very heterogeneous tissue, where approximately half of its nuclei reside outside of muscle fibres,<sup>S38</sup> it is essential to distinguish myonuclei from those of neighbouring mononuclear cells. Whereas most of studies presented above did not report the localization of nuclei,<sup>27,30,S20,S26,S28</sup> some animal studies detected TUNEL-positive nuclei outside (mononucleated cells)<sup>31,S27</sup> and inside<sup>S27</sup> the muscle fibre, suggesting the existence of myonuclear death. The multinucleated nature of the muscle fibre also implies that the death of a single myonucleus does not mean the destruction of the entire fibre. Accordingly, the whole number of fibres is maintained in skeletal muscle of human cachectic patients<sup>S3,S39</sup> and cancer mice.<sup>31</sup> Therefore, myonuclear death during cancer cachexia may trigger individual nuclei decay segmentally along the fibre, which may, over an extended period of time, locally weaken the transcriptional capacity of the fibre and contribute to atrophy and muscle dysfunction.

### Muscle fibre microenvironment

Pioneering studies at the beginning of the 20th century<sup>35</sup> and later<sup>S40</sup> reported an increased number of nuclei in the vicinity of the sarcolemma. More recently, inflammatory cells,<sup>36</sup> macrophages and fibro-adipogenic progenitors cells<sup>37</sup> have been identified in muscle of cachectic cancer patients. A higher number of activated satellite cells,<sup>38</sup> activated stem cells,<sup>38</sup> undifferentiated cells<sup>S41,S42</sup> and inflammatory cells<sup>27,39,40,S20,S26,S28</sup> have also been detailed in muscle of cachectic cancer mice.

The presence of inflammatory cells and macrophages, together with an augmented cytokine content (see below), suggests the persistence of unresolved inflammation in cachectic skeletal muscle, which may alter the properties of myogenic precursors.<sup>S43</sup> It has thus been shown that although progenitor cells do commit to the myogenic programme, they do not completely differentiate.<sup>38</sup> Muscle frailty may also cause localized episodes of muscle regeneration, as exhibited in cachectic cancer patients<sup>37,S44</sup> and in cancer mice,<sup>27,39,S20,S28</sup> and also illustrated by the presence of centralized

myonuclei.<sup>27,37,39,S20,S28,S44</sup> Furthermore, skeletal muscle of cachectic cancer mice regenerates less efficiently after freeze clamping-induced<sup>38,S45</sup> or cardiotoxin-induced<sup>S46</sup> muscle injury. Therefore, cancer cachexia is associated with an altered regenerative capacity of skeletal muscle that could lead to a reduction in the renewal of myonuclei by myogenic precursor cells, whose activity would be further impaired by unresolved inflammation, thus ultimately contributing to decrease the transcriptional capacity of the fibre.

### Ribosomal content

Reductions in skeletal muscle rRNA level,<sup>16,20</sup> 28S rRNA level<sup>S47</sup> and ribosomal S6 protein content<sup>41</sup> have been reported in skeletal muscle of tumour-bearing animals, along with defective transcriptional activity of polymerase I,<sup>S48</sup> the enzyme responsible for ribosomal gene transcription and a drop in ribosomal gene transcription.<sup>42</sup> Ribophagy (selective ribosome degradation by the autophagy–lysosome pathway) would be also increased.<sup>42</sup> Consequently, ribosomal gene transcription and protein machineries appear to be compromised in muscle during cancer cachexia. Clearly, the mechanisms regulating skeletal muscle ribosomal content warrants further research to determine the functional relevance of the translational capacity in controlling muscle wasting during cancer cachexia.

## Proteolytic capacity during cancer cachexia

### Ubiquitin–proteasome system


Skeletal muscle proteins targeted for degradation are marked by the 26S proteasome complex through an ATP-dependent ubiquitination process (Figure 2). The covalent attachment of a chain of ubiquitin molecules to the substrate protein involves a three-step enzymatic cascade driven by E1 (ubiquitin-activating enzymes), E2 (ubiquitin-conjugating enzymes) and E3 (ubiquitin-ligase enzymes). Ubiquitinated proteins are then docked to the proteasome for degradation.<sup>S49</sup> The activation of the ubiquitin–proteasome system has largely been inferred from increased expression of the E3 ligases, MuRF1 (*TRIM63*) and MAFbx/Atrogin1 (*FBXO32*) in multiple atrophy conditions.<sup>S50</sup> MuRF1 targets sarcomeric proteins (actin, myosin heavy chain, troponin) for degradation,<sup>S51–S53</sup> whereas MAFbx targets MyoD and the translational enhancer EIF3F for degradation.<sup>S54,S55</sup>

Although some studies show increased transcript and protein levels of MuRF1,<sup>36,43,S37,S56,S57</sup> and MAFbx,<sup>36,S37,S57</sup> in skeletal muscle of cachectic cancer patients, a majority of investigations report unchanged expression of

MuRF1,<sup>11,44–50,S37,S58</sup> and MAFbx,<sup>11,43–45,47–50,S37,S58</sup> in muscle of cachectic cancer patients compared with non-cachectic patients or healthy subjects (Figure 2). A decrease in MAFbx expression has even been documented.<sup>46</sup> By contrast, increased ubiquitin mRNA<sup>S1,52,S59,S60</sup> and protein<sup>37,52</sup> levels, ubiquitinated proteins<sup>36,S56</sup> and mRNA and protein levels of proteasome subunits<sup>11,53</sup> have been frequently reported in skeletal muscle of cancer patients, even if some discrepancies still persist.<sup>36,50,54,S61</sup> The activity of the ubiquitin–proteasome system also correlates with the severity of body mass loss.<sup>S59</sup> Finally, transcriptomic analyses provide divergent results with either increased,<sup>S62</sup> unchanged<sup>S58</sup> or decreased<sup>45</sup> expression of genes related to the ubiquitin–proteasome system in muscle of cachectic cancer patients.

Animal models consistently highlight an augmentation in the mRNA level of MAFbx,<sup>18,21,41,43,44,55–77,S17,S26,S33,S34,S57,S63–S87</sup> and MuRF1,<sup>18,21,41,43,44,56–62,64,65,67–69,71–76,78,79,S17,S26,S33,S34,S57,S64–S68,S70–S83,S85,S87–S90</sup> in skeletal muscle of cachectic cancer animals (Figure 2). This has also been confirmed at the protein level.<sup>18,80–83,S20,S42,S57,S69,S82,S85,S91–S94</sup> Ubiquitin mRNA<sup>23,24,32,56,59,62,73,75,84,85,S17–S19,S57,S71,S72,S95–S98</sup> and protein<sup>82,S86</sup> levels, the content of ubiquitinated proteins<sup>18,21,27,29,59,75,76,82,83,S20,S26,S66,S67,S86,S99</sup> and the transcript<sup>18,20,24,28,32,62,73,85,S17–S20,S26,S95,S96</sup> and protein<sup>S100,S101</sup> levels of several proteasome subunits are also increased in muscle of cachectic cancer mice. Transcriptomic studies also feature an up-regulation of genes related to the ubiquitin–proteasome system in muscle of cachectic cancer animals.<sup>33,34,55,73,78,86–88,S70,S72,S92,S102,S103</sup> Consistent with these observations, proteasome enzyme activity is elevated in muscle of cachectic cancer mice.<sup>28,89,S75,S100,S104</sup>

Regardless of a lesser number of investigations in human cancer patients, the analysis of MuRF1 and MAFbx expression presented above suggests that the ubiquitin–proteasome system does not contribute to muscle mass loss in cancer patients, which is in line with the observation that there is currently no clear evidence of an increase in muscle proteolysis rate in cachectic cancer patients. This would thus give prominence to a major interspecies difference. However, a careful examination of the data reveals that whereas the picture is effectively contrasted when looking at MuRF1 and MAFbx, a much more nuanced picture appears when looking at ubiquitin mRNA level, proteasome subunit mRNA level, ubiquitinated proteins and ubiquitin–proteasome activity (Figure 2). Therefore, when considering all markers of the ubiquitin–proteasome system, and not solely MuRF1 and MAFbx, currently available data rather indicate that the ubiquitin–proteasome system would be activated in human cachectic cancer patients. One may also consider the function of other E3 ligases such as MUSA1/Fbxo30<sup>S105</sup> and SMART/Fbxo21,<sup>S106</sup> all required for skeletal muscle atrophy. Although one study reports a decrease in MUSA1 transcript level in muscle of tumour-bearing mice,<sup>S66</sup> other studies show that MUSA1<sup>21,43,71,73</sup> and SMART<sup>43</sup> transcript levels are increased.



**Figure 2** Comparative analysis of the transcript and protein levels of components of the ubiquitin–proteasome system in cachectic skeletal muscle of cancer patients and animals. Proteins are targeted for degradation by the 26S proteasome through covalent attachment of a chain of ubiquitin molecules. The E1 ubiquitin-activating enzyme hydrolyses ATP to bind ubiquitin. E2 ubiquitin-conjugating enzymes receive ubiquitin from E1 and brings it to the E3 ubiquitin-ligase enzymes, which catalyse the transfer of the ubiquitin from E2 to the substrate. This reaction is the rate-limiting step of the ubiquitination process. The ubiquitinated protein is then docked to the proteasome for degradation. One gene encodes the E1 enzyme, whereas one hundred genes encode the E2 enzymes and almost one thousand genes the E3 enzymes. Significant variations are reported in red (increase) or blue (decrease). Unchanged levels are reported in white.

Expression of MUSA1 is also heightened in skeletal muscle of cachectic cancer patients.<sup>43</sup> Expression of Fbxo31, a novel E3 ligase that is induced during denervation and fasting,<sup>5106</sup> is also elevated in muscle of cachectic cancer mice.<sup>71,73</sup> It now remains to determine which substrates are targeted for degradation by these E3 ligases and to what extent they are associated with the regulation of the ubiquitin–proteasome

system. Finally, the kinetic of cancer cachexia is very different between patients and animals. In tumour-bearing mice, the subcutaneous injection of cancer cells induces a rapid and violent tumour burden in a couple of days/weeks and a fast and important depletion of muscle compartment. This allows for a well-controlled and accurate kinetic analysis of the ubiquitin–proteasome system. This is also true for genetic models

of cancer cachexia, even if they display a greater heterogeneity in disease progression and appearance of cachexia-associated symptoms. By contrast, the complexity of the clinical context in cancer patients, the difficulty of determining the onset of the tumour surge and the onset of cachexia and the lower rate of skeletal muscle depletion render the kinetic analysis of the ubiquitin–proteasome system during the cachectic syndrome particularly challenging.

### Autophagy–lysosome system

The autophagy–lysosome system is responsible for eliminating long-lived proteins and large supramolecular structures, including dysfunctional mitochondria.<sup>549</sup> Proteins and organelles to be degraded are engulfed during the formation of a double-membrane structure called the autophagosome, which then fuse with lysosomes allowing acidic proteolytic degradation of their contents by cathepsins. Because autophagy is constantly active to remove damaged proteins and organelles, a defect in autophagy will result in muscle functional impairment,<sup>5107</sup> but excessive autophagy will also contribute to muscle mass loss.<sup>5108,5109</sup> Therefore, a tight regulation of the autophagy–lysosome system is necessary for skeletal muscle homeostasis.

The protein level of autophagy-related genes such as *ATG5*, *ATG7*, *Beclin1* and *LC3B* is increased in skeletal muscle of cachectic cancer patients,<sup>30,47,68,90,539,556</sup> together with the number of autophagosomes,<sup>556</sup> suggesting an increase in autophagosome formation. The number of autophagosomes results from a dynamic equilibrium between autophagosome formation and autophagosome docking and fusion with lysosomes. Therefore, an accumulation of autophagosomes can be also interpreted as a default in autophagosome clearance. The multidomain protein p62/SQSTM1 is a cargo adaptor protein involved in selectively targeting protein aggregates to autophagosomes.<sup>5110</sup> Because p62 is constantly removed by autophagy, a rise in p62 protein content is a good marker of disturbances in autophagosome turnover. p62 protein content<sup>68,90,556</sup> and p62 aggregates<sup>37</sup> are increased in skeletal muscle of cachectic cancer patients, an observation that is consistent with a disrupted clearance of autophagosomes and suggests that defects in lysosomal rejuvenation/biogenesis and proteolytic capacity could be involved in impaired autophagosome turnover. Nevertheless, the mRNA levels of *TFEB*,<sup>90</sup> a master regulator of lysosome biogenesis,<sup>5111</sup> and cathepsin D<sup>54</sup> are unchanged in muscle of cachectic cancer patients, whereas the mRNA level of cathepsin B<sup>54</sup> is even increased. Although very limited, these data suggest that disturbances in autophagosome turnover may primarily be due to induction of pathways that promote autophagosome formation that are not properly matched to those promoting autophagosome clearance. Of note, some studies do not report

any difference in the mRNA<sup>44,45,49</sup> and protein<sup>36</sup> levels of autophagy markers nor in autophagosome number,<sup>36</sup> suggesting that autophagy can be also properly balanced in muscle of cachectic cancer patients.

Expression of autophagy-related genes is increased in skeletal muscle of cachectic cancer animals.<sup>27,41,44,67,68,72,73,77,79–81,91,526,528,566,578,588,5112</sup> This has also been confirmed by transcriptomic analysis.<sup>73,86,88,570,572</sup> Cathepsin expression<sup>20,32,62,69,77,79,92,517,581,595</sup> and activity<sup>25,26,29,595</sup> as well as lysosomal proteolysis,<sup>18,29,77,595</sup> are also increased. Together, these findings indicate that autophagy and the clearance of autophagosomes are activated in skeletal muscle of cachectic cancer mice. However, one should note that some studies mention either unchanged<sup>24</sup> or even decreased<sup>77,84</sup> cathepsin activity and unchanged lysosomal proteolysis<sup>24,89</sup> in muscle of cachectic cancer animals. Furthermore, an accumulation of p62 in muscle of cachectic cancer animals,<sup>68,77,80,91,526,5112</sup> with an increase in autophagosome number,<sup>68</sup> suggests a disequilibrium between autophagosome formation and clearance.

The E3 ligase TRAF6 mediates the conjugation of Lys63-linked polyubiquitin chains to ULK1 and BECLIN1,<sup>5113</sup> a post-translational modification that regulates their activity. TRAF6 expression is up-regulated in skeletal muscle of cancer patients<sup>52</sup> and in tumour-bearing mice.<sup>566,588</sup> Muscle-restricted ablation of *TRAF6* also preserves muscle mass in cancer mice<sup>588</sup> and restores LC3B and BECLIN1 expression to control levels,<sup>588</sup> suggesting an important role for TRAF6 during cancer cachexia. To what extent TRAF6 may impact the proteolytic capacity of the autophagy–lysosome system is currently unknown.

Animal studies showed marked alterations in mitochondrial function and network in cachectic muscle.<sup>93</sup> Therefore, the role of mitophagy in controlling mitochondriostasis needs to be questioned. The core mitophagy process is led by the kinase PINK1, which phosphorylates the E3 ligase PARKIN,<sup>5113</sup> whereas other proteins (BNIP3, NIX) are involved in ubiquitin-independent mitophagy.<sup>5113</sup> The protein level of PINK1 and PARKIN is either unchanged<sup>90</sup> or decreased<sup>5114</sup> in skeletal muscle of cachectic cancer patients, whereas the mRNA and protein levels of BNIP3 and NIX are unchanged.<sup>90</sup> In muscle of cachectic cancer animals, PINK1 protein level is either unchanged<sup>94</sup> or increased,<sup>91</sup> whereas that of PARKIN remains unchanged.<sup>94</sup> Consequently, human and animal data may suggest that mitophagy would be either unchanged or decreased. Mitochondrial fusion and fission proteins are essential in regulating organelle dynamic. The expression of fusion proteins is diminished (MFN2)<sup>5114</sup> or unchanged (OPA1),<sup>30,5114</sup> in muscle of cachectic cancer patients, whereas the mRNA level encoding the fission protein FIS1 is augmented,<sup>30,5114</sup> Expression of mitochondrial fusion proteins (MFN1, MFN2, OPA1) is unchanged<sup>79,94</sup> or reduced<sup>95,536</sup> in skeletal muscle of cachectic cancer mice, whereas expression of FIS1 is unchanged<sup>94,578</sup> or elevated.<sup>94,95,536</sup> Therefore, the

fission of mitochondria would be favoured in cachectic muscle, which, together with unchanged or decreased mitophagy, would lead to an accumulation of dysfunctional organelles. Clearly, the data currently available suggest that our view of the mechanisms involved in mitophagy during cancer cachexia is still fragmented and needs further exploration.

### Calpains

The calpain family consists of a group of 15 calcium-activated cysteine proteases.<sup>5115</sup> The ubiquitously expressed calpain-1 and calpain-2 as well as the muscle-specific isoform calpain-3 are expressed in skeletal muscle. Calpains have originally been described as facilitators of protein turnover by releasing myofilaments from myofibrils.<sup>5115,5116</sup>

In human, one study demonstrates increased calpain activity in skeletal muscle of non-weight-losing cancer patients,<sup>5117</sup> whereas another one does not show any difference.<sup>5118</sup> Although former experiments indicated that calcium-dependent proteolysis would not be involved in muscle proteolysis in cachectic cancer animals,<sup>20,24</sup> a more recent examination shows that calcium-dependent proteolysis is activated in tumour-bearing mice.<sup>521</sup> Regarding calpain activity *per se*, investigations show unchanged<sup>23,575,5119</sup> or increased activity<sup>533,5100,5104</sup> in muscle of cachectic cancer mice. This last result agrees with reports showing heightened expression of calpain isoforms<sup>24,27,32,62,517,521,571,572,574</sup> and decreased expression<sup>521,533</sup> and activity<sup>5119</sup> of calpastatin, the endogenous specific inhibitor of calpain. Therefore, considering the major impact of cancer cachexia on skeletal muscle structure and the remodelling function attributed to calpains,<sup>5115</sup> a deeper understanding of calpain function during cancer cachexia would be beneficial.

## Insulin/IGF1–AKT–mTOR pathway

### Insulin and IGF1

Insulin and IGF1 activate a cascade of phosphorylation that coordinately regulate protein synthesis and degradation (Figure 3). IGF1 is mainly produced by the liver under the control of pituitary-secreted growth hormone and accounts for 70–80% of serum IGF1.<sup>5120</sup> However, liver IGF1 only contributes approximately to 30% of adult body size,<sup>5121</sup> indicating that IGF1 originating from other tissues, including skeletal muscle, also contributes to the regulation of tissue growth. Among the different IGF1 isoforms expressed in skeletal muscle, IGF1-Ea, and to a lesser extent IGF1-Eb (IGF1-Ec in human), are the most powerful in increasing muscle mass and force in mice.<sup>5122</sup>

Data show that the driving force of the pathway is reduced in cachectic muscle. Circulating IGF1<sup>5123</sup> and muscle IGF1-Ec transcript level<sup>46</sup> are decreased in cachectic cancer patients. The circulating level of insulin,<sup>26,83,5124</sup> and IGF1,<sup>56,76,83,564</sup> as well as muscle IGF1-Ea<sup>56,563</sup> and IGF1-Eb,<sup>18,55,515,594</sup> transcript levels, and muscle IGF1 protein content<sup>564</sup> are also decreased in cachectic cancer animals.


### The pathway

Insulin/IGF1 signalling impinges a crucial regulatory step in the pathway, that is, the reaction catalysed by the serine/threonine protein kinase AKT. Under its phosphorylated active form, AKT indirectly activates the kinase mTOR. mTOR and then signals to the translation machinery via the activation of S6K, which controls the ribosomal protein S6, and the inhibition of 4EBP1, which hinders translation initiation by sequestering EIF4E.<sup>5125</sup> AKT also phosphorylates and inhibits GSK3, thus relieving GSK3-dependent inhibition of EIF2B.<sup>5126</sup>

AKT protein level<sup>50,96</sup> and the phosphorylation of GSK3,<sup>96</sup> mTOR<sup>96</sup> and S6K<sup>50,96</sup> are decreased in skeletal muscle of cachectic cancer patients compared with non-cachectic cancer patients, along with the phosphorylated inactive form of the translational repressor 4EBP1<sup>50</sup> (Figure 3). Transcriptomic analyses also show that the expression of genes involved in protein anabolism is down-regulated<sup>45,5127</sup> or altered<sup>562</sup> in skeletal muscle of cachectic cancer patients. Therefore, both transcriptional and post-translational events regulate the pathway. In animal models, a large majority of studies reports an inhibition of the pathway, as shown by the decrease in the phosphorylation of AKT,<sup>526,564,565,591</sup> mTOR,<sup>18,515,520,591,5100,5128</sup> S6K,<sup>18,21,41,82,515,520,566,591,5128</sup> S6,<sup>21,41,82,566,594</sup> and 4EBP1.<sup>18,41,515,594,5100,5128</sup> These events are also associated with weight loss,<sup>18,515,5128</sup> attesting that the greater is the inhibition of the pathway, the greater is the extent of cachexia. Microarray analyses also highlight a transcriptional regulation of the pathway in skeletal muscle.<sup>33</sup> Consequently, evidence based on global transcript and biochemical analyses indicate that the activity of the insulin/IGF1–AKT–mTOR pathway is reduced in cachectic muscle of cancer patients and cancer animals.

However, some discrepancies exist between studies both in human and animals. Some human investigations reveal unchanged<sup>11,30,36,49,50</sup> or even increased AKT phosphorylation<sup>47</sup> in skeletal muscle of cachectic cancer patients, together with unchanged phosphorylation of GSK3,<sup>47,50</sup> mTOR,<sup>47,50</sup> S6K<sup>11,47,50</sup> and 4EBP1<sup>11,47</sup> (Figure 3). Similarly, AKT phosphorylation remains unchanged,<sup>57,58,520,563,566,567</sup> or even increased,<sup>18,97,594</sup> in muscle of cachectic cancer animals. A few works also describe unchanged phosphorylation of S6K,<sup>39,58</sup> S6<sup>97</sup> and GSK3,<sup>568</sup> and sometimes even increased phosphorylation of S6K,<sup>57,563</sup> 4EBP1<sup>591</sup> and





**Figure 3** Comparative analysis of the regulation of the insulin/IGF1–AKT–mTOR pathway in clinical and preclinical studies. Upon receptor activation, IRS1 promotes the phosphorylation of phosphatidylinositol 4,5-bisphosphate into phosphatidylinositol 3,4,5-triphosphate at the plasma membrane by recruiting the kinase PI3K. Phospholipid phosphorylation promotes AKT recruitment and activation by PDK1. AKT positively or negatively regulates multiple targets including mTOR from the mTORC1 complex (not represented) and FOXO transcription factors. AMPK, whose activity is increased by energy stress, is another important modulator of the pathway. Black arrows indicate post-translational regulation. Red arrows indicate transcriptional regulation. Significant variations are reported in red (increase) or blue (decrease). Unchanged levels are reported in white.

GSK3.<sup>563</sup> Divergences between studies are important to consider. First, an activation of the pathway, which is at first counter-intuitive, could be interpreted as a compensatory response aimed at limiting the magnitude of muscle mass loss. Second, these results also strongly suggest that the pathway can be turned on or off depending on the clinical/experimental context (cancer type, severity of the disease and progression through the disease) and that a temporal regulation of the pathway may occur during the time course of the disease.

Is the insulin/IGF1–AKT–mTOR pathway constantly repressed to maintain the driving force of cancer cachexia, or are there episodes of reactivation of the pathway that succeed/alternate to episodes of repression? The data presented above hint at the second possibility would be an interesting alternative to consider. Finally, these data also suggest that regulatory influences coming not only from insulin/IGF1 but from other sources may converge to the pathway to modulate its activity.<sup>S125</sup> For instance, the accumulation of amino

acids released from catabolized proteins may mitigate the down-regulation of mTOR.<sup>5129</sup> Conversely, AMPK, a metabolic sensor that down-regulates mTOR signaling,<sup>5130</sup> is activated in muscle of cachectic cancer mice<sup>18,97,515,566,594</sup> and may thus further exacerbate mTOR inhibition.

### FOXO transcription factors

AKT inhibits the FOXO family of transcription factors by phosphorylation.<sup>5131</sup> FOXO regulate the expression of MuRF1 and MAFbx, as well as that of genes of the autophagy-lysosome system.<sup>5106,5108,5109,5132,5133</sup> AKT thus allows a coordinated regulation of protein synthesis and degradation.

FOXO-related genes are up-regulated in skeletal muscle of cachectic cancer patients,<sup>37</sup> and the transcript levels of FOXO1 and FOXO3 negatively correlate with body mass and muscle mass.<sup>98</sup> The ratio of the phosphorylated-to-total forms of FOXO1<sup>36</sup> and FOXO3<sup>36,96</sup> is diminished in skeletal muscle of cachectic cancer patients (*Figure 3*), suggesting heightened FOXO nuclear translocation and transcriptional activity. Importantly, the reduction in FOXO1 and FOXO3 phosphorylation reported in Puig-Vilanova et al.<sup>36</sup> was accompanied by an increase in MuRF1 and MAFbx protein content, which is consistent with a FOXO-dependent regulation of these E3 ligase. Similarly, unchanged transcript level of MuRF1 and MAFbx was associated with unchanged phosphorylated-to-total forms of FOXO1 and FOXO3 in muscle of cachectic cancer patients.<sup>47</sup>

In animal models of cancer cachexia, FOXO3,<sup>18,75,515,526,594</sup> phosphorylation is lowered, which agrees with increased FOXO3 nuclear localization<sup>83</sup> and transcriptional activity<sup>69</sup> reported by other studies (*Figure 3*). Furthermore, microarray analysis of FOXO-regulated transcripts<sup>34</sup> and motif analysis of promoter sequences<sup>570</sup> identify FOXO as a transcription factor involved in muscle atrophy during cancer cachexia. This concurs with the observation showing that blocking FOXO prevents muscle fibre atrophy and spares force deficits in tumour-bearing mice.<sup>34</sup> Nevertheless, a promoter analysis applied to skeletal muscle of cancer patients shows that weight-loss associated genes have only fewer FOXO binding sites,<sup>558</sup> highlighting that the molecular characteristics of skeletal muscle from cachectic cancer patients may be different from those of preclinical models.

Finally, some animal studies report unchanged FOXO1<sup>526</sup> and FOXO3<sup>41</sup> phosphorylation, whereas others even report increased FOXO1<sup>27,563</sup> phosphorylation and decreased FOXO1 DNA binding activity.<sup>563</sup> This may be interpreted as a compensatory mechanism to limit the extent of muscle mass loss. These data also remind us that additional FOXO-regulatory mechanisms exist, including phosphorylation by FOXO-activating kinases, such as AMPK, acetylation, ubiquitination and methylation.<sup>5131</sup> This adds another level of com-

plexity to the FOXO code and may dictate the transcription of different FOXO target genes.

## Endoplasmic reticulum stress and unfolded protein response

The endoplasmic reticulum (ER) is an organelle involved in the folding, maturation and trafficking of newly synthesized proteins.<sup>5134,5135</sup> ER stress, which leads to an abnormal accumulation of unfolded or misfolded proteins, is perceived by three key transducers of the unfolded protein response (UPR). ATF6, IRE1 $\alpha$  and PERK act as a surveillance system to relieve ER stress and regulate proteostasis.<sup>5136</sup> Although adaptive UPR contributes to skeletal muscle homeostasis, its prolonged or exacerbated activation leads to muscle atrophy.<sup>5137</sup>

The expression of several ER stress markers (GRP78, HSP60, HSP70, calnexin and calreticulin mRNA; PDIA3 and PI3K mRNA and protein) and the UPR (ATF6, XBP1 and CHOP mRNA and protein) is up-regulated in skeletal muscle of cachectic cancer patients,<sup>537</sup> as well as the ratio of the phosphorylated-to-total forms of PERK and eIF2 $\alpha$ .<sup>537</sup> Together, these data indicate the existence of an ER stress and the activation of the UPR in muscle of cachectic cancer patients. Nevertheless, supplementary investigations are needed to obtain reliable clinical data regarding the implication of ER stress and the UPR in cancer cachexia. In animal models of cancer cachexia, the induction of ER stress and all three arms of the UPR has been demonstrated.<sup>566</sup> These findings were recently corroborated by several animal studies,<sup>5138–5141</sup> thus providing robust evidence of ER stress and UPR activation during cancer cachexia. One should note, however, that a study mentions unchanged expression of multiple markers of ER stress and the UPR.<sup>5112</sup> This also agrees with repressed phosphorylation of eIF2 $\alpha$  reported in skeletal muscle of cachectic cancer rodents.<sup>563</sup>

A relevant question raised by these data is the function of ER stress and UPR activation during cancer cachexia. Under experimental ER stress, PERK-mediated eIF2 $\alpha$  phosphorylation contributes to stoppage of translation.<sup>5142</sup> The UPR represses the insulin/IGF1–AKT–mTOR pathway and activates autophagy.<sup>5143</sup> Furthermore, ATF4 transcription factor (PERK arm), whose expression is increased in cachectic skeletal muscle,<sup>78</sup> promotes muscle atrophy by modulating a subset of atrogenes like Gadd45 $\alpha$ , which induces the remodelling of chromatin to repress genes involved in anabolic signalling and to activate pro-atrophy genes.<sup>5144</sup> During cancer cachexia, the activation of ER stress and the UPR is associated with a noticeable inhibition of the insulin/IGF1–AKT–mTOR pathway and increased AMPK activity.<sup>566</sup> Therefore, an interplay between ER stress, the UPR, the insulin/IGF1–AKT–mTOR pathway, AMPK and the

expression of atrogenes may contribute to down-regulate protein synthesis and activate proteolysis. Intriguingly, the use of 4-phenylbutyrate (ER stress pan-inhibitor) or the targeted ablation of PERK in skeletal muscle aggravates the deleterious effects of cancer cachexia in tumour-bearing mice,<sup>566,5138</sup> suggesting that ER stress and the UPR (PERK arm) could be also viewed as a response originally aimed at limiting the effects of cancer cachexia.

## Oxidative stress

Oxidative stress (OxS) is due to an excessive production of reactive oxygen species (ROS) and reactive nitrogen species, together with impaired antioxidant defence, which results in an accumulation of oxidized and damaged proteins, organelles, membranes and DNA.<sup>5145</sup> In skeletal muscle, OxS reduces muscle strength and triggers atrophy.<sup>5146</sup>

OxS is increased in skeletal muscle of cachectic cancer patients, as shown by the elevation in total protein carbonylation<sup>36</sup> and malondialdehyde-protein adducts,<sup>36,5147</sup> and the reduced expression of genes encoding antioxidant proteins,<sup>525</sup> such as superoxide dismutase 2, glutamate-cysteine ligase (involved in glutathione synthesis) and Nrf2 (a transcription factor that regulates the expression of multiple cytoprotective genes). However, an increase in the protein level and activity of the antioxidant enzymes, superoxide dismutase 1 and 2, has also been reported in skeletal muscle of cachectic cancer patients,<sup>36</sup> which may illustrate the induction of a compensatory mechanism to alleviate increased ROS production.

OxS is also increased in skeletal muscle of cachectic cancer mice, as indicated by higher levels of ROS,<sup>91,94,575,5148</sup> lipid peroxidation<sup>40,63,99,100,575,5149</sup> and protein carbonylation,<sup>27,40,63,99,100,520,567,582,591,5150</sup> as well as by the increase in oxidized-to-reduced glutathione ratio,<sup>63,91,574,575,5112,5149,5151</sup> the decrease in glutathione peroxidase activity<sup>5148</sup> and the depletion of reduced glutathione and antioxidant peptides.<sup>5152</sup> The expression of antioxidant enzymes has been documented to be either decreased<sup>27,55,520,528,575,5148</sup> unchanged<sup>27,91,94,95,99,100,528,567,5148</sup> or even increased,<sup>40,55,91,100,528,567,574</sup> suggesting that the main factor responsible for augmenting OxS in muscle of cachectic cancer rodents is an increased ROS production. Transcriptomic analyses in mice also highlight increased transcriptional response of OxS-related genes.<sup>33,55,78</sup> Importantly, OxS would be more significant in Type II myofibres<sup>100</sup> and would also precede muscle mass loss,<sup>94</sup> suggesting that OxS could be a precocious event. Altogether, these data indicate that cachectic skeletal muscle is subjected to a chronic OxS that mainly results from increased ROS production associated or not with lowered antioxidant defence.

This obviously raises the question of the origin of ROS and the mechanisms linking OxS to muscle wasting during cancer cachexia. The main sources of ROS in skeletal muscle are complexes I and III of the mitochondrial respiratory chain and NADH oxidases, but also ER and peroxisomes where enzymatic complexes generate ROS.<sup>5145</sup> Sources of ROS are probably multiple during cancer cachexia, but previously reported alterations in muscle mitochondrial metabolism of cachectic cancer mice,<sup>79,93,101,5153</sup> should significantly contribute to ROS production by increasing leakage of electrons and the subsequent formation of superoxide anion. OxS integrates signal transduction pathways by regulating post-translational modifications of proteins through the redox regulation of the thiol side chain of cysteine amino acids.<sup>5154</sup> Redox proteomic studies have revealed that several hundred proteins contain reactive and potentially modulatory cysteine residues,<sup>5155</sup> including transcription factors (FOXO and NF- $\kappa$ B) and kinases (AMPK and mTOR), which functional implication in the loss of muscle mass during cancer cachexia we have demonstrated. Furthermore, the wide spectrum of action of OxS on other biomolecules (lipid and DNA) also suggests that OxS may target other pathways that contribute to muscle wasting during cancer cachexia.

## Inflammatory cytokines and downstream pathways

### *Circulating and skeletal muscle inflammatory cytokines*

Systemic inflammation is a well-described feature of cancer cachexia. Increased circulating level of C-reactive protein (CRP) is associated with weight loss in cancer patients.<sup>4,47,51,102,103,54,5156,5157</sup> The circulating levels of pro-inflammatory cytokines such as IL1 $\beta$ ,<sup>5158</sup> IL4,<sup>104</sup> IL6,<sup>47,50,51,104,105,5159–5163</sup> IL8,<sup>47,104,105,5160,5162,5163</sup> IL10,<sup>105</sup> IFN $\gamma$ <sup>104</sup> and TNF $\alpha$ <sup>104,5158,5162,5164</sup> are also elevated in cachectic cancer patients. Similar observations have been done in mice models of cancer cachexia for the circulating levels of CRP,<sup>591</sup> IL1 $\beta$ ,<sup>75,582,591</sup> IL6,<sup>18,39,43,55,72,74,75,80,87,95,105,106,58,515,542,573,582,585,5165–5167</sup> IL10,<sup>55</sup> IL11,<sup>87</sup> IFN $\gamma$ ,<sup>55,80,5168</sup> TNF $\alpha$ <sup>22,55,75,80,87,519,582,585,591,5124,5167</sup> and Tweak.<sup>591</sup>

The shock wave of this systemic inflammatory response is largely perceived by skeletal muscle. A large majority of human studies shows increased protein contents of IL1 $\beta$ ,<sup>36</sup> IL6,<sup>50,5169</sup> IFN $\gamma$ ,<sup>36</sup> TNF $\alpha$ <sup>5169</sup> and Tweak<sup>591</sup> in skeletal muscle of cachectic cancer patients. Only one work reports unchanged IL6 and TNF $\alpha$  protein levels.<sup>36</sup> IL1 $\beta$ ,<sup>592</sup> IL6,<sup>564,585</sup> IFN $\gamma$ <sup>27</sup> and TNF $\alpha$ <sup>528,564,585</sup> protein levels are also augmented in muscle of cachectic cancer mice, whereas one report shows unchanged IL1 $\beta$ , IL6 and TNF $\alpha$  protein levels.<sup>27</sup> One

may also consider that skeletal muscle itself can produce cytokines. However, a majority of investigations reports unchanged transcript levels (IL4,<sup>S170</sup> IL6<sup>S25</sup> and TNF $\alpha$ <sup>S0S25</sup>) in muscle of cachectic cancer patients, whereas one study reports an increase in IL6 mRNA level.<sup>50</sup> The picture is more contrasted in animal models of cancer cachexia with a fairly equivalent number of experiments describing increased (IL1 $\beta$ ,<sup>S8,S92</sup> IL6,<sup>21,43,58,106,S8,S74,S77,S92,S94</sup> and TNF $\alpha$ <sup>S92</sup>) or unchanged (IL6,<sup>95,S67,S69</sup> TNF $\alpha$ <sup>58,S67,S74</sup> and Tweak<sup>41</sup>) mRNA levels in muscle of cachectic cancer mice. Therefore, these data suggest that although non-muscle tissues (host immune system, tumour cells) seem to mainly contribute to skeletal muscle inflammation, it could also be partly supported by an increase in cytokine production by skeletal muscle. Corollary, the contribution of skeletal muscle fibre, resident and/or recruited mononucleated cells in producing cytokines needs to be evaluated.

### IL1 $\beta$ /TNF $\alpha$ -NF- $\kappa$ B pathway

Omic studies in human<sup>S62,S127</sup> and animals<sup>33,34,55,87,88,S92,S103,S171</sup> indicate that cachectic skeletal muscle is subjected to a persistent activation of a pro-inflammatory cytokine signalling. NF- $\kappa$ B transcription factor relays information from various cytokines (mainly IL1 $\beta$  and TNF $\alpha$ ). Five NF- $\kappa$ B transcription factors [p65 (Rel A), Rel B, c-Rel, p52, p50] are expressed in skeletal muscle.<sup>S172</sup> Activation of cytosolic NF- $\kappa$ B occurs when the I $\kappa$ B kinase phosphorylates I $\kappa$ B, resulting in its ubiquitination and proteasomal degradation. This allows NF- $\kappa$ B to translocate into the nucleus and regulate the expression of target genes. NF- $\kappa$ B activation in mouse skeletal muscle overexpressing I $\kappa$ B kinase leads to severe muscle wasting associated with increased MuRF1 expression and ubiquitin-proteasome proteolysis.<sup>S173</sup> Here, we will mainly discuss about the canonical p65 pathway, whose activation is associated with pathological conditions.

The phosphorylation level of NF- $\kappa$ Bp65 is similar in skeletal muscle of cachectic cancer patients compared with non-cachectic patients.<sup>S39</sup> However, mRNA level of NF- $\kappa$ B1 (precursor of p50)<sup>50</sup> and NF- $\kappa$ Bp65 protein content<sup>36,S169</sup> are increased compared with healthy controls, suggesting a precocious activation of NF- $\kappa$ B pathway before the development of cancer cachexia. A majority of animal experiments reports an increase in the phosphorylated active form of NF- $\kappa$ B,<sup>27,57,74,82,97,S8,S20,S26,S67,S94</sup> the phosphorylated inactive form of I $\kappa$ B,<sup>41</sup> the nuclear localization of NF- $\kappa$ B,<sup>S92</sup> NF- $\kappa$ B DNA binding<sup>S88,S89,S92,S101,S173</sup> and transcriptional activity<sup>27</sup> in cachectic muscle. The phosphorylated active form of NF- $\kappa$ Bp65 also negatively correlates with body mass loss and muscle force in cachectic cancer mice.<sup>S8</sup> Although still debated,<sup>88</sup> motif analysis of promoter sequences also identified NF- $\kappa$ B as a transcription factor involved in muscle atrophy during cancer cachexia.<sup>S70</sup> Of note, studies report un-

changed NF- $\kappa$ Bp65 phosphorylation<sup>41,88,S75</sup> and NF- $\kappa$ B DNA binding activity.<sup>32,88,S98,S174</sup>

### IL6-JAK-STAT3 pathway

The JAK-STAT pathway is activated by IFN $\alpha$ , IFN $\beta$ , IFN $\gamma$ , IL2 and IL6.<sup>S175</sup> Upon cytokine binding, activated JAK tyrosine kinase phosphorylates STAT proteins, which translocate into the nucleus to regulate the expression of target genes.<sup>S176</sup> STAT transcriptional activation contributes to muscle wasting by indirect activation of myostatin expression and expression of MAFbx and MuRF1.<sup>S177,S178</sup>


Limited information in patients indicate that the phosphorylation level of STAT3 is similar in cachectic compared with non-cachectic muscle.<sup>S39</sup> By contrast, numerous evidence of IL6-JAK-STAT3 pathway activation have been reported in preclinical models. Circulating IL6 level correlates with the development of cachexia in tumour-bearing mice.<sup>95,S166</sup> STAT3 activation<sup>74,82,95,97,106-108,S8,S15,S69,S82,S94</sup> and the expression of STAT3 target genes<sup>87</sup> are increased in skeletal muscle of cachectic cancer mice. The active form of STAT3 negatively correlates with body mass and muscle force,<sup>S8</sup> and media conditioned with serum of tumour-bearing mice activate STAT3 in C2C12 myotubes.<sup>S178</sup> *In silico* analysis of transcriptome data also reveals STAT3 as a transcription factor involved in muscle atrophy during cancer cachexia.<sup>S70</sup> Finally, hyperactivation of STAT3 in cancer mice exacerbates weight and muscle mass losses compared with control cancer mice.<sup>109</sup> Interestingly, recent results show that the IL6-JAK-STAT3 pathway contributes to the regulation of Noggin, an inhibitor of the anti-catabolic BMP-SMAD1/5/8 signalling pathway<sup>43</sup> (see below).

## TGF- $\beta$ signalling pathways

### Myostatin-SMAD2/3 signalling pathway

Myostatin is a TGF- $\beta$  superfamily member acting as a master negative regulator of skeletal muscle growth during embryonic and postnatal development in animals<sup>S179,S180</sup> and human<sup>S181</sup> and also in adulthood.<sup>110</sup> Myostatin binds to activin type IIB receptors (ActRIIB and ActRIIA) leading to the recruitment of TGF- $\beta$  Type I receptors (ALK4 and ALK5)<sup>S182</sup> and activation of SMAD2 and SMAD3 proteins (Figure 4). Once activated by phosphorylation, SMAD2/3 recruit SMAD4 to regulate the transcription of target genes.<sup>S183</sup> Myostatin also inhibits the insulin/IGF1-AKT-mTOR pathway.<sup>111</sup>

Surprisingly, myostatin circulating level is reduced in cachectic cancer patients compared with healthy controls<sup>S184,S185</sup> and decreased<sup>4</sup> or unchanged<sup>S185</sup> when compared with non-cachectic cancer patients (Figure 4). Myostatin mRNA



**Figure 4** Comparative analysis of the regulation of myostatin/activin A signalling in cachectic skeletal muscle of cancer patients and in animal models of cancer cachexia. Myostatin and activin A bind to activin Type II receptors (ActRIIB/IIA) that activate Type I receptors (ALK4/5/7), which phosphorylate and induce SMAD2/3 to form a complex with SMAD4 and translocate into the nucleus. Myostatin and activin A binding is modulated by the inhibitory action of follistatin. Significant variations are reported in red (increase) or blue (decrease). Unchanged levels are reported in white.

level is also unchanged<sup>11,47,48,50</sup> or diminished<sup>46</sup> in skeletal muscle of cachectic cancer patients. Accordingly, myostatin protein level,<sup>36</sup> as well as the phosphorylation of SMAD2 and SMAD3,<sup>47,S39</sup> is unchanged in muscle of cachectic cancer patients. Circulating<sup>4</sup> and muscle transcript<sup>48</sup> levels of follistatin, an inhibitor of myostatin and activin signalling, are also unchanged. Therefore, these studies indicate that myostatin signalling is not activated in skeletal muscle of cachectic cancer patients, even if this conclusion must be nuanced in the light of investigations showing that genes related to TGF- $\beta$  signalling<sup>37</sup> are up-regulated in skeletal muscle of cachectic cancer patients and that the transcript level of ActRIIB negatively correlates with muscle mass in cancer patients.<sup>98</sup>

How to reconcile the fact that myostatin is a master negative regulator of muscle mass in human<sup>S181</sup> and the observa-

tion that myostatin expression and signalling are not activated in muscle of cachectic cancer patients? One may first consider that as myostatin is produced by skeletal muscle, myostatin circulating level can thus simply be lowered as a consequence of the reduction in muscle mass during cancer cachexia. Furthermore, the data described above do not exclude the possibility that myostatin expression had increased earlier during the disease when muscle mass had not started to decrease yet. In support of this hypothesis, a strict temporal regulation of myostatin expression has been demonstrated in mice models of atrophy.<sup>112,S186</sup> Kinetic analysis of its expression during the time course of cachexia would allow to answer this question.

Animal studies show that the circulating level of myostatin is either elevated<sup>S91</sup> or unchanged<sup>75,S187</sup> (Figure 4). Although some studies report unchanged myostatin expression

in skeletal muscle of cachectic cancer rodents,<sup>21,41,65,80,S80,S97</sup> a majority of works report an increase in both myostatin mRNA<sup>61,62,75,113,S80,S187,S188</sup> and protein<sup>27,S26,S68,S91,S187,S188</sup> levels. Downstream, the mRNA level of ActRIIB<sup>S97</sup> and the phosphorylation of SMAD2<sup>75,80</sup> and SMAD3<sup>74,S91</sup> are also increased in muscle of cachectic cancer mice. Muscle follistatin mRNA level is either unchanged<sup>113,S68</sup> or decreased,<sup>S80,S97</sup> which is also consistent with increased myostatin activity. However, a reduction in ActRIIB mRNA level,<sup>41</sup> unchanged ALK4/5 mRNA level<sup>41</sup> and SMAD2/3 phosphorylation<sup>41,80,106</sup> have also been reported in tumour-bearing mice. Consequently, and even if this cannot be generalized to all studies, myostatin signalling is commonly activated during cancer cachexia in rodents.

### Activin A–SMAD2/3 signalling pathway

Activin A is another TGF- $\beta$  family member that binds to ActRIIA and ActRIIB and activate SMAD2/3 signalling with comparable potencies and efficacy as myostatin does.<sup>S189</sup> Increasing circulating activin A levels in mice increases E3 ligase expression, inhibits the insulin/IGF1–AKT–mTOR–pathway and reduces muscle mass and function.<sup>S190</sup>

Activin A circulating level is consistently elevated in cachectic cancer patients,<sup>4,43,104,105,114,S185,S191</sup> as well as in cachectic cancer mice<sup>43,74,75,80,105,106,115,116,S192</sup> (Figure 4). By contrast, studies indicate that skeletal muscle activin A mRNA level is either decreased<sup>21,106,S192</sup> or unchanged,<sup>41,43</sup> whereas activin A protein level is increased.<sup>S192</sup> Therefore, circulating activin A may thus come from another source than skeletal muscle, in particular tumour cells.<sup>21,75,116,S76,S192–S194</sup> Importantly, while myostatin circulating level is higher in mouse than in human<sup>S189</sup> and is the main negative regulator of muscle mass in mouse,<sup>S87</sup> activin A circulating level is higher in human than in mouse.<sup>S189</sup> Therefore, activin A could play a more prominent role in cachectic cancer patients than myostatin.

### BMP–SMAD1/5/8 signalling pathway

BMP (BMP7, BMP13 and BMP14) bind to BMP Type II receptor (BMPRII), ActRIIB or ActRIIA and promote the recruitment and activation of Type I receptors BMPRIA (ALK3), BMPRIIB (ALK6) or ActRIA (ALK2). This triggers the phosphorylation of SMAD1, SMAD5 and SMAD8, which, together with SMAD4, regulate the expression of target genes. As SMAD4 is shared by SMAD1/5/8 and SMAD2/3, BMP–SMAD1/5/8 and myostatin–SMAD2/3 pathways operate in parallel, and in opposition. Increased BMP–SMAD1/5/8 pathway activity in muscle induces hypertrophy,<sup>S105,S195</sup> whereas its inhibition causes muscle atrophy<sup>S105,S195</sup> and abolishes the hypertrophic phe-

notype of myostatin-deficient mice,<sup>S105</sup> indicating that this pathway is dominant over myostatin signalling.

Diminished BMP signalling and augmented expression of the BMP inhibitor Noggin are observed in skeletal muscle of cancer patients and mildly cachectic mice.<sup>43</sup> Importantly, both IL6 and activin A trigger the expression of Noggin,<sup>43</sup> a BMP–SMAD1/5/8 pathway inhibitor. BMP signalling inhibition is also associated with neuromuscular junction impairment.<sup>43</sup> Therefore, perturbed BMP signalling appears to be a critical pathogenic mechanism regulating muscle mass and function in cancer patients and animals.

## Glucocorticoid signalling

Glucocorticoids are steroid hormones produced by the adrenal glands under the control of the hypothalamic–pituitary axis. The hypothalamus secretes CRH, which stimulates the secretion of ACTH by the anterior pituitary gland. ACTH then binds its receptor on the adrenal cortex to activate the biosynthesis and release of glucocorticoids. Glucocorticoids bind the nuclear receptor NR3C1 of target tissues to activate or inhibit the transcription of multiple target genes.<sup>S196</sup> Glucocorticoids are well known to exert strong catabolic effects on skeletal muscle by activating the expression of multiple genes involved in proteolysis while inhibiting those involved in proteosynthesis.<sup>S197,S198</sup>

Circulating glucocorticoid level is increased in cachectic cancer patients<sup>117,S199,S200</sup> and in cachectic cancer mice.<sup>22,23,26,85,86,118,S124,S201–S203</sup> Hypothalamic CRH mRNA level<sup>67</sup> and pituitary ACTH secretion<sup>S204</sup> are also increased in animals, as well as the adrenal gland mass in cachectic cancer patients<sup>S205</sup> and animals.<sup>25,S204,S206</sup> More recently, our group established that the hypothalamic–pituitary–adrenal axis was activated in cachectic cancer mice, along with increased corticosterone level in serum and muscle, and increased skeletal muscle expression of glucocorticoid-responsive genes.<sup>119</sup> Interestingly, the analysis of transcriptomic data also reveals an increase in the expression of multiple glucocorticoid-responsive genes in muscle of cachectic cancer mice.<sup>33,34,44,55,73,78,86–88,S70,S72,S83,S92,S102,S103</sup> Therefore, a neuroendocrine mechanism that involves the hypothalamic–pituitary–adrenal axis may also contribute to the transcriptional regulation of skeletal muscle catabolism.

## Therapeutic perspectives

A brief description of the effects of therapeutic strategies in preclinical models is presented below.

## IGF1

IGF1 treatment reduces weight loss and improves outcome in a rat model of cancer cachexia.<sup>S207</sup> Anamorelin is a ghrelin agonist that increases the production of growth hormone from the pituitary gland and stimulates the liver to secrete IGF1.<sup>S208,S209</sup> Anamorelin has proven efficacy to limit the extent of cancer cachexia in human,<sup>S210,S211</sup> suggesting potential positive effects on the regulation of insulin/IGF1–AKT–mTOR pathway in skeletal muscle. However, care should be taken as IGF1 may promote tumour growth, even if there was no clear trend of increased tumour progression due to anamorelin.<sup>S210</sup>

## ER stress and the UPR

Targeted ablation of XBP1 (IRE1 $\alpha$  arm of the UPR) in skeletal muscle reduces muscle mass loss in tumour-bearing mice,<sup>S139</sup> whereas targeted ablation of PERK worsened muscle wasting.<sup>S138</sup> Different arms of the UPR may thus be implicated differently and provide different signalling outcomes.

## Oxidative stress

Administration of different antioxidants ( $\alpha$ -tocopherol, dehydroepiandrosterone, cocktail of catechins, quercetin and vitamin C) have provided contrasted results. Some studies show a sparing effect on muscle mass loss,<sup>S75</sup> reduced expression of E3 ligases<sup>S63,S75</sup> and a restoration of MyoD and myogenin expression,<sup>S75</sup> whereas another investigation reports an acceleration of cachexia by increasing muscle atrophy and promoting tumour growth.<sup>S76</sup>

## TNF $\alpha$ –NF- $\kappa$ B

Therapeutic strategies invalidating TNF $\alpha$  receptor,<sup>S18,S19,S31</sup> as well as the injection of anti-TNF $\alpha$  antibody,<sup>22</sup> failed to provide beneficial results. However, the pharmacological inhibition of NF- $\kappa$ B has proven its efficiency in cancer cachectic animals.<sup>27</sup> Besides, selective pharmacological inhibition of iNOS, which is a downstream effector of NF- $\kappa$ B pathway and is highly expressed in cachectic muscle,<sup>120</sup> ameliorates cancer cachexia in mouse.<sup>120</sup>

## IL6–JAK–STAT3

Cancer mice lacking IL6<sup>107,109,S69</sup> do not develop cachexia. Injection of an anti-IL6 antibody<sup>S166</sup> or an anti-IL6 receptor antibody<sup>18</sup> prevents cachexia progression in tumour-bearing animals. STAT3 inhibition also reduces muscle wasting in tumour-bearing mice.<sup>S178,S212</sup>

## Myostatin/activin A–SMAD1/3

Cancer cachexia is blocked by myostatin gene invalidation,<sup>41</sup> the administration of a myostatin antisense RNA,<sup>S188</sup> a myostatin antibody,<sup>31</sup> a soluble form of ActRIIB,<sup>21,62,75,113,S17,S192</sup> an ALK4/5 receptor antagonist,<sup>66</sup> and by the administration of IMB0901 (myostatin signalling inhibitor).<sup>S93</sup> AAV-targeted inhibition of myostatin and activin A also prevents muscle wasting in tumour-bearing mice.<sup>S87</sup>

## BMP–SMAD1/5/8

Increasing BMP signalling in skeletal muscle of tumour-bearing mice by gene delivery or pharmacological means can prevent muscle wasting.<sup>43</sup>

## Glucocorticoids

Ablation of adrenal glands does not attenuate cachexia<sup>26,S213</sup> in tumour-bearing animals. However, adrenalectomy itself induces weight loss.<sup>S214</sup> The steroid inhibitor RU486 shows contrasted results, with some positive effects on the attenuation of body weight loss<sup>S202</sup> or not.<sup>118,S201</sup> However, RU486 also exerts anti-progestogenic and anti-androgenic effects, which may mitigate the potential anti-catabolic effects of glucocorticoid inhibition. Muscle wasting is abrogated in muscle-specific glucocorticoid receptor knockout mice inoculated with LLC cells.<sup>61</sup> This spatially targeted tissue approach suggests that targeting glucocorticoids through the hypothalamic–pituitary–adrenal axis by specific molecular tools may be promising in preclinical models of cancer cachexia.

## Conclusion and future directions

This analysis of the literature highlights several points and redraw the contours of some accepted ideas.

Our analysis shows the existence of species-dependent molecular and biochemical responses. This is an essential factor to consider when evaluating the relevance of preclinical data to the clinical field. This also raises the question of the preclinical models used for research that should reproduce as closely as possible the complexity of the clinical context. Important factors are the model used (syngeneic ectopic/orthotopic graft, human tumour xenograft, genetic engineered mouse), the rate of disease progression and tumour growth, the age of the animals, the presence of metastasis or not and the use of additional chemotherapy or not.<sup>S215</sup>

There are still uncertainties whether protein degradation rate and MuRF1 and MAFbx expression are increased in skeletal muscle of cachectic cancer patients. This remains a key issue, even though data suggest that the ubiquitin–protea-

some system would be activated in muscle of human cancer patients.

Activin A may play a more pre-eminent role in cachectic cancer patients than myostatin, the reverse being true in animals.

The regulation of the molecular mechanisms involved in cancer cachexia is not a linear process during the natural history of the disease. A temporal analysis of the mechanisms involved in muscle proteostasis should be performed in pre-clinical models. In cancer patients, due to the complexity of the clinical context and the impossibility to determine the onset of tumour growth, a time-course analysis of muscle atrophy and muscle function will help to more accurately determine the cachectic state of the patients throughout the course of the disease.

Emerging evidence indicate that muscle fibre microenvironment contributes to cancer cachexia. Whether or not this is quantitatively important in determining muscle wasting remains to investigate.

One major unsolved question is the influence of gender on the molecular and biological expression of cancer cachexia. This is a key issue that needs to be addressed.

Finally, although substantial progress has been made in developing therapeutic strategies in preclinical models, treatments of cancer cachexia are not available in the clinic. Considering the numerous molecular mechanisms that are coordinately regulated within a specific time frame during the disease, multi-targeted strategies could be more effective in addressing the diversity and complexity of cancer cachexia.

## References

1. Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition and classification of cancer cachexia: an international consensus. *Lancet Oncol* 2011;**12**:489–495.
2. Baracos VE, Martin L, Korc M, Guttridge DC, Fearon KCH. Cancer-associated cachexia. *Nat Rev Dis Primers* 2018;**4**:17105.
3. Cohn SH, Gartenhaus W, Sawitsky A, Rai K, Zanzi I, Vaswani A, et al. Compartmental body composition of cancer patients by measurement of total body nitrogen, potassium, and water. *Metabolism* 1981;**30**:222–229.
4. Loumaye A, de Barys M, Nachit M, Lause P, Frateur L, van Maanen A, et al. Role of activin A and myostatin in human cancer cachexia. *J Clin Endocrinol Metab* 2015;**100**:2030–2038.
5. Prado CM, Sawyer MB, Ghosh S, Lieffers JR, Esfandiari N, Antoun S, et al. Central tenet of cancer cachexia therapy: do patients with advanced cancer have exploitable anabolic potential? *Am J Clin Nutr* 2013;**98**:1012–1019.
6. Argiles JM, Stemmler B, Lopez-Soriano FJ, Busquets S. Inter-tissue communication in cancer cachexia. *Nat Rev Endocrinol* 2018;**15**:9–20.
7. Weerink LBM, van der Hoorn A, van Leeuwen BL, de Bock GH. Low skeletal muscle mass and postoperative morbidity in surgical oncology: a systematic review and meta-analysis. *J Cachexia Sarcopenia Muscle* 2020;**11**:636–649.
8. Anker MS, Holcomb R, Muscaritoli M, von Haehling S, Haverkamp W, Jatoi A, et al. Orphan disease status of cancer cachexia in the USA and in the European Union: a systematic review. *J Cachexia Sarcopenia Muscle* 2019;**10**:22–34.
9. Emery PW, Edwards RH, Rennie MJ, Souhami RL, Halliday D. Protein synthesis in muscle measured in vivo in cachectic patients with cancer. *Br Med J (Clin Res Ed)* 1984;**289**:584–586.
10. Dworzak F, Ferrari P, Gavazzi C, Maiorana C, Bozzetti F. Effects of cachexia due to cancer on whole body and skeletal muscle protein turnover. *Cancer* 1998;**82**:42–48.
11. Williams JP, Phillips BE, Smith K, Atherton PJ, Rankin D, Selby AL, et al. Effect of tumour burden and subsequent surgical resection on skeletal muscle mass and protein turnover in colorectal cancer patients. *Am J Clin Nutr* 2012;**96**:1064–1070.
12. Lundholm K, Bennegard K, Eden E, Svaninger G, Emery PW, Rennie MJ. Efflux of 3-methylhistidine from the leg in cancer patients who experience weight loss. *Cancer Res* 1982;**42**:4807–4811.
13. Shaw JH, Humberstone DA, Douglas RG, Koea J. Leucine kinetics in patients with benign disease, non-weight-losing cancer, and cancer cachexia: studies at the whole-body and tissue level and the response to nutritional support. *Surgery* 1991;**109**:37–50.
14. MacDonald AJ, Johns N, Stephens N, Greig C, Ross JA, Small AC, et al. Habitual myofibrillar protein synthesis is normal in patients with upper GI cancer cachexia. *Clin Cancer Res* 2015;**21**:1734–1740.
15. Beck SA, Smith KL, Tisdale MJ. Anticachectic and antitumor effect of eicosapentaenoic acid and its effect on protein turnover. *Cancer Res* 1991;**51**:6089–6093.

## Acknowledgements

The authors certify that they comply with the ethical guidelines for publishing in the *Journal of Cachexia, Sarcopenia and Muscle*.<sup>S215</sup>

## Funding

A.M. was financially supported by the Ministère de l'Enseignement Supérieur de la Recherche et de l'Innovation. This work was supported by La Ligue contre le cancer (Comité de la Loire) et la Fondation ARC pour la Recherche sur le Cancer.

## Conflict of interest

A.M., Y.S.G. and D.F. declare that they have no conflict of interest.

## Online supplementary material

Additional supporting information may be found online in the Supporting Information section at the end of the article.



16. Emery PW, Lovell L, Rennie MJ. Protein synthesis measured in vivo in muscle and liver of cachectic tumor-bearing mice. *Cancer Res* 1984;**44**:2779–2784.
17. Smith KL, Tisdale MJ. Increased protein degradation and decreased protein synthesis in skeletal muscle during cancer cachexia. *Br J Cancer* 1993;**67**:680–685.
18. White JP, Baynes JW, Welle SL, Kostek MC, Matesic LE, Sato S, et al. The regulation of skeletal muscle protein turnover during the progression of cancer cachexia in the Apc (Min/+) mouse. *PLoS ONE* 2011;**6**:e24650.
19. Samuels SE, Knowles AL, Tilignac T, Debiton E, Madelmont JC, Attaix D. Higher skeletal muscle protein synthesis and lower breakdown after chemotherapy in cachectic mice. *Am J Physiol Regul Integr Comp Physiol* 2001;**281**:R133–R139.
20. Baracos VE, DeVivo C, Hoyle DH, Goldberg AL. Activation of the ATP-ubiquitin-proteasome pathway in skeletal muscle of cachectic rats bearing a hepatoma. *Am J Physiol Endocrinol Metab* 1995;**268**:E996–E1006.
21. Nissinen TA, Hentila J, Penna F, Lampinen A, Lautaoja JH, Fachada V, et al. Treating cachexia using soluble ACVR2B improves survival, alters mTOR localization, and attenuates liver and spleen responses. *J Cachexia Sarcopenia Muscle* 2018;**9**:514–529.
22. Costelli P, Carbo N, Tessitore L, Bagby GJ, Lopez-Soriano FJ, Argiles JM, et al. Tumor necrosis factor- $\alpha$  mediates changes in tissue protein turnover in a rat cancer cachexia model. *J Clin Invest* 1993;**92**:2783–2789.
23. Costelli P, Garcia-Martinez C, Llovera M, Carbo N, Lopez-Soriano FJ, Agell N, et al. Muscle protein waste in tumor-bearing rats is effectively antagonized by a beta 2-adrenergic agonist (clenbuterol). Role of the ATP-ubiquitin-dependent proteolytic pathway. *J Clin Invest* 1995;**95**:2367–2372.
24. Temparis S, Asensi M, Taillandier D, Arousseau E, Larbaud D, Obled A, et al. Increased ATP-ubiquitin-dependent proteolysis in skeletal muscles of tumor-bearing rats. *Cancer Res* 1994;**54**:5568–5573.
25. Tessitore L, Costelli P, Bonetti G, Baccino FM. Cancer cachexia, malnutrition, and tissue protein turnover in experimental animals. *Arch Biochem Biophys* 1993;**306**:52–58.
26. Tessitore L, Costelli P, Baccino FM. Pharmacological interference with tissue hypercatabolism in tumour-bearing rats. *Biochem J* 1994;**299**:71–78.
27. Chacon-Cabrera A, Fermoselle C, Urtreger AJ, Mateu-Jimenez M, Diamant MJ, de Kier Joffe ED, et al. Pharmacological strategies in lung cancer-induced cachexia: effects on muscle proteolysis, autophagy, structure, and weakness. *J Cell Physiol* 2014;**229**:1660–1672.
28. Khal J, Wyke SM, Russell ST, Hine AV, Tisdale MJ. Expression of the ubiquitin-proteasome pathway and muscle loss in experimental cancer cachexia. *Br J Cancer* 2005;**93**:774–780.
29. Lorite MJ, Thompson MG, Drake JL, Carling G, Tisdale MJ. Mechanism of muscle protein degradation induced by a cancer cachectic factor. *Br J Cancer* 1998;**78**:850–856.
30. de Castro GS, Simoes E, Lima J, Ortiz-Silva M, Festuccia WT, Tokeshi F, et al. Human cachexia induces changes in mitochondria, autophagy and apoptosis in the skeletal muscle. *Cancers (Basel)* 2019;**11**:1264.
31. Murphy KT, Chee A, Gleeson BG, Naim T, Swiderski K, Koopman R, et al. Antibody-directed myostatin inhibition enhances muscle mass and function in tumor-bearing mice. *Am J Physiol Regul Integr Comp Physiol* 2011;**301**:R716–R726.
32. Busquets S, Figueras MT, Fuster G, Almendro V, Moore-Carrasco R, Ametller E, et al. Anticachectic effects of formoterol: a drug for potential treatment of muscle wasting. *Cancer Res* 2004;**64**:6725–6731.
33. Nosacka RL, Delitto AE, Delitto D, Patel R, Judge SM, Trevino JG, et al. Distinct cachexia profiles in response to human pancreatic tumours in mouse limb and respiratory muscle. *J Cachexia Sarcopenia Muscle* 2020;**11**:820–837.
34. Judge SM, Wu CL, Beharry AW, Roberts BM, Ferreira LF, Kandarian SC, et al. Genome-wide identification of FoxO-dependent gene networks in skeletal muscle during C26 cancer cachexia. *BMC Cancer* 2014;**14**:997.
35. Jewesbury RC, Topley WW. Pathological changes in voluntary muscles in general diseases. *Proc R Soc Med* 1912;**5**:161–178.
36. Puig-Vilanova E, Rodriguez DA, Lloreta J, Ausin P, Pascual-Guardia S, Broquetas J, et al. Oxidative stress, redox signaling pathways, and autophagy in cachectic muscles of male patients with advanced COPD and lung cancer. *Free Radic Biol Med* 2015;**79**:91–108.
37. Judge SM, Nosacka RL, Delitto D, Gerber MH, Cameron ME, Trevino JG, et al. Skeletal muscle fibrosis in pancreatic cancer patients with respect to survival. *JNCI Cancer Spectr* 2018;**2**:pk043.
38. He WA, Berardi E, Cardillo VM, Acharyya S, Aulino P, Thomas-Ahner J, et al. NF- $\kappa$ B-mediated Pax7 dysregulation in the muscle microenvironment promotes cancer cachexia. *J Clin Invest* 2013;**123**:4821–4835.
39. Mehl KA, Davis JM, Berger FG, Carson JA. Myofiber degeneration/regeneration is induced in the cachectic ApcMin/+ mouse. *J Appl Physiol* 2005;**99**:2379–2387.
40. Salazar-Degracia A, Busquets S, Argiles JM, Lopez-Soriano FJ, Barreiro E. Formoterol attenuates increased oxidative stress and myosin protein loss in respiratory and limb muscles of cancer cachectic rats. *PeerJ* 2017;**5**:e4109.
41. Gallot YS, Durieux AC, Castells J, Desgeorges MM, Vernus B, Plantureux L, et al. Myostatin gene inactivation prevents skeletal muscle wasting in cancer. *Cancer Res* 2014;**74**:7344–7356.
42. Kim HG, Huot JR, Pin F, Guo B, Bonetto A, Nader GA. Reduced rDNA transcription diminishes skeletal muscle ribosomal capacity and protein synthesis in cancer cachexia. *FASEB J* 2021;**35**:e21335.
43. Sartori R, Hagg A, Zampieri S, Armani A, Winbanks CE, Viana LR, et al. Perturbed BMP signaling and denervation promote muscle wasting in cancer cachexia. *Sci Transl Med* 2021;**13**.
44. Talbert EE, Cuitino MC, Ladner KJ, Rajasekera PV, Siebert M, Shakya R, et al. Modeling human cancer-induced cachexia. *Cell Rep* 2019;**28**:1612–1622.
45. Gallagher IJ, Stephens NA, MacDonald AJ, Skipworth RJ, Husi H, Greig CA, et al. Suppression of skeletal muscle turnover in cancer cachexia: evidence from the transcriptome in sequential human muscle biopsies. *Clin Cancer Res* 2012;**18**:2817–2827.
46. Bonetto A, Penna F, Aversa Z, Mercantini P, Baccino FM, Costelli P, et al. Early changes of muscle insulin-like growth factor-1 and myostatin gene expression in gastric cancer patients. *Muscle Nerve* 2013;**48**:387–392.
47. Op den Kamp CM, Langen RC, Snepvangers FJ, de Theije CC, Schellekens JM, Laugs F, et al. Nuclear transcription factor kappa B activation and protein turnover adaptations in skeletal muscle of patients with progressive stages of lung cancer cachexia. *Am J Clin Nutr* 2013;**98**:738–748.
48. D'Orlando C, Marzetti E, Francois S, Lorenzi M, Conti V, di Stasio E, et al. Gastric cancer does not affect the expression of atrophy-related genes in human skeletal muscle. *Muscle Nerve* 2014;**49**:528–533.
49. Stephens NA, Skipworth RJ, Gallagher IJ, Greig CA, Guttridge DC, Ross JA, et al. Evaluating potential biomarkers of cachexia and survival in skeletal muscle of upper gastrointestinal cancer patients. *J Cachexia Sarcopenia Muscle* 2015;**6**:53–61.
50. Murton AJ, Maddocks M, Stephens FB, Marimuthu K, England R, Wilcock A. Consequences of late-stage non-small-cell lung cancer cachexia on muscle metabolic processes. *Clin Lung Cancer* 2017;**18**:e1–e11.
51. DeJong CH, Busquets S, Moses AG, Schrauwen P, Ross JA, Argiles JM, et al. Systemic inflammation correlates with increased expression of skeletal muscle ubiquitin but not uncoupling proteins in cancer cachexia. *Oncol Rep* 2005;**14**:257–263.
52. Sun YS, Ye ZY, Qian ZY, Xu XD, Hu JF. Expression of TRAF6 and ubiquitin mRNA in skeletal muscle of gastric cancer patients. *J Exp Clin Cancer Res* 2012;**31**:81.
53. Khal J, Hine AV, Fearon KC, DeJong CH, Tisdale MJ. Increased expression of pro-

- teasome subunits in skeletal muscle of cancer patients with weight loss. *Int J Biochem Cell Biol* 2005;**37**:2196–2206.
54. Jagoe RT, Redfern CP, Roberts RG, Gibson GJ, Goodship TH. Skeletal muscle mRNA levels for cathepsin B, but not components of the ubiquitin-proteasome pathway, are increased in patients with lung cancer referred for thoracotomy. *Clin Sci (Lond)* 2002;**102**:353–361.
  55. Constantinou C, Fontes de Oliveira CC, Mintzopoulos D, Busquets S, He J, Kesarwani M, et al. Nuclear magnetic resonance in conjunction with functional genomics suggests mitochondrial dysfunction in a murine model of cancer cachexia. *Int J Mol Med* 2011;**27**:15–24.
  56. Costelli P, Muscaritoli M, Bossola M, Penna F, Reffo P, Bonetto A, et al. IGF-1 is downregulated in experimental cancer cachexia. *Am J Physiol Regul Integr Comp Physiol* 2006;**291**:R674–R683.
  57. Acharyya S, Butchbach ME, Sahenk Z, Wang H, Saji M, Carathers M, et al. Dystrophin glycoprotein complex dysfunction: a regulatory link between muscular dystrophy and cancer cachexia. *Cancer Cell* 2005;**8**:421–432.
  58. Murphy KT, Struk A, Malcontenti-Wilson C, Christophi C, Lynch GS. Physiological characterization of a mouse model of cachexia in colorectal liver metastases. *Am J Physiol Regul Integr Comp Physiol* 2013;**304**:R854–R864.
  59. Acharyya S, Ladner KJ, Nelsen LL, Damrauer J, Reiser PJ, Swoap S, et al. Cancer cachexia is regulated by selective targeting of skeletal muscle gene products. *J Clin Invest* 2004;**114**:370–378.
  60. Braun TP, Zhu X, Szumowski M, Scott GD, Grossberg AJ, Levasseur PR, et al. Central nervous system inflammation induces muscle atrophy via activation of the hypothalamic-pituitary-adrenal axis. *J Exp Med* 2011;**208**:2449–2463.
  61. Braun TP, Grossberg AJ, Krasnow SM, Levasseur PR, Szumowski M, Zhu XX, et al. Cancer- and endotoxin-induced cachexia require intact glucocorticoid signaling in skeletal muscle. *FASEB J* 2013;**27**:3572–3582.
  62. Busquets S, Toledo M, Orpi M, Massa D, Porta M, Capdevila E, et al. Myostatin blockage using actRIIB antagonism in mice bearing the Lewis lung carcinoma results in the improvement of muscle wasting and physical performance. *J Cachexia Sarcopenia Muscle* 2012;**3**:37–43.
  63. Guarnier FA, Cecchini AL, Suzukawa AA, Maragno AL, Simao AN, Gomes MD, et al. Time course of skeletal muscle loss and oxidative stress in rats with Walker 256 solid tumor. *Muscle Nerve* 2010;**42**:950–958.
  64. Julienne CM, Dumas JF, Goupille C, Pinault M, Berri C, Collin A, et al. Cancer cachexia is associated with a decrease in skeletal muscle mitochondrial oxidative capacities without alteration of ATP production efficiency. *J Cachexia Sarcopenia Muscle* 2012;**3**:265–275.
  65. Kir S, Komaba H, Garcia AP, Economopoulos KP, Liu W, Lanske B, et al. PTH/PTHrP receptor mediates cachexia in models of kidney failure and cancer. *Cell Metab* 2016;**23**:315–323.
  66. Levolger S, Wiemer EAC, van Vugt JLA, Huisman SA, van Vledder MG, van Damme-van Engel S, et al. Inhibition of activin-like kinase 4/5 attenuates cancer cachexia associated muscle wasting. *Sci Rep* 2019;**9**:9826.
  67. Michaelis KA, Zhu X, Burfeind KG, Krasnow SM, Levasseur PR, Morgan TK, et al. Establishment and characterization of a novel murine model of pancreatic cancer cachexia. *J Cachexia Sarcopenia Muscle* 2017;**8**:824–838.
  68. Pigna E, Berardi E, Aulino P, Rizzuto E, Zampieri S, Carraro U, et al. Aerobic exercise and pharmacological treatments counteract cachexia by modulating autophagy in colon cancer. *Sci Rep* 2016;**6**:26991.
  69. Reed SA, Sandesara PB, Senf SM, Judge AR. Inhibition of FoxO transcriptional activity prevents muscle fiber atrophy during cachexia and induces hypertrophy. *FASEB J* 2011;**26**:987–1000.
  70. Schwarzkopf M, Coletti D, Sassoon D, Marazzi G. Muscle cachexia is regulated by a p53-PW1/Peg3-dependent pathway. *Genes Dev* 2006;**20**:3440–3452.
  71. Shakri AR, Zhong TJ, Ma W, Coker C, Kim S, Calluori S, et al. Upregulation of ZIP14 and altered zinc homeostasis in muscles in pancreatic cancer cachexia. *Cancers (Basel)* 2019;**12**.
  72. Talbert EE, Metzger GA, He WA, Guttridge DC. Modeling human cancer cachexia in colon 26 tumor-bearing adult mice. *J Cachexia Sarcopenia Muscle* 2014;**5**:321–328.
  73. Wang G, Biswas AK, Ma W, Kandpal M, Coker C, Grandgenett PM, et al. Metastatic cancers promote cachexia through ZIP14 upregulation in skeletal muscle. *Nat Med* 2018;**24**:770–781.
  74. Winbanks CE, Murphy KT, Bernardo BC, Qian H, Liu Y, Sepulveda PV, et al. Smad7 gene delivery prevents muscle wasting associated with cancer cachexia in mice. *Sci Transl Med* 2016;**8**:348ra398.
  75. Zhou X, Wang JL, Lu J, Song Y, Kwak KS, Jiao Q, et al. Reversal of cancer cachexia and muscle wasting by ActRIIB antagonism leads to prolonged survival. *Cell* 2010;**142**:531–543.
  76. Huot JR, Pin F, Narasimhan A, Novinger LJ, Keith AS, Zimmers TA, et al. ACVR2B antagonism as a countermeasure to multi-organ perturbations in metastatic colorectal cancer cachexia. *J Cachexia Sarcopenia Muscle* 2020;**11**:1779–1798.
  77. Penna F, Costamagna D, Pin F, Camperi A, Fanzani A, Chiarpotto EM, et al. Autophagic degradation contributes to muscle wasting in cancer cachexia. *Am J Pathol* 2013;**182**:1367–1378.
  78. Lecker SH, Jagoe RT, Gilbert A, Gomes M, Baracos V, Bailey J, et al. Multiple types of skeletal muscle atrophy involve a common program of changes in gene expression. *FASEB J* 2004;**18**:39–51.
  79. Penna F, Ballaro R, Martinez-Cristobal P, Sala D, Sebastian D, Busquets S, et al. Autophagy exacerbates muscle wasting in cancer cachexia and impairs mitochondrial function. *J Mol Biol* 2019;**431**:2674–2686.
  80. Marino FE, Risbridger G, Gold E. Activin-beta modulates cachexia by repressing the ubiquitin-proteasome and autophagic degradation pathways. *J Cachexia Sarcopenia Muscle* 2015;**6**:365–380.
  81. Zhang G, Liu Z, Ding H, Zhou Y, Doan HA, Sin KWT, et al. Tumor induces muscle wasting in mice through releasing extracellular Hsp70 and Hsp90. *Nat Commun* 2017;**8**:589.
  82. Hardee JP, Counts BR, Gao S, VanderVeen BN, Fix DK, Koh HJ, et al. Inflammatory signalling regulates eccentric contraction-induced protein synthesis in cachectic skeletal muscle. *J Cachexia Sarcopenia Muscle* 2018;**9**:369–383.
  83. Koutnik AP, Poff AM, Ward NP, DeBlasi JM, Soliven MA, Romero MA, et al. Ketone bodies attenuate wasting in models of atrophy. *J Cachexia Sarcopenia Muscle* 2020;**11**:973–996.
  84. Llovera M, Garcia-Martinez C, Agell N, Marzabal M, Lopez-Soriano FJ, Argiles JM. Ubiquitin gene expression is increased in skeletal muscle of tumour-bearing rats. *FEBS Lett* 1994;**338**:311–318.
  85. Carbo N, Lopez-Soriano J, Costelli P, Busquets S, Alvarez B, Baccino FM, et al. Interleukin-15 antagonizes muscle protein waste in tumour-bearing rats. *Br J Cancer* 2000;**83**:526–531.
  86. Goncalves MD, Hwang SK, Pauli C, Murphy CJ, Cheng Z, Hopkins BD, et al. Fenofibrate prevents skeletal muscle loss in mice with lung cancer. *Proc Natl Acad Sci U S A* 2018;**115**:E743–E752.
  87. Bonetto A, Aydogdu T, Kunzevitzky N, Guttridge DC, Khuri S, Koniaris LG, et al. STAT3 activation in skeletal muscle links muscle wasting and the acute phase response in cancer cachexia. *PLoS One* 2011;**6**:e22538.
  88. Cornwell EW, Mirbod A, Wu CL, Kandarian SC, Jackman RW. C26 cancer-induced muscle wasting is IKKbeta-dependent and NF-kappaB-independent. *PLoS One* 2014;**9**:e87776.
  89. Llovera M, Garcia-Martinez C, Agell N, Lopez-Soriano FJ, Argiles JM. Muscle wasting associated with cancer cachexia is linked to an important activation of the ATP-dependent ubiquitin-mediated proteolysis. *Int J Cancer* 1995;**61**:138–141.
  90. Aversa Z, Pin F, Lucia S, Penna F, Verzaro R, Fazi M, et al. Autophagy is induced in the skeletal muscle of cachectic cancer patients. *Sci Rep* 2016;**6**:30340.
  91. Ballaro R, Penna F, Pin F, Gomez-Cabrera MC, Vina J, Costelli P. Moderate exercise improves experimental cancer cachexia by modulating the redox homeostasis. *Cancers (Basel)* 2019;**11**:285.

92. Deval C, Mordier S, Obled C, Bechet D, Combaret L, Attaix D, et al. Identification of cathepsin L as a differentially expressed message associated with skeletal muscle wasting. *Biochem J* 2001;**360**: 143–150.
93. Martin A, Freyssen D. Phenotypic features of cancer cachexia-related loss of skeletal muscle mass and function: lessons from human and animal studies. *J Cachexia Sarcopenia Muscle* 2021;**12**: 252–273.
94. Brown JL, Rosa-Caldwell ME, Lee DE, Blackwell TA, Brown LA, Perry RA, et al. Mitochondrial degeneration precedes the development of muscle atrophy in progression of cancer cachexia in tumour-bearing mice. *J Cachexia Sarcopenia Muscle* 2017;**8**:926–938.
95. White JP, Baltgalvis KA, Puppa MJ, Sato S, Baynes JW, Carson JA. Muscle oxidative capacity during IL-6-dependent cancer cachexia. *Am J Physiol Regul Integr Comp Physiol* 2011;**300**:R201–R211.
96. Schmitt TL, Martignoni ME, Bachmann J, Fechtner K, Friess H, Kinscherf R, et al. Activity of the Akt-dependent anabolic and catabolic pathways in muscle and liver samples in cancer-related cachexia. *J Mol Med (Berl)* 2007;**85**:647–654.
97. Puppa MJ, Murphy EA, Fayad R, Hand GA, Carson JA. Cachectic skeletal muscle response to a novel bout of low-frequency stimulation. *J Appl Physiol* 1985;**2014**: 1078–1087.
98. Johns N, Stretch C, Tan BH, Solheim TS, Sorhaug S, Stephens NA, et al. New genetic signatures associated with cancer cachexia as defined by low skeletal muscle index and weight loss. *J Cachexia Sarcopenia Muscle* 2017;**8**: 122–130.
99. Barreiro E, de la Puente B, Busquets S, Lopez-Soriano FJ, Gea J, Argiles JM. Both oxidative and nitrosative stress are associated with muscle wasting in tumour-bearing rats. *FEBS Lett* 2005;**579**: 1646–1652.
100. Marin-Corral J, Fontes CC, Pascual-Guardia S, Sanchez F, Olivan M, Argiles JM, et al. Redox balance and carbonylated proteins in limb and heart muscles of cachectic rats. *Antioxid Redox Signal* 2010;**12**:365–380.
101. Neyroud D, Nosacka RL, Judge AR, Hepple RT. Colon 26 adenocarcinoma (C26)-induced cancer cachexia impairs skeletal muscle mitochondrial function and content. *J Muscle Res Cell Motil* 2019;**40**: 59–65.
102. O’Gorman P, McMillan DC, McArdle CS. Longitudinal study of weight, appetite, performance status, and inflammation in advanced gastrointestinal cancer. *Nutr Cancer* 1999;**35**:127–129.
103. Scott HR, McMillan DC, Forrest LM, Brown DJ, McArdle CS, Milroy R. The systemic inflammatory response, weight loss, performance status and survival in patients with inoperable non-small cell lung cancer. *Br J Cancer* 2002;**87**:264–267.
104. Lerner L, Hayes TG, Tao N, Krieger B, Feng B, Wu Z, et al. Plasma growth differentiation factor 15 is associated with weight loss and mortality in cancer patients. *J Cachexia Sarcopenia Muscle* 2015;**6**: 317–324.
105. Lerner L, Tao J, Liu Q, Nicoletti R, Feng B, Krieger B, et al. MAP 3K11/GDF15 axis is a critical driver of cancer cachexia. *J Cachexia Sarcopenia Muscle* 2016;**7**: 467–482.
106. Chen JL, Walton KL, Qian H, Colgan TD, Hagg A, Watt MJ, et al. Differential effects of IL6 and activin A in the development of cancer-associated cachexia. *Cancer Res* 2016;**76**:5372–5382.
107. Baltgalvis KA, Berger FG, Pena MM, Davis JM, Muga SJ, Carson JA. Interleukin-6 and cachexia in ApcMin/+ mice. *Am J Physiol Regul Integr Comp Physiol* 2008;**294**: R393–R401.
108. Segatto M, Fittipaldi R, Pin F, Sartori R, Dae Ko K, Zare H, et al. Epigenetic targeting of bromodomain protein BRD4 counteracts cancer cachexia and prolongs survival. *Nat Commun* 2017;**8**:1707.
109. Miller A, McLeod L, Alhayyani S, Szczepny A, Watkins DN, Chen W, et al. Blockade of the IL-6 trans-signalling/STAT3 axis suppresses cachexia in Kras-induced lung adenocarcinoma. *Oncogene* 2017;**36**: 3059–3066.
110. Durieux AC, Amirouche A, Banzet S, Koulmann N, Bonnefoy R, Padeloup M, et al. Ectopic expression of myostatin induces atrophy of adult skeletal muscle by decreasing muscle gene expression. *Endocrinology* 2007;**148**:3140–3147.
111. Amirouche A, Durieux AC, Banzet S, Koulmann N, Bonnefoy R, Mouret C, et al. Down-regulation of akt/mammalian target of rapamycin signaling pathway in response to myostatin overexpression in skeletal muscle. *Endocrinology* 2009;**150**: 286–294.
112. Morel J, Palao JC, Castells J, Desgeorges M, Busso T, Molliex S, et al. Regulation of Akt-mTOR, ubiquitin-proteasome and autophagy-lysosome pathways in locomotor and respiratory muscles during experimental sepsis in mice. *Sci Rep* 2017;**7**:10866.
113. Benny Klimek ME, Aydogdu T, Link MJ, Pons M, Koniaris LG, Zimmers TA. Acute inhibition of myostatin-family proteins preserves skeletal muscle in mouse models of cancer cachexia. *Biochem Biophys Res Commun* 2010;**391**: 1548–1554.
114. Paajanen J, Ilonen I, Lauri H, Jarvinen T, Sutinen E, Ollila H, et al. Elevated circulating activin A levels in patients with malignant pleural mesothelioma are related to cancer cachexia and reduced response to platinum-based chemotherapy. *Clin Lung Cancer* 2020;**21**:e142–e150.
115. Matzuk MM, Finegold MJ, Mather JP, Krummen L, Lu H, Bradley A. Development of cancer cachexia-like syndrome and adrenal tumors in inhibin-deficient mice. *Proc Natl Acad Sci U S A* 1994;**91**: 8817–8821.
116. Parajuli P, Kumar S, Loumaye A, Singh P, Eragamreddy S, Nguyen TL, et al. Twist1 activation in muscle progenitor cells causes muscle loss akin to cancer cachexia. *Dev Cell* 2018;**45**:e716.
117. Flint TR, Janowitz T, Connell CM, Roberts EW, Denton AE, Coll AP, et al. Tumor-induced IL-6 reprograms host metabolism to suppress anti-tumor immunity. *Cell Metab* 2016;**24**:672–684.
118. Llovera M, Garcia-Martinez C, Costelli P, Agell N, Carbo N, Lopez-Soriano FJ, et al. Muscle hypercatabolism during cancer cachexia is not reversed by the glucocorticoid receptor antagonist RU38486. *Cancer Lett* 1996;**99**:7–14.
119. Martin A, Castells J, Allibert V, Emerit A, Zolotoff C, Cardot-Ruffino V, et al. Hypothalamic-pituitary-adrenal axis activation and glucocorticoid-responsive gene expression in skeletal muscle and liver of Apc mice. *J Cachexia Sarcopenia Muscle* 2022;**16**:1686–1703.
120. Sadek J, Hall DT, Colalillo B, Omer A, Tremblay AK, Sanguin-Gendreau V, et al. Pharmacological or genetic inhibition of iNOS prevents cachexia-mediated muscle wasting and its associated metabolism defects. *EMBO Mol Med* 2021;**13**: e13591.

## Supplementary references

- S1. Weber MA, Krakowski-Roosen H, Schroder L, Kinscherf R, Krix M, Kopp-Schneider A et al. Morphology, metabolism, microcirculation, and strength of skeletal muscles in cancer-related cachexia. *Acta Oncol* 2009;**48**:116-124.
- S2. Stephens NA, Gray C, MacDonald AJ, Tan BH, Gallagher IJ, Skipworth RJ et al. Sexual dimorphism modulates the impact of cancer cachexia on lower limb muscle mass and function. *Clin Nutr* 2012;**31**:499-505.
- S3. Weber MA, Kinscherf R, Krakowski-Roosen H, Aulmann M, Renk H, Kunkele A et al. Myoglobin plasma level related to muscle mass and fiber composition: a clinical marker of muscle wasting? *J Mol Med (Berl)* 2007;**85**:887-896.
- S4. Vanhoutte G, van de Wiel M, Wouters K, Sels M, Bartolomeeussen L, De Keersmaecker S et al. Cachexia in cancer: what is in the definition? *BMJ Open Gastroenterol* 2016;**3**:e000097.
- S5. Roeland EJ, Ma JD, Nelson SH, Seibert T, Heavey S, Revta C et al. Weight loss versus muscle loss: re-evaluating inclusion criteria for future cancer cachexia interventional trials. *Support Care Cancer* 2017;**25**:365-369.
- S6. Narasimhan A, Greiner R, Bathe OF, Baracos V, Damaraju S. Differentially expressed alternatively spliced genes in skeletal muscle from cancer patients with cachexia. *J Cachexia Sarcopenia Muscle* 2018;**9**:60-70.
- S7. Jackson KM, Cole CL, Dunne RF. From bench to bedside: updates in basic science, translational and clinical research on muscle fatigue in cancer cachexia. *Curr Opin Clin Nutr Metab Care* 2021;**24**:216-222.
- S8. VanderVeen BN, Hardee JP, Fix DK, Carson JA. Skeletal muscle function during the progression of cancer cachexia in the male Apc(Min/+) mouse. *J Appl Physiol (1985)* 2018;**124**:684-695.
- S9. Huang DD, Wang SL, Zhuang CL, Zheng BS, Lu JX, Chen FF et al. Sarcopenia, as defined by low muscle mass, strength and physical performance, predicts complications after surgery for colorectal cancer. *Colorectal Dis* 2015;**17**:O256-264.
- S10. Jung HW, Kim JW, Kim JY, Kim SW, Yang HK, Lee JW et al. Effect of muscle mass on toxicity and survival in patients with colon cancer undergoing adjuvant chemotherapy. *Support Care Cancer* 2015;**23**:687-694.
- S11. Tan BH, Brammer K, Randhawa N, Welch NT, Parsons SL, James EJ et al. Sarcopenia is associated with toxicity in patients undergoing neo-adjuvant chemotherapy for oesophago-gastric cancer. *Eur J Surg Oncol* 2015;**41**:333-338.
- S12. Mir O, Coriat R, Blanchet B, Durand JP, Boudou-Rouquette P, Michels J et al. Sarcopenia predicts early dose-limiting toxicities and pharmacokinetics of sorafenib in patients with hepatocellular carcinoma. *PLoS One* 2012;**7**:e37563.
- S13. Wendrich AW, Swartz JE, Bril SI, Wegner I, de Graeff A, Smid EJ et al. Low skeletal muscle mass is a predictive factor for chemotherapy dose-limiting toxicity in patients with locally advanced head and neck cancer. *Oral Oncol* 2017;**71**:26-33.
- S14. Smith HJ, Greenberg NA, Tisdale MJ. Effect of eicosapentaenoic acid, protein and amino acids on protein synthesis and degradation in skeletal muscle of cachectic mice. *Br J Cancer* 2004;**91**:408-412.
- S15. White JP, Puppa MJ, Gao S, Sato S, Welle SL, Carson JA. Muscle mTORC1 suppression by IL-6 during cancer cachexia: a role for AMPK. *Am J Physiol Endocrinol Metab* 2013;**304**:E1042-1052.
- S16. Lima M, Sato S, Enos RT, Baynes JW, Carson JA. Development of an UPLC mass spectrometry method for measurement of myofibrillar protein synthesis: application to

- analysis of murine muscles during cancer cachexia. *J Appl Physiol (1985)* 2013;**114**:824-828.
- S17. Toledo M, Busquets S, Penna F, Zhou X, Marmonti E, Betancourt A et al. Complete reversal of muscle wasting in experimental cancer cachexia: Additive effects of activin type II receptor inhibition and beta-2 agonist. *Int J Cancer* 2016;**138**:2021-2029.
- S18. Llovera M, Garcia-Martinez C, Lopez-Soriano J, Agell N, Lopez-Soriano FJ, Garcia I et al. Protein turnover in skeletal muscle of tumour-bearing transgenic mice overexpressing the soluble TNF receptor-1. *Cancer Lett* 1998;**130**:19-27.
- S19. Llovera M, Garcia-Martinez C, Lopez-Soriano J, Carbo N, Agell N, Lopez-Soriano FJ et al. Role of TNF receptor 1 in protein turnover during cancer cachexia using gene knockout mice. *Mol Cell Endocrinol* 1998;**142**:183-189.
- S20. Chacon-Cabrera A, Mateu-Jimenez M, Langohr K, Fermoselle C, Garcia-Arumi E, Andreu AL et al. Role of PARP activity in lung cancer-induced cachexia: Effects on muscle oxidative stress, proteolysis, anabolic markers, and phenotype. *J Cell Physiol* 2017;**232**:3744-3761.
- S21. Pin F, Minero VG, Penna F, Muscaritoli M, De Tullio R, Baccino FM et al. Interference with Ca(2+)-dependent proteolysis does not alter the course of muscle wasting in experimental cancer cachexia. *Front Physiol* 2017;**8**:213.
- S22. Rosa-Caldwell ME, Benson CA, Lee DE, Brown JL, Washington TA, Greene NP et al. Mitochondrial function and protein turnover in the diaphragm are altered in LLC tumor model of cancer cachexia. *Int J Mol Sci* 2020;**21**:7841.
- S23. Bossola M, Mirabella M, Ricci E, Costelli P, Pacelli F, Tortorelli AP et al. Skeletal muscle apoptosis is not increased in gastric cancer patients with mild-moderate weight loss. *Int J Biochem Cell Biol* 2006;**38**:1561-1570.
- S24. Busquets S, Deans C, Figueras M, Moore-Carrasco R, Lopez-Soriano FJ, Fearon KC et al. Apoptosis is present in skeletal muscle of cachectic gastro-intestinal cancer patients. *Clin Nutr* 2007;**26**:614-618.
- S25. Brzezczynska J, Johns N, Schilb A, Degen S, Degen M, Langen R et al. Loss of oxidative defense and potential blockade of satellite cell maturation in the skeletal muscle of patients with cancer but not in the healthy elderly. *Aging (Albany NY)* 2016;**8**:1690-1702.
- S26. Salazar-Degracia A, Busquets S, Argiles JM, Bargallo-Gispert N, Lopez-Soriano FJ, Barreiro E. Effects of the beta2 agonist formoterol on atrophy signaling, autophagy, and muscle phenotype in respiratory and limb muscles of rats with cancer-induced cachexia. *Biochimie* 2018;**149**:79-91.
- S27. Baltgalvis KA, Berger FG, Pena MM, Mark Davis J, White JP, Carson JA. Activity level, apoptosis, and development of cachexia in Apc(Min/+) mice. *J Appl Physiol (1985)* 2010;**109**:1155-1161.
- S28. Salazar-Degracia A, Blanco D, Vila-Ubach M, de Biurrun G, de Solorzano CO, Montuenga LM et al. Phenotypic and metabolic features of mouse diaphragm and gastrocnemius muscles in chronic lung carcinogenesis: influence of underlying emphysema. *J Transl Med* 2016;**14**:244.
- S29. Belizario JE, Lorite MJ, Tisdale MJ. Cleavage of caspases-1, -3, -6, -8 and -9 substrates by proteases in skeletal muscles from mice undergoing cancer cachexia. *Br J Cancer* 2001;**84**:1135-1140.
- S30. Figueras M, Busquets S, Carbo N, Barreiro E, Almendro V, Argiles JM et al. Interleukin-15 is able to suppress the increased DNA fragmentation associated with muscle wasting in tumour-bearing rats. *FEBS Lett* 2004;**569**:201-206.

- S31. Carbo N, Busquets S, van Royen M, Alvarez B, Lopez-Soriano FJ, Argiles JM. TNF- $\alpha$  is involved in activating DNA fragmentation in skeletal muscle. *Br J Cancer* 2002;**86**:1012-1016.
- S32. van Royen M, Carbo N, Busquets S, Alvarez B, Quinn LS, Lopez-Soriano FJ et al. DNA fragmentation occurs in skeletal muscle during tumor growth: A link with cancer cachexia? *Biochem Biophys Res Commun* 2000;**270**:533-537.
- S33. Lin XY, Chen SZ. Calpain inhibitors ameliorate muscle wasting in a cachectic mouse model bearing CT26 colorectal adenocarcinoma. *Oncol Rep* 2017;**37**:1601-1610.
- S34. Wang Q, Li C, Peng X, Kang Q, Deng D, Zhang L et al. Combined treatment of carfilzomib and z-VAD-fmk inhibits skeletal proteolysis and apoptosis and ameliorates cancer cachexia. *Med Oncol* 2015;**32**:100.
- S35. Antunes D, Padrao AI, Maciel E, Santinha D, Oliveira P, Vitorino R et al. Molecular insights into mitochondrial dysfunction in cancer-related muscle wasting. *Biochim Biophys Acta* 2014;**1841**:896-905.
- S36. White JP, Puppa MJ, Sato S, Gao S, Price RL, Baynes JW et al. IL-6 regulation on skeletal muscle mitochondrial remodeling during cancer cachexia in the ApcMin/+ mouse. *Skelet Muscle* 2012;**2**:14.
- S37. Barreiro E, Salazar-Degracia A, Sancho-Munoz A, Gea J. Endoplasmic reticulum stress and unfolded protein response profile in quadriceps of sarcopenic patients with respiratory diseases. *J Cell Physiol* 2019;**234**:11315-11329.
- S38. Schmalbruch H, Hellhammer U. The number of nuclei in adult rat muscles with special reference to satellite cells. *Anat Rec* 1977;**189**:169-175.
- S39. Johns N, Hatakeyama S, Stephens NA, Degen M, Degen S, Friauff W et al. Clinical classification of cancer cachexia: phenotypic correlates in human skeletal muscle. *PLoS One* 2014;**9**:e83618.
- S40. Marin OS, Denny-Brown D. Changes in skeletal muscle associated with cachexia. *Am J Pathol* 1962;**41**:23-39.
- S41. Mu X, Agarwal R, March D, Rothenberg A, Voigt C, Tebbets J et al. Notch Signaling Mediates Skeletal Muscle Atrophy in Cancer Cachexia Caused by Osteosarcoma. *Sarcoma* 2016;**2016**:3758162.
- S42. Penna F, Costamagna D, Fanzani A, Bonelli G, Baccino FM, Costelli P. Muscle wasting and impaired myogenesis in tumor bearing mice are prevented by ERK inhibition. *PLoS ONE* 2010;**5**:e13604.
- S43. Bentzinger CF, Wang YX, Dumont NA, Rudnicki MA. Cellular dynamics in the muscle satellite cell niche. *EMBO Rep* 2013;**14**:1062-1072.
- S44. Zampieri S, Doria A, Adami N, Biral D, Vecchiato M, Savastano S et al. Subclinical myopathy in patients affected with newly diagnosed colorectal cancer at clinical onset of disease: evidence from skeletal muscle biopsies. *Neurol Res* 2010;**32**:20-25.
- S45. Coletti D, Aulino P, Pigna E, Barteri F, Moresi V, Annibali D et al. Spontaneous physical activity downregulates Pax7 in cancer cachexia. *Stem Cells Int* 2016;**2016**:6729268.
- S46. Inaba S, Hinohara A, Tachibana M, Tsujikawa K, Fukada SI. Muscle regeneration is disrupted by cancer cachexia without loss of muscle stem cell potential. *PLoS One* 2018;**13**:e0205467.
- S47. Lautaoja JH, Lalowski M, Nissinen TA, Hentila J, Shi Y, Ritvos O et al. Muscle and serum metabolomes are dysregulated in colon-26 tumor-bearing mice despite amelioration of cachexia with activin receptor type 2B ligand blockade. *Am J Physiol Endocrinol Metab* 2019;**316**:E852-E865.
- S48. Goodlad GA, Clark CM. Response of skeletal muscle RNA polymerases I and II to tumour growth. *Biochim Biophys Acta* 1988;**950**:296-302.

- S49. Sandri M. Protein breakdown in muscle wasting: Role of autophagy-lysosome and ubiquitin-proteasome. *Int J Biochem Cell Biol* 2013;**45**:2121-2129.
- S50. Bodine SC, Baehr LM. Skeletal muscle atrophy and the E3 ubiquitin ligases MuRF1 and MAFbx/atrogen-1. *Am J Physiol Endocrinol Metab* 2014;**307**:E469-484.
- S51. Kedar V, McDonough H, Arya R, Li HH, Rockman HA, Patterson C. Muscle-specific RING finger 1 is a bona fide ubiquitin ligase that degrades cardiac troponin I. *Proc Natl Acad Sci U S A* 2004;**101**:18135-18140.
- S52. Clarke BA, Drujan D, Willis MS, Murphy LO, Corpina RA, Burova E et al. The E3 Ligase MuRF1 degrades myosin heavy chain protein in dexamethasone-treated skeletal muscle. *Cell Metab* 2007;**6**:376-385.
- S53. Polge C, Heng AE, Jarzaguat M, Ventadour S, Claustre A, Combaret L et al. Muscle actin is polyubiquitinated in vitro and in vivo and targeted for breakdown by the E3 ligase MuRF1. *FASEB J* 2011;**25**:3790-3802.
- S54. Tintignac LA, Lagirand J, Batonnet S, Sirri V, Leibovitch MP, Leibovitch SA. Degradation of MyoD mediated by the SCF (MAFbx) ubiquitin ligase. *J Biol Chem* 2005;**280**:2847-2856.
- S55. Csibi A, Cornille K, Leibovitch MP, Poupon A, Tintignac LA, Sanchez AM et al. The translation regulatory subunit eIF3f controls the kinase-dependent mTOR signaling required for muscle differentiation and hypertrophy in mouse. *PLoS ONE* 2010;**5**:e8994.
- S56. Zhang Y, Wang J, Wang X, Gao T, Tian H, Zhou D et al. The autophagic-lysosomal and ubiquitin proteasome systems are simultaneously activated in the skeletal muscle of gastric cancer patients with cachexia. *Am J Clin Nutr* 2020;**111**:570-579.
- S57. Yuan L, Han J, Meng Q, Xi Q, Zhuang Q, Jiang Y et al. Muscle-specific E3 ubiquitin ligases are involved in muscle atrophy of cancer cachexia: an in vitro and in vivo study. *Oncol Rep* 2015;**33**:2261-2268.
- S58. Stephens NA, Gallagher IJ, Rooyackers O, Skipworth RJ, Tan BH, Marstrand T et al. Using transcriptomics to identify and validate novel biomarkers of human skeletal muscle cancer cachexia. *Genome Med* 2010;**2**:1.
- S59. Bossola M, Muscaritoli M, Costelli P, Grieco G, Bonelli G, Pacelli F et al. Increased muscle proteasome activity correlates with disease severity in gastric cancer patients. *Ann Surg* 2003;**237**:384-389.
- S60. Bossola M, Muscaritoli M, Costelli P, Bellantone R, Pacelli F, Busquets S et al. Increased muscle ubiquitin mRNA levels in gastric cancer patients. *Am J Physiol Regul Integr Comp Physiol* 2001;**280**:R1518-1523.
- S61. Taskin S, Stumpf VI, Bachmann J, Weber C, Martignoni ME, Friedrich O. Motor protein function in skeletal abdominal muscle of cachectic cancer patients. *J Cell Mol Med* 2014;**18**:69-79.
- S62. Skorokhod A, Bachmann J, Giese NA, Martignoni ME, Krakowski-Roosen H. Real-time imaging cDNA-AFLP transcript profiling of pancreatic cancer patients: Egr-1 as a potential key regulator of muscle cachexia. *BMC Cancer* 2012;**12**:265.
- S63. Penna F, Bonetto A, Muscaritoli M, Costamagna D, Minero VG, Bonelli G et al. Muscle atrophy in experimental cancer cachexia: is the IGF-1 signaling pathway involved? *Int J Cancer* 2010;**127**:1706-1717.
- S64. Zhuang P, Zhang J, Wang Y, Zhang M, Song L, Lu Z et al. Reversal of muscle atrophy by Zhimu and Huangbai herb pair via activation of IGF-1/Akt and autophagy signal in cancer cachexia. *Support Care Cancer* 2016;**24**:1189-1198.
- S65. Asp ML, Tian M, Wendel AA, Belury MA. Evidence for the contribution of insulin resistance to the development of cachexia in tumor-bearing mice. *Int J Cancer* 2010;**126**:756-763.

- S66. Bohnert KR, Gallot YS, Sato S, Xiong G, Hindi SM, Kumar A. Inhibition of ER stress and unfolding protein response pathways causes skeletal muscle wasting during cancer cachexia. *FASEB J* 2016;**30**:3053-3068.
- S67. Assi M, Derbre F, Lefeuvre-Orfila L, Rebillard A. Antioxidant supplementation accelerates cachexia development by promoting tumor growth in C26 tumor-bearing mice. *Free Radic Biol Med* 2016;**91**:204-214.
- S68. Bonetto A, Penna F, Minero VG, Reffo P, Bonelli G, Baccino FM et al. Deacetylase inhibitors modulate the myostatin/follistatin axis without improving cachexia in tumor-bearing mice. *Curr Cancer Drug Targets* 2009;**9**:608-616.
- S69. Baltgalvis KA, Berger FG, Pena MM, Davis JM, White JP, Carson JA. Muscle wasting and interleukin-6-induced atrogen-I expression in the cachectic Apc ( Min/+ ) mouse. *Pflugers Arch* 2009;**457**:989-1001.
- S70. Fernandez GJ, Ferreira JH, Vechetti IJ, Jr., de Moraes LN, Cury SS, Freire PP et al. MicroRNA-mRNA co-sequencing identifies transcriptional and post-transcriptional regulatory networks underlying muscle wasting in cancer cachexia. *Front Genet* 2020;**11**:541.
- S71. Fontes-Oliveira CC, Busquets S, Toledo M, Penna F, Paz Aylwin M, Sirisi S et al. Mitochondrial and sarcoplasmic reticulum abnormalities in cancer cachexia: altered energetic efficiency? *Biochim Biophys Acta* 2013;**1830**:2770-2778.
- S72. Fontes-Oliveira CC, Busquets S, Fuster G, Ametller E, Figueras M, Oliván M et al. A differential pattern of gene expression in skeletal muscle of tumor-bearing rats reveals dysregulation of excitation-contraction coupling together with additional muscle alterations. *Muscle Nerve* 2014;**49**:233-248.
- S73. Hall DT, Griss T, Ma JF, Sanchez BJ, Sadek J, Tremblay AMK et al. The AMPK agonist 5-aminoimidazole-4-carboxamide ribonucleotide (AICAR), but not metformin, prevents inflammation-associated cachectic muscle wasting. *EMBO Mol Med* 2018;**10**:e8307.
- S74. Ham DJ, Murphy KT, Chee A, Lynch GS, Koopman R. Glycine administration attenuates skeletal muscle wasting in a mouse model of cancer cachexia. *Clin Nutr* 2014;**33**:448-458.
- S75. Mastrocola R, Reffo P, Penna F, Tomasinelli CE, Boccuzzi G, Baccino FM et al. Muscle wasting in diabetic and in tumor-bearing rats: role of oxidative stress. *Free Radic Biol Med* 2008;**44**:584-593.
- S76. Matsuyama T, Ishikawa T, Okayama T, Oka K, Adachi S, Mizushima K et al. Tumor inoculation site affects the development of cancer cachexia and muscle wasting. *Int J Cancer* 2015;**137**:2558-2565.
- S77. Murphy KT, Chee A, Trieu J, Naim T, Lynch GS. Importance of functional and metabolic impairments in the characterization of the C-26 murine model of cancer cachexia. *Dis Model Mech* 2012;**5**:533-545.
- S78. Pin F, Busquets S, Toledo M, Camperi A, Lopez-Soriano FJ, Costelli P et al. Combination of exercise training and erythropoietin prevents cancer-induced muscle alterations. *Oncotarget* 2015;**6**:43202-43215.
- S79. Re Cecconi AD, Forti M, Chiappa M, Zhu Z, Zingman LV, Cervo L et al. Musclin, a myokine induced by aerobic exercise, retards muscle atrophy during cancer cachexia in mice. *Cancers (Basel)* 2019;**11**:
- S80. Roberts EW, Deonaraine A, Jones JO, Denton AE, Feig C, Lyons SK et al. Depletion of stromal cells expressing fibroblast activation protein-alpha from skeletal muscle and bone marrow results in cachexia and anemia. *J Exp Med* 2013;**210**:1137-1151.
- S81. Roberts BM, Ahn B, Smuder AJ, Al-Rajhi M, Gill LC, Beharry AW et al. Diaphragm and ventilatory dysfunction during cancer cachexia. *FASEB J* 2013;**27**:2600-2610.



- S82. Shukla SK, Markov SD, Attri KS, Vernucci E, King RJ, Dasgupta A et al. Macrophages potentiate STAT3 signaling in skeletal muscles and regulate pancreatic cancer cachexia. *Cancer Lett* 2020;**484**:29-39.
- S83. Tseng YC, Kulp SK, Lai IL, Hsu EC, He WA, Frankhouser DE et al. Preclinical investigation of the novel histone deacetylase inhibitor AR-42 in the treatment of cancer-induced cachexia. *J Natl Cancer Inst* 2015;**107**:d1v274.
- S84. Tzika AA, Fontes-Oliveira CC, Shestov AA, Constantinou C, Psychogios N, Righi V et al. Skeletal muscle mitochondrial uncoupling in a murine cancer cachexia model. *Int J Oncol* 2013;**43**:886-894.
- S85. Zhang L, Tang H, Kou Y, Li R, Zheng Y, Wang Q et al. MG132-mediated inhibition of the ubiquitin-proteasome pathway ameliorates cancer cachexia. *J Cancer Res Clin Oncol* 2013;**139**:1105-1115.
- S86. Aulino P, Berardi E, Cardillo VM, Rizzuto E, Perniconi B, Ramina C et al. Molecular, cellular and physiological characterization of the cancer cachexia-inducing C26 colon carcinoma in mouse. *BMC Cancer* 2010;**10**:363.
- S87. Chen JL, Walton KL, Hagg A, Colgan TD, Johnson K, Qian H et al. Specific targeting of TGF-beta family ligands demonstrates distinct roles in the regulation of muscle mass in health and disease. *Proc Natl Acad Sci U S A* 2017;**114**:E5266-E5275.
- S88. Paul PK, Gupta SK, Bhatnagar S, Panguluri SK, Darnay BG, Choi Y et al. Targeted ablation of TRAF6 inhibits skeletal muscle wasting in mice. *J Cell Biol* 2010;**191**:1395-1411.
- S89. Shadfar S, Couch ME, McKinney KA, Weinstein LJ, Yin X, Rodriguez JE et al. Oral resveratrol therapy inhibits cancer-induced skeletal muscle and cardiac atrophy in vivo. *Nutr Cancer* 2011;**63**:749-762.
- S90. Damrauer JS, Stadler ME, Acharyya S, Baldwin AS, Couch ME, Guttridge DC. Chemotherapy-induced muscle wasting: association with NF-kappaB and cancer cachexia. *Eur J Transl Myol* 2018;**28**:7590.
- S91. Padrao AI, Oliveira P, Vitorino R, Colaco B, Pires MJ, Marquez M et al. Bladder cancer-induced skeletal muscle wasting: disclosing the role of mitochondria plasticity. *Int J Biochem Cell Biol* 2013;**45**:1399-1409.
- S92. Bae T, Jang J, Lee H, Song J, Chae S, Park M et al. Paeonia lactiflora root extract suppresses cancer cachexia by down-regulating muscular NF-kappaB signalling and muscle-specific E3 ubiquitin ligases in cancer-bearing mice. *J Ethnopharmacol* 2020;**246**:112222.
- S93. Liu D, Qiao X, Ge Z, Shang Y, Li Y, Wang W et al. IMB0901 inhibits muscle atrophy induced by cancer cachexia through MSTN signaling pathway. *Skelet Muscle* 2019;**9**:8.
- S94. Puppa MJ, Gao S, Narsale AA, Carson JA. Skeletal muscle glycoprotein 130's role in Lewis lung carcinoma-induced cachexia. *FASEB J* 2014;**28**:998-1009.
- S95. Fujita J, Tsujinaka T, Yano M, Ebisui C, Saito H, Katsume A et al. Anti-interleukin-6 receptor antibody prevents muscle atrophy in colon-26 adenocarcinoma-bearing mice with modulation of lysosomal and ATP-ubiquitin-dependent proteolytic pathways. *Int J Cancer* 1996;**68**:637-643.
- S96. Busquets S, Garcia-Martinez C, Alvarez B, Carbo N, Lopez-Soriano FJ, Argiles JM. Calpain-3 gene expression is decreased during experimental cancer cachexia. *Biochim Biophys Acta* 2000;**1475**:5-9.
- S97. Busquets S, Toledo M, Marmonti E, Orpi M, Capdevila E, Betancourt A et al. Formoterol treatment downregulates the myostatin system in skeletal muscle of cachectic tumour-bearing rats. *Oncol Lett* 2012;**3**:185-189.

- S98. Moore-Carrasco R, Busquets S, Almendro V, Palanki M, Lopez-Soriano FJ, Argiles JM. The AP-1/NF-kappaB double inhibitor SP100030 can revert muscle wasting during experimental cancer cachexia. *Int J Oncol* 2007;**30**:1239-1245.
- S99. Solomon V, Baracos V, Sarraf P, Goldberg AL. Rates of ubiquitin conjugation increase when muscles atrophy, largely through activation of the N-end rule pathway. *Proc Natl Acad Sci U S A* 1998;**95**:12602-12607.
- S100. Russell ST, Siren PM, Siren MJ, Tisdale MJ. Attenuation of skeletal muscle atrophy in cancer cachexia by D-myo-inositol 1,2,6-triphosphate. *Cancer Chemother Pharmacol* 2009;**64**:517-527.
- S101. Eley HL, Russell ST, Tisdale MJ. Attenuation of muscle atrophy in a murine model of cachexia by inhibition of the dsRNA-dependent protein kinase. *Br J Cancer* 2007;**96**:1216-1222.
- S102. Blackwell TA, Cervenka I, Khatri B, Brown JL, Rosa-Caldwell ME, Lee DE et al. Transcriptomic analysis of the development of skeletal muscle atrophy in cancer-cachexia in tumor-bearing mice. *Physiol Genomics* 2018;**50**:1071-1082.
- S103. Shum AM, Fung DC, Corley SM, McGill MC, Bentley NL, Tan TC et al. Cardiac and skeletal muscles show molecularly distinct responses to cancer cachexia. *Physiol Genomics* 2015;**47**:588-599.
- S104. Costelli P, Bossola M, Muscaritoli M, Grieco G, Bonelli G, Bellantone R et al. Anticytokine treatment prevents the increase in the activity of ATP-ubiquitin- and Ca(2+)-dependent proteolytic systems in the muscle of tumour-bearing rats. *Cytokine* 2002;**19**:1-5.
- S105. Sartori R, Schirwis E, Blaauw B, Bortolanza S, Zhao J, Enzo E et al. BMP signaling controls muscle mass. *Nat Genet* 2013;**45**:1309-1318.
- S106. Milan G, Romanello V, Pescatore F, Armani A, Paik JH, Frasson L et al. Regulation of autophagy and the ubiquitin-proteasome system by the FoxO transcriptional network during muscle atrophy. *Nat Commun* 2015;**6**:6670.
- S107. Masiero E, Agatea L, Mammucari C, Blaauw B, Loro E, Komatsu M et al. Autophagy is required to maintain muscle mass. *Cell Metab* 2009;**10**:507-515.
- S108. Mammucari C, Milan G, Romanello V, Masiero E, Rudolf R, Del Piccolo P et al. FoxO3 controls autophagy in skeletal muscle in vivo. *Cell Metab* 2007;**6**:458-471.
- S109. Zhao J, Brault JJ, Schild A, Cao P, Sandri M, Schiaffino S et al. FoxO3 coordinately activates protein degradation by the autophagic/lysosomal and proteasomal pathways in atrophying muscle cells. *Cell Metab* 2007;**6**:472-483.
- S110. Sanchez-Martin P, Komatsu M. p62/SQSTM1 - steering the cell through health and disease. *J Cell Sci* 2018;**131**:
- S111. Sardiello M, Palmieri M, di Ronza A, Medina DL, Valenza M, Gennarino VA et al. A gene network regulating lysosomal biogenesis and function. *Science* 2009;**325**:473-477.
- S112. Hentila J, Nissinen TA, Korkmaz A, Lensu S, Silvennoinen M, Pasternack A et al. Activin receptor ligand blocking and cancer have distinct effects on protein and redox homeostasis in skeletal muscle and liver. *Front Physiol* 2018;**9**:1917.
- S113. Grumati P, Dikic I. Ubiquitin signaling and autophagy. *J Biol Chem* 2018;**293**:5404-5413.
- S114. Marzetti E, Lorenzi M, Landi F, Picca A, Rosa F, Tanganelli F et al. Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. *Exp Gerontol* 2017;**87**:92-99.
- S115. Hyatt HW, Powers SK. The role of calpains in skeletal muscle remodeling with exercise and inactivity-induced atrophy. *Int J Sports Med* 2020;**41**:994-1008.
- S116. Huang J, Forsberg NE. Role of calpain in skeletal-muscle protein degradation. *Proc Natl Acad Sci U S A* 1998;**95**:12100-12105.

- S117. Smith IJ, Aversa Z, Hasselgren PO, Pacelli F, Rosa F, Doglietto GB et al. Calpain activity is increased in skeletal muscle from gastric cancer patients with no or minimal weight loss. *Muscle Nerve* 2011;**43**:410-414.
- S118. Tardif N, Klaude M, Lundell L, Thorell A, Rooyackers O. Autophagic-lysosomal pathway is the main proteolytic system modified in the skeletal muscle of esophageal cancer patients. *Am J Clin Nutr* 2013;**98**:1485-1492.
- S119. Costelli P, De Tullio R, Baccino FM, Melloni E. Activation of Ca(2+)-dependent proteolysis in skeletal muscle and heart in cancer cachexia. *Br J Cancer* 2001;**84**:946-950.
- S120. Sjogren K, Liu JL, Blad K, Skrtic S, Vidal O, Wallenius V et al. Liver-derived insulin-like growth factor I (IGF-I) is the principal source of IGF-I in blood but is not required for postnatal body growth in mice. *Proc Natl Acad Sci U S A* 1999;**96**:7088-7092.
- S121. Stratikopoulos E, Szabolcs M, Dragatsis I, Klinakis A, Efstratiadis A. The hormonal action of IGF1 in postnatal mouse growth. *Proc Natl Acad Sci U S A* 2008;**105**:19378-19383.
- S122. Ascenzi F, Barberi L, Dobrowolny G, Villa Nova Bacurau A, Nicoletti C, Rizzuto E et al. Effects of IGF-1 isoforms on muscle growth and sarcopenia. *Aging Cell* 2019;**18**:e12954.
- S123. Simons JP, Schols AM, Buurman WA, Wouters EF. Weight loss and low body cell mass in males with lung cancer: relationship with systemic inflammation, acute-phase response, resting energy expenditure, and catabolic and anabolic hormones. *Clin Sci (Lond)* 1999;**97**:215-223.
- S124. Tessitore L, Costelli P, Baccino FM. Humoral mediation for cachexia in tumour-bearing rats. *Br J Cancer* 1993;**67**:15-23.
- S125. Liu GY, Sabatini DM. mTOR at the nexus of nutrition, growth, ageing and disease. *Nat Rev Mol Cell Biol* 2020;**21**:183-203.
- S126. Bodine SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R et al. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol* 2001;**3**:1014-1019.
- S127. Stretch C, Aubin JM, Mickiewicz B, Leugner D, Al-Manasra T, Tobola E et al. Sarcopenia and myosteatosis are accompanied by distinct biological profiles in patients with pancreatic and periampullary adenocarcinomas. *PLoS One* 2018;**13**:e0196235.
- S128. Eley HL, Russell ST, Tisdale MJ. Effect of branched-chain amino acids on muscle atrophy in cancer cachexia. *Biochem J* 2007;**407**:113-120.
- S129. Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L et al. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science* 2008;**320**:1496-1501.
- S130. Bolster DR, Crozier SJ, Kimball SR, Jefferson LS. AMP-activated protein kinase suppresses protein synthesis in rat skeletal muscle through down-regulated mammalian target of rapamycin (mTOR) signaling. *J Biol Chem* 2002;**277**:23977-23980.
- S131. Calissi G, Lam EW, Link W. Therapeutic strategies targeting FOXO transcription factors. *Nat Rev Drug Discov* 2021;**20**:21-38.
- S132. Sandri M, Sandri C, Gilbert A, Skurk C, Calabria E, Picard A et al. Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy. *Cell* 2004;**117**:399-412.
- S133. Stitt TN, Drujan D, Clarke BA, Panaro F, Timofeyeva Y, Kline WO et al. The IGF-1/PI3K/Akt pathway prevents expression of muscle atrophy-induced ubiquitin ligases by inhibiting FOXO transcription factors. *Mol Cell* 2004;**14**:395-403.

- S134. Bohnert KR, McMillan JD, Kumar A. Emerging roles of ER stress and unfolded protein response pathways in skeletal muscle health and disease. *J Cell Physiol* 2018;**233**:67-78.
- S135. Almanza A, Carlesso A, Chintla C, Creedican S, Doultzinos D, Leuzzi B et al. Endoplasmic reticulum stress signalling - from basic mechanisms to clinical applications. *FEBS J* 2019;**286**:241-278.
- S136. Hetz C, Zhang K, Kaufman RJ. Mechanisms, regulation and functions of the unfolded protein response. *Nat Rev Mol Cell Biol* 2020;
- S137. Gallot YS, Bohnert KR. Confounding roles of ER stress and the unfolded protein response in skeletal muscle atrophy. *Int J Mol Sci* 2021;**22**:
- S138. Gallot YS, Bohnert KR, Straughn AR, Xiong G, Hindi SM, Kumar A. PERK regulates skeletal muscle mass and contractile function in adult mice. *FASEB J* 2019;**33**:1946-1962.
- S139. Bohnert KR, Goli P, Roy A, Sharma AK, Xiong G, Gallot YS et al. The Toll-Like receptor/MyD88/XBP1 signaling axis mediates skeletal muscle wasting during cancer cachexia. *Mol Cell Biol* 2019;**39**:e00184-00119.
- S140. Qiu L, Chen W, Wu C, Yuan Y, Li Y. Exosomes of oral squamous cell carcinoma cells containing miR-181a-3p induce muscle cell atrophy and apoptosis by transmissible endoplasmic reticulum stress signaling. *Biochem Biophys Res Commun* 2020;**533**:831-837.
- S141. Straughn AR, Kelm NQ, Kakar SS. Withaferin A and ovarian cancer antagonistically regulate skeletal muscle mass. *Front Cell Dev Biol* 2021;**9**:636498.
- S142. Harding HP, Zhang Y, Ron D. Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature* 1999;**397**:271-274.
- S143. Qin L, Wang Z, Tao L, Wang Y. ER stress negatively regulates AKT/TSC/mTOR pathway to enhance autophagy. *Autophagy* 2010;**6**:239-247.
- S144. Ebert SM, Dyle MC, Kunkel SD, Bullard SA, Bongers KS, Fox DK et al. Stress-induced skeletal muscle Gadd45a expression reprograms myonuclei and causes muscle atrophy. *J Biol Chem* 2012;**287**:27290-27301.
- S145. Le Moal E, Pialoux V, Juban G, Groussard C, Zouhal H, Chazaud B et al. Redox control of skeletal muscle regeneration. *Antioxid Redox Signal* 2017;**27**:276-310.
- S146. Dobrowolny G, Aucello M, Rizzuto E, Beccafico S, Mammucari C, Boncompagni S et al. Skeletal muscle is a primary target of SOD1G93A-mediated toxicity. *Cell Metab* 2008;**8**:425-436.
- S147. Ramamoorthy S, Donohue M, Buck M. Decreased Jun-D and myogenin expression in muscle wasting of human cachexia. *Am J Physiol Endocrinol Metab* 2009;**297**:E392-401.
- S148. Sullivan-Gunn MJ, Campbell-O'Sullivan SP, Tisdale MJ, Lewandowski PA. Decreased NADPH oxidase expression and antioxidant activity in cachectic skeletal muscle. *J Cachexia Sarcopenia Muscle* 2011;**2**:181-188.
- S149. Padilha CS, Borges FH, Costa Mendes da Silva LE, Frajacomo FTT, Jordao AA, Duarte JA et al. Resistance exercise attenuates skeletal muscle oxidative stress, systemic pro-inflammatory state, and cachexia in Walker-256 tumor-bearing rats. *Appl Physiol Nutr Metab* 2017;**42**:916-923.
- S150. Gilliam LA, Lark DS, Reese LR, Torres MJ, Ryan TE, Lin CT et al. Targeted overexpression of mitochondrial catalase protects against cancer chemotherapy-induced skeletal muscle dysfunction. *Am J Physiol Endocrinol Metab* 2016;**311**:E293-301.
- S151. Ushmorov A, Hack V, Droge W. Differential reconstitution of mitochondrial respiratory chain activity and plasma redox state by cysteine and ornithine in a model of cancer cachexia. *Cancer Res* 1999;**59**:3527-3534.

- S152. Der-Torossian H, Wysong A, Shadfar S, Willis MS, McDunn J, Couch ME. Metabolic derangements in the gastrocnemius and the effect of Compound A therapy in a murine model of cancer cachexia. *J Cachexia Sarcopenia Muscle* 2013;**4**:145-155.
- S153. Fermoselle C, Garcia-Arumi E, Puig-Vilanova E, Andreu AL, Urtreger AJ, de Kier Joffe ED et al. Mitochondrial dysfunction and therapeutic approaches in respiratory and limb muscles of cancer cachectic mice. *Exp Physiol* 2013;**98**:1349-1365.
- S154. Dickinson BC, Chang CJ. Chemistry and biology of reactive oxygen species in signaling or stress responses. *Nat Chem Biol* 2011;**7**:504-511.
- S155. Weerapana E, Wang C, Simon GM, Richter F, Khare S, Dillon MB et al. Quantitative reactivity profiling predicts functional cysteines in proteomes. *Nature* 2010;**468**:790-795.
- S156. Deans DA, Tan BH, Wigmore SJ, Ross JA, de Beaux AC, Paterson-Brown S et al. The influence of systemic inflammation, dietary intake and stage of disease on rate of weight loss in patients with gastro-oesophageal cancer. *Br J Cancer* 2009;**100**:63-69.
- S157. Abbass T, Dolan RD, Laird BJ, McMillan DC. The relationship between imaging-based body composition analysis and the systemic inflammatory response in patients with cancer: a systematic review. *Cancers (Basel)* 2019;**11**:1304.
- S158. Scheede-Bergdahl C, Watt HL, Trutschnigg B, Kilgour RD, Haggarty A, Lucar E et al. Is IL-6 the best pro-inflammatory biomarker of clinical outcomes of cancer cachexia? *Clin Nutr* 2012;**31**:85-88.
- S159. Op den Kamp CM, Gosker HR, Lagarde S, Tan DY, Snepvangers FJ, Dingemans AM et al. Preserved muscle oxidative metabolic phenotype in newly diagnosed non-small cell lung cancer cachexia. *J Cachexia Sarcopenia Muscle* 2015;**6**:164-173.
- S160. Krzystek-Korpacka M, Matusiewicz M, Diakowska D, Grabowski K, Blachut K, Kustrzeba-Wojcicka I et al. Impact of weight loss on circulating IL-1, IL-6, IL-8, TNF-alpha, VEGF-A, VEGF-C and midkine in gastroesophageal cancer patients. *Clin Biochem* 2007;**40**:1353-1360.
- S161. Pettersen K, Andersen S, Degen S, Tadini V, Grosjean J, Hatakeyama S et al. Cancer cachexia associates with a systemic autophagy-inducing activity mimicked by cancer cell-derived IL-6 trans-signaling. *Sci Rep* 2017;**7**:2046.
- S162. Pfitzenmaier J, Vessella R, Higano CS, Noteboom JL, Wallace D, Jr., Corey E. Elevation of cytokine levels in cachectic patients with prostate carcinoma. *Cancer* 2003;**97**:1211-1216.
- S163. Penafuerte CA, Gagnon B, Sirois J, Murphy J, MacDonald N, Tremblay ML. Identification of neutrophil-derived proteases and angiotensin II as biomarkers of cancer cachexia. *Br J Cancer* 2016;**114**:680-687.
- S164. Bossola M, Muscaritoli M, Bellantone R, Pacelli F, Cascino A, Sgadari A et al. Serum tumour necrosis factor-alpha levels in cancer patients are discontinuous and correlate with weight loss. *Eur J Clin Invest* 2000;**30**:1107-1112.
- S165. Shum AM, Mahendradatta T, Taylor RJ, Painter AB, Moore MM, Tsoli M et al. Disruption of MEF2C signaling and loss of sarcomeric and mitochondrial integrity in cancer-induced skeletal muscle wasting. *Aging (Albany NY)* 2012;**4**:133-143.
- S166. Strassmann G, Fong M, Kenney JS, Jacob CO. Evidence for the involvement of interleukin 6 in experimental cancer cachexia. *J Clin Invest* 1992;**89**:1681-1684.
- S167. Zhou W, Jiang ZW, Tian J, Jiang J, Li N, Li JS. Role of NF-kappaB and cytokine in experimental cancer cachexia. *World J Gastroenterol* 2003;**9**:1567-1570.
- S168. Penna F, Busquets S, Pin F, Toledo M, Baccino FM, Lopez-Soriano FJ et al. Combined approach to counteract experimental cancer cachexia: eicosapentaenoic acid and training exercise. *J Cachexia Sarcopenia Muscle* 2011;**2**:95-104.

- S169. Dillon EL, Volpi E, Wolfe RR, Sinha S, Sanford AP, Arrastia CD et al. Amino acid metabolism and inflammatory burden in ovarian cancer patients undergoing intense oncological therapy. *Clin Nutr* 2007;**26**:736-743.
- S170. Prokopchuk O, Steinacker JM, Nitsche U, Otto S, Bachmann J, Schubert EC et al. IL-4 mRNA is downregulated in the liver of pancreatic cancer patients suffering from cachexia. *Nutr Cancer* 2017;**69**:84-91.
- S171. Gilabert M, Calvo E, Airolidi A, Hamidi T, Moutardier V, Turrini O et al. Pancreatic cancer-induced cachexia is Jak2-dependent in mice. *J Cell Physiol* 2014;**229**:1437-1443.
- S172. Hunter RB, Stevenson E, Koncarevic A, Mitchell-Felton H, Essig DA, Kandarian SC. Activation of an alternative NF-kappaB pathway in skeletal muscle during disuse atrophy. *FASEB J* 2002;**16**:529-538.
- S173. Cai D, Frantz JD, Tawa NE, Jr., Melendez PA, Oh BC, Lidov HG et al. IKKbeta/NF-kappaB activation causes severe muscle wasting in mice. *Cell* 2004;**119**:285-298.
- S174. Costelli P, Muscaritoli M, Bossola M, Moore-Carrasco R, Crepaldi S, Grieco G et al. Skeletal muscle wasting in tumor-bearing rats is associated with MyoD down-regulation. *Int J Oncol* 2005;**26**:1663-1668.
- S175. Schindler C, Levy DE, Decker T. JAK-STAT signaling: from interferons to cytokines. *J Biol Chem* 2007;**282**:20059-20063.
- S176. Moresi V, Adamo S, Berghella L. The JAK/STAT pathway in skeletal muscle pathophysiology. *Front Physiol* 2019;**10**:500.
- S177. Zhang L, Pan J, Dong Y, Twardy DJ, Dong Y, Garibotto G et al. Stat3 activation links a C/EBPdelta to myostatin pathway to stimulate loss of muscle mass. *Cell Metab* 2013;**18**:368-379.
- S178. Silva KA, Dong J, Dong Y, Schor N, Twardy DJ et al. Inhibition of Stat3 activation suppresses caspase-3 and the ubiquitin-proteasome system, leading to preservation of muscle mass in cancer cachexia. *J Biol Chem* 2015;**290**:11177-11187.
- S179. McPherron AC, Lawler AM, Lee SJ. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 1997;**387**:83-90.
- S180. McPherron AC, Lee SJ. Double muscling in cattle due to mutations in the myostatin gene. *Proc Natl Acad Sci U S A* 1997;**94**:12457-12461.
- S181. Schuelke M, Wagner KR, Stolz LE, Hubner C, Riebel T, Komen W et al. Myostatin mutation associated with gross muscle hypertrophy in a child. *N Engl J Med* 2004;**350**:2682-2688.
- S182. Lee SJ, Lehar A, Liu Y, Ly CH, Pham QM, Michaud M et al. Functional redundancy of type I and type II receptors in the regulation of skeletal muscle growth by myostatin and activin A. *Proc Natl Acad Sci U S A* 2020;**117**:30907-30917.
- S183. Han HQ, Zhou X, Mitch WE, Goldberg AL. Myostatin/activin pathway antagonism: Molecular basis and therapeutic potential. *Int J Biochem Cell Biol* 2013;**45**:2333-2347.
- S184. Breitbart A, Scharf GM, Duncker D, Widera C, Gottlieb J, Vogel A et al. Highly specific detection of myostatin prodomain by an immunoradiometric sandwich assay in serum of healthy individuals and patients. *PLoS One* 2013;**8**:e80454.
- S185. Talar-Wojnarowska R, Wozniak M, Borkowska A, Olakowski M, Malecka-Panas E. Clinical significance of activin A and myostatin in patients with pancreatic adenocarcinoma and progressive weight loss. *J Physiol Pharmacol* 2020;**71**:105-112.
- S186. Hill NE, Saeed S, Phadke R, Ellis MJ, Chambers D, Wilson DR et al. Detailed characterization of a long-term rodent model of critical illness and recovery. *Crit Care Med* 2015;**43**:e84-96.

- S187. Costelli P, Muscaritoli M, Bonetto A, Penna F, Reffo P, Bossola M et al. Muscle myostatin signalling is enhanced in experimental cancer cachexia. *Eur J Clin Invest* 2008;**38**:531-538.
- S188. Liu CM, Yang Z, Liu CW, Wang R, Tien P, Dale R et al. Myostatin antisense RNA-mediated muscle growth in normal and cancer cachexia mice. *Gene Ther* 2008;**15**:155-160.
- S189. Latres E, Mastaitis J, Fury W, Miloscio L, Trejos J, Pangilinan J et al. Activin A more prominently regulates muscle mass in primates than does GDF8. *Nat Commun* 2017;**8**:15153.
- S190. Chen JL, Walton KL, Winbanks CE, Murphy KT, Thomson RE, Makanji Y et al. Elevated expression of activins promotes muscle wasting and cachexia. *FASEB J* 2014;**28**:1711-1723.
- S191. Lerner L, Gyuris J, Nicoletti R, Gifford J, Krieger B, Jatoi A. Growth differentiating factor-15 (GDF-15): A potential biomarker and therapeutic target for cancer-associated weight loss. *Oncol Lett* 2016;**12**:4219-4223.
- S192. Zhong X, Pons M, Poirier C, Jiang Y, Liu J, Sandusky GE et al. The systemic activin response to pancreatic cancer: implications for effective cancer cachexia therapy. *J Cachexia Sarcopenia Muscle* 2019;**10**:1083-1101.
- S193. Otani T, Minami S, Yamoto M, Umesaki N. Production of activin A in hyperplasia and adenocarcinoma of the human endometrium. *Gynecol Oncol* 2001;**83**:31-38.
- S194. Wildi S, Kleeff J, Maruyama H, Maurer CA, Buchler MW, Korc M. Overexpression of activin A in stage IV colorectal cancer. *Gut* 2001;**49**:409-417.
- S195. Winbanks CE, Chen JL, Qian H, Liu Y, Bernardo BC, Beyer C et al. The bone morphogenetic protein axis is a positive regulator of skeletal muscle mass. *J Cell Biol* 2013;**203**:345-357.
- S196. Kuo T, Lew MJ, Mayba O, Harris CA, Speed TP, Wang JC. Genome-wide analysis of glucocorticoid receptor-binding sites in myotubes identifies gene networks modulating insulin signaling. *Proc Natl Acad Sci U S A* 2012;**109**:11160-11165.
- S197. Bodine SC, Furlow JD. Glucocorticoids and Skeletal Muscle. *Adv Exp Med Biol* 2015;**872**:145-176.
- S198. Schakman O, Kalista S, Barbe C, Loumaye A, Thissen JP. Glucocorticoid-induced skeletal muscle atrophy. *Int J Biochem Cell Biol* 2013;**45**:2163-2172.
- S199. Knapp ML, al-Sheibani S, Riches PG, Hanham IW, Phillips RH. Hormonal factors associated with weight loss in patients with advanced breast cancer. *Ann Clin Biochem* 1991;**28**:480-486.
- S200. Cala MP, Agullo-Ortuno MT, Prieto-Garcia E, Gonzalez-Riano C, Parrilla-Rubio L, Barbas C et al. Multiplatform plasma fingerprinting in cancer cachexia: a pilot observational and translational study. *J Cachexia Sarcopenia Muscle* 2018;**9**:348-357.
- S201. Rivadeneira DE, Naama HA, McCarter MD, Fujita J, Evoy D, Mackrell P et al. Glucocorticoid blockade does not abrogate tumor-induced cachexia. *Nutr Cancer* 1999;**35**:202-206.
- S202. Russell ST, Tisdale MJ. The role of glucocorticoids in the induction of zinc-alpha2-glycoprotein expression in adipose tissue in cancer cachexia. *Br J Cancer* 2005;**92**:876-881.
- S203. Tanaka Y, Eda H, Tanaka T, Udagawa T, Ishikawa T, Horii I et al. Experimental cancer cachexia induced by transplantable colon 26 adenocarcinoma in mice. *Cancer Res* 1990;**50**:2290-2295.
- S204. Crespigio J, Weidmann R, Macioszek M, de Oliveira J, de Souza M, Pignatelli D. Impaired glucocorticoid synthesis in cancer cachexia-anorexia syndrome in an experimental model. *Annals of Clinical & Experimental Metabolism* 2016;**1**:1008.

- S205. Sarason EL. Adrenal cortex in systemic disease: a morphological study. *Arch Intern Med* 1943;**71**:702-712.
- S206. Ball HA, Samuels LT. Adrenal weights in tumor-bearing rats. *Proc Soc Exp Biol Med* 1938;**38**:441-443.
- S207. Schmidt K, von Haehling S, Doehner W, Palus S, Anker SD, Springer J. IGF-1 treatment reduces weight loss and improves outcome in a rat model of cancer cachexia. *J Cachexia Sarcopenia Muscle* 2011;**2**:105-109.
- S208. Garcia JM, Polvino WJ. Pharmacodynamic hormonal effects of anamorelin, a novel oral ghrelin mimetic and growth hormone secretagogue in healthy volunteers. *Growth Horm IGF Res* 2009;**19**:267-273.
- S209. Pietra C, Takeda Y, Tazawa-Ogata N, Minami M, Yuanfeng X, Duus EM et al. Anamorelin HCl (ONO-7643), a novel ghrelin receptor agonist, for the treatment of cancer anorexia-cachexia syndrome: preclinical profile. *J Cachexia Sarcopenia Muscle* 2014;**5**:329-337.
- S210. Temel JS, Abernethy AP, Currow DC, Friend J, Duus EM, Yan Y et al. Anamorelin in patients with non-small-cell lung cancer and cachexia (ROMANA 1 and ROMANA 2): results from two randomised, double-blind, phase 3 trials. *Lancet Oncol* 2016;**17**:519-531.
- S211. Takayama K, Katakami N, Yokoyama T, Atagi S, Yoshimori K, Kagamu H et al. Anamorelin (ONO-7643) in Japanese patients with non-small cell lung cancer and cachexia: results of a randomized phase 2 trial. *Support Care Cancer* 2016;**24**:3495-3505.
- S212. Bonetto A, Aydogdu T, Jin X, Zhang Z, Zhan R, Puzis L et al. JAK/STAT3 pathway inhibition blocks skeletal muscle wasting downstream of IL-6 and in experimental cancer cachexia. *Am J Physiol Endocrinol Metab* 2012;**303**:E410-421.
- S213. Svaninger G, Gelin J, Lundholm K. Tumor-host wasting not explained by adrenal hyperfunction in tumor-bearing animals. *J Natl Cancer Inst* 1987;**79**:1135-1141.
- S214. Gosselin C, Cabanac M. Adrenalectomy lowers the body weight set-point in rats. *Physiol Behav* 1997;**62**:519-523.
- S215. Ballaro R, Costelli P, Penna F. Animal models for cancer cachexia. *Curr Opin Support Palliat Care* 2016;**10**:281-287.