

Modeling the HDO cycle with the LMD Mars GCM

Margaux Vals, Loïc Rossi, Franck Montmessin, François Forget, Ehouarn Millour, Kevin Olsen, Anna Fedorova, Alexander Trokhimovskiy, Oleg Korablev

▶ To cite this version:

Margaux Vals, Loïc Rossi, Franck Montmessin, François Forget, Ehouarn Millour, et al.. Modeling the HDO cycle with the LMD Mars GCM. Europlanet Science Congress 2020, Sep 2020, Online, Unknown Region. pp.EPSC2020-680, 10.5194/epsc2020-680. hal-04492906

HAL Id: hal-04492906

https://hal.science/hal-04492906

Submitted on 6 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Europlanet Science Congress 2020

Virtual meeting 21 September – 9 October 2020

Home > EPSC2020-680

EPSC Abstracts

Vol. 14, EPSC2020-680, 2020, updated on 09 Jan 2024

https://doi.org/10.5194/epsc2020-680 Europlanet Science Congress 2020

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Modeling the HDO cycle with the LMD Mars GCM

Margaux Vals et al.

Abstract

HDO and the D/H ratio are important in order to understand Mars past and present climate, in particu through ages of the Martian water cycle. We present here further modeling developments of the HD (Forget et al. 1999), in continuation of the work presented by L. Rossi in session TP16 ("Modeling of th Storm on the martian HDO cycle."). These improvements are led with the perspective of compar provided by the Atmospheric Chemistry Suite (ACS) on board the ESA/Roscosmos Trace Gas Orbiter.

Introduction

The D/H ratio observed in a planetary atmosphere is a proxy for the ratio of the current water reservo of the planet. The current D/H ratio measured in the Martian atmosphere is at least five that of the Vie (SMOW) (Owen et al. 1988, Encrenaz et al. 2018, Krasnopolsky 2015, Villanueva et al. 2015). This hig derived from the HDO/H_2O abundance ratio, is a precious indicator of the large escape of water fron time. Apart from the mass difference between both isotopes, the differential escape of H and D comes of H_2O over HDO (Cheng et al. 1999) and the Vapor Pressure Isotope Effect (VPIE) that produ condensation (Krasnopolsky, 2000, Bertaux et al. 2001, Fouchet et al. 2000).

Modeling HDO

Although the version of the model used by L. Rossi et al. comprises the main last developments of the of the water cycle (no radiative effect of clouds, simple conversion from vapour to ice to reach the sa consistent comparison with the previous study led by Montmessin et al. 2005. We present here the rewith the complete representation of the water cycle, including the activation of the radiative effect (referring to the parametrization of the different processes of formation of the clouds as nucleation of the water ice growth, dust scavenging end supersaturation) implemented in the model by Navarro et al. 20 have been proved to considerably improve the representation of the water cycle in comparison to availa

1 sur 3 06/03/2024 17:50

The reinforcement of the Hadley circulation by the radiative effect of clouds, so as the involvement obviously affect the transport of HDO and its vertical distribution, which directly impacts the D/H ratio i a GCM simulation run over three martian years reveal a persistent behaviour of the D/H cycle. The D/H is close to the one observed by Montmessin et al. 2005, and also by L. Rossi in the abstract subremphasis of the latitudinal gradient appearing during the Northern Hemisphere Summer, probably of Hadley cell generated by the radiative effect of clouds (see Figure 1). The D/H ratio cycle of the ice phase between what happens during the first and the second part of the martian year. In particular, the high the tropics during the Southern Hemisphere spring and summer, while the corresponding amounts of the (respectively around 10~ppb and 5000~ppb), have to be further investigated (see Figure 1). The vertical also impacted by the modified structure of the clouds led by the integration of the microphysics (see Figure 1).




Figure 1: Zonal mean of the integrated column of D/H vapor and ice ratio over a martian year at the k Ls (Solar Longitude). GCM simulation run with the radiative active water ice clouds and the microphysic




Figure 2: Zonal mean cross sections of HDO vapor, HDO ice, D/H vapor and D/H ice during southern st

Conclusion

While the results of the simple physics of clouds allow a straightforward analysis of the fractionation regarding H_2O , the last simulations obtained with the microphysics of clouds account for a more reacycle and will offer the best basis for interpretation and comparison with the observations $\frac{1}{2} \frac{1}{2} \frac{$

CO Meeting Organizer EPSC 2020 around 120~km and they don't include the parametrizations of the thermospil Gonzalez-Galindo et al. 2009) and the photochemistry (Lefèvre et al. 2004, 2008). The ultimate goal of a complete representation of the deuterium cycle, from its source as HDO in the lower atmosphere thermosphere.

Results of the last achievements will be presented at the conference.

Acknowledgements

ExoMars is the space mission of ESA and Roscosmos. The ACS experiment is led by IKI Space Research acknowledges funding by Roscosmos and CNES. Science operations of ACS are funded by Roscosmos a funded by Federal agency of science organizations (FANO). M.V. acknowledges support from the DIM AC region in support for the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the research (Domaine d'Intérêt Majeur, Astrophysique et Conditions d'Apparition of the condition of th

How to cite: Vals, M., Rossi, L., Montmessin, F., Forget, F., Millour, E., Olsen, K., Fedorova, A., Trokhimo Modeling the HDO cycle with the LMD Mars GCM, Europlanet Science Congress 2020, online, 21 Sep-9 https://doi.org/10.5194/epsc2020-680, 2020.

Contact

Imprint

Contact

Data protection

© 2019 Copernicus GmbH Header image © Europlanet Society



3 sur 3 06/03/2024 17:50