



HAL
open science

Enzymatic and Metabolic Diagnostic of Nitrogen Deficiency in *Arabidopsis thaliana* Wassileskija Accession

T Lemaitre, Laure Gaufichon, Stephanie Boutet-Mercey, Aurélie Christ, Céline Masclaux-daubresse

► To cite this version:

T Lemaitre, Laure Gaufichon, Stephanie Boutet-Mercey, Aurélie Christ, Céline Masclaux-daubresse. Enzymatic and Metabolic Diagnostic of Nitrogen Deficiency in *Arabidopsis thaliana* Wassileskija Accession. *Plant and Cell Physiology*, 2008, 49, pp.1056 - 1065. 10.1093/pcp/pcn081 . hal-04495814

HAL Id: hal-04495814

<https://hal.science/hal-04495814>

Submitted on 27 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enzymatic and metabolic diagnostic of nitrogen deficiency in *Arabidopsis thaliana*
Wassileskija accession

Thomas Lemaître[‡], Laure Gaufichon[‡], Stéphanie Boutet-Mercey, Aurélie Christ and Céline Masclaux-Daubresse*

Unité de Nutrition Azotée des Plantes, UNAP, UR511, INRA, Route de Saint Cyr, F-78000 Versailles

* corresponding author : Céline Masclaux-Daubresse
phone +33 (0)1 30 83 30 88
fax +33 (0)1 30 83 30 96
mail masclaux@versailles.inra.fr

‡ These two authors contributed equally to the paper

Running title:

Nitrogen deficiency in *Arabidopsis*

Key words

Nitrogen limitation, nitrogen remobilisation, glutamine synthetase, nitrate reductase, amino acid, seed composition, asparagine, glutamine

Abbreviations

LN: Low nitrogen nutrition (cultivated upon)

HN: High nitrogen nutrition (cultivated upon)

DW: dry weight

FW: fresh weight

DAS: days after sowing

GABA: γ -aminobutyric acid

NRE: Nitrogen remobilization efficiency

%P: partitioning

RSA: Relative Specific Allocation

Financial supports

The ¹⁵N tracing experiment was performed thanks to the financial support received from the ANR Génoplante ARCOLE program (ANR-05-GPLA-032-05).

Dr Thomas Lemaître was Attaché Temporaire d'Enseignement et de Recherche (ATER), supported by the University of Versailles Saint-Quentin en Yvelines (UVSQ).

Abstract

Adaptation to steady-state low-nutrient availability was investigated by comparing the *Wassileskija* (WS) accession of *Arabidopsis thaliana* grown on 2 mM nitrate or 10 mM nitrate. Low nitrogen conditions led to a limited rosette biomass and seed yield. The latter was mainly due to reduced seed number while seed weight was less affected. However, harvest index was lower upon high nitrate compared to limited nitrate condition. Under nitrogen limiting conditions, nitrate reductase activity was decreased while glutamine synthetase activity was increased due to a higher accumulation of the cytosolic enzyme. The level of nitrogen remobilisation to the seeds was higher under low nitrogen and the vegetative parts of the plants remaining after seed production stored very low residual nitrogen. Through promoting nitrogen remobilisation and recycling pathways, nitrogen limitation modified plant and seed compositions. Rosette leaves contained more sugars and less free amino acids when grown under nitrogen limiting conditions. Compared to high nitrogen, their levels of proline, asparagine and glutamine were decreased. The seed amino acid composition reflected that of the rosette leaves, thus suggesting that phloem loading for seed filling was poorly selective. The major finding of this report was that together with decreasing biomass and yield, nitrogen limitation triggers large modifications in vegetative products and seed quality.

Introduction

The availability of nitrogen and water are two major limiting factors in plant growth (Lea and Azevedo 2006). Plants have a fundamental dependence on inorganic nitrogenous fertilizers, and their quantity, added to soil worldwide annually, is increasing every year (Good et al. 2004). Nitrogen has a major impact on seed and forage production while the quality of these products is strongly influenced by nitrogen availability. Together with yield, one important agronomical trait is the percentage of proteins contained in cereal grains that determines seed nutritional quality, and is defined by the N harvest index (NIH, as N in grain/total N uptake).

To date, most plant cultivars bred for crops have been selected under non-limiting conditions (Bänziger et al. 1997; Presterl et al. 2003) for productivity and seed yield. Numerous studies have reported a negative correlation between grain protein concentration and yield (Beninati and Busch 1992) and references cited therein), thus suggesting that breeding cultivars for a higher yield have resulted in decreasing the NIH.

Nitrogen, as a commercial fertilizer, is one of the most expensive nutrients to supply and it represents the major cost in plant production (Singh 2005). Moreover, crop plants are only able to convert 30-40% of this applied nitrogen to useful food products. This has led to an important ecological problem due to lost nitrogen polluting soils and water. The possibility of lowering fertilizer-input and breeding plants with better nitrogen use efficiency is a major aim (Lea and Azevedo 2006; Hirel et al. 2007). Indeed, improving nitrogen use efficiency, defined as the yield of grain per unit of available nitrogen in the soil, is now the central point of a many physiological and agronomical studies. However, limiting nitrogen input in the field will certainly have other effects other than limiting dry matter production and yield, and will probably also modify the quality of useful food products.

The effect of low nitrogen availability on plant biomass, nitrate uptake, root architecture, has already been widely studied (Lea and Azevedo 2007; Drew and Saker 1976; Brouwer 1962; Van der Werf and Nagel 1996). We know that plants modify their root architecture, changing the lateral-root/primary-root ratio and decreasing the shoot/root ratio at the same time, to forage the soil for nutrients. The influence of nitrogen limitation on the fine-tuning of plant metabolism is less documented. Attention has been mainly paid to plant responses to nitrogen starvation; an abiotic stress that can be encountered by all plants, even in the field, several times during their development cycles (Scheible et al. 2004; Wang et al. 2000). On the other hand, adaptation to steady-state low-nutrient availability is poorly documented (see Forde and Lea 2007 for a review). The available results on the comparison of plant metabolism under low and high nitrogen nutrition are often fragmentary and difficult to compare and analyse since they have been obtained on different plant species, tissues or by using different growth conditions (Fritz et al. 2006; Tercé-Laforgue et al. 2004).

It would be useful to obtain a complete picture of the modifications in plant metabolism when nitrogen is poor or abundant. Transcriptomics is a powerful tool to help analyse and compare the phenotypes of plants and it has been used with success to detect changes and adaptations brought about in mutants and stressed plants, such as phosphate-starved, nitrogen-starved or sugar-induced plants. Enzymatic or metabolic analyses can be used as a diagnostic tool for nutritional problems and to help decipher the effects of a mutation on the ability of plants to use nutrients, such as nitrogen for example. In this report we have compared the metabolism traits of the rosette leaves of *Arabidopsis thaliana* plants grown under limiting or under full nitrate nutrition and characterised the metabolic pathways favoured by an adaptation of plants to low nitrogen conditions. It was found that nitrate nutrition changed considerably metabolite levels and enzyme activities. The data show that the fine-tuning of

plant metabolism is sensitive to nitrate levels and that plant-material quality depends on nitrogen availability. This report provides a metabolic diagnostic of nitrogen deficiency in *Arabidopsis thaliana* that should facilitate the detection or analysis of nitrogen-related problems.

Results and Discussion

Development, biomass and yield is affected by low nitrogen nutrition

Many *Arabidopsis* genotypes show a large variation in traits related to development and fitness between low (2mM nitrate) and high (10mM nitrate) nitrogen growth conditions (Loudet et al., 2003). We observed that nitrogen limitation delayed flowering time from 60 DAS to 66 DAS in the WS accession analysed in this work. After 43 DAS, the fresh weight (FW) of the rosette was significantly higher under HN than under LN, and this difference increased to 6-fold at flowering time (Figure 1A). The dry weight (DW) of the seeds and of the dried harvested vegetative remains (rosette, flowering stems and silique envelopes) that correspond to the stover in crop plants, were respectively 2.3 and 2.9 times higher for HN grown plants than for LN grown plants (Figure 1B). Therefore, HN nutrition appeared to be more valuable for the vegetative organs compared to seed production and harvest index (seeds DW.(whole plant DW)⁻¹) was slightly lower in HN than in LN conditions. Nitrogen availability did not influence single seed DW (Figure 1C) but it increased seed yield more than 2-fold, as a result of higher seed number under HN conditions (Figure 1D). This also suggests that plant fitness is better under HN conditions, whereas single seed biomass is not affected.

Nitrogen nutrition modifies the activities of nitrogen assimilation and nitrogen recycling enzymes

Since, nitrogen limitation did not affect the total soluble protein concentration in rosette leaves (Figure 2A), measured enzyme activities were expressed with respect to protein content (nmol.min⁻¹.mg⁻¹protein).

Nitrogen allocation in the plant depends on several steps of plant metabolism that determine the physiology of a plant. In *Arabidopsis* it is assumed that most of the nitrate, taken up *via* transporters in the roots, is reduced and assimilated in the leaves. Beside nitrate

uptake efficiency, nitrate reduction is the main limiting point for nitrogen assimilation. Nitrate reductase (NR, EC.1.7.1.1) is known to be positively regulated by nitrate availability (Vincentz et al. 1993). Indeed, the total NR activity was higher in HN than in LN grown plants during the rosette lifespan, except at 29 DAS when rosette biomass was similar for HN and LN plants and at 64 DAS when leaf senescence of HN grown plants had reduced NR activity to the same level as LN plants (Figure 2B). NR activity decreased with plant ageing under both conditions however, this decrease occurred earlier in LN plants compared to HN plants. *In vivo* NR activity depends on the activation state of the enzyme, which is controlled through NR phosphorylation. The activation state (see Material and Methods) was high (more than 50%) during the time course of our experiments in both LN and HN rosettes (Figure 2C). At 43 DAS and 48 DAS, the activation state was significantly higher in LN rosettes compared to HN. This suggests that the lower NR activity of LN rosettes was due to a lower NR protein content rather than to an inhibition of the enzyme by phosphorylation.

When nitrate and nitrite are reduced, the ammonium produced is assimilated by glutamine synthetase (GS, EC.6.3.1.2) isoforms. Surprisingly the *in vitro* GS activity was significantly higher in the LN rosettes compared to those from HN plants (this was seen after 43 DAS, Figure 2D). Antibodies raised against a conserved peptide domain of GS1 and GS2 allowed us to detect both isoenzymes using western blots and protein extracts from rosettes grown under HN or LN conditions (Figure 2E). GS1 was clearly detected in all the LN rosette extracts and GS1 amount was increased with rosette ageing. At 64 DAS, the GS1 amount in LN rosettes appeared as high as the GS2 amount. This was not the case for the HN rosette extract. In HN rosettes, the GS1 content detected was clearly lower than the GS2 content. Moreover GS1 decreased with leaf ageing and was almost undetectable at 64 DAS in HN rosettes. Loading HN and LN extracts on the same gel (data not shown) confirmed that GS2 amount was similar in HN and LN rosettes and did not change with ageing. It also confirmed that GS1 amount was higher in LN rosettes than in HN rosettes. The increase of GS1 in LN rosettes with ageing and the decrease of GS1 in the HN rosette with ageing could explain that higher GS activity was measured in LN rosettes.

GS1 isoforms are well known markers of N-remobilisation and senescence (Masclaux et al. 2000; Martin et al. 2005). Five genes in Arabidopsis that are differentially regulated encode GS1. The different GS1 isoforms certainly have different location and roles in nitrogen assimilation and recycling (see Masclaux-Daubresse et al. 2008, for a review). In other plant species, several GS1 isoforms have indeed been detected in different plant tissues such as phloem or mesophyll cells depending on leaf development (Brugière et al. 2000;

Masclaux et al. 2000). The low level and decrease of GS1 in HN rosette extracts suggests that the GS1 isoforms present in young HN rosette are mainly dedicated to nitrogen primary assimilation. The increase of GS1 proteins during LN rosette ageing confirms GS1 as a senescence-related marker in *Arabidopsis* when nitrogen is limited (Diaz et al. 2008). Together with the early decrease of NR activity in the LN rosettes, the high GS1 amount suggests that LN rosettes exhibit senescence-related symptoms early during their development and mobilise nitrogen efficiently. This is consistent with the idea that N-limitation is an exogenous senescence-triggering factor that facilitates N-remobilisation at the whole plant level (Wingler et al. 2004).

Nitrogen deficiency stimulates nitrogen remobilisation to the seeds

Since GS1 proteins were more abundant and appeared earlier in LN rosettes compared to HN rosettes, the effect of LN on N-remobilisation was analysed by ^{15}N tracing experiments to measure the amount of nitrogen mobilised from the rosettes to the seeds. Labelling of the rosettes was performed at the vegetative stage (40 and 42 DAS), 20 days before flower bud emergence and 80 days before seed and vegetative part harvest. This allowed us to assume that all of the provided $^{15}\text{NO}_3^-$ had been assimilated and incorporated into organic molecules, before remobilisation took place.

At seed maturity, the dry vegetative remains and the total seeds were harvested for analysis. ^{15}N enrichment, total nitrogen concentration and dry matter of the vegetative remains and seeds compartments were measured allowing the calculation of the ^{15}N enrichment, total nitrogen concentration and dry matter in the whole plant (vegetative remains + seeds). The partitioning of ^{15}N (P% ^{15}N), nitrogen (P%N) and dry matter (P%DM) in the seeds was calculated as the ratio [seeds / (seeds + vegetative remains)]. The relative specific allocation (RSA) of ^{15}N in the different compartments (seeds, vegetative remains and whole plant) was also calculated and it is a representation of tissue enrichment (Table 2, (Deléens et al. 1994)).

Results showed that nitrogen limitation had no significant effect on P%DM and that seeds represented about 28% of the whole plant dry matter in both 2 mM and 10 mM nitrate grown plants. However, nitrogen limitation strongly modified nitrogen allocation to seeds (P%N) since more than 65 % of the total nitrogen absorbed by the plant during its lifespan was located to the seed compartment of the LN plants, versus 25% for HN plants. This result is coherent with the very low N% remaining in the vegetative remains of LN plants compared

to HN plants, whereas the seed N% are similar in LN and HN seeds. The P%N and %P%¹⁵N were similar for LN plants thus resulting in a $RSA_{seeds}:RSA_{whole-plant}$ close to 1. This underpins a close relationship between $RSA_{vegetative-remains}$ and RSA_{seeds} suggesting that when *Arabidopsis* plants are cultivated with a 2 mM nitrate nutrition, N-remobilisation occurred in a way that seems to follow the model described by Gallais et al. (2006) for maize. This model proposes that quasi equality to 1 observed for the ratio $RSA_{seeds}:RSA_{whole-plant}$ would have existed very early during development when N-remobilisation and N-uptake were both active. There would be no dilution or concentration of ¹⁵N in the seeds because the amino acids from newly assimilated N are mixed with amino acids from proteolysis for synthesis of new proteins that will be further hydrolysed to support protein neosynthesis. This recycling process would be maintained until amino acid flux to the seeds. Under a nitrogen limitation, proteolysis supporting rapid protein turn over and continuous N-recycling for protein neosynthesis, together with a rapid and efficient nitrogen remobilisation to the seeds after flowering could explain the observed $RSA_{seeds}:RSA_{whole-plant}$ close to 1. However, whereas this equality was independent of environmental conditions in maize (Gallais et al. 2006), this does not appear to be the case in *Arabidopsis* since we observed $RSA_{seeds} > RSA_{whole-plant}$ for HN grown plants (Table 2). This suggests that in the seeds of HN plants, the amount of nitrogen coming from N-remobilisation is higher than the amount of unlabelled nitrogen assimilated after flowering and during grain maturation. This is also consistent with the fact that P%N is lower than P%¹⁵N in HN plant seeds whereas P%N equals P%¹⁵N in LN seeds. The comparison of P%¹⁵N / P%DM between plants is a good way to estimate and compare the NRE when plants do not show the same biomass or sink-source development. Indeed this ratio normalises ¹⁵N flux to seeds taking into account the respective sizes of sink and source compartments. The P%¹⁵N / P%DM ratio shows that NRE in LN plants is 1.7 fold that of HN plants, and it confirms that nitrogen deficiency stimulates plant nitrogen remobilisation (Schulze et al. 1994). Altogether these results shows that whereas NRE is higher in LN plants than in HN plants, the amount of nitrogen coming from N-remobilisation is higher in the HN seeds than in the LN seeds. This surprising feature might be due to HN plants absorbing high quantities of unlabelled nitrogen, before flowering, after flowering and during seed filling, that was stored in leaf rosettes, possibly as nitrate, rather than incorporated into organic compounds. This unlabelled and non-remobilised nitrogen would dilute the rosette ¹⁵N pool thus decreasing the vegetative remains RSA and at the same time increase the ratio $RSA_{seeds}:RSA_{whole-plant}$.

Nitrogen nutrition modifies metabolite concentrations in rosette leaves

Since the nitrogen pools in rosettes and seeds seem to be differentially influenced by N-assimilation and N-remobilisation depending on HN and LN growth conditions, it was hypothesized that metabolite quality and quantity would be modified. As expected, the total nitrogen concentration was higher (2 to 4 times) in the rosettes of HN grown plants (Figure 3A). This difference was mainly due to a higher absorption of nitrate by the HN plants that resulted in the accumulation of nitrate and amino acids, since the ammonium and soluble protein concentrations were similar in rosettes of HN and LN plants (Figure 3B and 2A). The total ^{15}N remaining in the whole plant is related to plant nitrate uptake and transport if we assume that no efflux has occurred. This showed that rosette $^{15}\text{NO}_3^-$ absorption at 40 and 42 DAS was ten times higher under HN ($0.2451 \pm 0.0730 \text{ mg.plant}^{-1}$, $n=4$) than under LN nutrition ($0.0219 \pm 0.0036 \text{ mg.plant}^{-1}$, $n=4$). Nitrate content in HN rosettes was high (50-100 nmol. mg^{-1} FW) and increased with rosette ageing (Figure 3C). On the other hand, nitrate content in LN rosettes was very low (around 5 nmol.mg^{-1} FW) during plant development except at 29 DAS when no difference of biomass could be detected between HN and LN rosettes. However, the nitrate concentration was significantly higher in HN rosettes than in LN rosettes at all time points, 29 DAS included. Whereas nitrate concentration was lower in LN rosettes, the nitrite content appeared slightly higher than in HN rosettes. This difference was significant at 43, 50 and 64 DAS (Figure 3D). Ammonium concentrations were found to be similar in HN and LN rosettes thus suggesting an efficient assimilation into amino acids in both LN and HN plants. However, the total free amino acid concentration was 2-fold higher in HN rosettes (Figure 3E). Sugar accumulation in plant tissues is a well-known consequence of nitrogen deficiency (Tercé-Laforgue et al. 2004). Hexoses and sucrose were almost undetectable in the HN rosettes whereas their concentration increased steadily from 10 and 2 nmol.mg^{-1} FW at 29 DAS, up to 50 and 10 nmol.mg^{-1} FW respectively at 64 DAS in LN rosettes. Starch was detected in both LN and HN plants (Figure 3F). At 29 DAS starch levels were not different in LN and HN rosette, after which starch dramatically increased in LN rosettes, peaking at 50-57 DAS, just before flowering (Figure 3F). On the other hand, HN rosettes did not show an increase in starch levels during the development of the plants. Under low nitrogen nutrition conditions, LN plants become nitrogen-deficient, and the imbalance between carbon fixation and nitrogen assimilation leads to starch accumulation. The carbohydrate accumulation in LN rosettes could contribute to the promotion of “senescence-like” symptoms such as GS1 early induction (Masclaux-Daubresse et al. 2005).

Individual amino acid proportions are modified by nitrogen availability in a similar manner in the seeds and in the rosette leaves.

Individual amino acid contents were determined in rosettes and seeds of LN and HN grown plants using anion exchange chromatography. All of the individual amino acid levels were higher in HN grown rosettes (Table 1) when compared to LN grown rosettes; glutamine, proline and aspartate were especially higher. However, since the total free amino acid concentration was higher in the HN grown rosettes (Figure 3E), individual amino acid concentrations mainly reflected the global increase in total amino acid content. Glutamate occupies a central position in amino acid metabolism since this is the main α -amino group donor for the biosynthesis of glutamine through the GS pathway and for the biosynthesis of other amino acids through the action of a wide variety of amino transferases (Forde and Lea 2007). A good picture of the α -amino transfer network is given by the analysis of the proportion of each amino acid among the total amino acid pool. The relative amount (as % of the total amino acid content) allowed us to determine amino acid pools favoured by plant metabolism when plants have been cultivated under LN or HN conditions (Figure 4). Except at 29 DAS (Figure 4A), the relative amounts of serine, glutamate, alanine, valine and GABA were higher in the rosettes of LN grown plants whereas the relative amounts of threonine, asparagine, glutamine, arginine and proline were higher in the rosettes of plants grown in nitrate rich conditions. These trends and the difference between LN and HN were exacerbated with ageing (Figure 4A-F). Under HN growing conditions, rosettes preferentially accumulated glutamine, asparagine, arginine and proline. Glutamine is the former product obtained from ammonium assimilation. Glutamine is dedicated to transport in the phloem sap and can accumulate to buffer glutamate concentration and detoxify ammonium. Arg molecule contains four nitrogen atoms and acts as a nitrogen reservoir. Proline which is preferentially synthesized in young leaf tissues can also accumulate to high concentration, especially in young leaf tissues where nitrogen assimilation is high. Asparagine build-up was observed in several water-stressed plants and has been proposed as a transient nitrogen storage metabolite in sugar-starved maize roots, in seeds during dehydration and germination (Martinelli et al. 2007 and references therein). In high nitrogen conditions, the increase of asparagine that parallels the increase of glutamine stress the importance of this molecule as a temporary N

storage compound in annuals (Lea et al. 2007).

Since the relative amount of each amino acid was different between LN and HN rosettes, a possible influence on seed composition was investigated. Amino acid levels were 2-fold higher in HN seeds than in LN seeds (30 ± 5 versus 14 ± 2 nmol.mg⁻¹FW; n=4). The total nitrogen concentration and the total protein content were also significantly higher in HN seeds than in LN seeds (4.46 ± 0.17 versus 3.31 ± 0.13 , n=6, [mg nitrogen.100 mg⁻¹FW] and 220 ± 8 versus 171 ± 16 , n=3, [μ g protein. mg⁻¹FW] respectively). However, it can be noted that HN seeds preferentially accumulated free amino acids rather than proteins, compared to LN seeds. Indeed, the concentration of each individual amino acid, except glutamate, was higher in HN seeds compared to LN seeds. Glutamate content was similar in LN and HN seeds, whereas glutamine and asparagine levels were 19 and 6 times higher in HN seeds. Arginine, proline and alanine were 3-fold higher (Figure 5A). The relative individual amino acid amounts were calculated for HN and LN seeds (Figure 5B). HN seeds preferentially accumulated asparagine, glutamine, alanine, arginine and proline, thus showing similarities between the relative amino acid amounts found in HN rosettes. Our results suggest that the individual amino acid proportions in the seeds are directly related to the individual amino acid proportions in the rosette leaves.

Diagnostic of nitrogen deficiency main effects on Arabidopsis thaliana metabolism and physiology

It is known from a long time that nitrogen availability strongly influences plant yield and biomass. However, comparisons of nutrition effects on plant physiology and metabolism have remained fragmentary and poorly documented.

The most striking results showed that limiting nitrogen-nutrition not only influenced plant development, yield and biomass, but also modified nitrogen metabolism and metabolite composition. N-limiting conditions promoted early nitrogen remobilisation and recycling allowing *per se* efficient plant economy. More than 68% of the rosette nitrogen was remobilised to the seeds under LN conditions, versus 36 % under HN conditions. Since at the same time as nitrogen was efficiently remobilised from LN rosettes, N-assimilation was limited by poor nitrogen availability, the total nitrogen remaining in the LN vegetative remains at the end of the plant cycle was very low (six times less than in the HN vegetative

remains). In addition, seed %N ($\text{mg}\cdot\text{mg}^{-1}$ DW) was less affected and only 1.4 fold lower in LN seeds compared to HN seeds. Therefore, the main problematic effect of LN nutrition is the large decrease of yield.

In accordance with a positive effect of nitrogen limitation on N-recycling and remobilisation to the seeds, N-limitation has been seen to control senescence-related enzyme activities by inducing the early synthesis of GS1 and a premature decrease of NR activity (Masclaux et al. 2000). However, other premature senescence symptoms such as leaf yellowing had not been reported; by contrast, leaf redness was increased by LN as a result of anthocyanin accumulation (data not shown). As a result of the changes in N-assimilation:N-remobilisation balance and in N-allocation, it was found that the quality and the quantity of nitrogenous compounds was modified in both seeds and vegetative organs. The HN rosettes accumulated large amounts of nitrate and amino acids but protein concentration was similar to LN rosettes. Accumulation of amino acids in HN rosettes compared to LN rosettes, resulted in a preferential increase in storage molecules such as amides, arginine and proline. Seed composition was also modified with HN seeds accumulating 2-fold more amino acids than LN seeds. However, in contrast to HN rosettes, HN seeds also accumulated proteins. The nature of the amino acids stored into seeds was similar to those stored in the old rosettes (after 50 DAS), and HN seeds preferentially accumulated amides, arginine and proline compared to LN seeds. For both LN and HN plants, the seed amino acid composition paralleled rosette amino acid composition. These observations lead to questions concerning the mechanisms involved in amino acid flux to the seeds (van der Graaff et al. 2006; Masclaux-Daubresse et al. 2008). The fact that amino acid patterns in the phloem sap of several species resemble those in the cytosol of source cells indicate that the phloem loading process would be mainly unspecific (Lohaus and Moellers 2000). This could explain that loading seeds with phloem amino acid results in similar amino acid compositions in both source and sink compartments.

An important feature of this study is that a modification of fertilizer quantity will result in major changes in plant product quantities and yields as well as in their nutritional quality.

Material and Methods

Plant material

Seeds of *Arabidopsis thaliana* accession WS were stratified for 48 h in 0.1 % agar solution (in

water) in the dark at 4°C before sowing. Homogeneous germination occurred 2 days after sowing (DAS). Three times per week, the pots were watered (by immersion of their base) in a solution containing 2 mM nitrate or 10 mM nitrate. Phosphate and sulfate were present at the same concentration (0.25 mM), as well as magnesium (0.25 mM) and sodium (0.20 mM) ions. The watering solution also contained 2.75 mM potassium, 0.5 mM calcium, and 0.7 mM chloride ions. The pH of the watering solutions remained between 5.1 and 5.5. The homogeneous vegetative plant material was grown in a growth chamber under controlled conditions (Loudet et al. 2002). The experimental unit was a small pot (long=60 mm, large=65 mm, high=60 mm) containing six plants positioned in a circle for the nitrate limiting condition (2 mM nitrate) or a single plant per pot for the nitrate pletoric condition (10 mM). The plants were maintained under short days with a photoperiod of 8 h throughout the culture. Day and night temperatures were regulated at 21°C and 17°C, respectively. Light was provided by 20 mercury-vapour bulbs, ensuring a photosynthetic photon flux density of approximately 160 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

Plant growth and development on 10 mM and 2 mM nitrate was similar until 21 days after sowing. The first harvesting time was 29 DAS, after which rosettes were harvested every 7 days. Four distinct batches of rosettes constituted four independent biological repeats. At each harvesting time, four different bulks of rosettes (four biological repeats) were harvested between 10:00 and 11:00 am, and stored at -80°C until use.

Metabolite extraction and analysis, total nitrogen and protein content determination

Amino acids and ammonium were determined after extraction in a solution of 2% 5-sulfosalicylic acid (50 mg FW mL⁻¹). Ammonium was determined according to the Berthelot method (Berthelot 1859). Total amino acid content was assayed by the Rosen colorimetric method using Gln as a reference (Rosen 1957). Amino acid composition was determined in pooled samples extracted from equal dry weights of three individual plant repeats by ion-exchange chromatography using the AminoTac JLC-500/V amino acid analyzer according to the instructions of the manufacturer (JEOL [Europe], Croissy-sur-Seine, France).

Carbohydrates were extracted from aliquots (60 mg FW) of frozen plant material using a three step ethanol-water procedure. The first step consisted of a 30 min extraction at 80°C using 600 μl of 80% ethanol, the second step consisted of a 20 min extraction at 80°C using 300 μl of 50% ethanol and the third step completed the extraction by using 300 μl of water at 80°C

for 20 min. The supernatant and pellets of each extraction step were obtained after centrifugation (15 min, 13000 g, 4°C) and pooled. The sucrose and hexose levels were determined from the ethanolic extract using a commercially available kit (R-Biopharm, Glasgow, Scotland). Starch content was determined from pellets using the same kit. Nitrate was measured according to the method of Cataldo et al. (1975) in the same extract used to measure the nitrate reductase activity. Nitrite was determined from the blank of the nitrate reductase assay (see below).

Total nitrogen was determined using oven-dried plant material and the Dumas combustion method with an NA 1500CN Fisons Instrument (Thermoquest, Runcorn, Cheshire, UK) analyser.

Protein concentration was determined using a commercially available kit (Coomassie Protein assay reagent, BioRad, Hercules, California, USA) and BSA as a standard. Seed protein concentrations were determined by measuring the amino acid levels of NaOH hydrolysates according to Baud et al. (2007).

Enzymatic assays and western blots

Enzymes were extracted from frozen leaf material stored at -80°C . Soluble protein content was determined in crude leaf extracts. Nitrate reductase (NR) maximal extractable activity and activation state were measured as described by Ferrario-Méry et al. (1998). The activation state of NR is defined as the ratio of the activity measured in the presence of 10 mM MgCl_2 divided by the activity in the presence of 5 mM EDTA, and is expressed as a percentage. Glutamine synthetase (GS) was measured according to the method of O'Neal and Joy (1973) (O'Neal and Joy 1973). GS1 and GS2 isoforms were separated by SDS-PAGE according to Masclaux et al. (2000). Antibodies were raised in rabbits against the synthetic peptide AYGEGNERRLTG by Eurogentec (Seraing, Belgium) and they detected both GS1 and GS2 isoenzymes.

^{15}N tracing

Plants were grown in sand under similar conditions as described previously and at 56 DAS, plants were transferred to long day conditions (light/dark cycle 16 h/8 h) to induce flowering. At the two ^{15}N -uptake time points 40 DAS and 42 DAS, the normal watering solution was replaced by a solution that had the same nutrient composition except that $^{14}\text{NO}_3$ was replaced

by $^{15}\text{NO}_3$ to give 10% enrichment. After labelling, plant roots and sand were rinsed with deionised water and unlabelled nutritive solution was used for the rest of the culture cycle. Plants were harvested at the end of their cycle when seeds were mature and the rosette dry. Samples were separated to give (i) “dry vegetative remains” [rosette + stem + cauline leaves + siliques], and (ii) total seeds. Four replicates (plants) were harvested and the experiment was carried out twice.

For all the experiments, unlabelled samples were harvested in order to determine ^{15}N natural abundance. After the drying and weighing of each plant part, material was ground to obtain homogenous fine powder. A subsample of 1000 to 2000 μg was carefully weighed in tin capsules to determine total N content and ^{15}N abundance using an elemental analyzer (roboprep CN, PDZ Europa Scientific Ltd, Crewe, UK) coupled to an isotoper ratio mass spectrometer (Twenty-twenty, PDZ Europa Scientific Ltd, Crewe, UK) calibrated measuring natural abundance. The ^{15}N abundance of samples was calculated as atom percent and defined as $A\% = 100 \cdot (^{15}\text{N}) / (^{15}\text{N} + ^{14}\text{N})$ for labeled plant samples and for unlabelled plant controls (Acontrol% was ca. 0.3660). The ^{15}N enrichment of the sample i was then defined as $(A_i\% - A_{\text{control}}\%)$. The absolute quantity of ^{15}N contained in the sample i was defined as $Q_i = \text{DW}_i \cdot (A_i\% - A_{\text{control}}\%) \cdot \%N_i$ with $\%N_i$ the proportion of nitrogen in the sample i. The Q value of the whole plant (wp) can be measured and $Q_{\text{wp}} = \sum Q_i$. The partition P% of ^{15}N in the organ i was calculated as $(\%P^{15}\text{N in } i) = [(A_i\% - A_{\text{control}}\%) \cdot \text{DW}_i \cdot \%N_i] / [(A_{\text{wp}}\% - A_{\text{control}}\%) \cdot \text{DW}_{\text{wp}} \cdot \%N_{\text{wp}}] = Q_i / Q_{\text{wp}}$. The relative specific allocation of ^{15}N was calculated as $(A_i\% - A_{\text{control}}\%) / (A_{\text{nutritive solution}}\% - A_{\text{control}}\%)$.

Acknowledgements

Authors thanks Dr Michael Hodges (IBP, University of Orsay, France) for manuscript proof reading.

References

- Bänziger, M, Betran, FJ and Lafitte, HR (1997) Efficiency of high nitrogen environment for improving maize for low nitrogen environment. *Crop Science* 37: 1103-1109
- Baud, S, Wuilleme, S, Dubreucq, B, de Almeida, A, Vuagnat, C, Lepiniec, L, Miquel, M and Rochat, C (2007) Function of plastidial pyruvate kinases in seeds of *Arabidopsis thaliana*. *The Plant Journal* 52: 405-419
- Beninati, NF and Busch, RH (1992) Grain protein inheritance and nitrogen uptake and redistribution in a spring wheat cross. *Crop Science* 32: 1471-1475
- Berthelot, MPE (1859) Violet d'aniline. Report, *chimie appliquée* 1: 284

- Brouwer, R (1962) Nutrient influences on the distribution of the dry matter in the plants. *Netherlands Journal of Agricultural Sciences* 10: 399-408
- Brugière, N, Dubois, F, Masclaux, C, Sangwan, RS, and Hirel, B (2000) Immunolocalization of glutamine synthetase in senescing tobacco (*Nicotiana tabacum* L.) leaves suggests that ammonia assimilation is progressively shifted to the mesophyll cytosol. *Planta* 211: 519-527
- Cataldo, DA, Haroon, M, Schrader, LE and Young, VL (1975) Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commun. Soil Science and Plant Analysis* 6: 71-80
- Deléens, E, Cliquet, JB and Prioul, J-L (1994) Use of ¹³C and ¹⁵N plant label near natural abundance for monitoring carbon and nitrogen partitioning. *Australian Journal of Plant Physiology* 21: 133-146
- Diaz, C, Lemaître, T, Christ, A, Azzopardi, M, Kato, Y, Sato, F, Morot-Gaudry, JF, Le Dily, F and Masclaux-Daubresse, C (2008) Nitrogen recycling and remobilisation are differentially controlled by leaf senescence and development stage in *Arabidopsis thaliana* under low nitrogen nutrition doi: 10.1104/pp.108.119040 (in press)
- Drew, MC and Saker, LR (1976) Nutrient supply and the growth of the seminal root system in barley. II. Localised compensatory changes in lateral root growth and the rate of nitrate uptake when nitrate is restricted to only one part of the root system. *Journal of Experimental Botany* 26: 79-90
- Forde, BG and Lea, PJ (2007) Glutamate in plants: metabolism, regulation, and signalling. *Journal of Experimental Botany* 58: 2339-2358
- Fritz, C, Mueller, C, Matt, P, Feil, R and Stitt, M (2006) Impact of the C-N status on the amino acid profile in tobacco source leaves. *Plant, Cell and Environment* 29: 2055-2076
- Gallais, A, Coque, M, Quillere, I, Prioul, JL and Hirel, B (2006) Modelling postsilking nitrogen fluxes in maize (*Zea mays*) using N-15-labelling field experiments. *New Phytologist* 172: 696-707
- Good, AG, Shrawat, AK and Muench, DG (2004) Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends in Plant Science* 9: 597-605
- Hirel, B, Le Gouis, J, Ney, B and Gallais, A (2007) The challenge of improving nitrogen use efficiency in crop plants: towards a more central role for genetic variability and quantitative genetics within integrated approaches. *Journal of Experimental Botany* 58: 2369-2387
- Lea, PJ, Azevedo, RA (2006) Nitrogen use efficiency 1. Uptake of nitrogen from the soil. *Annals of Applied Biology* 149: 243-247
- Lea, PJ, Azevedo, RA (2007) Nitrogen use efficiency 2. Amino acid metabolism. *Annals of Applied Biology* 151: 269-275
- Lea, PJ, Sodek, L, Parry, MAJ, Shewry, PR and Halford, NG (2007) Asparagine in plants. *Annals of Applied Biology* 150: 1-26
- Lohaus, G, Moellers, C (2000) Phloem transport of amino acids in two *Brassica napus* L. genotypes and one *B. carinata* genotype in relation to their seed protein content. *Planta* 211: 833-840
- Loudet, O, Chaillou, S, Camilleri, C, Bouchez, D and Daniel-Vedele, F (2002) Bay-0 x Shahdara recombinant inbred line population: a powerful tool for the genetic dissection of complex traits in *Arabidopsis*. *Theoretical and Applied Genetics* 104: 1173-1184
- Martin, A, Belastegui-Macadam, X, Quillere, I, Floriot, M, Valadier, MH, Pommel, B, Andrieu, B, Donnison, I and Hirel, B (2005) Nitrogen management and senescence in

- two maize hybrids differing in the persistence of leaf greenness: agronomic, physiological and molecular aspects. *New Phytologist* 167: 483-492
- Masclaux, C, Valadier, M, Brugière, N, Morot-Gaudry, J and Hirel, B (2000) Characterization of the sink/source transition in tobacco (*Nicotiana tabacum* L.) shoots in relation to nitrogen management and leaf senescence. *Planta* 211: 510-518
- Masclaux-Daubresse, C, Carrayol, E and Valadier, M-H (2005) The two nitrogen mobilisation- and senescence-associated GS1 and GDH genes are controlled by C and N metabolites. *Planta* 221: 58-588
- Masclaux-Daubresse, C, Reisdorf-Cren, M, and Orsel, M (2008) Leaf nitrogen remobilization for plant development and grain filling. *Plant Biology* (in press)
- Martinelli, T, Whittaker, A, Bochicchio, A, Vazzana, C, Suzuki, A, Masclaux-Daubresse, C (2007) Amino acid pattern and glutamate metabolism during dehydration stress in the “resurrection” plant *Sporobolus stapfianus*. A comparison between desiccation-sensitive and desiccation-tolerant leaves. *Journal of Experimental Botany* 58: 3037-3046
- O'Neal, D and Joy, KD (1973) Glutamine synthetase of pea leaves. I. Purification, stabilisation and pH optima. *Archive of Biochemistry and Biophysics* 159: 113-122
- Presterl, T, Seitz, G, Landbeck, M, Thiemt, W, Schmidt, W and Geiger, HH (2003) Improving nitrogen use efficiency in European maize: estimation of quantitative parameters. *Crop Science* 43: 1259-1265
- Rosen, H (1957) A modified ninhydrin colorimetric analysis for amino acids. *Archive of Biochemistry and Biophysics* 67: 10-15
- Scheible, WR, Morcuende, R, Czechowski, T, Fritz, C, Osuna, D, Palacios-Rojas, N, Schindelasch, D, Thimm, O, Udvardi, MK and Stitt, M (2004) Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of *Arabidopsis* in response to nitrogen. *Plant Physiology* 136: 2483-2499
- Schulze, W, Schulze, ED, Stadler, J, Heilmeyer, H, Stitt, M and Mooney, HA (1994) Growth and reproduction of *Arabidopsis thaliana* in relation to storage of starch and nitrate in the wild-type and in starch-deficient and nitrate-uptake-deficient mutants. *Plant, Cell and Environment* 17: 795-809
- Singh, U (2005) Integrated nitrogen fertilization for intensive and sustainable agriculture. In Basra, A, Goyal, S and Tishner, R, eds, *Enhancing the efficiency of nitrogen utilisation in plants*. *Journal of Crop Improvement*, Vol 15. Food Product Press, The Haworth press Inc., pp 213-257
- Tercé-Laforgue, T, Mäck, G and Hirel, B (2004) New insights towards the function of glutamate dehydrogenase revealed during source-sink transition of tobacco (*Nicotiana tabacum*) plants grown under different nitrogen regimes. *Physiologia Plantarum* 120: 220-228
- van der Graaff, E, Schwacke, R, Schneider, A, Desimone, M, Flugge, UI and Kunze, R (2006) Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence. *Plant Physiology* 141: 776-792
- Van der Werf, A and Nagel, OW (1996) Carbon allocation to shoots and roots in relation to nitrogen supply is mediated by cytokinins and sucrose. *Opinion in Plant Soil* 185: 21-32
- Vincentz, M, Moureaux, T, Leydecker, M-T, Vaucheret, H and Caboche, M (1993) Regulation of nitrate and nitrite reductase expression in *Nicotiana plumbaginifolia* leaves by nitrogen and carbon metabolites. *The Plant Journal* 3: 315-324
- Wang, R, Guegler, K, LeBrie, ST and Crawford, NM (2000) Genomic analysis of a nutrient response in *Arabidopsis* reveals diverse expression patterns and novel metabolic and potential regulatory genes induced by nitrate. *The Plant Cell* 12: 1491-1509

Wingler, A, Marès, M and Pourtau, N (2004) Spatial patterns and metabolic regulation of photosynthetic parameters during leaf senescence. *New Phytologist* 161: 781-78

Tables

Table 1: Individual amino acid composition in the rosettes of plants grown on 10 mM and 2 mM nitrate during 29, 36, 50, 57 and 64 DAS. Free amino acids were separated by chromatography and determined as nmol.(mg FW)⁻¹. Results were obtained after pooling the amino acid samples from four plants on the basis of equal FW used in the extraction.

nmol . (mg FW) ⁻¹	29 DAS		36 DAS		50 DAS		57 DAS		64 DAS	
	10 mM	2 mM								
ASP	1.672	1.133	1.210	0.403	2.177	1.039	1.830	0.969	1.385	1.010
THR	0.376	0.249	0.232	0.107	0.591	0.248	0.831	0.216	0.778	0.194
SER	0.584	0.510	0.406	0.353	0.906	0.873	1.148	0.867	1.260	0.757
ASN	0.596	0.336	0.559	0.087	0.930	0.081	0.772	0.093	1.048	0.102
GLU	3.270	2.188	2.071	0.732	3.167	2.539	3.130	2.413	2.855	2.509
GLN	1.919	1.108	1.631	0.328	2.978	0.483	2.348	0.685	2.115	0.908
GLY	0.371	0.183	0.300	0.027	0.176	0.080	0.134	0.131	0.179	0.116
ALA	1.167	0.919	0.610	0.211	0.814	0.390	0.832	0.529	0.556	0.509
VAL	0.180	0.143	0.139	0.099	0.253	0.328	0.256	0.378	0.254	0.370
ARG	0.026	0.026	0.026	0.027	0.118	0.038	0.203	0.028	0.525	0.025
PRO	0.393	0.169	0.328	0.112	1.342	0.211	1.588	0.198	1.441	0.189

Table 2 : Comparison of RSA and P% of samples from plants grown on 10 mM and 2 mM nitrate. Partitioning P% is the ratio between the total-seed compartment and the whole plant. P%¹⁵N (g ¹⁵N in seeds . g⁻¹ ¹⁵N in the whole plant), P%N (g N in seeds . g⁻¹ N in the whole plant) and P%DM (g dry matter of seeds . g⁻¹ dry matter of the whole plant). P% and RSA are the mean of four plants and standard errors are presented. The experiment was repeated twice giving similar results and differences between 10 mM and 2 mM. A single experiment is presented.

	LN (2 mM nitrate)	HN (10 mM nitrate)
P%DM	30.1% ± 2.2%	25.4% ± 1.9%
P%N	65.5% ± 3.2%	25.3% ± 3.0%
P% ¹⁵ N	68.7% ± 1.8%	35.8% ± 7.8%
P% ¹⁵ N / P%DM	2.29 ± 0.17	1.40 ± 0.22
RSA _{seeds}	0.90 ± 0.10	0.83 ± 0.07
RSA _{seeds} / RSA _{wholeplant}	1.03 ± 0.03	1.19 ± 0.09
N% seeds (mg. (100 mg DW) ⁻¹)	3.19 ± 0.06	4.34 ± 0.11
N% vegetative remains (mg. (100 mg DW) ⁻¹)	0.72 ± 0.04	4.39 ± 0.23

Legends

Figure 1: Comparison of biomass (A ; B ; C) and seed production (D) between plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Standard errors and means of 4 individual plants are presented. # indicates a non significant difference between white and black bars with $P \geq 0.05$.

Figure 2: Comparison of protein concentration (A), nitrate reductase activity (NR ; B), nitrate reductase activation state (C), glutamine synthetase activity (GS ; D) and GS protein content (E) in rosettes of plants grown in low nitrate (black bars) or in high nitrate (white bars) conditions. Standard error and mean of 4 individual plants are presented. # indicates a non significant difference between white and black bars with $P \geq 0.05$. For protein separation before western blot, equal amount of soluble protein was loaded to each lane (10 μg).

Figure 3: Comparison of total nitrogen content ($\text{mg} \cdot (100 \text{ mg DW})^{-1}$; A), ammonium concentration ($\text{nmol} \cdot (\text{mg FW})^{-1}$; B), nitrate ($\text{nmol} \cdot (\text{mg FW})^{-1}$; C), nitrite ($\text{nmol} \cdot (\text{mg FW})^{-1}$; D), free amino acids ($\text{nmol} \cdot (\text{mg FW})^{-1}$; E) and starch ($\text{nmol} \cdot (\text{mg FW})^{-1}$; F) in the rosettes of plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Standard errors and means of 4 individual plants are presented. # indicates a non significant difference between white and black bars with $P \geq 0.05$.

Figure 4: Comparison of individual amino acid proportions (in % of total amino acid) between extracts of rosettes grown under nitrate limiting (black bars) or plethoric (white bars) conditions for 29, 36, 43, 50, 57 and 64 days. Individual amino acids were quantified using ion-exchange chromatography using pooled samples extracted from 4 plants and mixed on the basis of equal total free amino acid content.

Figure 5: Comparison of individual amino acid concentrations (A, $\text{nmol} \cdot \text{mg}^{-1} \text{ DW}$) and proportions (B, in % of total amino acid) between seeds of plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Individual amino acids were quantified using ion-exchange chromatography using pooled samples extracted from seeds of 4 plants and mixed on the basis of equal total free amino acid content.




Figure 1: Comparison of biomass (A ; B ; C) and seed production (D) between plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Standard errors and means of 4 individual plants are presented. # indicates a non significant difference between white and black bars with $P \geq 0.05$.




Figure 2: Comparison of protein concentration (A), nitrate reductase activity (NR ; B), nitrate reductase activation state (C), glutamine synthetase activity (GS ; D) and GS protein content (E) in rosettes of plants grown in low nitrate (black bars) or in high nitrate (white bars) conditions. Standard error and mean of 4 individual plants are presented. # indicates a non significant difference between white and black bars with $P \geq 0.05$. For protein separation before western blot, equal amount of soluble protein was loaded to each lane ($10 \mu\text{g}$).




Figure 3: Comparison of total nitrogen content (mg.(100 mg DW)⁻¹ ; A), ammonium concentration (nmol.(mg FW)⁻¹; B), nitrate (nmol.(mg FW)⁻¹; C), nitrite (nmol.(mg FW)⁻¹; D), free amino acids (nmol.(mg FW)⁻¹; E) and starch (nmol.(mg FW)⁻¹; F) in the rosettes of plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Standard errors and means of 4 individual plants are presented. # indicates a non significant difference between white and black bars with P ≥ 0.05.




Figure 4: Comparison of individual amino acid proportions (in % of total amino acid) between extracts of rosettes grown under nitrate limiting (black bars) or plethoric (white bars) conditions for 29, 36, 43, 50, 57 and 64 days. Individual amino acids were quantified using ion-exchange chromatography using pooled samples extracted from 4 plants and mixed on the basis of equal total free amino acid content.




Figure 5: Comparison of individual amino acid concentrations (A, nmol. mg⁻¹ DW) and proportions (B, in % of total amino acid) between seeds of plants grown under nitrate-limiting (black bars) or plethoric (white bars) conditions. Individual amino acids were quantified using ion-exchange chromatography using pooled samples extracted from seeds of 4 plants and mixed on the basis of equal total free amino acid content.