

### The ornamentation steps of the Bull Rotunda of the Lascaux cave give new insights into the Upper Palaeolithic natural life cycle

Ina Reiche, Colette Vignaud, Emilie Chalmin, Michel Menu, Jean-Michel

Geneste

#### ▶ To cite this version:

Ina Reiche, Colette Vignaud, Emilie Chalmin, Michel Menu, Jean-Michel Geneste. The ornamentation steps of the Bull Rotunda of the Lascaux cave give new insights into the Upper Palaeolithic natural life cycle. Archaeometry, inPress, 10.1111/arcm.12960. hal-04498458

### HAL Id: hal-04498458 https://hal.science/hal-04498458

Submitted on 11 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Archaeometry

# archaeo**metry**

#### The ornamentation steps of the Bull Rotunda of the Lascaux cave give new insights into the Upper Palaeolithic natural life cycle

| Journal:  | Archaeometry  |
|---|---|
| Manuscript ID   | ARCH-09-0737.R1   |
| Wiley - Manuscript type:  | Original Article  |
| Date Submitted by the<br>Author:                                    | n/a   |
| Complete List of Authors:   | Reiche, Ina; Chimie ParisTech, Insitut de Recherche de Chimie-Paris<br>(IRCP); CNRS, FR 3506 CNRS - Centre de recherche et de restauration<br>des musées de France<br>Vignaud, Colette; Centre de Recherche et de Restauration des Musées de<br>France,<br>Chalmin , Emilie; Université Savoie Mont-Blanc, EDYTEM<br>Menu, Michel; Centre de Recherche et de Restauration des musées de<br>France; The Cyprus Institute, APAC Labs<br>Geneste, Jean-Michel; PACEA   |
| Keywords:   | Lascaux cave, creation steps, manganese and iron oxides, SEM-EDX, TEM-EDX-SAED, Palaeolithic paintings, natural life cycle  |
| Abstract:   | Although the ornamentation of the Lascaux cave seems relatively<br>homogeneous in style, the analysis by scanning and transmission<br>electron microscopy of 32 micro-samples from the figures of the Hall of<br>the Bulls (Bull Rotunda) and one desquamated sample from the soil<br>highlighted different paint pots. The black and red paint matters with<br>their associated mineralogical phases were extensively characterized.<br>Considering previous stylistic and superimposition studies, we were<br>eventually able to confirm five creation steps of monothematic figures<br>("assemblages") based on the chemical and mineralogical characteristics<br>Further interpretations in terms of the rhythm and temporal framework<br>of the Hall of the Bulls (Bull Rotunda) of the Lascaux cave could be<br>reinforced. Some particular representations such as the black cave bear<br>and one of the cross shaped signs were shown to be likely added at later<br>moments by different hands because their paint pots were different from<br>those of the Bull Rotunda and similar to that of other representations in<br>deeper rooms of the Lascaux cave. We also found that the paint matter<br>of the Bull Rotunda was different than those of other Palaeolithic cave<br>sites on a more regional scale indicating a very local sourcing. |
|   |   |
| lote: The following files were s<br>You must view these files (e.g. | ubmitted by the author for peer review, but cannot be converted to PDF. movies) online.   |
| igure7_R.svg  |   |
|   |   |

| 1        |                         |
|----------|-------------------------|
| 2        |                         |
| 3        |                         |
| 4        |                         |
| 5        | SCHOLARONE <sup>™</sup> |
| 6        | Manuscripts             |
| 7        | Manascripts             |
| 8        |                         |
| 9        |                         |
| 10       |                         |
| 11       |                         |
| 12       |                         |
| 13       |                         |
| 14       |                         |
| 15       |                         |
| 16       |                         |
| 17       |                         |
| 18       |                         |
| 19<br>20 |                         |
| 20       |                         |
| 22       |                         |
| 22 23    |                         |
| 24       |                         |
| 25       |                         |
| 26       |                         |
| 27       |                         |
| 28       |                         |
| 29       |                         |
| 30       |                         |
| 31       |                         |
| 32       |                         |
| 33       |                         |
| 34       |                         |
| 35       |                         |
| 36       |                         |
| 37       |                         |
| 38       |                         |
| 39       |                         |
| 40       |                         |
| 41       |                         |
| 42       |                         |
| 43<br>44 |                         |
| 44<br>45 |                         |
| 45       |                         |
| 40       |                         |
| 47 48    |                         |
| 49       |                         |
| 50       |                         |
| 51       |                         |
| 52       |                         |
| 53       |                         |
| 54       |                         |
| 55       |                         |
| 56       |                         |
| 57       |                         |
| 58       |                         |
| 59       |                         |
| 60       |                         |

### The ornamentation steps of the Bull Rotunda of the Lascaux cave give new insights into the Upper Palaeolithic natural life cycle

Ina Reiche<sup>1,2\*</sup>, Colette Vignaud<sup>3</sup>, Emilie Chalmin<sup>4</sup>, Michel Menu<sup>5</sup>, Jean-Michel Geneste<sup>6</sup>

1 Fédération de recherche New AGLAE, FR3506 CNRS/Ministère de la Culture, Palais du Louvre, Paris, France.

2 PSL Université, Chimie Paristech, Institut de recherche de Chimie Paris, UMR 8247 CNRS, PCMTH, Paris, France

3 former address: Laboratoire du Centre de recherche et de restauration des musées de France, C2RMF-UMR171 CNRS, Palais du Louvre, Paris, France, present address: Colette.vignaud@noos.fr

4 EDYTEM, UMR CNRS 5204, Université Savoie Mont-Blanc, Le Bourget-du-Lac, France

5 former address: Centre de recherche et de restauration des musées de France, C2RMF, Palais du Louvre, Paris, France, present address: APAC Labs, The Cyprus Institute, Nicosia, Cyprus

6 PACEA UMR 5199 CNRS/Ministère de la culture & Université de Bordeaux, Bordeaux, France

\* corresponding author: ina.reiche@chimieparistech.psl.eu

#### Abstract

Although the ornamentation of the Lascaux cave seems relatively homogeneous in style, the analysis by scanning and transmission electron microscopy of 32 micro-samples from the figures of the *Hall of the Bulls (Bull Rotunda)* and one desquamated sample from the soil highlighted different paint pots. The black and red paint matters with their associated mineralogical phases were extensively characterized. Considering previous stylistic and superimposition studies, we were eventually able to confirm five creation steps of monothematic figures ("assemblages") based on the chemical and mineralogical characteristics. Further interpretations in terms of the rhythm and temporal framework of the *Hall of the Bulls (Bull Rotunda)* of the Lascaux cave could be reinforced. Some particular representations such as the black cave bear and one of the cross shaped signs were shown to be likely added at later moments by different hands because their paint pots were different from those of the Bull Rotunda and similar to that of other representations in deeper rooms of the Lascaux cave. We also found that the paint matter of the Bull Rotunda was different than those of other Palaeolithic cave sites on a more regional scale indicating a very local sourcing.

#### Introduction. Previous work on the Lascaux cave and paint samples

Electron microscopy can bring to light invisible information about rock art. The Lascaux cave is one of the most famous decorated caves in the world dating back to the Magdalenian period. Its rock art has been studied in detail by N. Aujoulat (Aujoulat, 2004). Lascaux's paintings and graphical entities seem to have been created in relatively short period of time in all parts of the cave (Leroi-Gourhan and Allain, 1979), likely a few centuries at the height of the temperate interstadial of Lascaux. There are many instances

 of enrichments, repainting and retouching of the parietal decorations, but there is an obvious unity of style in Lascaux. Recent radiocarbon dates of five reindeer scraps from the main sectors of the cave (Axial gallery, *Passage*, *Nef*, *Puits*) place the occupation of the cave around 17,600 uncal. BP (meaning 21,500-21,000 cal. BP) (Ducasse and Langlais, 2019) but there is still no date of paintings.

The results obtained from the physicochemical analysis of 17 micro-samples (GLAS26, GLAS45-46, GLAS50-55, GLAS62, GLAS101-107) of isolated scenes at different locations of the Lascaux cave have already been reported elsewhere following a strategy of panels or coherent scenes such as the scène du Puits (Aujoulat et al., 2002), blasons (Chalmin et al., 2004) and the bisons adossés (Vignaud et al., 2006) as well as two isolated figures, horse and deer (Chalmin, 2003). The different scenes were related in stylistic accordance, method of application and paint matter. It was brought to light that paint matter was deliberately prepared as a function of a specific use. As an example, Lascaux' artists used different paint matters for the *blasons*' rectangles, each with a singular shade (Chalmin et al., 2004). Beyond these preparation specific features, it turned out that there was a large palette of manganese (Mn) oxide minerals and various qualities of haematite that allowed painters to obtain subtle colour hues (Aujoulat et al., 2002; Chalmin et al., 2004; Vignaud et al., 2006). Haematite is less discriminative than the black Mn oxide based phases in distinguishing prehistoric figures. Scanning Electron Microscopy (SEM) observations on microscale only allow the identification of Mn oxide agglomerates. For the discrimination of different mineralogical phases, Transmission Electron Microscopy (TEM) coupled with an Energy Dispersive Analysing system (EDS) and Selected Area Electron Diffraction (SAED) is required. These new criteria allow new insights into the different paint recipes of the prehistoric artists of Lascaux.

Despite the homogeneity of the style in Lascaux, previous results indicated that different paint matters and preparations allow the identification of groupings ("assemblages") that contribute to a better understanding of the various execution sequences of representations. Furthermore, technological information on raw materials, preparation, applications and also possibly economic activities (choice of raw materials, diversity, specific selection) can be obtained.

The *Bull Rotunda* is an enigmatic chamber and its decoration seems to be among the first one to be executed in the Lascaux cave. It is 18 m long, 7 m large and 4-5 m high with a surface of 117 m<sup>2</sup> (Aujoulat, 2004) (Fig. 1.) The quality of the wall is not suitable for engraving but favourable for painting because of a homogeneous very indurated white carbonate concretion with relatively large grains (from 5 to 20 mm in diameter) and high reflective power that covers the entire substrate rock in the *Rotunda*. There is a change in the colour of the substrate that materializes a baseline used as an imaginary soil for most of the figures in the room. Only a few gaps in the carbonate concretion, scale imprints and corroded beaches disturb the homogeneity of this covering and there is the presence of macro-cupules in the lower part of the panel. Among the animals, the *horses* are the most numerous (N=17) then *aurochs* (bulls) and *cows* (N=11) and finally *deer* (N=5). There is also a single *bison* (N=1), a *unicorn* (N=1) and a *cave bear* (N=1) (35) (Fig. 2 and 3 (Aujoulat, 2004)). The animal representation of the *unicorn* (mysterious animal actually represented with two horns) is unique by the way. There are also other representations of rare animal species such as a *cave bear*. The *bear* is superimposed on the initially

executed large ventral line of the first bull on the right side in the *Rotunda* (Fig. 3, 1D101). It was discovered thanks to its muzzle and its ears, which went over the ventral line of the *black bull*.

Animals are either monochrome (black or red) or bichrome with different colour associations (black and red, brown or yellow). Some figures present different colour gradients such as the *brown horse* 1G204 with colours ranging from light brown to black and the *deer* 1G214 with colours from yellow to black. Basically, large *black bulls* and *red cows* are found in the Rotunda: 3.5 m (L) X 3.0 m (H) for bull 1G203 (Fig. 2), 4.6 m (L) X 2.8 m (H) for bull 1G205 (Fig. 2) and 5.6 m (L) X ca. 3 mm (H) for bull 1D101 (Fig. 3). They are the largest representations found in all European Palaeolithic art. However, the size contrasts strongly between certain species of animals. *Horses* are bichrome and smaller in size (Aujoulat, 2004).

Different methods of paint application were identified: painting with a brush, a stamp, spraying and/or stencil drawing. While their study and interpretation are not in the focus of the present work, we may mention, as an example, the black dorsal lines of the *bulls* were executed using a series of very close punctuations avoiding any pointillism effect. The regularity of point diameters indicates a similar or even the same tool was used. A consistent application technique was also used for the *red cow* representations. In the lower part of the figures, spraying was used, whereas in the upper part of the figures, paint was applied with a brush, due to the (in)accessibility of the wall. More details are reported elsewhere (Aujoulat, 2004).

A. Laming-Emperaire suggested in 1959 that the identification of groups in different parts of the cave ("assemblages") corresponded to a conscious order in the distribution of animals, as well as a combination of size, species and direction of animals (Laming-Emperaire, 1959). N. Aujoulat also set the sequence of execution steps in the *Bull Rotunda* in the 1980-90's, each composed of different monothematic animal representations. According to stylistic and macro-photographical observations of the walls, he proposed five steps (Aujoulat, 2004): step 1 corresponded to a series of monochromatic *horses*, step 2 to a series of *bichrome and polychrome horses* including the *unicorn*, step 3 to a series of *black bulls*, step 4 to a series of *red cows* and step 5 finally to a series of *deer* (step 5). J.-M. Geneste further interpreted this sequence in 2021 on the basis of stylistic and superimposition studies in terms of the natural life cycle of the Upper Palaeolithic hunter societies, and proposed the different steps of monothematic animal depictions to be related to the natural rhythm of the animal herds in the prehistoric landscape, e.g. seasons (Geneste, 2021).

#### Aim of this work

In the present work we discuss the consistency of the hypothesized execution steps in terms of different paint pots, especially the colouring phases, used for the different animal series in the *Bull Rotunda*. Electron microscopic analyses and observations of micro-samples taken from different paintings in the emblematic *Bull Rotunda* of the Lascaux cave allow the characterisation of these paint pots. We can infer from different paint pots different execution steps under the following conditions: first, different paint matters can be identified in the executed prehistoric graphical entities; second, the paint matter is characteristic at the scale of the micro-samples for each figure, and can thus be

considered representative enough of the figures on the wall; and third, it can be postulated that one single hand or a very small group of artists created in the cave a series of stylistically similar figures, and completed them in a short period of time (nearly simultaneously), when the same or a similar type of components is found in the micro-samples corresponding to a particular series of prehistoric figures.

Based on this, it was assumed that series of figures could be defined according to their chemical and mineralogical characteristics as observed by electron microscopy. Additionally, we suggested that it was possible to attribute, from a stylistically, chemically and mineralogically homogeneous group of prehistoric figures of the same colour, to a consistent step of execution. However, it is not possible to determine the duration or the number of artists involved in a particular execution step (although sometimes individual hands of prehistoric artists could be stylistically identified (Fritz and Tosello, 2015)) nor the time lapse between different execution steps.

By identifying different steps of execution in the *Bull Rotunda* we intented to reinforce the interpretation of the ornamentation steps as related to the temporal natural life cycle of the Upper Palaeolithic hunter-gatherer society, e.g. seasons.

#### Studied material and methods

This study summarized the results of the analyses of 33 micro-samples taken from different figures in the Bull Rotunda of the Lascaux cave, including the desquamated sample collected from the soil. Three field trips were authorized to take micro-samples: 25 micro-samples were taken in 1996 (GLAS1-25), 3 in 1999 (GLAS61-63) and 4 in 2005 (GLAS117-120) as well as one desquamated sample from the soil (LAS121). The two field trips in 1996 and 1999 took place before the closing of the cave, while the one in 2005 occurred after. Micro-samples of black, red and yellow graphical entities were taken using a sterilized scalpel. On the left wall, we counted 21 samples from ten animal figures (sample GLAS 1-2: black unicorn 1G201, GLAS3-6: bichrome horse G202, GLAS7-8: bichrome horns of the bull 1G203, GLAS10: brown horse 1G204, GLAS11: red deer 1G212, GLAS12-13: bichrome deer 1G214 and a black horse, numbered here 1G206bis, GLAS15: red cow 1G206, GLAS16: small black horse, GLAS17-18: black bull 1G205) and two signs (GLAS9: 1G223 (could also correspond to the muzzle of a hidden animal) and GLAS 14:1G226). On the right, twelve samples from five animal figures (GLAS 22-23, 118: black bull 1D101, GLAS20: black bull 1D102, GLAS24: red cow 1D103, GLAS26: red calf 1D108, GLAS25,117: black bear 1D107 and two signs (GLAS 19: cross 1D112, GLAS21: 1D114) were studied, as well as a desquamated sample (LAS121) laying on the floor (Fig. 1, Table 1) and GLAS119 corresponding to the red spot on the bench seat.

We used optical microscopy (OM), SEM-EDS, TEM-EDS and TEM-SAED to characterize the large set of paint micro-samples on the micro- and the nanoscale. The sample LAS121 was prepared as a cross-section inserted in polyester resin and then polished. The other samples were studied without resin inclusion by OM and SEM. Only when the paint matter was concentrated enough in the micro-sample was a sample prepared for TEM observations, which requires invasive preparation.

#### Optical microscopy

The paint cross-section and the micro-samples were examined under a light microscope (NIKON Eclipse LV 100ND) and photographed without any preparation.

#### SEM-EDS

The micro-samplings were deposited without any preliminary preparation directly into a metallic cup to avoid charge phenomena, nickel was chosen in order to avoid interfering with detected elements in the EDS spectrum .

The cross-section is carbon-coated to enable charge evacuation.

For this study, a Jeol JSM-840 SEM and a Philips XL 30 CP SEM were used, both equipped with an EDX Link Isis 300 analysis system. An acceleration voltage of 20 kV of the electrons with a beam current of about 100  $\mu$ A was generally used.

#### TEM- EDS and TEM-SAED

Each selected sample was ground in an agate mortar. The obtained powder was dispersed ultrasonically in ethanol and then a drop of this preparation was deposited on a carbon-coated copper (Cu) or gold (Au) grid. The TEM observations were carried out at 200 kV on the sample grid using a JEOL 2000FX microscope equipped with a Link Isis EDS. Using a double tilted object holder, electron diffraction (SAED) is carried out and the crystallographic structure of the poly-crystals was then recorded. If individual crystals were large enough, a single-crystal diffraction pattern could be recorded on each one. Knowing its chemical composition, the single-crystal identity and its zone axis is identified using the Carine software®.

#### Results

#### Microscale morphological and chemical characteristics

Optical microscope observations of the micro-samples enabled us to evaluate the thickness, the colour and the texture of the pigment. We were also able to extract the paint matter from the wall support. The thickness of the micro-samples was approx. between 50 and 500  $\mu$ m. This thickness allowed good access to the distribution of each component in the paint matter and avoided disturbance from potential components coming from the limestone support or various posterior encrusts (Fig. 4a-d).

The cross section LAS121 corresponded to a red paint sandwiched between a thin outer layer of transparent calcium carbonate and a thick layer of calcium (Ca) carbonate from the wall support. The paint sample was actually a peeling off of the wall well below the paint. The paint layer in the cross section contained a thin layer of about 5 to 10  $\mu$ m of inhomogeneous clay rich in iron. Punctually, iron (Fe) oxides were visible in the form of platelets of 1 to 2  $\mu$ m in the electron micrograph (Fig. 4b and 5a).

SEM observations of the micro-samples showed micro-morphological characteristics in addition to chemical analysis (EDS) (Table 1). All black samples were agglomerates composed of different types of manganese (Mn) oxides, some containing additionally aluminosilicates material. Chemically, three types of Mn oxides are found: pure Mn oxide, Barium (Ba)-containing Mn oxide and Mn oxide with variable amounts of Ba and potassium (K). Different morphologies can be identified, but not different crystallographic structures. For example, the *cave bear* sample GLAS117 showed three morphologies of

Mn oxides: simple Mn oxide or hydroxide (fig. 5c) and at higher magnification fines needles of a Ba-bearing Mn oxide as well as cubic Ca carbonate (Fig 5d).

Aluminosilicates were present in most of the black paints; quartz was more rare. The presence of a phase combining calcium (Ca) and phosphorus (P) could be noticed in only two samples (black punctuations GLAS21 and *cave bear* GLAS25). In black signs mercury (Hg) was also as small droplets. This was likely contemporary pollution, as the temperature of the cave was controlled using Hg-thermometers, that were frequently broken inside the cave. Morphologically, platelet-like crystals, sheets, sticks, needle-shaped, filaments, balls and spherical grains as well as "desert roses" and meshes could be described in these Mn oxide blocs (Table 1). The morphology corresponding to the crystallization modus did not, however, give any information about the types of crystallographic phase present in the Mn oxides.

Red paints were composed of iron (Fe) oxides of different crystal shapes. Fibrous crystals are found as well as needles, platelet-like crystals and spherical grains. Chemically, other natural phases such as quartz, aluminosilicates and calcium carbonate were observed besides Fe oxide. Particularly, one Ca and P-based phase (probably hydroxyapatite) was also identified in most samples. Further minor elements like K and titanium (Ti) were detected but cannot be attributed to a specific compound.

The yellow paint sample was composed of a mixture of iron oxide, a high amount of calcium carbonate possibly from the wall support or from recrystallization process, some aluminosilicates and a Ca-P based compound, likely hydroxyapatite.

Brown paints were composed of mixtures of Fe and Mn oxides in addition to aluminosilicates, calcium carbonate, quartz and calcium sulphate (likely gypsum). The latter phase possibly originated from weathering processes. Interestingly, the presence of a Ca and P-based phase was observed for the red flat paints (filling). Manganese oxide crystal agglomerates were needle- or fibre-shaped and Fe oxides were sheet-like.

Further discrimination of the crystallographic structures present in the paint samples is not possible. Therefore, nanoscale observations using TEM-EDS and TEM-SAED were required.

#### Nanoscale morphological and crystallographic characteristics

TEM observations allowed the determination of the nano-morphological characteristics at the scale of individual crystals. Chemical analysis by EDS on individual crystals allows identifying characteristic minor elements associated to a special crystal in the phase mixtures. After chemical identification of the individual crystal, crystallographic phase and crystal orientation (zone axis) identification was performed on the individual crystal electron diffraction patterns (SAED).

All black samples were either identified as cryptomelane (simple non-stoichiometric Mn oxide:  $[(Ba,K)Mn_8O_{16}].xH_2O)$  alone or mixtures of cryptomelane with hollandite  $(BaMn_8O_{16}.xH_2O)$  or todorokite ( $[(Ba,Ca,K,Na)Mn_6O_{12}].H_2O$ ) (Fig.6 a-c). These crystallographic phases correspond to Mn oxides with variable amounts of Ba and K, which is consistent with the SEM-EDS observations. From a morphological point of view, there were generally longitudinal single-crystals, and special meshes were found for the

todorokite phase. The presence of K could directly be linked to the Mn oxide phase cryptomelane and not to the aluminosilicates as it was identified on individual crystals. In some cases, the K-bearing Mn oxide was prevalent with respect to the Ba-containing Mn oxide (Table 1).

Particular phases were observed in the *cave bear* (1D107) paint samples (GLAS25 and GLAS117) that showed a mixture of pyrolusite (MnO<sub>2</sub>), manganite (MnOOH) and romanechite (Ba<sub>2</sub>Mn<sub>5</sub>O<sub>10</sub>.xH<sub>2</sub>O) (Fig. 7a-c) and the black sample GLAS19 corresponding to the *cross-shaped sign* 1D112, which was composed of a mixture of cryptomelane and magnetite (Fe<sub>3</sub>O<sub>4</sub>) (Fig. 7d).

TEM-SAED provided evidence of haematite crystals in every red paint sample. Most of the haematite was poorly crystallized without characteristic shape and was mixed within aluminosilicates, but two types of lamellar crystals (lamellar sheets with hexagonal shape (type I) and lamellar sheets of trapezoidal shape (type II)), could be found and were chosen for electron diffraction. Both crystal types showed more or less regular features and variable dimensions. Hexagonal type I crystals were present in the sample GLAS24 of the *red cow* 1D103, in GLAS9 corresponding to the *signs* 1G223, in GLAS11 corresponding to the *red deer* 1G212 (size range of 100 to 250 nm) as well as in GLAS63 (Fig 8a) corresponding to the *red cow's* horn 1G206. Type II haematite was found in GLAS26 for the *red calf* 1D108 (Fig. 8b).

For the most part, haematite crystals could be found in association with hydroxyapatite crystals in red, yellow and brown flat paints of the *Bull Rotunda* such as in the *red cow* (1G206) and in the *red cow* 1D108 (GLAS26, Fig. 8c). By contrast, red lines or dots such as the red spot GLAS119, the *signs* 1G223 (GLAS9) and the red lines underlining the black horns of the *big bull* 1G203 (GLAS 8) as well as the desquamated sample LAS121 did not contain any hydroxyapatite. There were also a few black samples (the black outline of the *black and yellow deer* 1G214 (GLAS12), the black dots 1D114 (GLAS21) and the *cave bear's* ear (GLAS25), that contained hydroxyapatite, but its presence seems to be linked to yellow, brown or red paints underlaid or located very close to the black samples.

TEM-EDS also showed the local presence of a titanium (Ti) oxide phase associated with haematite crystals in the sample GLAS26 of the *red cow* 1D108 and the sample GLAS11 of the *red deer* 1G212. The type of Ti oxide phase could not be determined because it was not crystalline enough.

The yellow samples were not studied extensively using TEM but some electron diffraction patterns confirmed the presence of the Fe oxyhydroxide (goethite FeOOH) structure in good agreement with the yellow colour.

Brown paints contain haematite and cryptomelane in the form of lamellar sheets and longitudinal crystals, respectively. The presence of one Fe oxide and one Mn oxide phase was consistent with SEM-EDS analyses. The presence of hydroxyapatite nanocrystals was also observed in brown paints by TEM.

#### Discussion. New information in the ornamentation steps of the Bull Rotunda

## Chemical-mineralogical grouping of figures corresponding to one execution step at the Bull Rotunda

The TEM-EDS-SAED results were grouped according to the postulated creation steps in order to check their consistency in terms of chemical and crystallographic phases present in the corresponding paint pot:

#### 1. Step of monochromatic horses

Only one sample was studied of step 1 (GLAS16) by SEM, as the homogeneity of the figures this step was not questioned. Its black corresponded to a Ba-bearing Mn oxide in the form of rolled needles. The additional presence of aluminosilicates was detected.

#### 2. Step of bichrome and polychrome horses including the unicorn

Nine samples of this creation step were studied corresponding to black, red and brown colours. All black matter corresponds to a Ba-bearing Mn oxide either in the form of sheets, needles, spherical grains or plates according to SEM. TEM showed the presence of cryptomelane. A very unique figure in the *Bull Rotunda* is the black *unicorn* corresponding with the samples GLAS1 and GLAS2. A Ba-bearing Mn oxide was identified in the two samples with a very similar morphology corresponding both to the horn and the beard of the *unicorn*, but the paint matter was applied in two different application modes. One implies a brush and the other spraying or blowing. This difference in application mode could support the separation of the horns from the rest of the body of the *unicorn*. However, the close association of the horns continues more than 15 cm along with the lines of the head (Aujoulat, 2004). The same type of paint matter found in the horn and the beard of the beard of the *unicorn* reinforces this coherence and association of these elements in one figure despite different modes of application.

A red flat paint sample (GLAS 5) coming from the *big horse* 1G202 were also studied. The sample showed iron oxide in the form of haematite that was associated with an aluminosilicate and small amounts of a Ca-P phase, likely hydroxyapatite. We indicated the red colour in the *horse* 1G202 in this execution step with 2\* in Table 1 because it cannot be ascertained by chemical and mineralogical characterization that this colour was added during this execution step.

Brown colour was studied in the *brown horse* 1G204 (GLAS10, 61 and 62). Its paint matter was dominated by the presence of aluminosilicates rich in Fe and Mn. The Ca-P-based phase (hydroxyapatite) was also identified in the brown paint. The shape of the hydroxyapatite nanocrystals in the sample GLAS 62 indicated that they could have originated from reindeer antler in this particular case (Chadefaux et al., 2008).

It is possible that the Ca-P-based phase, hydroxyapatite, was deliberately added in order to change the consistence of the paint matter. It can also originate from the use of antler tools in the paint preparation. Taphonomy might also be an explanation for the formation of hydroxyapatite on cave walls in karst environments (Chalmin et al., 2017; Chalmin and Reiche, 2013). Finally, hydroxyapatite can be naturally associated to haematite from natural origin (Cornell and Schwertmann, 2003).

If hydroxyapatite originated from a natural mixture with haematite, it would likely have been found in all samples, which was not observed here. It was found systematically in the red and brown flat paints but was absent in the red drawing lines. Therefore, its presence seems to be related to red and brown paints applied in thin layers. Indeed, haematite nanocrystals might be intentionally diluted in other media because of its strong colouring properties giving rise to lighter red and brown colours. The systematic presence of hydroxyapatite in the red and brown flat paints indicates likely a similar geological origin of the paint matter and strengthens the hypothesis of the homogeneity of the paint preparation of this step and hence the synchronicity of the creation of these figures.

#### 3. Step of the black bulls

Ten samples corresponding to figures executed in the step 3 were examined.

The black paint matter generally corresponded to Ba-bearing Mn oxides with various morphologies including longitudinal single crystals, rolls of needles, rods, "desert roses", sheets and spherical grains. Their crystallographic phases corresponded to cryptomelane and/or hollandite. The chemical and crystallographic composition of this *black bulls* was very consistent. It was also characterized by the absence of a Ca-P phase. The *black sign* 1G226 in *big bull* 1G205 as well as the *black punctuations* 1D114 in the body of the *black bulls* of this step and were thus confirmed to belong to the same ornamentation step.

However, the *cross-shaped sign* 1D112 was revealed to be different in composition with respect to the two *black bulls* 1D101 and 1D102 represented beside it. Magnetite was detected along with cryptomelane in this black sign (GLAS 19). The cross-shaped sign 1D112, should thus be attributed to another creation step and is consequently indicated as such (3) in Table 1.

Todorokite, another crystallographic Mn oxide phase, was found with cryptomelane and hollandite in one sample of the abdomen line of the *black bull* 1D101 (GLAS 23). Its presence in GLAS 23 is probably linked to a repaint on the ventral bull strip evidenced by macrophotography.

Besides black paints, haematite was found in the red lines on the horns of the *black bull* 1G203 and dots (GLAS 8). They were either hexagonal single crystals type I or sheets. They also contained aluminosilicates rich in K and Ti. As the third step was consistently executed with black paint matter, it is questionable if the red lines of the horns belong to this step. Therefore, the samples of the red horns were indicated as 3\* in Table 1. They might be related to the creation step 5 as described below. The paint matter found in the signs 1G223 (GLAS 9) on the left-hand side of the *Bull Rotunda* in front of the bull 1G203 cannot be directly related to the haematite type found in the red lines of the bull 1G203 and might belong to another execution step. They were also indicated as step 5 in Table 1.

4. Step of the red cows and red dots on black bull

Nine red samples were studied from the figures of step 4. The red colour was related to the presence of haematite in different shapes as observed by TEM. Lamellar sheets with hexagonal single crystals type I and trapezoidal single crystals type II were found with

polycrystalline clusters (hyper-dispersed nanocrystals). These morphologies were common habits for haematite of sedimentary origin (Cornell and Schwertmann, 2003). Haematite was associated with quartz, Na- and K-rich aluminosilicates, calcium carbonate and in a quasi-systematic manner to a Ca-P-based (hydroxyapatite) phase. In the sample GLAS 26 corresponding to the red calf 1D108, hydroxyapatite with the characteristic features of that of reindeer antler (Chadefaux et al., 2008) was identified along with haematite type II. The chemical and mineralogical composition of this creation step is very consistent. Interestingly, the same association of Fe oxide-based (haematite) colouring matter with the presence of a Ca-P-phase (hydroxyapatite) was observed as for the second step of *bi- and polychrome horses*. This strengthened the observation of the systematic presence of hydroxyapatite in red and brown flat paints.

The red desquamated sample LAS121 found on the soil showed a very similar mineralogical composition, which is consistent with that of *red cows* 1D108 et 1D103. Therefore, this sample can be associated to the creation step 4.

The sample GLAS119 from the bench contained polycrystalline haematite and haematite in sheets as well as an important amount of Al-rich clay, but no calcite. The bench composition was thus different from that of the wall.

#### 5. Step of *bichrome deer*

Three samples of the creation step 5 were investigated (GLAS11-13). They showed a Babased Mn oxide and haematite or goethite. The Ba-bearing Mn oxide showed fibrous plates or filaments, whereas haematite single crystals corresponded to type I. The haematite phase was rich in Ti. The presence of Ti in haematite crystals indicated ilmenite or another Ti-Fe based mineral  $Fe_xTi_YO_z$ , which could be helpful to identify the raw material source as some magmatic rock. Some aluminosilicates, calcium carbonate and systematically a Ca-P-phase (hydroxyapatite) were found in the paint matter. The presence of aluminosilicates and calcium carbonate could be linked to the cretaceous limestone substrate. This step seemed also consistent with regard to its chemical and mineralogical composition of the figures. As mentioned above, the mineralogical features of the red horns of the *black bull* 1G203 (GLAS8, step 3\*) corresponding to hexagonal single crystals type I or sheets were similar to the samples from the *bichrome deer* (GLAS 11) and can thus be related to this execution step.

#### Other step: isolated cave bear

The *cave bear* figure sampled in GLAS25 and GLAS117 was hidden in the path of the ventral strip of the bull 1D101. The idea of a separate, probably later, creation is supported by the different chemical and mineralogical composition of the cave bear paint matter with respect to the large *black bull* 1D101 (GLAS 22 and 118, step 3). This *bull* was executed with a mixture of two different types of black Ba-bearing Mn oxides (cryptomelane and/or hollandite) whereas the *cave bear* contained a different mixture of a pure Mn oxide and hydroxide (pyrolusite and manganite) in the form of needles in mikado-shape and Babearing Mn oxide (romanechite). The particular composition of the *cave bear* excluded its correspondence to any of the five creation steps of the *Bull Rotunda*.

Figure 9 summarizes the different creation steps in terms of the chemical and mineralogical composition of the paint pots observed in different representations of the *Bull Rotunda* of the Lascaux cave.

### b) Interpretation of the execution sequence of the *Bull Rotunda* in terms of temporality of the living

The cave of Lascaux was probably much frequented, but was not a settlement site. The subsistence hunter-gatherers had camps likely at different locations with respect to their settlement sites. At the camps, they crossed and hunted the animal herds at different seasons. The construction of the *Bull Rotunda* ornamentation in several creation steps of monothematic animal depictions could be related to the natural rhythm of transhumance in the cold Palaeolithic landscape, e.g. seasons (Aujoulat, 2004; Geneste, 2021).

Each creation step translates the perception of the temporality of the living and likely represents each season. The *horses* (steps 1 and 2) represent the end of winter and the start of spring. The series of *black bulls* (step 3) and *red cows* (step 4) corresponds to summer, and the *deer* represents autumn (step 5). Lastly, the *cave bear* might represent winter as hibernating animal in these caves. The insulated figure, namely the *unicorn*, is difficult to interpret in terms of living temporality, although the *unicorn* could be associated with the *horse* series (steps 1 and 2).

## c) Comparison with previous results obtained on insulated scenes of other rooms in the Lascaux cave and with other cave sites at a regional scale

The electron microscopy study of the black paint matter used for the *Bull Rotunda* confirmed the presence of Ba-bearing Mn oxides in the form of hollandite, cryptomelane (with Ba and K) or todorokite. These pigments were already identified in other representations of the Lascaux cave (e.g. *scène du Puits, blasons, bisons adossés*), where additionally pyrolusite (pure Mn oxide), romanechite (with Ba), manganite and magnetite were found in single representations. The presence of pyrolusite and manganite was confirmed in the isolated figure of the *cave bear* of the *Bull Rotunda*. Its composition (a mixture of pyrolusite, manganite and romanechite) was similar to that of the falling horse and to the dark pigment blocks of *Diverticule axial* of the Lascaux cave. Pyrolusite and romanechite were the main components of some dark blocks found during excavations in particular in *crayons* no. 53 and 6 (Chalmin et al., 2004). This might indicate that the cave bear was executed at another moment, possibly at the same time when the falling horse of the *Diverticule Axial* was created. It should be noted that there were two other engraved, but not painted *cave bear* representations in the *Passage* of the Lascaux cave (Man-Estier, 2009).

Furthermore, there is a parallel between presence of todorokite identified in GLAS23 of the ventral stripe of *black bull* 1D101 and another figure (*blasons*) within the ornamentation of the *Nef* of the Lascaux cave and the colouring blocs found on the soil (Chalmin et al., 2006). Pyrolusite and todorokite were found in the rectangle 43 of the *blasons*. Romanechite was mixed with todorokite in the rectangle 47. Cryptomelane and todorokite were found in the rectangles 49 and 52. Therefore, todorokite, a special crystallographic phase identified in the repainted ventral stripe of the *black bull* is not unknown in other

rooms of the Lascaux cave, a fact that suggests a possible creation of this overpaint at another moment as the execution step of the *blasons*.

The black *cross-shaped sign* 1D112 was also revealed to be different in its composition to the two nearby *black bulls* 1D101 and 1D102. The presence of magnetite in the *sign* 1D112, its particular paint matter of two mixed Fe and Mn oxide-based black mineral phases, corresponds to that found in the right *bison* (GLAS103 and 105) of the *bisons adossés* in the *Nef* of the Lascaux cave (Vignaud et al., 2006). This, too, might also have been a later addition to the panel.

Concerning the red paints, different morphologies of haematite were already observed in the red paints of the *blasons* and *bisons adossés*. Strikingly, the presence of the Ca-P phase is observed in a quasi-systematic way in the Fe oxide-based paints of the *Bull Rotunda* but Ca-P was absent in the desquamated red sample LAS121 collected on the soil. Its presence in addition to the paint matter seems to be a characteristic marker of the paint preparation involved in the execution steps of the *Bull Rotunda*.

At a regional scale, the black Mn-based paint matter identified by non-invasive methods such as portable X-ray fluorescence and micro-Raman analyses in other Upper Palaeolithic decorated cave sites such as that of Rouffignac and Font-de-Gaume in Dordogne showed a different mineralogical composition with respect to those found in this study in Lascaux. According to our current knowledge, the black matters of Rouffignac and Font-de-Gaume are mixtures of romanechite and pyrolusite (Gay et al., 2020; Trosseau et al., 2021), whereas that of Lascaux showed mainly K-rich and cryptomelane-based Mn oxide. The collected paint matter of each site likely came from different sources and should not be considered on a more regional scale (in the range of some ten to hundred km) because there was a lot of choice in the region.

#### Conclusions

Scanning and transmission electron microscopy of 32 micro-samples of the figures of the Bull Rotunda of the Lascaux cave allowed the extensive study of the black and red paint matters with their associated mineralogical phases at micro- and nanoscale. While the red paint matter only contained one pigment, haematite, and the differentiation can be performed by the associated crystallographic phases, black paint matters could be distinguished thanks to the presence of different crystallographic Mn oxide phases such as cryptomelane and hollandite. These phases were shown to be Ba- and/or K-bearing Mn oxides. On this basis it was possible to confirm the five consistent creation steps of the ornamentation of the *Bull Rotunda* in the Lascaux cave proposed by previous stylistic and superimposition studies. Our new results also allowed corroborating the hypothesis to read the figures in terms of the natural life cycle of the Upper Palaeolithic hunter-gatherer society, e.g. seasons.

Some particular representations such as the black *cave bear* and one of the *cross-shaped signs* were highlighted, and understood to likely have been added at posterior moments and by other "artists" operating in deeper rooms as the *Nef* and the *Diverticule axial* of this

extraordinary cave. This was shown by the parallels in terms of paint pot composition with figures from deeper galleries in the Lascaux cave.

A comparison of the black paint matter of the *Bull Rotunda* (mainly cryptomelane, except for the *cave bear*) with that of other Palaeolithic cave sites in the Dordogne such as Rouffignac and Font-de-Gaume (generally pyrolusite and romanechite) revealed the use of different pigments in these caves indicating distinct colouring matter provenience.

#### Acknowledgments

We gratefully acknowledge the support and discussions with Hélène Salomon during sampling on site and after. She is also thanked for having provided the SEM micrographs. Françoise Pillier is thanked for her support during TEM observations and sample preparation. Anne Maigret, photographer at the C2RMF, is kindly thanked for scanning the transmission electron micrograph negatives. Nicolette is kindly acknowledged for the revision of the English language. The DRAC Nouvelle Aquitaine is acknowledged for the support of this study. Norbert Aujoulat who passed away is especially acknowledged for his fundamental work and understanding of Lascaux cave art. We dedicate this publication to his memory.

#### References

Aujoulat, N., 2004. Lascaux: le geste, l'espace et le temps, Arts rupestres. Seuil, Paris.

Aujoulat, N., Chalmin, E., Vignaud, C., Menu, M., 2002. Lascaux : les pigments noirs de la Scène du puits. 10es journées de la Section française de l'institut international de conservation (SFIIC) L'art avant l'histoire. Conservation de l'Art Préhistorique, Paris, 23-24 mai 2002, p. 5-14.

Chadefaux, C., Vignaud, C., Menu, M., Reiche, I., 2008. Multianalytical Study of Palaeolithic reindeer antler. Discovery of Antler traces in Lascaux pigments by TEM. Archaeometry 50, 516–534. https://doi.org/10.1111/j.1475-4754.2008.00373.x

Chalmin, E., 2003. Caractérisation des oxydes de manganèse et usage des pigments noirs au Paléolithic supérieur (PhD Thesis). Université de Marne la Vallée.

Chalmin, E., Hœrlé, S., Reiche, I., 2017. Taphonomy on the Surface of the Rock Wall: Rock-Paint-Atmosphere Interactions, in: David, B., McNiven, I.J. (Eds.), The Oxford Handbook of the Archaeology and Anthropology of Rock Art. Oxford University Press. https://doi.org/10.1093/oxfordhb/9780190607357.013.47

Chalmin, E., Menu, M., Pomiès, M.-P., Vignaud, C., Aujoulat, N., Geneste, J.-M., 2004. Les blasons de Lascaux. L'Anthropologie 108, 571–592. https://doi.org/10.1016/j.anthro.2004.12.001

Chalmin, E., Reiche, I., 2013. Synchrotron X-Ray Microanalysis and Imaging of Synthetic Biological Calcium Carbonate in Comparison With Archaeological Samples Originating from the Large Cave of Arcy-sur-Cure (28000-24500 BP, Yonne, France). Microsc. Microanal. 19, 1523–1534. https://doi.org/10.1017/S1431927613013342

Chalmin, E., Vignaud, C., Salomon, H., Farges, F., Susini, J., Menu, M., 2006. Minerals discovered in paleolithic black pigments by transmission electron microscopy and micro-

| 1<br>2                                       |   |
|--|---|
| 2<br>3<br>4<br>5                             | X-ray absorption near-edge structure. Appl. Phys. A 83, 213–218.<br>https://doi.org/10.1007/s00339-006-3510-7   |
| 6<br>7<br>8                                  | Cornell, R.M., Schwertmann, U., 2003. The Iron Oxides: Structure, Properties, Reactions, Occurences and Uses, 1st ed. Wiley. https://doi.org/10.1002/3527602097   |
| 9<br>10<br>11<br>12                          | Ducasse, S., Langlais, M., 2019. Twenty years on, a new date with Lascaux.<br>Reassessing the chronology of the cave's Paleolithic occupations through new 14C AMS dating. Paléo 130–147. https://doi.org/10.4000/paleo.4558  |
| 13<br>14<br>15                               | Fritz, C., Tosello, G., 2015. From Gesture to Myth: Artists' techniques on the walls of Chauvet Cave. Palethnologie. https://doi.org/10.4000/palethnologie.876  |
| 16<br>17<br>18<br>19<br>20                   | Gay, M., Plassard, F., Müller, K., Reiche, I., 2020. Relative chronology of Palaeolithic drawings of the Great Ceiling, Rouffignac cave, by chemical, stylistic and superimposition studies. J. Archaeol. Sci. Rep. 29, 102006.<br>https://doi.org/10.1016/j.jasrep.2019.102006 |
| 21<br>22<br>23<br>24                         | Geneste, JM., 2021. Les grandes lignes d'un monde mythique insoupçonné. Des rythmes, de la temporalité et de la place du sujet dans l'art rupestre paléolithique. Afr. Stud. 9–27.  |
| 25<br>26<br>27                               | Laming-Emperaire, A., 1959. Lascaux : Paintings and Engravings, Harmondsworth. ed. Pelican Books, Middlesex, Engl. ; Baltimore, Maryland.   |
| 28<br>29<br>30                               | Leroi-Gourhan, A., Allain, J. (Eds.), 1979. Lascaux inconnu, Supplément à "Gallia préhistoire." Editions de Centre national de la recherche scientifique, Paris.  |
| 31<br>32<br>33                               | Man-Estier, E., 2009. Les ursidés au naturel et au figuré pendant la préhistoire (Sciences de l'Homme et Société.). Muséum National d'Histoire Naturelle, Paris, France.  |
| 34<br>35<br>36                               | Trosseau, A., Maigret, A., Coquinot, Y., Reiche, I., 2021. <i>In situ</i> XRF study of black colouring matter of the Palaeolithic figures in the Font-de-Gaume cave. J. Anal. At. Spectrom. 36, 2449–2459. https://doi.org/10.1039/D1JA00202C                                   |
| 37<br>38<br>39<br>40<br>41<br>42<br>43       | Vignaud, C., Salomon, H., Chalmin, E., Geneste, JM., Menu, M., 2006. Le groupe des « bisons adossés » de Lascaux. Étude de la technique de l'artiste par analyse des pigments. L'Anthropologie, Art préhistorique 110, 482–499. https://doi.org/10.1016/j.anthro.2006.07.008    |
| 44<br>45                                     | Data availability statement   |
| 46<br>47<br>48<br>49                         | The data are available through request to the corresponding author.   |
| 50<br>51                                     | Figure and table captions   |
| 52<br>53<br>54<br>55<br>56<br>57<br>58<br>59 | Figure 1. Lascaux cave map with location of the left and right wall of <i>Bull Rotonda</i> . © <i>Bassier</i>   |
| 60   |   |

Figure 2. Left wall of *Bull Rotonda* with location of the microsamples (GLASXX). Numbering 1G10X corresponds to each entity: black monochrome figures are indicated in black, red monochrome figures are indicated in red and bichrome figures are indicated in green. © I. Reiche, CNRS, adapted from N. Aujoulat.

Figure 3. Right wall of *Bull Rotonda* with location of the micro-samples (GLASXX). LAS121 corresponds to a desquamated sample from the ground. Numbering 1D10X corresponds to each entity: black monochrome figures are indicated in black and red monochrome figures are indicated in red. © I. Reiche, CNRS, adapted from N. Aujoulat.

Figure 4. Optical micrographs: (a and b) red desquamation LAS121 sample, (b) prepared as a cross section; (c and d) black samples: (c) *big bull* 1D101 GLAS22 and (d) *cave bear* GLAS 117. © C. Vignaud

Figure 5. Scanning electron micrographs (SEM in BSE mode): (a and b) cross section las 121 sample, (b) detail on punctual red pigment; (c and d) *cave bear* GLAS117, Mn oxide, Ba-bearing Mn oxide and Ca oxide, (d) detail on Ba-bearing Mn oxide. ©E. Chalmin and H. Salomon

Figure 6. Transmission electron micrographs of black phases detected in the black *big bull* 1D101 samples GLAS 22 and 23. (a and b) GLAS 22 on the its back line, (c) GLAS 23 on its black ventral strip: (a) hollandite, monoclinic, zone axis [0-11], (b) cryptomelane, monoclinic, zone axis [001], (c) todorokite, monoclinic, characteristic meshwork crystals. © E. Chalmin and C. Vignaud


Figure 7. Transmission electron micrographs with corresponding electron diffraction patterns of the black phases detected in the *cave bear* sample GLAS 117 (a, b and c) and in the *cross* 1D112 sample GLAS19 (d): (a) manganite, monoclinic, zone axis [2-10], (b) romanechite, monoclinic, zone axis [021], (c) pyrolusite, tetragonal, zone axis [12-1], (d) magnetite Fe<sub>3</sub>O<sub>4</sub>, face-centered cubic, zone axis [10-1]. © E. Chalmin and C. Vignaud

Figure 8. Transmission electron micrographs with corresponding electron diffraction patterns of the red phases detected in the *red cow* 1G206 and the *red calf* 1D108: a) haematite crystal of hexagonal shape in GLAS 63 corresponding to the *red cow* (1G206) and corresponding electron diffraction (trigonal), b) Haematite crystal of trapezoidal shape found in micro-samples GLAS 26 corresponding to the *red calf* 1D108, c) Haematite nanocrystals mixed within hydroxyapatite in GLAS 26 and corresponding TEM-EDX spectrum indicating the presence of a Ca-P-based phase (hydroxyapatite) and Fe oxide trapped in an aluminosilicate matrix. © C. Vignaud

Figure 9. Schematic overview of the different creation steps in terms of the chemical and mineralogical composition of the paint pots observed in different representations of the *Bull Rotunda* of the Lascaux cave. © Conception: I. Reiche, C. Vignaud and E. Chalmin; Realization: E. Chalmin

Table 1. Summary of the morphological, mineralogical and chemical features at the microand at nanoscale of the studied micro-samples of the *Bull Rotunda*. If the creation step is shown in brackets it means that its attribution to this step has changed according to the determined mineralogical composition. If an asterisk is indicated at the number of the execution step it means that this attribution is questionable and cannot be ascertained.

| 1 2                  |  |
|----------------------|--|
| 3<br>4<br>5          |  |
| 5<br>6<br>7          |  |
| 8<br>9               |  |
| 10<br>11             |  |
| 12<br>13             |  |
| 14<br>15             |  |
| 16<br>17<br>18       |  |
| 19<br>20             |  |
| 21<br>22             |  |
| 23<br>24             |  |
| 25<br>26             |  |
| 27<br>28<br>29       |  |
| 30<br>31             |  |
| 32<br>33             |  |
| 34<br>35             |  |
| 35<br>36<br>37<br>38 |  |
| 39<br>40             |  |
| 41<br>42             |  |
| 43<br>44             |  |
| 45<br>46<br>47       |  |
| 48<br>49             |  |
| 50<br>51             |  |
| 52<br>53             |  |
| 54<br>55<br>56       |  |
| 50<br>57<br>58       |  |
| 59<br>60             |  |





60




Figure 1. Lascaux cave map with location of the left and right wall of Bull Rotonda. © Bassier 137x97mm (118 x 118 DPI)




Figure 2. Left wall of Bull Rotonda with location of the microsamples (GLASXX). Numbering 1G10X corresponds to each entity: black monochrome figures are indicated in black, red monochrome figures are indicated in red and bichrome figures are indicated in green. © I. Reiche, CNRS, adapted from N. Aujoulat.

275x139mm (59 x 59 DPI)




Figure 3. Right wall of Bull Rotonda with location of the micro-samples (GLASXX). LAS121 corresponds to a desquamated sample from the ground. Numbering 1D10X corresponds to each entity: black monochrome figures are indicated in black and red monochrome figures are indicated in red. © I. Reiche, CNRS, adapted from N. Aujoulat.

117x103mm (118 x 118 DPI)




Figure 4. Optical micrographs: (a and b) red desquamation LAS121 sample, (b) prepared as a cross section; (c and d) black samples: (c) big bull 1D101 GLAS22 and (d) cave bear GLAS 117. © C. Vignaud

748x517mm (118 x 118 DPI)




Figure 5. Scanning electron micrographs (SEM in BSE mode): (a and b) cross section las 121 sample, (b) detail on punctual red pigment; (c and d) cave bear GLAS117, Mn oxide, Ba-bearing Mn oxide and Ca oxide, (d) detail on Ba-bearing Mn oxide. ©E. Chalmin and H. Salomon

758x521mm (59 x 59 DPI)




Figure 6. Transmission electron micrographs of black phases detected in the black big bull 1D101 samples GLAS 22 and 23. (a and b) GLAS 22 on the its back line, (c) GLAS 23 on its black ventral strip: (a) hollandite, monoclinic, zone axis [0-11], (b) cryptomelane, monoclinic, zone axis [001], (c) todorokite, monoclinic, characteristic meshwork crystals. © E. Chalmin and C. Vignaud

206x209mm (59 x 59 DPI)




Figure 8. Transmission electron micrographs with corresponding electron diffraction patterns of the red phases detected in the red cow 1G206 and the red calf 1D108: a) haematite crystal of hexagonal shape in GLAS 63 corresponding to the red cow (1G206) and corresponding electron diffraction (trigonal), b) Haematite crystal of trapezoidal shape found in micro-samples GLAS 26 corresponding to the red calf 1D108, c) Haematite nanocrystals mixed within hydroxyapatite in GLAS 26 and corresponding TEM-EDX spectrum indicating the presence of a Ca-P-based phase (hydroxyapatite) and Fe oxide trapped in an aluminosilicate matrix. © C. Vignaud

84x128mm (96 x 96 DPI)




Figure 9. Schematic overview of the different creation steps in terms of the chemical and mineralogical composition of the paint pots observed in different representations of the Bull Rotunda of the Lascaux cave. © Conception: I. Reiche, C. Vignaud and E. Chalmin; Realization: E. Chalmin

756x505mm (118 x 118 DPI)

**Table 1.** Summary of the morphological, mineralogical and chemical features at the micro- and at nanoscale of the studied micro-samples of the Bull Rotunda. If the creation step is shown in brackets it means that its attribution to this step has changed according to the determined mineralogical composition. If an asterisk is indicated at the number of the execution step it means that this attribution is questionable and cannot be ascertained.

| Hypothe-<br>sized<br>creation<br>step | Figure                     | Sample<br>number | Main colour /<br>rendering             | Other phases or<br>elements                      | Inferred<br>creation<br>step |
|---------------------------------------|----------------------------|------------------|--|--|------------------------------|
| Results indicated                     | Type of<br>paint<br>matter |                  | Crystal morphology                     | Additional minerals                              |                              |
| 1                                     | Small<br>black<br>horse    | GLAS16           | Black / filling                        |  |                              |
|                                       | Mixed Mn o<br>Ba           | xide with        | Needle rolls                           | Aluminosilicates                                 | 1                            |
| 2                                     | Unicorn 1G                 | 6201<br>GLAS1    | Black horn /<br>continuous outline     |  |                              |
|                                       | Mixed Mn o<br>Ba           |                  | Sheets 10µm x 1µm                      | Aluminosilicates                                 | 2                            |
|                                       |                            | GLAS2            | Black beard /<br>tight punctuations    |  |                              |
|                                       | Mixed Mn o<br>Ba and K     | xide with        | Sheets 10µm x 1µm                      | Aluminosilicates                                 | 2                            |
| 2                                     | Big horse 1                | IG202<br>GLAS3   | Black back leg /<br>continuous outline |  |                              |
|                                       | Mixed Mn o<br>Ba           | xide with        | Spherical grains of 0.5µm and sticks   | Aluminosilicates<br>CI+Hg                        |                              |
|                                       |                            | GLAS6            | Black neckline / tigh punctuations     | t  |                              |
|                                       | Mixed Mn o<br>Ba           | xide with        | Spherical grains of<br>0.5µm to 1µm    | Silicate<br>(absence of Al)                      | 2                            |
|                                       |                            | GLAS4            | Red neckline / contin<br>outline       | nuous  | 2*                           |
|                                       | Fe oxide                   |                  | Sheets and<br>plaquettes               | Aluminosilicates and<br>Mn traces                |                              |
|                                       |                            | GLAS5            | Red flat colour / fillir               | ng   | 2*                           |
|                                       | Fe oxide                   |                  | Sheets and plaquettes                  | Ca/P phase,<br>aluminosilicates and<br>Mn traces |                              |
| 2                                     | Brown hors<br>GLAS10       | se 1G204         | Brown / continuous outline             |  |                              |

|   | Fe oxide +               | Mn oxide       | Fe and Mn enriched aluminosilicates                               | Ca/P phase,<br>Aluminosilicates,  | 2 |
|---|--------------------------|----------------|---|---|---|
|   |                          |                |   | calcium carbonate, calcium sulfate  |   |
|   |                          | GLAS61         | Proximal end of the I<br>line of the back /<br>continuous outline | black   |   |
|   | Mixed Mn c<br>Ba         | oxide with     | Needles of 0.5 to<br>2.5µm length                                 | Aluminosilicates,<br>quartz, hair included<br>in Mn oxide, CI+Hg<br>(pollution) | 2 |
|   |                          | GLAS62         | Brown distal end of I   | back  |   |
|   | Fe oxide +               | Mn oxide       | Sheets Fe/O, fibres<br>Mn/O                                       | Ca/P phase<br>Aluminosilicates  |   |
|   | Haematite,<br>cryptomela | ne             | Single crystals and<br>sticks of 200 to 400<br>nm                 | Ca/P phase in the<br>form of<br>hydroxyapatite likely<br>from reindeer antler   |   |
| 3 | Big bull<br>1G203        | GLAS7          | Black horn /<br>continuous outline                                |   |   |
|   | Mixed Mn c<br>Ba         | oxide with     | Sheets inf. 1µm   | Aluminosilicates  | : |
| 3 |                          | GLAS8          | Red horn / tight punctuations                                     |   |   |
|   | Fe oxide                 |                | Sheets 1-10µm   | Aluminosilicates + K<br>and Ti  | ł |
| 3 | Signs 1G2                | 23 GLAS9       | Red dots and lines / punctuations                                 |   |   |
|   | Fe oxide                 |                | Fibres  | Aluminosilicates, <b>quartz</b> , K, Ti   |   |
|   | Haematite                |                | Hexagonal single crys<br>type I                                   | tals -  | Į |
| 3 | Black sign<br>the body o |                | Black / punctuations  |   |   |
|   | Mixed Mn o<br>Ba and K   | oxide with     | "Desert roses"  | Ti  |   |
|   | Cryptomela<br>hollandite | ne and         | Longitudinal single crystals                                      | TiO <sub>2</sub>  | 3 |
| 3 | Big bull 1G              | 6205<br>GLAS17 | Black eye / tight<br>punctuations                                 |   |   |
|   | Mixed Mn c<br>Ba         | oxide with     |   | Aluminosilicates<br>grains 0.1µm piled<br>up 0.5µm                              |   |

| 2<br>3         | 1 |
|----------------|---|
| 4              |   |
| 5<br>6<br>7    |   |
| 8<br>9         |   |
| 10<br>11       |   |
| 12<br>13       |   |
| 13<br>14<br>15 |   |
| 16<br>17       |   |
| 17<br>18<br>19 |   |
| 20<br>21       |   |
| 22<br>23       |   |
| 23<br>24<br>25 |   |
| 26<br>27       |   |
| 28<br>29       |   |
| 30<br>31       |   |
| 32<br>33       |   |
| 34<br>35       |   |
| 36<br>37       |   |
| 38<br>39       |   |
| 40<br>41       |   |
| 42<br>43       |   |
| 44<br>45       |   |
| 46<br>47       |   |
| 48<br>49       |   |
| 50<br>51       |   |
| 52<br>53       |   |
| 54<br>55       |   |
| 56<br>57       |   |
| 58<br>59       |   |
| 60             |   |

|   |                          | GLAS18                                 | Black back / continuo<br>outline | ous   |   |
|---|--------------------------|--|----------------------------------|---|---|
|   | Mixed Mn o<br>Ba         | xide with                              | Rolls of needles 2 x<br>0.1µm    | Aluminosilicates + Al                                 |   |
|   | Cryptomela<br>hollandite | ne and                                 | Longitudinal single crystals     | -   | 3   |
| 3 | muzzle of 1              | 12 in front of<br>ID102<br>GLAS19      | the Black / co                   | ontinuous outline                                     |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Needles, rolls and sticks        | aluminosilicates +<br>quartz, Hg                      | (3)<br>posterior<br>to the                |
|   | Cryptomela<br>magnetite  | ne and                                 | Single crystals                  | -   | decora-<br>tion of the<br>Bull<br>Rotunda |
| 3 | Big bull 1D<br>GLAS20    | 102                                    | Black chamfer facing             | eye / punctuations                                    |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Spherical grains and sticks      | aluminosilicates,<br>quartz                           |   |
|   | Cryptomela               | ne                                     | Longitudinal single crystals     | -   | 3   |
| 3 |                          | tuations 1D <sup>.</sup><br>101 GLAS21 |                                  |   |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Spherical grains and sticks      | Ca/P phase,<br>aluminosilicates and<br>Hg (pollution) |   |
|   | Cryptomela               | ne                                     | Longitudinal single crystals     | -   | 3   |
| 3 | Big bull 1D<br>GLAS22    | 101                                    | Black back / continuo            | ous outline   |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Rolls                            | aluminosilicates +Al,<br>Hg                           |   |
|   | Cryptomela               | ne and hollar                          | dite                             | -   | 3   |
| 3 |                          | GLAS23                                 | Black abdomen /<br>punctuations  | outside the bear                                      |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Rolls and sticks                 | aluminosilicates +Al,<br>Hg                           |   |
|   | Todorokite               |  | Meshes                           | -   | 3*  |
| 3 |                          | GLAS118                                | Black abdomen /<br>punctuations  | outside the bear                                      |   |
|   | Mixed Mn o<br>Ba         | xide with                              | Sheets 1µm and needles           | Calcium carbonate                                     |   |
|   | Cryptomela               | ne                                     | Rare small crystals              | -   | 3*  |

| 1              |
|----------------|
| 2              |
| 3              |
| 4              |
| 5<br>6         |
| 7              |
| 8              |
| 9              |
| 10<br>11       |
| 12             |
| 13             |
| 14             |
| 15<br>16       |
| 17             |
| 18             |
| 19             |
| 20<br>21       |
| 22             |
| 23             |
| 24             |
| 25<br>26       |
| 27             |
| 28             |
| 29             |
| 30<br>31<br>32 |
| 32             |
| 33             |
| 34<br>35       |
| 36             |
| 37             |
| 38             |
| 39             |
| 40<br>41       |
| 42             |
| 43             |
| 44<br>45       |
| 45<br>46       |
| 47             |
| 48             |
| 49<br>50       |
| 50<br>51       |
| 52             |
| 53             |
| 54<br>55       |
| 55<br>56       |
| 57             |
| 58             |
| 59<br>60       |

| 4 | Big red cov        | w 1G206<br>GLAS15       | Red flat colour / filling                              | I  |   |
|---|--------------------|-------------------------|--|--|---|
|   | Fe oxide           |                         | Poor in oxide  | Ca/P phase   | 4 |
|   |                    | GLAS63                  | Red left horn / filling                                |  |   |
|   | Fe oxide           |                         | Needles and small plates                               | Quartz   |   |
|   | Haematite          |                         | Hexagonal single<br>crystals type I (and<br>II), 1.2µm | -  | 4 |
| 4 | Red cow 1I         | D103<br>GLAS24          | Red tail / continuous o                                | outline  |   |
|   | Fe oxide           |                         | Spherical grains<br>0.1µm                              | Ca/P phase,<br>aluminosilicates  |   |
|   | Haematite          |                         | Lamellar sheets type I                                 |  | 4 |
|   |                    | GLAS120                 | Red / filling  |  |   |
|   | Fe oxide           |                         | K-bearing alumina-<br>silicates with Fe                | Calcite with<br>pigment in the<br>fissures   | 4 |
| 4 | Red calf 1D        | 0108<br>GLAS26          | Red / filling  |  |   |
|   | Fe oxide           |                         |  | <b>Ca/P phase</b> ,<br>Alumino-silicates +<br>Fe, Ti, K                                    | 4 |
|   | Haematite          |                         | Sheets type II and rods                                | Hydroxy-<br>apatite likely<br>from reindeer<br>antler                                      | 4 |
| 4 | Red spot o<br>seat | n bench<br>GLAS119      | Red spot under t<br>red cow / punctua                  |  |   |
|   | Fe oxide           |                         |  | many Fe grains<br>entrapped in K-<br>bearing<br>aluminosilicates and<br>locally Na-bearing |   |
|   | Hematite           |                         | Hematite in sheets and polycrystalline hematite        | -  | 4 |
| 4 |                    | amation: flal<br>LAS121 | ke on the ground                                       | Under the red cow  |   |
|   | Fe oxide           |                         | Porous calcite wall                                    | aluminosilicates<br>rich in Fe, K, Cl  |   |
|   | Hematite           |                         | Hematite in sheets and different crystal dimension     |  | 4 |
| 5 | Two-heade<br>1G212 | d deer<br>GLAS11        | Red / continuous outli                                 | ine  |   |

|       | Fe oxide                  |                       | Fibrous plates <1µm                     | Ca/P phase a few aluminosilicates, Ti                   |                                |
|-------|---------------------------|-----------------------|---|---|--------------------------------|
|       | Hematite                  |                       | Single crystals type I,<br>100-250nm    | Ті  | 5                              |
| 5     |                           | yellow deer<br>GLAS12 | Black outline / contine                 | uous outline  |                                |
|       | Mixed Mn o<br>Ba          | xide with             | Filaments 1µmx0.1µm                     | Ca/P phase  | 5                              |
|       |                           | GLAS13                | Yellow body / filling                   |   |                                |
|       | Fe oxide                  |                       | Few material                            | Ca/P phase,<br><b>Ca carbonate</b><br>aluminosili-cates | 5                              |
| other | Bear<br>1D107             | GLAS25                | Black ear / punctuatio                  | on  |                                |
|       | Mn oxide ar<br>Mn oxide w |                       | Grains and rods                         | Ca/P phase,<br>aluminosili-cates, Mg                    | posterior<br>to the<br>decora- |
|       | Pyrolusite, and romane    |                       | Longitudinal single crystals            |   | tion of the<br>Bull<br>Rotunda |
|       |                           | GLAS117               | Black muzzle / punctu                   | lation  |                                |
|       | Mn oxide ar<br>Mn oxide w |                       | Prisms of Mn oxide<br>and needles Mn-Ba | big prisms of calcite<br>and round quartz<br>grains     | posterior<br>to the<br>decora- |
|       | Pyrolusite a romanechite  |                       | Needles in mikado-sha single crystals   | pe and longitudinal                                     | tion of the<br>Bull<br>Rotunda |
|       |                           |                       |   | 2   | <u>.</u>                       |