



HAL
open science

Convergence of the solutions of the discounted Hamilton–Jacobi equation

Andrea Davini, Albert Fathi, Renato Iturriaga, Maxime Zavidovique

► **To cite this version:**

Andrea Davini, Albert Fathi, Renato Iturriaga, Maxime Zavidovique. Convergence of the solutions of the discounted Hamilton–Jacobi equation. *Inventiones Mathematicae*, 2016, pp.29-55. 10.1007/s00222-016-0648-6 . hal-04502592

HAL Id: hal-04502592

<https://hal.science/hal-04502592>

Submitted on 13 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONVERGENCE OF THE SOLUTIONS OF THE DISCOUNTED EQUATION

ANDREA DAVINI, ALBERT FATHI, RENATO ITURRIAGA AND MAXIME ZAVIDOVIQUE

ABSTRACT. We consider a continuous coercive Hamiltonian H on the cotangent bundle of the compact connected manifold M which is convex in the momentum. If $u_\lambda : M \rightarrow \mathbb{R}$ is the viscosity solution of the discounted equation

$$\lambda u_\lambda(x) + H(x, d_x u_\lambda) = c(H),$$

where $c(H)$ is the critical value, we prove that u_λ converges uniformly, as $\lambda \rightarrow 0$, to a specific solution $u_0 : M \rightarrow \mathbb{R}$ of the critical equation

$$H(x, d_x u) = c(H).$$

We characterize u_0 in terms of Peierls barrier and projected Mather measures.

1. INTRODUCTION

The so called *ergodic approximation* is a technique introduced in [20] to show the existence of viscosity solutions to an equation of the kind

$$H(x, d_x u) = c, \tag{1.1}$$

where c is a real constant and H , the *Hamiltonian*, is a continuous function defined on $\mathbb{T}^k \times \mathbb{R}^k$, where $\mathbb{T}^k = \mathbb{R}^k / \mathbb{Z}^k$ is the canonical flat torus.

In fact, the arguments in [20] work as well for a Hamiltonian defined on the cotangent bundle of a compact manifold. Therefore in the sequel $H : T^*M \rightarrow \mathbb{R}$ will be a given continuous function, called the Hamiltonian, where T^*M is the cotangent bundle of M , compact connected manifold without boundary.

The method in [20] to find solutions of (1.1) is to perturb the Hamiltonian by adding a term consisting of u multiplied by a positive parameter λ to obtain the *discounted equation*

$$\lambda u(x) + H(x, d_x u) = 0. \tag{1.2}$$

This equation obeys a maximum principle, and therefore it has a unique solution $u_\lambda : M \rightarrow \mathbb{R}$. The idea is then to study the behavior of u_λ when the *discount factor* λ tends to zero. When the Hamiltonian $H(x, p)$ is coercive in p , uniformly with respect to x , the functions λu_λ are equi-bounded and the u_λ are equi-Lipschitz. Furthermore, the functions $-\lambda u_\lambda$ uniformly converge on M , as λ tends to 0, to a constant $c(H)$, henceforth termed *critical value*. By adding a suitable constant to each function u_λ , we obtain an equi-bounded and equi-Lipschitz family of functions $\hat{u}_\lambda : M \rightarrow \mathbb{R}$ satisfying, for each $\lambda > 0$,

$$H(x, d_x \hat{u}_\lambda) = -\lambda u_\lambda(x)$$

Date: Submitted Version 12 Mars 2014.

2010 Mathematics Subject Classification. 35B40, 37J50, 49L25.

Key words and phrases. asymptotic behavior of solutions, weak KAM Theory, viscosity solutions, optimal control.

Work supported by ANR-07-BLAN-0361-02 KAM faible & ANR-12-BS01-0020 WKBHJ.

in the viscosity sense. By the Ascoli–Arzelà Theorem and the stability of the notion of viscosity solution, we derive that the functions \hat{u}_λ uniformly converge, *along subsequences* as λ goes to 0, to global viscosity solutions on M of the *critical equation*

$$H(x, d_x u) = c(H) \quad \text{in } M. \quad (1.3)$$

This is also the sole equation of the family (1.1) that admits solutions. Solutions, subsolutions and supersolutions of (1.3) will be termed *critical* in the sequel.

Due to the lack of a uniqueness result for the critical equation, it is not clear at this point that limits of \hat{u}_λ along different subsequences yield the same solution of (1.3). In this paper, we address the problem when H is convex in the momentum.

Theorem 1.1. *Let $H : T^*M \rightarrow \mathbb{R}$ be a continuous Hamiltonian, which is coercive, and convex in the momentum. For $\lambda > 0$, denote by $u_\lambda : M \rightarrow \mathbb{R}$ the unique continuous viscosity solution of*

$$\lambda u_\lambda + H(x, d_x u_\lambda) = c(H), \quad (1.4)$$

where $c(H)$ is the critical value of H . The family u_λ converges as $\lambda \rightarrow 0$, to a single critical solution u_0 .

Note that we have replaced 0 in the second member of (1.2) by the critical constant $c(H)$. With this choice, the solutions of (1.4) are uniformly bounded independently of λ , see Corollary 4.5 below. Note also that the solution of

$$\lambda u + H(x, d_x u) = c$$

is $u_\lambda + c/\lambda$, where u_λ is the solution of (1.2). Therefore there is at most one c for which the family of solutions are bounded, independently of λ .

In fact, as we will see, without loss of generality we can assume in Theorem 1.1 that H is superlinear. In that case, by Fenchel’s formula, the Hamiltonian H has a conjugated Lagrangian $L : TM \rightarrow \mathbb{R}$ which is superlinear and convex in the fibers of the tangent bundle. We can then apply weak KAM theory—or rather its extension to general Lagrangians, see the appendices to this paper—to characterize u_0 in terms of the Peierls barrier and of projected Mather measures—defined respectively by equation (B.3) and Definition B.21 in §B.2.

Proposition 1.2. *The function $u_0 = \lim_{\lambda \rightarrow 0} u_\lambda$, obtained in Theorem 1.1 above, can be characterized in either of the following two ways:*

- (i) *it is the largest critical subsolution $u : M \rightarrow \mathbb{R}$ such that $\int_M u d\mu \leq 0$ for every projected Mather measure μ ,*
- (ii) *it is the infimum over all projected Mather measure μ of the functions h_μ defined by $h_\mu(x) = \int_M h(y, x) d\mu(y)$, where h is the Peierls barrier.*

The theorem and proposition above extend the results of Renato Iturriaga and Hector Sánchez-Morgado [19], where the convergence is proved for a Tonelli Hamiltonian under the assumption that the Aubry set consists of a finite number of hyperbolic fixed point of the Lagrangian flow. In [18], Diogo Gomes found some constraints on the possible accumulation points of u_λ in terms of a concept of generalized Mather measures.

Since we could prove the results without any regularity assumptions on H , we need to establish firmly the weak KAM theory beyond its previous scopes to general continuous superlinear Lagrangians convex in the fibers, with a particular emphasis on the Mather measures. This is done in two appendices. Most of the results in these appendices appear here for the first time in this generality.

2. PRELIMINARIES

In this work, we will denote by M a compact connected smooth manifold without boundary of dimension m . It will be convenient to endow M with an auxiliary C^∞ Riemannian metric. The associated Riemannian distance on M will be denoted by d . We denote by TM the tangent bundle and by $\pi : TM \rightarrow M$ the canonical projection. A point of TM will be denoted by (x, v) with $x \in M$ and $v \in T_x M = \pi^{-1}(x)$. In the same way, a point of the cotangent bundle T^*M will be denoted by (x, p) , with $x \in M$ and $p \in T_x^*M$ a linear form on the vector space $T_x M$. We will denote by $p(v)$ the value of the linear form $p \in T_x^*M$ evaluated at $v \in T_x M$, and by $\|v\|_x$ the norm of v at the point x . We will use the same notation $\|p\|_x$ for the dual norm of a form $p \in T_x^*M$.

On a smooth manifold like M , there is an intrinsic notion of measure zero set: a subset Z of M is said to be of measure zero, if for every smooth coordinate patch $\varphi : U \rightarrow \mathbb{R}^m$, the image $\varphi(U \cap Z)$ has Lebesgue measure 0 in \mathbb{R}^m . Note that Z has measure zero in this sense if and only if it has measure 0 for the *Riemannian volume measure* associated to a Riemannian metric. We say that a property holds *almost everywhere* (*a.e.* for short) on M if it holds up to a set of measure zero as defined above.

Since M is compact, we can endow the space $C(M, \mathbb{R})$ of continuous real function on M with the *sup-norm*

$$\|u\|_\infty := \sup_{x \in M} |u(x)|, \quad u \in C(M, \mathbb{R}).$$

We will say that κ is a *Lipschitz constant* for $u \in C(M, \mathbb{R})$ if it satisfies $u(x) - u(y) \leq \kappa d(x, y)$ for every $x, y \in M$. Any such a function will be termed *Lipschitz*, or κ -*Lipschitz* if we want to specify the Lipschitz constant. The space of real valued Lipschitz functions on M will be denoted by $\text{Lip}(M, \mathbb{R})$. If the function $u : M \rightarrow \mathbb{R}$ is differentiable at a point $x \in M$, we will denote by $d_x u$ its derivative (called also differential). Rademacher's theorem states that a (locally) Lipschitz function has a derivative almost everywhere.

Given a continuous function u on M , we will call *subtangent* (respectively, *supertangent*) of u at y a function ϕ of class C^1 in a neighborhood U of y such that $u - \phi$ has a local minimum (resp., maximum) at y . Its differential $d_y \phi$ will be called a *subdifferential* (resp. *superdifferential*) of u at y , respectively. The set of sub and superdifferentials of u at y will be denoted $D^-u(y)$ and $D^+u(y)$, respectively. The function ϕ will be furthermore termed *strict subtangent* (resp., *strict supertangent*) if $u - \phi$ has a *strict* local minimum (resp., maximum) at y . Any subtangent (resp., supertangent) ϕ of u can be always assumed strict at y without affecting $d_y \phi$ by possibly replacing it with $\phi - d^2(y, \cdot)$ (resp. $\phi + d^2(y, \cdot)$).

We recall that u is differentiable at y if and only if $D^+u(y)$ and $D^-u(y)$ are both nonempty. In this instance, $D^+u(y) = D^-u(y) = \{d_y u\}$. We refer the reader to [7] or [13] for the proofs.

Let $G \in C(\mathbb{R} \times T^*M)$ and let us consider the following Hamilton–Jacobi equation:

$$G(u(x), x, d_x u) = 0 \quad \text{in } M. \tag{2.1}$$

Let $u \in C(M, \mathbb{R})$. We will say that u is a *viscosity subsolution* of (2.1) if

$$G(u(x), x, p) \leq 0 \quad \text{for every } p \in D^+u(x) \text{ and } x \in M.$$

We will say that u is a *viscosity supersolution* of (2.1) if

$$G(u(x), x, p) \geq 0 \quad \text{for every } p \in D^-u(x) \text{ and } x \in M.$$

We will say that u is a *viscosity solution* if it is both a sub and a supersolution. In the sequel, solutions, subsolutions and supersolutions will be always meant in the viscosity sense, hence the adjective *viscosity* will be omitted. Moreover, they will be implicitly assumed continuous, with no further specification.

It is easily seen, by Rademacher's theorem, that a Lipschitz-continuous subsolution is also an almost everywhere subsolution. The converse is not true, in general. However, when G is convex in p , the following holds, see any of the references [2, 3, 13, 24]:

Proposition 2.1. *Assume $G \in C(\mathbb{R} \times T^*M)$ such that $G(u, x, \cdot)$ is convex in T_x^*M for every fixed $u \in \mathbb{R}$ and $x \in M$. Let $u \in \text{Lip}(M, \mathbb{R})$. The following facts are equivalent:*

- (i) $G(u(x), x, p) \leq 0$ for every $p \in D^+u(x)$ and $x \in M$;
- (ii) $G(u(x), x, p) \leq 0$ for every $p \in D^-u(x)$ and $x \in M$;
- (iii) $G(u(x), x, p) \leq 0$ for every $p \in \partial^c u(x)$ and $x \in M$;
- (iv) $G(u(x), x, d_x u) \leq 0$ for a.e. $x \in M$.

In (iii), we have used the Clarke derivative $\partial^c u(x)$ of the Lipschitz function u at the point x . For definition and properties of the Clarke derivative see §A.1.

3. CRITICAL AND DISCOUNTED HAMILTON–JACOBI EQUATIONS

In this paper, we will consider a continuous function $H : T^*M \rightarrow \mathbb{R}$, called the Hamiltonian, satisfying the following assumptions:

- (H1) (Convexity) For every $x \in M$, the map $p \mapsto H(x, p)$ is convex on T_x^*M .
- (H2) (Coercivity) $H(x, p) \rightarrow +\infty$ as $\|p\|_x \rightarrow +\infty$ uniformly in $x \in M$.

The coercivity condition will be actually reinforced as follows:

- (H2') (Superlinearity) $H(x, p)/\|p\|_x \rightarrow +\infty$ as $\|p\|_x \rightarrow +\infty$ uniformly in $x \in M$.

A Hamiltonian H satisfying (H1)–(H2') will be furthermore termed *Tonelli* if it is of class C^2 on T^*M , and $\partial^2 H / \partial p^2(x, p)$ is a strictly positive bilinear form for every $(x, p) \in T^*M$.

We will show below that, for our study, we can always reduce to the case of a superlinear Hamiltonian, without any loss of generality.

Conditions (H2) and (H2') are given in terms of the norm $\|\cdot\|_x$ associated with the Riemannian metric, but they do not actually depend on the particular choice of it for all Riemannian metrics are equivalent on a compact manifold. For $c \in \mathbb{R}$, we will consider the Hamilton-Jacobi equation

$$H(x, d_x u) = c. \tag{3.1}$$

Notice that any given C^1 function $u : M \rightarrow \mathbb{R}$ is a subsolution (resp. supersolution) of $H(x, d_x u) = c$ provided $c \geq \max_{x \in M} H(x, d_x u)$ (resp. $c \leq \min_{x \in M} H(x, d_x u)$). Moreover, the convexity and coercivity of H allow to give the following characterization of viscosity subsolutions of (3.1), see any of the references [2, 3, 13, 24].

Proposition 3.1. *Given a convex coercive Hamiltonian H on the compact manifold M , and $c \in \mathbb{R}$, a function $u : M \rightarrow \mathbb{R}$ is a viscosity subsolution of the Hamilton-Jacobi equation $H(x, d_x u) = c$ if and only if u is Lipschitz, and satisfies $H(x, d_x u) \leq c$, for almost every $x \in M$.*

Moreover, for any fixed $c \in \mathbb{R}$, the set of viscosity subsolutions of $H(x, d_x u) = c$ is equi-Lipschitz with a common Lipschitz constant κ_c given by

$$\kappa_c = \sup\{\|p\|_x \mid H(x, p) \leq c\}. \quad (3.2)$$

We define the *critical value* $c(H)$ as

$$c(H) = \inf\{a \in \mathbb{R} \mid \text{equation (3.1) admits subsolutions}\}. \quad (3.3)$$

By the Ascoli–Arzelà Theorem and the stability of the notion of viscosity subsolution, it is easily seen that such an infimum is attained, meaning that there are subsolutions also at the critical level. Moreover, $c(H)$ is the only real value c for which equation (3.1) admits solutions.

Here and in the sequel, we will assume $c(H) = 0$. This is not restrictive, since we can always reduce to this case by possibly replacing H with $H - c(H)$. We will hence refer to

$$H(x, d_x u) = 0 \quad (3.4)$$

as the *critical equation*. Correspondingly, solutions, subsolutions and supersolutions to (3.4) will be termed *critical* in the sequel.

We will be also interested in the discounted version of (3.4), that is the equation

$$\lambda u(x) + H(x, d_x u) = 0, \quad (3.5)$$

where $\lambda > 0$. The following holds:

Proposition 3.2. *Let $\lambda \geq 0$. Then any subsolution of (3.5) is Lipschitz–continuous and satisfies*

$$\lambda w(x) + H(x, d_x w) \leq 0, \text{ for a.e. } x \in M. \quad (3.6)$$

Proof. A subsolution w of (3.5) satisfies

$$H(x, d_x w) \leq \|\lambda w\|_\infty \quad \text{in } M$$

in the viscosity sense, hence it is Lipschitz continuous by the coercivity of H , see [3]. In particular, it satisfies the inequality (3.6) at every differentiability point, i.e. almost everywhere by Rademacher’s theorem. \square

The crucial difference between the critical equation (3.4) and the discounted equation (3.5) with $\lambda > 0$ is that the latter satisfies a strong comparison principle. In fact, we have

Theorem 3.3. *Let $\lambda > 0$. If v, u are a sub and a supersolution of (3.5), respectively, then $v \leq u$ in M . Moreover, there exists a unique solution of (3.5).*

This theorem is well known, see for instance [3]. For the reader’s convenience, we propose here a short proof by exploiting an approximation argument that works due to our assumption that H is convex in p .

Proof of Theorem 3.3. We want to prove that $\min_M(u - v) \geq 0$. We introduce the Hamiltonian $\hat{H}_{\lambda v}$ defined by

$$\hat{H}_{\lambda v}(x, p) = \lambda v(x) + H(x, p).$$

This Hamiltonian is continuous, convex, and coercive. Obviously v is a subsolution of the equation

$$\hat{H}_{\lambda v}(x, d_x v) = 0,$$

Therefore applying Theorem A.1, for every integer $n \geq 1$, we can find a C^1 function $w_n : M \rightarrow \mathbb{R}$ such that

$$\hat{H}_{\lambda v}(x, d_x w_n) = \lambda v(x) + H(x, d_x w_n) \leq \frac{1}{n} \text{ for every } x \in M, \quad (3.7)$$

and w_n converges uniformly to v on M . This uniform convergence implies that $\min_M(u - w_n)$ converges to $\min_M(u - v)$ as $n \rightarrow +\infty$. Let x_n be a point in M where $u - w_n$ attains its minimum. Then w_n is a subtangent to u at x_n . Since u is a supersolution of (3.5) we have

$$\lambda u(x_n) + H(x_n, d_{x_n} w_n) \geq 0,$$

and by subtracting inequality (3.7) with $x = x_n$, we end up with

$$-\frac{1}{n} \leq \lambda(u(x_n) - v(x_n)) \leq \lambda \min_M(u - w_n) + \lambda \|w_n - v\|_\infty.$$

By sending n to $+\infty$ we obtain the first assertion of the statement. This first part implies the uniqueness of the solution. The existence part follows by applying Perron's method, see [3], or the end of the proof of Theorem 4.4. \square

Next, we show that the solutions of (3.5) are equi-Lipschitz.

Proposition 3.4. *There exists a constant κ independent of $\lambda > 0$, such that the solution u_λ of (3.5) is Lipschitz with Lipschitz constant κ .*

Proof. We already know that u_λ is Lipschitz. We want to prove that its Lipschitz constant can be chosen independent of λ . Let us set $\beta = \max_{x \in M} |H(x, 0)|$. The function $w \equiv -\beta/\lambda$ is obviously a subsolution of (3.5). By Theorem 3.3, we must have $\lambda u_\lambda(x) \geq -\beta$ for every $x \in M$. Hence, we get

$$H(x, d_x u_\lambda) \leq -\lambda u_\lambda(x) \leq \beta, \text{ for a.e. } x \in M,$$

and u_λ is κ_β -Lipschitz by coercivity of H , with κ_β given by (3.2). This proves the proposition with $\kappa = \kappa_\beta$. \square

Note that $\beta = \max_{x \in M} |H(x, 0)| \geq c(H) = 0$. In fact, we know that there exists a solution $u : M \rightarrow \mathbb{R}$ of equation (3.4). At a minimum x_0 of u the constant function $w \equiv u(x_0)$ is a subtangent therefore $H(x_0, 0) \geq 0 = c(H)$, which implies $\beta \geq 0$. This remark together with Proposition 3.4 above shows that not all of H is relevant in order to study the discounted and critical equations. In particular, we may modify H outside the compact set $\{(x, p) \in T^*M : \|p\|_x \leq \kappa\}$, with $\kappa = \kappa_\beta$, to obtain a new Hamiltonian which is still continuous and convex, and satisfies the stronger growth condition (H2'). Since the κ -sublevel of the two Hamiltonians coincide, the solutions of the corresponding critical and discounted equations are the same.

In the remainder of the paper, without any loss of generality, we will therefore assume that H is superlinear in p , i.e. that satisfies condition (H2').

4. THE DISCOUNTED VALUE FUNCTION

We recall that we are assuming $c(H) = 0$, and also that H satisfies condition (H2'). We will denote by $L : TM \rightarrow \mathbb{R}$ the associated Lagrangian, see section (B.1).

For every $\lambda > 0$, we define the discounted value function $u_\lambda : M \rightarrow \mathbb{R}$ by

$$u_\lambda(x) = \inf_{\gamma} \int_{-\infty}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds, \quad (4.1)$$

where the infimum is taken over all absolutely continuous curves $\gamma :]-\infty, 0] \rightarrow M$, with $\gamma(0) = x$.

In this section we will show that u_λ defined as above is indeed the unique viscosity solution of the discounted Hamilton–Jacobi equation

$$\lambda u_\lambda + H(x, d_x u_\lambda) = 0 \quad \text{in } M. \quad (4.2)$$

First we derive some crucial information about the function u_λ defined by the variational formula (4.1).

Proposition 4.1. *The function u_λ defined by (4.1) satisfies the following properties:*

(i) *For every $\lambda > 0$*

$$\frac{\min_{TM} L}{\lambda} \leq u_\lambda(x) \leq \frac{L(x, 0)}{\lambda} \quad \text{for every } x \in M.$$

In particular, $\|\lambda u_\lambda\|_\infty \leq C_0$ for some positive constant C_0 independent of $\lambda > 0$.

(ii) *For every absolutely continuous curve $\gamma : [a, b] \rightarrow M$, we have*

$$e^{\lambda b} u_\lambda(\gamma(b)) - e^{\lambda a} u_\lambda(\gamma(a)) \leq \int_a^b e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds. \quad (4.3)$$

(iii) *There exists a positive constant κ , independent of $\lambda > 0$, such that*

$$u_\lambda(x) - u_\lambda(y) \leq \kappa d(x, y) \quad \text{for every } x, y \in M \text{ and } \lambda > 0,$$

that is, the functions $\{u_\lambda : \lambda > 0\}$ are equi-Lipschitz.

Proof. In (i), the first inequality comes from the fact that every absolutely continuous curve $\gamma : (-\infty, 0] \rightarrow M$ satisfies

$$\int_{-\infty}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds \geq \left(\min_{TM} L \right) \int_{-\infty}^0 e^{\lambda s} ds = \frac{\min_{TM} L}{\lambda}.$$

The second inequality follows by choosing, as a competitor, the steady curve identically equal to the point x .

To prove (ii), we first note that we can assume $b = 0$, since we can always reduce to this case by replacing γ with the curve $\gamma_{-b}(\cdot) := \gamma(\cdot + b)$ defined on the interval $[a - b, 0]$ and by dividing (4.3) by $e^{\lambda b}$. Note that a change of variables gives

$$\int_a^b e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds = e^{\lambda b} \int_{a-b}^0 e^{\lambda s} L(\gamma_{-b}(s), \dot{\gamma}_{-b}(s)) ds.$$

So, let $\gamma \in \text{AC}([a, 0]; M)$ be fixed. For every absolutely continuous curve $\xi : (-\infty, 0] \rightarrow M$ with $\xi(0) = \gamma(a)$, we define a curve $\xi_a : (-\infty, a] \rightarrow M$ by setting

$\xi_a(\cdot) := \xi(\cdot - a)$ and a curve $\eta := \xi_a \star \gamma : (-\infty, 0] \rightarrow M$ obtained by concatenation of ξ_a and γ . By definition of u_λ and arguing as above we get:

$$\begin{aligned} u_\lambda(\gamma(0)) &\leq \int_{-\infty}^0 e^{\lambda s} L(\eta, \dot{\eta}) ds = \int_{-\infty}^a e^{\lambda s} L(\xi_a, \dot{\xi}_a) ds + \int_a^0 e^{\lambda s} L(\gamma, \dot{\gamma}) ds \\ &= e^{\lambda a} \int_{-\infty}^0 e^{\lambda s} L(\xi, \dot{\xi}) ds + \int_a^0 e^{\lambda s} L(\gamma, \dot{\gamma}) ds. \end{aligned}$$

By minimizing with respect to all $\xi \in \text{AC}((-\infty, 0]; M)$ with $\xi(0) = \gamma(a)$ we get the assertion by definition of $u_\lambda(\gamma(a))$.

To prove (iii), pick $x, y \in M$ and let $\gamma : [-d(x, y), 0] \rightarrow M$ be the geodesic joining y to x parameterized by the arc-length. According to item (ii), we have

$$u_\lambda(x) - u_\lambda(y) \leq (-u_\lambda(y)) \left(1 - e^{-\lambda d(x, y)}\right) + \int_{-d(x, y)}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds.$$

Let $C_1 := \max \{L(z, v) : z \in M, \|v\|_z \leq 1\}$ and C_0 the constant given by item (i). We get

$$u_\lambda(x) - u_\lambda(y) \leq (\| \lambda u_\lambda \|_\infty + C_1) \frac{1 - e^{-\lambda d(x, y)}}{\lambda} \leq (C_0 + C_1) d(x, y),$$

where, for the last inequality, we have used the fact that, by concavity, $1 - e^{-h} \leq h$ for every $h \in \mathbb{R}$. \square

Next, we prove that the discounted value function satisfies the Dynamical Programming Principle.

Proposition 4.2. *Let $\lambda > 0$. For every $x \in M$ and $t > 0$*

$$u_\lambda(x) = \inf_{\gamma(0)=x} \left\{ e^{-\lambda t} u_\lambda(\gamma(-t)) + \int_{-t}^0 e^{\lambda s} L(\gamma, \dot{\gamma}) ds : \gamma \in \text{AC}([-t, 0]; M) \right\}. \quad (4.4)$$

Moreover, the above infimum is attained.

Proof. Fix $x \in M$ and $t > 0$. By part (ii) of Proposition 4.1 we immediately derive that $u_\lambda(x)$ is less or equal than the right-hand side term of (4.4).

Let us prove the opposite inequality. Let $\varepsilon > 0$ and choose an absolutely continuous curve $\gamma : (-\infty, 0] \rightarrow M$ with $\gamma(0) = x$ such that

$$u_\lambda(x) + \varepsilon > \int_{-\infty}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds.$$

Then

$$u_\lambda(x) + \varepsilon > \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds + e^{-\lambda t} \int_{-\infty}^{-t} e^{\lambda(s+t)} L(\gamma(s), \dot{\gamma}(s)) ds.$$

We now make a change of variables in the second integral. By setting $\xi(\cdot) := \gamma(\cdot - t)$ and by exploiting the definition of u_λ we end up with

$$\begin{aligned} u_\lambda(x) + \varepsilon &> \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds + e^{-\lambda t} \int_{-\infty}^0 e^{\lambda s} L(\xi(s), \dot{\xi}(s)) ds \\ &\geq \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds + e^{-\lambda t} u_\lambda(\gamma(-t)). \end{aligned}$$

The desired inequality follows as $\varepsilon > 0$ was arbitrarily chosen.

To prove the last assertion, we take minimizing sequence $\gamma_n : [-t, 0] \rightarrow M$ with $\gamma_n(0) = x$, i.e. such that

$$\lim_{n \rightarrow +\infty} e^{-\lambda t} u_\lambda(\gamma_n(-t)) + \int_{-t}^0 e^{\lambda s} L(\gamma_n(s), \dot{\gamma}_n(s)) ds = u_\lambda(x).$$

For n large enough, we have:

$$\int_{-t}^0 e^{\lambda s} L(\gamma_n(s), \dot{\gamma}_n(s)) ds \leq 1 + u_\lambda(\gamma_n(0)) - e^{-\lambda t} u_\lambda(\gamma_n(-t)) \leq 1 + 2\|u_\lambda\|_\infty.$$

According to Theorem B.2, the curves γ_n uniformly converge, up to subsequences, to an absolutely continuous curve $\gamma : [-t, 0] \rightarrow M$ with $\gamma(0) = x$ and satisfying

$$\int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds \leq \liminf_{n \rightarrow +\infty} \int_{-t}^0 e^{\lambda s} L(\gamma_n(s), \dot{\gamma}_n(s)) ds.$$

This readily implies that γ is a minimizer of (4.4). \square

We proceed to show that u_λ is the unique solution of the discounted Hamilton-Jacobi equation (4.2). We prove a preliminary result first.

Proposition 4.3. *Let $u \in C(M)$. Then u is a subsolution of (4.2) if and only if*

$$u(\gamma(0)) - e^{-\lambda t} u(\gamma(-t)) \leq \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds \quad (4.5)$$

for every curve $\gamma \in \text{AC}([-t, 0]; M)$ and every $t > 0$.

Proof. Let us first assume that u is a subsolution of (4.2). By Proposition 3.2, we know that u is Lipschitz continuous. Like in the proof of Theorem 3.3, applying Theorem A.1, we can find a sequence of C^1 functions $u_n : M \rightarrow \mathbb{R}$ such that $\|u_n - u\|_\infty \leq 1/n$ and

$$\lambda u(x) + H(x, d_x u_n) \leq \frac{1}{n}, \text{ for every } x \in M. \quad (4.6)$$

Therefore

$$\lambda u_n(x) + H(x, d_x u_n) \leq \frac{1 + \lambda}{n}, \text{ for every } x \in M. \quad (4.7)$$

Pick a curve $\gamma \in \text{AC}([-t, 0]; M)$. By exploiting the Fenchel inequality (B.2) of appendix B together with (4.7), we get

$$\begin{aligned} u_n(\gamma(0)) - e^{-\lambda t} u_n(\gamma(-t)) &= \int_{-t}^0 \frac{d}{ds} \left(e^{\lambda s} u_n(\gamma(s)) \right) ds \\ &= \int_{-t}^0 e^{\lambda s} \left(\lambda u_n(\gamma(s)) + \langle d_{\gamma(s)} u_n, \dot{\gamma}(s) \rangle \right) ds \\ &\leq \int_{-t}^0 e^{\lambda s} \left(\lambda u_n(\gamma(s)) + H(\gamma(s), d_{\gamma(s)} u_n) + L(\gamma(s), \dot{\gamma}(s)) \right) ds \\ &\leq \frac{1 + \lambda}{n} \int_{-t}^0 e^{\lambda s} ds + \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds \end{aligned}$$

with the last inequality obtained from (4.7). The assertion follows by sending n to $+\infty$.

Conversely, let us assume that (4.5) holds for any absolutely continuous curve. Arguing as in the proof of part (iii) of Proposition 4.1 we see that u is Lipschitz continuous. Let x be a differentiability point of u . Pick a vector $v \in T_x M$ and let

$\gamma \in C^1([-1, 0]; M)$ such that $(\gamma(0), \dot{\gamma}(0)) = (x, v)$. Then, for every $t \in (0, 1)$, we have

$$\frac{u(\gamma(0)) - e^{-\lambda t}u(\gamma(-t))}{t} \leq \frac{1}{t} \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds,$$

hence, letting $t \rightarrow 0$,

$$\lambda u(x) + \langle d_x u, v \rangle - L(x, v) \leq 0.$$

By taking the supremum of this inequality with respect to $v \in T_x M$ we conclude by Proposition B.1 that

$$\lambda u(x) + H(x, d_x u) \leq 0$$

for every differentiability point x of u , i.e. u is a subsolution of (4.2) by Proposition 2.1. \square

We are now ready to prove the announced result:

Theorem 4.4. *For every $\lambda > 0$, the discounted value function u_λ is the unique continuous viscosity solution of (4.2).*

Proof. We already know that u_λ is Lipschitz continuous on M . According to Propositions 4.1–(ii) and 4.3 we derive that u_λ is a subsolution of (4.2).

In order to prove that u_λ is a supersolution, we will show that it is maximal in the family of continuous subsolution of (4.2). Indeed, let $w \in C(M)$ be a subsolution of (4.2) and pick a point x in M . Fix $t > 0$ and let $\gamma : [-t, 0] \rightarrow M$ be a minimizer of (4.4). According to Proposition 4.3 we have

$$\begin{aligned} u_\lambda(x) &= e^{-\lambda t} u_\lambda(\gamma(-t)) + \int_{-t}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds \\ &\geq w(x) + e^{-\lambda t} (u_\lambda(\gamma(-t)) - w(\gamma(-t))) \geq w(x) - e^{-\lambda t} (\|u_\lambda\|_\infty + \|w\|_\infty). \end{aligned}$$

Sending $t \rightarrow +\infty$ we get $u_\lambda(x) \geq w(x)$, as it was claimed.

A standard argument now implies that u_λ is a supersolution. We will sketch it here for the reader's convenience. Were u_λ not a supersolution, there would exist a strict subgradient ϕ to u_λ at some point $y \in M$ with $\phi(y) = u_\lambda(y)$ such that

$$\lambda \phi(y) + H(y, d_y \phi) < 0.$$

Then we can find $r > 0$ and $\varepsilon > 0$ small enough so that the function w defined as

$$w = \max\{\phi + \varepsilon, u_\lambda\} \quad \text{in } B_r(y) \quad \text{and} \quad w = u_\lambda \quad \text{elsewhere}$$

is still a subsolution of (4.2). But this is in contradiction with the maximality of u_λ since $w(y) > u_\lambda(y)$ by construction.

The uniqueness part comes from Theorem 3.3. \square

As a consequence, we derive the following

Corollary 4.5. *The functions $\{u_\lambda : \lambda > 0\}$ are equi-Lipschitz and equi-bounded. In particular, $\|\lambda u_\lambda\|_\infty \rightarrow 0$ as $\lambda \rightarrow 0$.*

Proof. The equi-Lipschitz character of the functions u_λ has been already proved in Proposition 4.1. To see they are equi-bounded, take a solution u of (3.4)—recall that we are assuming $c(H) = 0$. By addition of suitable constants, we obtain two critical solutions \underline{u}, \bar{u} of equation (3.4) such that $\underline{u} \leq 0 \leq \bar{u}$ in M . It is easily seen that,

for every fixed $\lambda > 0$, \underline{u} and \bar{u} are, respectively, a sub and a supersolution of (3.5). By the comparison principle stated in Theorem 3.3 we derive

$$\underline{u} \leq u_\lambda \leq \bar{u} \quad \text{in } M \quad \text{for every } \lambda > 0,$$

as it was to be shown. \square

Here is now a first obvious case of convergence.

Proposition 4.6. *Suppose that the constants are critical subsolutions, or equivalently that $L + c(H) \geq 0$, then $u_\lambda \geq 0$, and $u_\lambda \nearrow$ uniformly as $\lambda \searrow 0$ to some solution of the critical equation $H(x, d_x u) = c(H)$.*

Proof. Replacing L by $L + c(H)$, we can assume $c(H) = 0$. By Corollary 4.5, the family $(u_\lambda)_{\lambda > 0}$ is equi-continuous and bounded. Moreover, any uniform accumulation point of u_λ is a solution of equation (3.4). Therefore it suffices to show that $u_\lambda \geq 0$, and $u_\lambda \nearrow$, as $\lambda \searrow 0$. Since $L \geq 0$, for $\lambda' \geq \lambda$, and $\gamma :] - \infty, 0] \rightarrow M$, we have

$$0 \leq \int_{-\infty}^0 e^{\lambda' s} L(\gamma(s), \dot{\gamma}(s)) ds \leq \int_{-\infty}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds.$$

taking the infimum over all $\gamma :] - \infty, 0] \rightarrow M$, with $\gamma(0) = x$ yields $0 \leq u_{\lambda'}(x) \leq u_\lambda(x)$. \square

We end this section by proving the existence of Lipschitz curves that realize the infimum in the definition of discounted value function.

Proposition 4.7. *Let $\lambda > 0$ and $x \in M$. Then there exists a curve $\gamma_x^\lambda : (-\infty, 0] \rightarrow M$ with $\gamma_x^\lambda(0) = x$ such that*

$$u_\lambda(x) = e^{-\lambda t} u_\lambda(\gamma_x^\lambda(-t)) + \int_{-t}^0 e^{\lambda s} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds \quad \text{for every } t > 0. \quad (4.8)$$

Moreover, there exists a constant $\alpha > 0$, independent of λ and x , such that $\|\dot{\gamma}_x^\lambda\|_\infty \leq \alpha$. In particular

$$u_\lambda(x) = \int_{-\infty}^0 e^{\lambda s} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds. \quad (4.9)$$

Proof. According to Proposition 4.2 we know that, for every $n \in \mathbb{N}$, there exists a curve $\xi_n : [-n, 0] \rightarrow M$ with $\xi_n(0) = x$ such that

$$u_\lambda(x) = e^{-\lambda n} u_\lambda(\xi_n(-n)) + \int_{-n}^0 e^{\lambda s} L(\xi_n(s), \dot{\xi}_n(s)) ds. \quad (4.10)$$

We claim that, for every $[a, b] \subset [-n, 0]$,

$$e^{\lambda b} u_\lambda(\xi_n(b)) - e^{\lambda a} u_\lambda(\xi_n(a)) = \int_a^b e^{\lambda s} L(\xi_n(s), \dot{\xi}_n(s)) ds. \quad (4.11)$$

Indeed, by taking into account Proposition 4.1–(ii) we have

$$\begin{aligned} \int_{-n}^0 e^{\lambda s} L(\xi_n, \dot{\xi}_n) ds &= \int_b^0 e^{\lambda s} L(\xi_n, \dot{\xi}_n) ds + \int_a^b e^{\lambda s} L(\xi_n, \dot{\xi}_n) ds + \int_{-n}^a e^{\lambda s} L(\xi_n, \dot{\xi}_n) ds \\ &\geq \left(u_\lambda(x) - e^{\lambda b} u_\lambda(\xi_n(b)) \right) + \left(e^{\lambda b} u_\lambda(\xi_n(b)) - e^{\lambda a} u_\lambda(\xi_n(a)) \right) \\ &\quad + \left(e^{\lambda a} u_\lambda(\xi_n(a)) - e^{-\lambda n} u_\lambda(\xi_n(-n)) \right) = u_\lambda(x) - e^{-\lambda n} u_\lambda(\xi_n(-n)). \end{aligned}$$

Now we remark that the first term in the above inequality is equal to the last one, according to (4.10), hence all inequalities must be equalities. This proves (4.11). By reasoning as in the proof of Proposition 4.2 and using a diagonal argument, we derive from Theorem B.2 that there exists an absolutely continuous curve $\gamma_x^\lambda : (-\infty, 0] \rightarrow M$ with $\gamma_x^\lambda(0) = x$ which is, up to extraction of a subsequence, the uniform limit of the curves ξ_n over compact subsets of $(-\infty, 0]$. Such curve satisfies

$$e^{\lambda b} u_\lambda(\gamma_x^\lambda(b)) - e^{\lambda a} u_\lambda(\gamma_x^\lambda(a)) = \int_a^b e^{\lambda s} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds. \quad (4.12)$$

for every $[a, b] \subset (-\infty, 0]$. To see this, it suffices to pass to the limit in (4.11). The equality holds also for the limit curve γ_x^λ by the lower semicontinuity of the functional \mathbb{L}^λ stated in Theorem B.2 and by Proposition 4.1-(ii). In particular, this proves assertion (4.8).

The fact that the curves γ_x^λ are equi-Lipschitz is a consequence of the fact that the functions u_λ are equi-Lipschitz, say κ -Lipschitz, according to Proposition 4.1. Indeed, by superlinearity of L , there exists a constant A_κ , depending on κ , such that

$$L(x, v) \geq (\kappa + 1)\|v\|_x - A_\kappa \quad \text{for every } (x, v) \in TM.$$

For every $a \in (-\infty, 0)$ and $h > 0$ small enough we get, from (4.12),

$$\begin{aligned} e^{\lambda(a+h)} u_\lambda(\gamma_x^\lambda(a+h)) - e^{\lambda a} u_\lambda(\gamma_x^\lambda(a)) &= \int_a^{a+h} e^{\lambda s} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds \\ &\geq e^{\lambda a} (\kappa + 1) \int_a^{a+h} \|\dot{\gamma}_x^\lambda(s)\|_{\gamma_x^\lambda(s)} ds - A_\kappa \int_a^{a+h} e^{\lambda s} ds \\ &\geq e^{\lambda a} \left(\kappa d(\gamma_x^\lambda(a), \gamma_x^\lambda(a+h)) + \int_a^{a+h} \|\dot{\gamma}_x^\lambda(s)\|_{\gamma_x^\lambda(s)} ds - A_\kappa \frac{e^{\lambda h} - 1}{\lambda} \right) \end{aligned} \quad (4.13)$$

On the other hand

$$\begin{aligned} e^{\lambda(a+h)} u_\lambda(\gamma_x^\lambda(a+h)) - e^{\lambda a} u_\lambda(\gamma_x^\lambda(a)) &\leq (e^{\lambda(a+h)} - e^{\lambda a}) u_\lambda(\gamma_x^\lambda(a+h)) + e^{\lambda a} \kappa d(\gamma_x^\lambda(a), \gamma_x^\lambda(a+h)) \\ &\leq e^{\lambda a} \left(C_0 \frac{e^{\lambda h} - 1}{\lambda} + \kappa d(\gamma_x^\lambda(a), \gamma_x^\lambda(a+h)) \right), \end{aligned} \quad (4.14)$$

where C_0 is the constant given by Proposition 4.1-(i). Plugging (4.14) into (4.13) and dividing by $h e^{\lambda a}$ we end up with

$$\frac{1}{h} \int_a^{a+h} \|\dot{\gamma}_x^\lambda(s)\|_{\gamma_x^\lambda(s)} ds \leq (A_\kappa + C_0) \frac{e^{\lambda h} - 1}{\lambda h}. \quad (4.15)$$

Sending $h \rightarrow 0$ we infer

$$\|\dot{\gamma}_x^\lambda(a)\|_{\gamma_x^\lambda(a)} \leq \alpha := (A_\kappa + C_0) \quad \text{for a.e. } a \in (-\infty, 0],$$

as it was to be shown. In particular, by sending $t \rightarrow +\infty$ in (4.8) we get (4.9) by the Dominated Convergence Theorem. \square

5. CONVERGENCE OF THE DISCOUNTED VALUE FUNCTIONS

In this section we will prove our main theorem, namely that the discounted value functions u_λ converge, as $\lambda \rightarrow 0$, to a particular solution u_0 of the critical equation (3.4).

To define u_0 , we consider the family \mathcal{F}_- of subsolutions $u : M \rightarrow \mathbb{R}$ of the critical equation (3.4) satisfying the following condition

$$\int_M u d\mu \leq 0 \text{ for every projected Mather measure } \mu. \quad (5.1)$$

For the concept of Mather measure see Definition B.21 in §B.4 below.

Note that, given any critical subsolution u , the function $u - \|u\|_\infty$ is in \mathcal{F}_- . Therefore \mathcal{F}_- is not empty.

Lemma 5.1. *The family \mathcal{F}_- is uniformly bounded from above, i.e.*

$$\sup\{u(x) \mid x \in M, u \in \mathcal{F}_-\} < +\infty.$$

Proof. The family of critical subsolutions is equi-Lipschitz. Call κ a common Lipschitz constant. Since the set of projected Mather measure μ is not empty, picking such a probability measure μ , for $u \in \mathcal{F}_-$, we have $\min u = \int_M \min u d\mu \leq \int_M u d\mu \leq 0$. Hence $\max u \leq \max u - \min u$. Since u is κ -Lipschitz, we also $\max u - \min u \leq \kappa \text{diam}(M) < +\infty$. \square

Therefore we can define $u_0 : M \rightarrow \mathbb{R}$ by

$$u_0 = \sup_{\mathcal{F}_-} u.$$

As the supremum of a family of viscosity subsolutions, we know that u_0 is itself a critical subsolution. We will obtain later that u_0 is a solution, see Theorem 5.6 below.

We now start to study the asymptotic behavior of the discounted value functions u_λ as $\lambda \rightarrow 0$ and the relation with u_0 . We will use the set $\mathfrak{M}_0(L)$ of Mather measures on TM , and the set $\mathfrak{M}_0(L)$ of projected Mather measures on M , see section B.4 for definition and properties. We begin with the following result:

Proposition 5.2. *Let $\lambda > 0$. Then, for every $\mu \in \mathfrak{M}_0(L)$, we have*

$$\int_M u_\lambda(x) d\mu(x) \leq 0.$$

In particular, if the functions u_{λ_n} uniformly converge to u for some sequence $\lambda_n \rightarrow 0$, then $u \leq u_0$ on M .

Proof. Like in the proof of Theorem 3.3, there exists a sequence $(w_n)_n$ of functions in $C^1(M)$ such that $\|u_\lambda - w_n\|_\infty \leq 1/n$ and

$$\lambda u_\lambda(x) + H(x, d_x w_n) \leq 1/n, \text{ for every } x \in M.$$

By the Fenchel inequality

$$L(x, v) + H(x, d_x w_n) \geq d_x w_n(v), \text{ for every } (x, v) \in TM,$$

Combining these two inequalities, yields

$$\lambda u_\lambda(x) + d_x w_n(v) \leq L(x, v) + \frac{1}{n}, \text{ for every } (x, v) \in TM. \quad (5.2)$$

Let us fix some $\tilde{\mu} \in \tilde{\mathfrak{M}}_0(L)$, and set $\mu = \pi_{\#}\tilde{\mu} \in \mathfrak{M}_0(L)$. Since μ is closed and minimizing, we have $\int_{TM} d_x w_n(v) d\tilde{\mu}(x, v) = 0$, and $\int_{TM} L(x, v) d\tilde{\mu}(x, v) = 0$. Therefore if we integrate (5.2), we obtain

$$\lambda \int_M u_\lambda(x) d\mu(x) \leq \frac{1}{n}.$$

Since $\lambda > 0$, letting $n \rightarrow \infty$, yields $\int_M u_\lambda(x) d\mu(x) \leq 0$. If u is the uniform limit of $(u_{\lambda_n})_n$ for some $\lambda_n \rightarrow 0$, we know that it is a solution of the critical equation (3.4). Moreover, it also has to satisfy $\int_M u(x) d\mu(x) \leq 0$ for every projected Mather measure μ . Therefore $u \in \mathcal{F}_-$ and $u \leq u_0$. \square

The next (and final) step is to show that $u \geq u_0$ in M whenever u is the uniform limit of $(u_{\lambda_n})_n$ for some $\lambda_n \rightarrow 0$. We need to introduce some tools and to prove some preliminary results first.

We will use for this the following way to construct closed measure. Suppose $\gamma :]-\infty, 0] \rightarrow M$ is a Lipschitz curve, and $\lambda > 0$. We define the measure $\tilde{\mu}_\gamma^\lambda$ on TM by

$$\begin{aligned} \int_{TM} f(x, v) d\tilde{\mu}_\gamma^\lambda &:= \int_{-\infty}^0 \frac{d}{ds} (e^{\lambda s}) f(\gamma(s), \dot{\gamma}(s)) ds \\ &= \lambda \int_{-\infty}^0 e^{\lambda s} f(\gamma(s), \dot{\gamma}(s)) ds, \end{aligned}$$

for every $f \in C_c(TM)$. It is not difficult to see that μ_γ^λ is a probability measure whose support is contained in the closure of

$$\{(\gamma(s), \dot{\gamma}(s)) \mid s \in]-\infty, 0] \text{ where } \gamma(s) \text{ is differentiable}\}$$

which is compact because it is contained in $\{(x, v) \in TM \mid \|v\|_x \leq \kappa\}$, where κ is a Lipschitz constant for γ .

Therefore if $\gamma_\lambda :]-\infty, 0] \rightarrow M, \lambda > 0$ is a family of equi-Lipschitz curves, the family of probability measures $\tilde{\mu}_{\gamma_\lambda}^\lambda, \lambda > 0$ is relatively compact in the weak topology on measures. Therefore for any sequence $\lambda_n \rightarrow 0$, we can extract a subsequence of $\tilde{\mu}_{\gamma_{\lambda_n}}^{\lambda_n}$ converging to a probability measure $\tilde{\mu}$ on TM . We now show that this measure is necessarily closed.

Proposition 5.3. *Suppose $\gamma_\lambda :]-\infty, 0] \rightarrow M, \lambda > 0$, is a family of equi-Lipschitz curves. If the measure $\tilde{\mu}$ on TM is the weak limit of $\tilde{\mu}_{\gamma_{\lambda_n}}^{\lambda_n}$ for some $\lambda_n \rightarrow 0$ then $\tilde{\mu}$ is closed.*

Proof. Call κ a common Lipschitz constant for the family of curves γ_λ , then both $\tilde{\mu}$, and the measures $\tilde{\mu}_{\gamma_{\lambda_n}}^{\lambda_n}$ have all support in the compact set $\{(x, v) \in TM \mid \|v\|_x \leq \kappa\}$. Therefore $\tilde{\mu}$ is a probability measure and $\int_{TM} \|v\|_x d\tilde{\mu}(x, v) \leq \kappa < +\infty$. Moreover, if $\varphi : M \rightarrow \mathbb{R}$ is C^1 , then the function $s \mapsto e^{\lambda s} \varphi(\gamma_\lambda(s))$ is Lipschitz on $]-\infty, 0]$ with derivative

$$s \mapsto \lambda e^{\lambda s} \varphi(\gamma_\lambda(s)) + e^{\lambda s} d_{\gamma_\lambda(s)} \varphi(\dot{\gamma}_\lambda(s)).$$

Hence

$$\varphi(\gamma_\lambda(0)) = \int_{-\infty}^0 e^{\lambda s} d_{\gamma_\lambda(s)} \varphi(\dot{\gamma}_\lambda(s)) ds + \int_{-\infty}^0 \lambda e^{\lambda s} \varphi(\gamma_\lambda(s)) ds.$$

Note that the left hand side is bounded by $\|\varphi\|_\infty$, and also

$$\left| \int_{-\infty}^0 \lambda e^{\lambda s} \varphi(\gamma_\lambda(s)) ds \right| \leq \int_{-\infty}^0 \lambda e^{\lambda s} \|\varphi\|_\infty ds \leq \|\varphi\|_\infty.$$

It follows that

$$\left| \int_{TM} d_x \varphi(v) d\tilde{\mu}_{\gamma_{\lambda_n}}^{\lambda_n} \right| = \left| \lambda_n \int_{-\infty}^0 e^{\lambda_n s} d_{\gamma_{\lambda_n}(s)} \varphi(\dot{\gamma}_{\lambda_n}(s)) ds \right| \leq 2\lambda_n \|\varphi\|_\infty \rightarrow 0,$$

as $\lambda_n \rightarrow 0$. Therefore, we obtain $\int_{TM} d_x \varphi(v) d\tilde{\mu} = \lim_{n \rightarrow +\infty} \int_{TM} d_x \varphi(v) d\tilde{\mu}_{\gamma_{\lambda_n}}^{\lambda_n} = 0$. \square

Suppose now that $x \in M$ is given. By Proposition 4.7, for every $\lambda > 0$, we can choose as $\gamma_x^\lambda : (-\infty, 0] \rightarrow M$ a Lipschitz curve satisfying $\gamma_x^\lambda(0) = x$, and

$$u_\lambda(x) = \int_{-\infty}^0 e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds.$$

By Proposition 4.7, we know that the family γ_x^λ is equi-Lipschitz. Therefore, we define $\tilde{\mu}_x^\lambda = \tilde{\mu}_{\gamma_x^\lambda}^\lambda$ by

$$\begin{aligned} \int_{TM} f(x, v) d\tilde{\mu}_x^\lambda &= \int_{-\infty}^0 \frac{d}{ds} (e^{\lambda s}) f(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds \\ &= \lambda \int_{-\infty}^0 e^{\lambda s} f(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds, \end{aligned}$$

for every $f \in C_c(TM)$.

Lemma 5.4. *The measures $\tilde{\mu}_x^\lambda, \lambda > 0$ defined above are all probability measures, whose supports are all contained in a common compact subset of TM . In particular, for any sequence $\lambda_n \rightarrow 0$, extracting a subsequence if necessary, we can assume that $\tilde{\mu}_x^{\lambda_n}$ converges to a probability measure on TM . This measure is a (closed) Mather measure.*

Proof. From what we have done above we know that the family $\tilde{\mu}_x^\lambda, \lambda > 0$ is weakly compact and any weak limit point is a closed measure. It remains to show with the notations of the lemma that $\int_{TM} L d\tilde{\mu} = 0$. We have

$$\begin{aligned} \int_{TM} L d\tilde{\mu} &= \lim_{n \rightarrow \infty} \int_{TM} L d\tilde{\mu}_x^{\lambda_n} \\ &= \lim_{n \rightarrow \infty} \int_{-\infty}^0 \frac{d}{ds} (e^{\lambda_n s}) L(\gamma_x^{\lambda_n}(s), \dot{\gamma}_x^{\lambda_n}(s)) ds \\ &= \lim_{n \rightarrow \infty} \lambda_n u_{\lambda_n}(x) \\ &= 0, \end{aligned}$$

where the last equality follows from the fact that $\lambda u_\lambda \rightarrow 0$, see Corollary 4.5 \square

The following lemma will be crucial in the proof of Proposition 1.2.

Lemma 5.5. *Let w be any critical subsolution. For every $\lambda > 0$ and $x \in M$*

$$u_\lambda(x) \geq w(x) - \int_{TM} w(y) d\tilde{\mu}_x^\lambda(y, v). \quad (5.3)$$

Proof. Let $\varepsilon > 0$. According to Theorem A.1, there exists a smooth function w_ε such that

$$\|w - w_\varepsilon\|_\infty < \varepsilon \text{ and } H(x, d_x w_\varepsilon) < \varepsilon, \text{ for every } x \in M.$$

By the Fenchel inequality, we have

$$\begin{aligned} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) &\geq d_{\gamma_x^\lambda(s)} w_\varepsilon(\dot{\gamma}_x^\lambda(s)) - H(\gamma_x^\lambda(s), d_{\gamma_x^\lambda(s)} w_\varepsilon) \\ &\geq d_{\gamma_x^\lambda(s)} w_\varepsilon(\dot{\gamma}_x^\lambda(s)) - \varepsilon \end{aligned}$$

for every $s < 0$. Using the definition of the curve γ_x^λ , see Proposition 4.7, we get

$$\begin{aligned} u_\lambda(x) &= e^{-\lambda t} u_\lambda(\gamma_x^\lambda(-t)) + \int_{-t}^0 e^{\lambda s} L(\gamma_x^\lambda(s), \dot{\gamma}_x^\lambda(s)) ds \\ &\geq e^{-\lambda t} u_\lambda(\gamma_x^\lambda(-t)) + \int_{-t}^0 e^{\lambda s} d_{\gamma_x^\lambda(s)} w_\varepsilon(\dot{\gamma}_x^\lambda(s)) ds - \varepsilon \int_{-t}^0 e^{\lambda s} ds \\ &= w_\varepsilon(x) - \int_{-t}^0 \frac{d}{ds} (e^{\lambda s}) w_\varepsilon(\gamma_x^\lambda(s)) ds \\ &\quad + e^{-\lambda t} \left(u_\lambda(\gamma_x^\lambda(-t)) - w_\varepsilon(\gamma_x^\lambda(-t)) \right) - \frac{\varepsilon}{\lambda} (1 - e^{-\lambda t}), \end{aligned}$$

where, for the last equality, we have used an integration by parts and the fact that $d_{\gamma_x^\lambda(s)} w_\varepsilon(\dot{\gamma}_x^\lambda(s)) = \frac{d}{ds} w_\varepsilon(\gamma_x^\lambda(s))$. Sending now $t \rightarrow +\infty$ we infer

$$u_\lambda(x) \geq w_\varepsilon(x) - \int_{-t}^0 \frac{d}{ds} (e^{\lambda s}) w_\varepsilon(\gamma_x^\lambda(s)) ds - \frac{\varepsilon}{\lambda} = w_\varepsilon(x) - \int_{TM} w_\varepsilon(y) d\tilde{\mu}_x^\lambda(y, v) - \frac{\varepsilon}{\lambda}.$$

The assertion follows by letting $\varepsilon \rightarrow 0$. \square

We are now ready to prove our main theorem:

Theorem 5.6. *The functions u_λ uniformly converge to u_0 on M as $\lambda \rightarrow 0$. In particular, as an accumulation point of u_λ , as $\lambda \rightarrow 0$, the function u_0 is a viscosity solution of (3.4).*

Proof. By Corollary 4.5 we know that the functions u_λ are equi-Lipschitz and equi-bounded, hence it is enough, by the Ascoli–Arzelà theorem, to prove that any converging subsequence has u_0 as limit.

Let $\lambda_n \rightarrow 0$ be such that u_{λ_n} uniformly converge to some $u \in C(M, \mathbb{R})$. We have seen in Proposition 5.2 that

$$u(x) \leq u_0(x) \quad \text{for every } x \in M.$$

To prove the opposite inequality, let us fix $x \in M$. Let w be a critical subsolution. By Proposition 5.5, we have

$$u_{\lambda_n}(x) \geq w(x) - \int_{TM} w(y) d\tilde{\mu}_x^{\lambda_n}(y, v).$$

By Lemma 5.4, extracting a further subsequence, we can assume that $\tilde{\mu}_x^{\lambda_n}$ converges weakly to a Mather measure $\tilde{\mu}$ whose projection on M is denoted by μ . Passing to the limit in the last inequality, we get

$$u(x) \geq w(x) - \int_{TM} w(y) d\mu(y),$$

where μ is a projected Mather measure. If we furthermore assume that $w \in \mathcal{F}_-$, the set of subsolutions satisfying (5.1), we obtain $\int_{TM} w(y) d\mu(y) \leq 0$, and $u \geq w$. Hence $u \geq u_0 = \sup_{w \in \mathcal{F}_-} w$. \square

6. ANOTHER FORMULA FOR THE LIMIT OF THE DISCOUNTED VALUE FUNCTIONS

In this section, we will give a characterization of u_0 as an infimum, using the Peierls barrier h , and the projected Mather measures. We define $\hat{u}_0 : M \rightarrow \mathbb{R}$ by

$$\hat{u}_0(x) = \min_{\mu \in \mathfrak{M}_0(L)} \int_{TM} h(y, x) d\mu(y), \text{ for every } x \in M, \quad (6.1)$$

where $\mathfrak{M}_0(L)$ is the set of projected Mather measures, see Definition B.21. We establish some properties of \hat{u}_0 .

Lemma 6.1. *The function \hat{u}_0 is a critical subsolution.*

Proof. We first remark that $\hat{u}_0 \geq \min_{M \times M} h > -\infty$, where the last strict inequality comes from the continuity of h . We then observe that the function $h_\mu : M \rightarrow \mathbb{R}, x \mapsto \int_{TM} h(y, x) d\mu(y)$ is a convex combination of the family of critical solutions $(h_y)_{y \in M}$, where $h_y(x) = h(y, x)$. By the convexity of H in the momentum p , it follows that each h_μ is a critical subsolution of (3.4). Again due to the convexity of H in the momentum p , a finite valued infimum of critical subsolutions is itself a critical subsolution. Therefore \hat{u}_0 is a critical subsolution. \square

Lemma 6.2. *We have $u_0 \leq \hat{u}_0$ everywhere on M .*

Proof. By the definitions of u_0 , and \hat{u}_0 , it suffices to show that $u \leq h_\mu$, for every critical subsolution u satisfying $\int_M u d\mu \leq 0$, where μ is a projected Mather measure on M . In fact, by Proposition B.4, we have

$$u(x) \leq u(y) + h(y, x),$$

for every $x, y \in M$. If we integrate with respect to y , we get $u(x) \leq \int_M u d\mu + h_\mu(x)$. But $\int_M u d\mu \leq 0$ by assumption. \square

Theorem 6.3. *We have $u_0 = \hat{u}_0$ everywhere on M . In particular, the function \hat{u}_0 is a critical solution.*

Proof. Since, by Lemma 6.2, we already know that $u_0 \leq \hat{u}_0$, we have to show the reverse inequality $u_0 \geq \hat{u}_0$. By Lemma 6.1, the function \hat{u}_0 is a subsolution of the critical Hamilton-Jacobi equation (3.4). Moreover, the function u_0 is a solution of (3.4). Therefore by Theorem B.6, it suffices to show that $\hat{u}_0 \leq u_0$ for every x in the projected Aubry set \mathcal{A} . Fix $x \in \mathcal{A}$, by part e) of Proposition B.4, the function $y \mapsto -h(y, x)$ is a critical subsolution. Hence the function $y \mapsto w(y) = -h(y, x) + \inf_{\mu \in \mathfrak{M}_0(L)} \int_M h(z, x) d\mu(z)$ is also a critical subsolution which satisfies condition (5.1). This implies $u_0 \geq w$ everywhere. In particular

$$u_0(x) \geq -h(x, x) + \inf_{\mu \in \mathfrak{M}_0(L)} \int_M h(z, x) d\mu(z).$$

Using $h(x, x) = 0$ for $x \in \mathcal{A}$, we get $u_0(x) \geq \inf_{\mu \in \mathfrak{M}_0(L)} \int_M h(y, x) d\mu(y) = \hat{u}_0(x)$. \square

We conclude this section with the case $L + c(H) \geq 0$.

Proposition 6.4. *Suppose that the constants are critical subsolutions, or equivalently that $L + c(H) \geq 0$, then the projected Mather set \mathcal{M} and the projected Aubry set \mathcal{A} are both equal to $\{x \in M \mid L(x, 0) + c(H) = 0\}$, and for every $x \in M$, we have $u_0(x) = \min\{h(y, x) \mid L(y, 0) + c(H) = 0\}$.*

Proof. Replacing L by $L + c(H)$, we can assume that $c(H) = 0$. Since the constant functions are C^1 critical subsolutions, by Lemma B.12, we have $\mathcal{A} \subset \{x \in M \mid L(x, 0) = 0\}$. Next we remark that the Dirac mass $\tilde{\delta}_{(x,0)}$ at $(x, 0)$ is a closed measure on TM for any $x \in M$. If $L(x, 0) = 0$, we have $\int_{TM} L d\tilde{\delta}_{(x,0)} = 0$. Therefore $\delta_{x,0}$ is a Mather minimizing measure. It follows that $\{x \in M \mid L(x, 0) = 0\} \subset \mathcal{M}$. Since $\mathcal{M} \subset \mathcal{A}$, we obtain $\mathcal{M} = \mathcal{A} = \{x \in M \mid L(x, 0) = 0\}$. If we set $u_1(x) = \min\{h(y, x) \mid L(y, 0) = 0\}$, since the Dirac mass at any y with $L(y, 0) = 0$ is a projected Mather measure, we obtain $u_0 = \hat{u}_0 \leq u_1$. On the other hand since the support of a Mather minimizing measure μ is contained in $\{y \in M \mid L(y, 0) = 0\}$, we get $\int_M h(y, x) d\mu(y) \geq \int_M u_1(x) d\mu(y) = u_1(x)$. Therefore $u_1 = u_0$. \square

APPENDIX A. CLARKE CALCULUS AND APPROXIMATION OF SUBSOLUTIONS

A.1. Clarke calculus. We will suppose in the sequel that N is a connected manifold (not necessarily compact). It will be useful to have a Riemannian metric g on N . For $(x, v) \in TN$, we will denote by $\|v\|_x$ the Riemannian norm $\sqrt{g_x(v, v)}$. For $(x, p) \in T^*N$, we will also denote by $\|p\|_x$ the norm of p obtained by duality from the Riemannian norm on T_xN . We will endow N with the distance d obtained from the Riemannian metric.

We will denote by δ (resp. δ^*) a distance defining the topology of TN (resp. T^*N). Replacing $\delta[(x, v), (x', v')]$ (resp. $\delta^*[(x, p), (x', p')]$), if necessary, by $\delta[(x, v), (x', v')] + d(x, x')$ (resp. $\delta^*[(x, p), (x', p')] + d(x, x')$), we will assume in the sequel that

$$\begin{aligned} \delta[(x, v), (x', v')] &\geq d(x, x'), \text{ for all } (x, v), (x', v') \in TN, \\ \delta^*[(x, p), (x', p')] &\geq d(x, x'), \text{ for all } (x, p), (x', p') \in T^*N. \end{aligned}$$

If $u : N \rightarrow \mathbb{R}$ is locally Lipschitz, recall that u is differentiable almost everywhere, by Rademacher's theorem. We will denote by $\partial^*u(y)$ the set of *reachable derivatives* (also called *reachable gradients*) of u at y , that is the set

$$\partial^*u(y) = \{p \in T_y^*N : (x_n, d_{x_n}u) \rightarrow (y, p) \text{ in } T^*M, u \text{ is differentiable at } x_n\}.$$

The *Clarke's generalized derivative* (or *gradient*) $\partial^c u(y)$ is the closed convex hull of $\partial^*u(y)$ in T_y^*N . The set $\partial^c u(y)$ contains both $D^+u(y)$ and $D^-u(y)$ defined in §2. In particular $d_y u \in \partial^c u(y)$ at any differentiability point y of u . Moreover, the set valued map $x \mapsto (x, \partial^c u(x))$ from N to T^*N is upper semicontinuous with respect to set inclusion. Recall that the upper semicontinuity means that for every open O subset of T^*N , the subset $\{x \in N \mid (x, \partial^c u(x)) \subset O\}$ is open in N . We refer the reader to [8] for a detailed treatment of the subject.

We denote by $\text{Graph}(\partial^c u)$, the graph of the set function $x \mapsto \partial^c u(x)$, i.e.

$$\text{Graph}(\partial^c u) = \{(x, p) \in T^*N \mid p \in \partial^c u(x)\}.$$

By the upper semi-continuity of $x \mapsto \partial^c u(x)$, this set is closed; moreover, for every compact subset $K \in N$, the set $\{(x, p) \in \text{Graph}(\partial^c u) \mid x \in K\}$ is compact.

For such a locally Lipschitz function $u : N \rightarrow \mathbb{R}$, it is convenient to introduce the Clarke gauges $\partial^+u, \partial^-u : TN \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} \partial^+u(x, v) &= \sup\{p(v) \mid p \in \partial^c u(x)\} = \sup\{p(v) \mid p \in \partial^*u(x)\} \\ \partial^-u(x, v) &= \inf\{p(v) \mid p \in \partial^c u(x)\} = \inf\{p(v) \mid p \in \partial^*u(x)\}. \end{aligned}$$

Note that both right equalities in the definitions above follow from the fact that $\partial^c u(x)$ is the convex envelop of $\partial^* u(x)$. Moreover, the sup and the inf in the definitions are attained by the compactness of $\partial^c u(x)$ and $\partial^* u(x)$.

Since $\partial^c(-u)(x) = -\partial^c u(x)$, we have $\partial^- u(x, v) = -\partial^+(-u)(x, v)$ —note that $\partial^+ u(x, v)$ is denoted by $u^\circ(x, v)$ in [8].

Since $x \mapsto \partial^c u(x)$ is upper semi-continuous with compact values, the function $\partial^+ u$ is upper semi-continuous, and the function $\partial^- u$ is lower semi-continuous.

A.2. Approximation of subsolutions by smooth almost subsolutions.

Theorem A.1. *Let $u : N \rightarrow \mathbb{R}$ be a locally Lipschitz function. Assume U is an open subset of N , and suppose that $H : T^*U \rightarrow \mathbb{R}$ is a continuous function such that*

$$H(x, p) < 0, \text{ for every } (x, p) \in \text{Graph}(\partial^c u), \text{ with } x \in U.$$

Then for every continuous function $\epsilon : U \rightarrow]0, +\infty[$, we can find a continuous $u_\epsilon : N \rightarrow \mathbb{R}$ such that

- 1) *the function u_ϵ is C^∞ on U , and $H(x, d_x u_\epsilon) < 0$ for every $x \in U$;*
- 2) *for every $x \in U$, we have $|u(x) - u_\epsilon(x)| \leq \epsilon(x)$;*
- 3) *the functions u and u_ϵ are equal on $N \setminus U$. Moreover, at a point $x \in N \setminus U$, the function u_ϵ is differentiable if and only if u is differentiable at x . Furthermore at such a point $x \in N \setminus U$ of differentiability, we have $d_x u_\epsilon = d_x u$.*

Although the following lemma is not necessary (we could use directly [15, Theorem 8.1]), it allows to reduce to the case where H is convex in the fibers.

Lemma A.2. *Under the assumptions of Theorem A.1, we can find a continuous function $\hat{H} : T^*U \rightarrow [-1, +\infty[$ such that*

- 1) *the function \hat{H} is convex in the fibers, i.e. for every $x \in U$, the map $p \rightarrow \hat{H}(x, p)$ is convex on the vector space T_x^*N ;*
- 2) *on the the intersection $\text{Graph}(\partial^c u) \cap T^*U$, the function \hat{H} is ≤ 0 ;*
- 3) *on the set $\{(x, p) \in T^*U \mid \hat{H}(x, p) \leq 1\}$, the function H is < 0 .*

Proof. Assume first that we could find an open cover $(U_i)_{i \in I}$ of U and a family of continuous functions $\hat{H}_i : T^*U_i \rightarrow \mathbb{R}$ such that 1) 2) and 3) are satisfied with U_i instead of U , for each $i \in I$. Choosing a partition of unity $(\varphi_i)_{i \in I}$ subordinated to the open cover $(U_i)_{i \in I}$ of U , we define $\hat{H} : T^*U \rightarrow \mathbb{R}$ by $\hat{H}(x, p) = \sum_{i \in I} \varphi_i(x) \hat{H}_i(x, p)$. It is obvious 1) and 2) are true. For 3), assume that we have $\hat{H}(x, p) \leq 1$. Since $\sum_{i \in I} \varphi_i(x) = 1$, and $\hat{H}(x, p) = \sum_{i \in I} \varphi_i(x) \hat{H}_i(x, p)$, we can find an $i \in I$ such that $\varphi_i(x) > 0$, and $\varphi_i(x) \hat{H}_i(x, p) \leq \varphi_i(x)$. This implies that $x \in U_i$, and $\hat{H}_i(x, p) \leq 1$. By the assumption on \hat{H}_i , we obtain $H(x, p) < 0$. This proves 3) for \hat{H} .

To finish the proof of the lemma, given an $x \in U$, we show that we can find an open neighborhood V_x of $x \in U$, and $\hat{H}_x : T^*V_x \rightarrow \mathbb{R}$ such that 1) 2) and 3) are satisfied by \hat{H}_x on V_x . Since this is a local statement, we can assume that $U = \mathbb{R}^n$, and $x = 0$. We endow \mathbb{R}^n with its usual Euclidean scalar product. We denote respectively by $\|\cdot\|_2, d_2$ the Euclidean norm and distance on \mathbb{R}^n . We also identify \mathbb{R}^n with its dual using the scalar product, Therefore $T^*U = \mathbb{R}^n \times \mathbb{R}^n$. Since H is < 0 on the compact subset $\{0\} \times \partial^c u(0)$, we can find an open neighborhood V of 0 in \mathbb{R}^n , and $\epsilon > 0$, such that H is still < 0 on $V \times \bar{V}_{2\epsilon}(\partial^c u(0))$, where $\bar{V}_\rho(\partial^c u(0)) = \{p \in \mathbb{R}^n \mid d_2(p, \partial^c u(0)) \leq \rho\}$, for $\rho \geq 0$. By the upper semi-continuity of $\partial^c u$, cutting down the neighborhood V of 0 if necessary, we can assume that

$\partial^c u(y) \subset V_\epsilon(\partial^c u(0))$, for every $y \in V$. It remains to define $\hat{H}_0 : V \times \mathbb{R}^n \rightarrow [-1, \infty[$ by

$$\hat{H}_0(x, p) = \frac{1}{\epsilon} d_2(p, \partial^c u(0)) - 1.$$

The convexity of \hat{H}_0 in p is a consequence of the convexity of $\partial^c u(0)$. Properties 2) and 3) follow from the choice of ϵ and V . \square

Proof of Theorem A.1. Set $A = N \setminus U$, and define $\hat{\epsilon} : N \rightarrow \mathbb{R}$ by

$$\hat{\epsilon}(x) = \begin{cases} \min(\epsilon(x), d^2(x, A)), & \text{if } x \in U, \\ 0, & \text{if } x \in A, \end{cases}$$

where $d^2(x, A) = \inf\{d(x, y)^2 \mid y \in A\}$, with d the Riemannian distance. Of course the function $\hat{\epsilon}$ is continuous on N , $\hat{\epsilon} \leq d^2(\cdot, A)$ everywhere on N .

Let $\hat{H} : TU \rightarrow \mathbb{R}$ be given by Lemma A.2. We can apply [13, Theorem 10.6] or [17, Theorem 6.2] to \hat{H} (see also [10]), which is convex in the fibers, to obtain a C^∞ function $u_\epsilon : U \rightarrow \mathbb{R}$ such that $\hat{H}(x, d_x u_\epsilon) \leq 1$ on U , and $|u_\epsilon(x) - u(x)| \leq \hat{\epsilon}(x)$, for every $x \in U$. By the choice of \hat{H} , and the properties of u_ϵ , conditions 1) and 2) of the theorem are satisfied. Since $\hat{\epsilon}(\cdot) \leq d^2(\cdot, A)$, the function $w = u_\epsilon - u$ can be extended by 0 on the closed set $A = N \setminus U$ to a continuous function on N . Moreover, for $x_0 \in A, x \in N$, we have $|w(x) - w(x_0)| = |w(x)| \leq d^2(x, A) \leq d(x, x_0)^2$, therefore w is differentiable at every point of A , with derivative 0. If we extend u_ϵ to N , by $u_\epsilon = w + u$, it is now obvious that condition 3) is satisfied. \square

The following known lemma is useful.

Lemma A.3. *Let $H : T^*U \rightarrow \mathbb{R}$ be a continuous function, then the function $\mathbb{H}(u) : U \rightarrow \mathbb{R}$ defined by*

$$\mathbb{H}(u)(x) = \sup\{H(x, p) \mid p \in \partial^c u(x)\}$$

is finite valued and upper semi-continuous. Therefore, for every lower semi-continuous function $\theta : U \rightarrow \mathbb{R}$, with $\mathbb{H}(u) < \theta$ everywhere on U , there exists a continuous function $\varphi : U \rightarrow \mathbb{R}$ such that $\mathbb{H}(u) < \varphi < \theta$. In particular, the function $\mathbb{H}(u)$ is the point-wise infimum of the family of continuous functions $\varphi : U \rightarrow \mathbb{R}$ such that $\varphi > \mathbb{H}(u)$ everywhere on U .

Proof. The fact that $\mathbb{H}(u)(x) < +\infty$ follows from the compactness of $\partial^c u(x)$. The fact that it is upper semi-continuous follows from the upper semi-continuity of $\partial^c u$. The rest depends only on the fact that $\mathbb{H}(u)$ is upper-semicontinuous. In fact, if $\psi, \theta : U \rightarrow \mathbb{R}$ are respectively upper and lower semi-continuous with $\psi < \theta$, then we can always find a continuous $\varphi : U \rightarrow \mathbb{R}$ with $\psi < \varphi < \theta$. This is known as the Baire insertion theorem. We recall the simple proof. For a given $x \in U$, we pick $t_x \in \mathbb{R}$ such that $\psi(x) < t_x < \theta(x)$. By semi-continuity, the set $\{y \mid \psi(y) < t_x < \theta(y)\}$ is open and contains x . Therefore we can find an open cover $(U_i)_{i \in I}$ of U , and a family $(t_i)_{i \in I}$ of real numbers such that $\psi < t_i < \theta$ on U_i . If we call $(\varphi_i)_{i \in I}$ a partition of unity subordinated to the open cover $(U_i)_{i \in I}$, and we define the continuous $\varphi = \sum_{i \in I} t_i \varphi_i$, it is easy to check that $\psi < \varphi < \theta$.

Let us prove the last statement, which is also true for any upper semi-continuous function ψ . We fix $x_0 \in U$. We define $K_{-1} = \emptyset, K_0 = \{x_0\}$, then we complete it to a sequence $K_n, n \geq -1$ of compact subsets of U , such that $K_n \subset \overset{\circ}{K}_{n+1}$, and $U = \cup_n K_n$. Since ψ is upper semi-continuous $c_n = \sup_{K_n} \psi$ is attained on the compact set K_n and is therefore finite. Note that $c_0 = \psi(x_0)$, and $c_{n+1} \geq c_n$. If

we define θ by $\theta = c_n$ on $K_n \setminus K_{n-1}$, for $n \geq 0$, we have $\theta \geq \psi$, and $\theta(x_0) = \psi(x_0)$. Moreover, the function θ is lower semi-continuous. In fact, if $x \in U$, and $n_x = \min\{n \geq 0 \mid x \in K_n\}$, then $\psi(x) = c_{n_x}$, and $\theta \geq c_{n_x}$, on the neighborhood $U \setminus K_{n_x-1}$ of x . By the previous part of the lemma, we can find a sequence of continuous functions $\varphi_m : U \rightarrow \mathbb{R}$ such that $\psi < \varphi_m < \theta + 1/m$. Since $\psi(x_0) < \varphi_m(x_0) < \theta(x_0) + 1/m = \psi(x_0) + 1/m$, the proof is finished. \square

Let $u : N \rightarrow \mathbb{R}$ be a locally Lipschitz function. Call $\rho : N \rightarrow \mathbb{R}$ a continuous function such that

$$\forall (x, p) \in \text{Graph}(\partial^c u), \|p\|_x < \rho(x).$$

Such a function ρ exists by the previous lemma. The following statement holds:

Corollary A.4. *Let u and ρ be as above. Then we can find a sequence $u_n : N \rightarrow \mathbb{R}$ of C^∞ functions, converging uniformly to u on N , such that $\|d_x u_n\|_x < \rho(x)$, for every $x \in N$, and every $n \geq 0$, and*

$$\partial^- u(x, v) \leq \liminf_{n \rightarrow +\infty} d_x u_n(v) \leq \limsup_{n \rightarrow +\infty} d_x u_n(v) \leq \partial^+ u(x, v),$$

for every $(x, v) \in TN$.

Proof. We apply Theorem A.1, with $U = N$, $\epsilon = 1/n$, and

$$H(x, p) = \max(\|p\|_x - \rho(x), \delta^*((x, p), \text{Graph}(\partial^c u)) - 1/n),$$

to obtain a C^∞ function $u_n : U \rightarrow \mathbb{R}$ such that $\|u_n - u\|_\infty \leq 1/n$, and satisfying

$$\|d_x u_n\|_x < \rho(x), \text{ and } \delta^*((x, d_x u_n), \text{Graph}(\partial^c u)) < \frac{1}{n}.$$

It remains to prove the last part of the corollary. Fix $(x, v) \in TN$, we prove the inequality

$$\limsup_{n \rightarrow +\infty} d_x u_n(v) \leq \partial^+ u(x, v).$$

We can find a subsequence n_i of the integers such that $d_x u_{n_i}(v)$ converges to $\limsup_{n \rightarrow +\infty} d_x u_n(v)$, as $i \rightarrow \infty$. Since $\delta^*((x, d_x u_n), \text{Graph}(\partial^c u)) < 1/n$, we can find $(x_n, p_n) \in \text{Graph}(\partial^c u)$ such that $\delta^*((x, d_x u_n), (x_n, p_n)) < 1/n$. Since $d(x_n, x) \leq \delta^*((x, d_x u_n), (x_n, p_n)) < 1/n$, we have $x_n \rightarrow x$. In particular, the set $K = \{x_n \mid n \in \mathbb{N}\} \cup \{x\}$ is a compact subset of N . Therefore, the sequence (x_n, p_n) is contained in the compact subset $\tilde{K} = \{(y, p) \in \text{Graph}(\partial^c u) \mid y \in K\}$. By possibly considering a subsequence, we can assume that (x_{n_i}, p_{n_i}) converges to (x, p) . Of course, we have $p \in \partial^c u(x)$, since $\text{Graph}(\partial^c u)$ is closed. By the choice of (x_n, p_n) , we conclude that $d_{x_{n_i}} u \rightarrow p$. Hence

$$\limsup_{n \rightarrow +\infty} d_x u_n(v) = \lim_{i \rightarrow \infty} d_x u_{n_i}(v) = p(v) \leq \partial^+ u(x, v).$$

The other inequality is proved similarly. \square

We will need to use Fatou's lemma for lim sup instead of lim inf. We recall and prove the statement.

Lemma A.5 (Fatou for lim sup). *Suppose (X, \mathcal{B}, μ) is a measure space, where \mathcal{B} is a σ -algebra on X , and μ is a measure defined on \mathcal{B} . Let $\varphi, \varphi_n : X \rightarrow \mathbb{R}, n \in \mathbb{N}$ be measurable functions such that*

$$|\varphi_n(x)| \leq \varphi(x) \text{ for } \mu\text{-almost every } x \in X, \text{ and every } n \geq 0.$$

If φ is integrable, and that for every $n \geq 0$, then all the function φ_n are integrable, and so is $\limsup \varphi_n$. Moreover, we have

$$\limsup \int_X \varphi_n d\mu \leq \int_X \limsup \varphi_n d\mu.$$

Proof. We will prove it as a consequence of Lebesgue's dominated convergence theorem, but in fact, the usual proof of Lebesgue's dominated convergence theorem contains a proof of this fact. Of course the domination condition implies that each φ_n is integrable. Note that domination condition can be written as

$$-\varphi \leq \varphi_n \leq \varphi, \mu\text{-a.e.}$$

Therefore, if we define $\psi_n = \sup_{m \geq n} \varphi_m$, we will also have the domination condition $|\psi_n| \leq \varphi$, μ -a.e., for every $n \geq 0$. It follows that each ψ_n is integrable. Moreover, since $\lim \psi_n = \limsup \varphi_n$, we obtain from Lebesgue's dominated convergence theorem that $\int_X \limsup \varphi_n d\mu = \lim \int_X \psi_n d\mu$. It remains to show that $\lim \int_X \psi_n d\mu \geq \limsup \int_X \varphi_n d\mu$, which follows easily from the inequality $\psi_n \geq \varphi_n$. \square

Corollary A.6. *Let $u : N \rightarrow \mathbb{R}$ be a locally Lipschitz function. For every absolutely continuous path $\gamma : [a, b] \rightarrow N$, we have*

$$\int_a^b \partial^- u(\gamma(t), \dot{\gamma}(t)) dt \leq u(\gamma(b)) - u(\gamma(a)) \leq \int_a^b \partial^+ u(\gamma(t), \dot{\gamma}(t)) dt.$$

Proof. We will prove the right hand side inequality. The left hand side inequality follows from the left hand-side applied to $-u$. Consider the sequence of functions u_n obtained in the previous Corollary A.4. Since the u_n are smooth, and γ is absolutely continuous, we have

$$u_n(\gamma(b)) - u_n(\gamma(a)) = \int_a^b d_{\gamma(t)} u_n(\dot{\gamma}(t)) dt.$$

Since u_n converges uniformly to u , and $\limsup_{n \rightarrow +\infty} d_{\gamma(t)} u_n(\dot{\gamma}(t)) \leq \partial^+ u(\gamma(t), \dot{\gamma}(t))$ on $[a, b]$, it suffices to show that

$$\int_a^b \limsup_{n \rightarrow +\infty} d_{\gamma(t)} u_n(\dot{\gamma}(t)) dt \geq \limsup_{n \rightarrow +\infty} \int_a^b d_{\gamma(t)} u_n(\dot{\gamma}(t)) dt.$$

But this follows from the \limsup version of Fatou's lemma A.5 applied to the sequence of functions $\varphi_n(t) = d_{\gamma(t)} u_n(\dot{\gamma}(t))$, once we find a common dominated function. Since γ is an absolutely continuous curve, the function $\psi(t) = \|\dot{\gamma}(t)\|_{\gamma(t)}$ is integrable over $[a, b]$. Using $\|d_x u_n\|_x < \rho(x)$, we obtain $|\varphi_n(t)| \leq C\psi(t)$, where $C = \sup_{t \in [a, b]} \rho(\gamma(t)) < +\infty$. This exactly the type of domination we are looking for. \square

Corollary A.7. *Let $u : N \rightarrow \mathbb{R}$ be a locally Lipschitz function, and $u_n : N \rightarrow \mathbb{R}$ be a sequence of functions given by Corollary A.4. Let $\tilde{\mu}$ be a Borel measure on TN for which the norm function $TN \rightarrow \mathbb{R}, (x, v) \mapsto \|v\|_x$ is $\tilde{\mu}$ -integrable on the set $T_K N = \{(x, v) \in TN \mid x \in K\}$, for every compact subset K of N . Then the functions $\partial^+ u : TN \rightarrow \mathbb{R}$ and $du_n : TN \rightarrow \mathbb{R}, (x, v) \mapsto d_x u_n(v), n \in \mathbb{N}$ are all $\tilde{\mu}$ -integrable on $T_K N$, with*

$$\limsup \int_{T_K N} d_x u_n(v) d\tilde{\mu}(x, v) \leq \int_{T_K N} \partial^+ u(x, v) d\tilde{\mu}(x, v).$$

Proof. By the choice of ρ , we have

$$\begin{aligned} |d_x u_n(v)| &\leq C\|v\|_x, \\ |\partial^+ u(x, v)| &\leq C\|v\|_x, \end{aligned}$$

where $C = \max_{x \in K} \rho(x) < +\infty$. Since $(x, v) \mapsto C\|v\|_x$ is integrable on $T_K N$, the rest of the proof is a consequence of the limsup version of Fatou's lemma A.5 and Corollary A.4, like in the previous proof. \square

APPENDIX B. AUBRY-MATHER AND WEAK KAM THEORIES FOR NON-SMOOTH LAGRANGIANS

In this appendix, we present the main results of weak KAM Theory we use. This material is well known in the case of a Tonelli Hamiltonian, see [5, 9, 14]. The lack of Hamiltonian and Lagrangian flows requires some different arguments, see [11–13]. Although given specifically for the torus the arguments of [11, 12] can be easily rephrased in our setting.

B.1. The Lagrangian and its action along curves. With any given continuous Hamiltonian H satisfying (H1)–(H2') we can associate a function $L : TM \rightarrow \mathbb{R}$ through the *Fenchel transform*, i.e. for $(x, v) \in TM$

$$L(x, v) := \sup_{p \in T_x^* M} p(v) - H(x, p). \quad (\text{B.1})$$

The function L is called the *Lagrangian* associated with the Hamiltonian H . It is continuous on TM and satisfies properties analogous to (H1) and (H2'), see Appendix A.2 in [7]. In particular, $L(x, \cdot)$ is superlinear in $T_x M$ for every fixed $x \in M$.

The *subdifferential*, in the sense of convex analysis, of the convex function $L(x, \cdot)$ at v is the set defined by

$$\partial_v L(x, v) := \{p \in T_x^* M : L(x, v') \geq L(x, v) + p(v' - v), \text{ for every } v' \in T_x M\}.$$

We recall that the set-valued map $(x, v) \mapsto (x, \partial L/\partial v(x, v))$ from TM to T^*M is upper semicontinuous with respect to set inclusion. Analogous definitions and results hold for H . We record for later use the following well known facts, see [23, Theorem 23.5].

Proposition B.1. *Let H and L be as above. The following inequality, called Fenchel inequality, holds*

$$L(x, v) + H(x, p) \geq p(v), \text{ for every } (x, v) \in TM \text{ and } (x, p) \in T^*M, \quad (\text{B.2})$$

and

$$H(x, p) = \sup_{v \in T_x M} \{p(v) - L(x, v)\} \quad \text{for every } (x, p) \in T^*M.$$

Furthermore, the following conditions on $v \in T_x M$ and $p \in T_x^* M$ are equivalent to each other:

- (i) $L(x, v) + H(x, p) = p(v)$;
- (ii) $p \in \partial_v L(x, v)$;
- (iii) $v \in \partial_p H(x, p)$.

Let J be a closed interval in \mathbb{R} . A curve $\gamma : J \rightarrow M$ is said to be *absolutely continuous* if, for every $\varepsilon > 0$, there exists $\delta > 0$ such that, for each family $\{(a_n, b_n) : n \in \mathbb{N}\}$ of pairwise disjoint intervals included in J , the following property holds:

$$\text{if } \sum_{n=1}^{+\infty} |b_n - a_n| < \delta \quad \text{then} \quad \sum_{n=1}^{+\infty} d(\gamma(a_n), \gamma(b_n)) < \varepsilon.$$

The family of absolutely continuous curves from J to M will be henceforth denoted by $\text{AC}(J; M)$. We will say that a sequence of curves γ_n in $\text{AC}(J; M)$ uniformly converge to $\gamma : J \rightarrow M$ if $\sup_{t \in J} d(\gamma_n(t), \gamma(t)) \rightarrow 0$ as $n \rightarrow +\infty$. The limit curve γ is continuous, but not absolutely continuous in general.

An absolutely continuous curve $\gamma : J \rightarrow M$ is differentiable at almost every point of J . We will denote by $\|\dot{\gamma}\|_\infty$ the essential supremum norm of the velocity field of the curve, i.e.

$$\|\dot{\gamma}\|_\infty := \text{ess sup}_{s \in J} \|\dot{\gamma}(s)\|_{\gamma(s)}$$

If there exists $\alpha > 0$ such that $d(\gamma(s), \gamma(t)) \leq \alpha |s - t|$ for every $s, t \in J$, then the curve γ will be termed *Lipschitz*, or α -*Lipschitz* when we want to specify the Lipschitz constant. This is equivalent to requiring $\|\dot{\gamma}\|_\infty \leq \alpha$ for M is a length space. When M is embedded in \mathbb{R}^k , it is well known that γ is absolutely continuous if and only if its distributional derivative $\dot{\gamma}$ belongs to $L^1(J; \mathbb{R}^k)$.

Let us now assume that J is bounded, i.e. of the form $[a, b]$ for some $a < b$ in \mathbb{R} . For every $\lambda \geq 0$, we define a functional \mathbb{L}^λ on $\text{AC}([a, b]; M)$ by setting

$$\mathbb{L}^\lambda(\gamma) := \int_a^b e^{\lambda s} L(\gamma(s), \dot{\gamma}(s)) ds, \quad \gamma \in \text{AC}([a, b]; M).$$

From the fact that L is bounded from below, it is easily seen that this integral is always well defined, with values in $\mathbb{R} \cup \{+\infty\}$. The following Tonelli–type theorem holds:

Theorem B.2. *Let $(\gamma_n)_n$ be a sequence in $\text{AC}([a, b]; M)$ such that*

$$\sup_{n \in \mathbb{N}} \mathbb{L}^\lambda(\gamma_n) < +\infty.$$

If a continuous curve $\xi : [a, b] \rightarrow M$ is the uniform limit of some subsequence $(\gamma_{n_k})_k$, then $\xi \in \text{AC}([a, b]; M)$ and

$$\mathbb{L}^\lambda(\xi) \leq \liminf_{k \rightarrow +\infty} \mathbb{L}^\lambda(\gamma_{n_k}).$$

Moreover, there exists a subsequence $(\gamma_{n_k})_k$ uniformly converging to a curve $\gamma \in \text{AC}([a, b]; M)$.

When M is contained in \mathbb{R}^k , the above theorem easily follows by making use of the Dunford–Pettis theorem, see [6, Theorems 2.11 and 2.12], and of standard semicontinuity results in the Calculus of Variations, see [6, Theorem 3.6]. To get the result in full generality, it suffices to show that we can always reduce to this case by localizing the argument and by reasoning in local charts, see for instance [14].

B.2. Weak KAM Theory. For every $t > 0$, we define a function $h_t : M \times M \rightarrow \mathbb{R}$ by setting

$$h_t(x, y) = \inf \left\{ \int_{-t}^0 L(\gamma, \dot{\gamma}) + c(H) ds : \gamma \in \text{AC}([-t, 0]; M), \gamma(-t) = x, \gamma(0) = y \right\}.$$

The quantity $h_t(x, y)$ is called the *minimal action* to go from x to y in time t .

The following characterization holds, see [14]:

Proposition B.3. *Let $w \in C(M, \mathbb{R})$. Then w is a critical subsolution if and only if*

$$w(x) - w(y) \leq h_t(y, x) \quad \text{for every } x, y \in M \text{ and } t > 0.$$

The *Peierls barrier* is the function $h : M \times M \rightarrow \mathbb{R}$ defined by

$$h(x, y) = \liminf_{t \rightarrow +\infty} h_t(x, y). \quad (\text{B.3})$$

It satisfies the following properties, see for instance [12]:

Proposition B.4. a) *The Peierls barrier h is finite valued and Lipschitz continuous.*

b) *If w is a critical subsolution, then*

$$w(x) - w(y) \leq h(y, x), \text{ for every } x, y \in M.$$

c) *For every $x, y, z \in M$ and $t > 0$, we have*

$$\begin{aligned} h(y, x) &\leq h(y, z) + h_t(z, x) \\ h(y, x) &\leq h_t(y, z) + h(z, x) \\ h(y, x) &\leq h(y, z) + h(z, x). \end{aligned}$$

d) *For every fixed $y \in M$, the function $h(y, \cdot)$ is a critical solution.*

e) *For every fixed $y \in M$, the function $-h(\cdot, y)$ is a critical subsolution.*

The projected *Aubry set* \mathcal{A} is the closed set defined by

$$\mathcal{A} := \{y \in M : h(y, y) = 0\}.$$

We use the terminology *projected Aubry set* to be coherent with the Tonelli case.

The following holds, see [13]:

Theorem B.5. *There exists a critical subsolution w which is of class C^1 and strict in $M \setminus \mathcal{A}$, i.e. satisfies*

$$H(x, d_x w) < c(H), \text{ for every } x \in M \setminus \mathcal{A}.$$

In particular, the projected Aubry set \mathcal{A} is nonempty.

When the Hamiltonian is locally Lipschitz in x and strictly convex in p , such a strict subsolution can be taken of class C^1 on the whole manifold M [16, 17], and even of class $C^{1,1}$ when H is Tonelli [4]. It is not known whether the result keeps holding in the case of a purely continuous Hamiltonian.

A consequence of Theorem B.5 is that \mathcal{A} is a uniqueness set for the critical equation. In fact, we have, see [13]:

Theorem B.6. *Let w, u be a critical sub and supersolution, respectively. If $w \leq u$ on \mathcal{A} , then $w \leq u$ on M . In particular, if two critical solutions coincide on the projected Aubry set \mathcal{A} , then they coincide on the whole manifold M .*

The next result provides a converse of Theorem B.5. In particular, it implies that \mathcal{A} is the set where the obstruction to the existence of strict critical subsolutions concentrates.

Proposition B.7. *Let $y \in \mathcal{A}$. For every critical subsolution w we have*

$$H(y, p) = c(H) \text{ for every } p \in D^-w(y).$$

Proof. Let w be a critical subsolution. By Proposition B.4–(ii), we have

$$w(x) \leq w(y) + h(y, x) =: u(x), \text{ for every } x \in M,$$

with equality holding for $x = y$ since $h(y, y) = 0$ by definition of the projected Aubry set. Then $D^-w(y) \subset D^-u(y)$, so the assertion follows from Proposition 2.1 and the fact that u is a critical solution. \square

Let us now consider the symmetric function δ_M defined by

$$\delta_M(x, y) = h(x, y) + h(y, x), \quad x, y \in M.$$

Proposition B.4 yields that δ_M is always nonnegative and satisfies the triangular inequality. Moreover, it induces a semidistance on \mathcal{A} , i.e. a function which fails to be a distance because the equality $\delta_M(x, y) = 0$ for $x, y \in \mathcal{A}$ does not necessarily imply that $x = y$.

To make δ_M a distance, we introduce a natural equivalence relation on \mathcal{A} , called the *Mather relation*, defined by $x \sim y$ if $h(x, y) + h(y, x) = 0$. Its equivalence classes are called the *Mather classes*. Two points x and y are in the same Mather class if and only if

$$h(x, y) + h(y, x) = 0.$$

Clearly, δ_M is a distance function in the quotient set \mathcal{A}/\sim formed by Mather classes. This quotient, endowed with the distance δ_M , is known as *Mather quotient*.

Proposition B.8. *Let $w : M \rightarrow \mathbb{R}$ be a critical subsolution. Then*

$$w(x) - w(y) = h(y, x)$$

for every $x, y \in \mathcal{A}$ in the same Mather class. Moreover, if $w_1, w_2 : M \rightarrow \mathbb{R}$ are critical subsolutions, we have

$$|(w_1 - w_2)(x) - (w_1 - w_2)(y)| \leq \delta_M(y, x) \quad \text{for every } x, y \in M.$$

In particular, $w_1 - w_2$ is constant on each Mather class.

Proof. By Proposition B.4–(ii) we have

$$\begin{aligned} w(x) - w(y) &\leq h(y, x) \\ w(y) - w(x) &\leq h(x, y). \end{aligned}$$

If we add these two inequalities, we get

$$w(x) - w(y) + w(y) - w(x) \leq h(y, x) + h(x, y) = \delta_M(y, x).$$

But the left hand side is zero, and so is the the right hand side if x, y are in the same Mather class. Hence the sum of the two inequalities is an equality, therefore both inequalities are equalities.

In the same way, from Proposition B.4–(ii) we get

$$\begin{aligned} w_1(x) - w_1(y) &\leq h(y, x) \\ w_2(y) - w_2(x) &\leq h(x, y). \end{aligned}$$

Adding and rearranging yields

$$(w_1 - w_2)(x) - (w_1 - w_2)(y) \leq \delta_M(x, y).$$

Since δ_M is symmetric, we get

$$|(w_1 - w_2)(x) - (w_1 - w_2)(y)| \leq \delta_M(x, y).$$

The rest of the proposition is obvious. \square

The following holds:

Theorem B.9. *Let $y \in \mathcal{A}$. Then there exists an absolutely continuous curve $\gamma : \mathbb{R} \rightarrow M$ with $\gamma(0) = y$ such that, for every $a, b \in \mathbb{R}$, with $a \leq b$, we have*

- (i) $\int_a^b L(\gamma(s), \dot{\gamma}(s)) + c(H) ds = h(\gamma(a), \gamma(b));$
- (ii) $h(\gamma(a), \gamma(b)) + h(\gamma(b), \gamma(a)) = 0.$

In particular, the curve γ is supported in the Mather class of y .

Proof. The existence of a curve satisfying (i) and (ii) above, in the case of a purely continuous Hamiltonian, is proved in [11], see also [12]. It is then clear by (ii) that $\gamma(t) \sim y$ for every $t \in \mathbb{R}$. \square

A curve satisfying assumptions (i) and (ii) in the above statement is called *static*.

Proposition B.10. *Any static curve $\gamma : \mathbb{R} \rightarrow M$ is α_0 -Lipschitz for some constant α_0 only depending on H . Moreover, for every critical subsolution $u : M \rightarrow \mathbb{R}$, and every $a, b \in \mathbb{R}$, with $a \leq b$, we have*

$$u(\gamma(b)) - u(\gamma(a)) = \int_a^b L(\gamma(s), \dot{\gamma}(s)) + c(H) ds.$$

Proof. Let us prove that γ is Lipschitz. The function $h(z, \cdot)$ is, for every fixed $z \in M$, a critical solution, in particular it is κ -Lipschitz continuous by Proposition 3.4. From property (i) of static curves, we derive

$$\int_a^{a+h} L(\gamma(s), \dot{\gamma}(s)) + c(H) ds \leq \kappa d(\gamma(a), \gamma(a+h)), \quad (\text{B.4})$$

for every $a \in \mathbb{R}$ and $h > 0$. By the superlinearity of L , there exists a constant A_κ such that

$$L(x, v) \geq (\kappa + 1)\|v\|_x - A_\kappa, \text{ for every } (x, v) \in TM.$$

We plug this inequality in (B.4) and we use the fact that the length of a curve is greater or equal than the distance between its extreme points. By dividing by h we end up with

$$\frac{1}{h} \int_a^{a+h} \|\dot{\gamma}(s)\|_{\gamma(s)} ds \leq A_\kappa - c(H). \quad (\text{B.5})$$

Sending $h \rightarrow 0$ we infer

$$\|\dot{\gamma}(a)\|_{\gamma(a)} \leq A_\kappa - c(H), \text{ for a.e. } a \in \mathbb{R},$$

as it was to be shown. To prove the last equality of the theorem, we observe that we have

$$\begin{aligned} u(\gamma(b)) - u(\gamma(a)) &\leq h(\gamma(a), \gamma(b)) = \int_a^b L(\gamma(s), \dot{\gamma}(s)) + c(H) ds, \\ u(\gamma(a)) - u(\gamma(b)) &\leq h(\gamma(b), \gamma(a)). \end{aligned}$$

If we add these two inequalities, we get the inequality $0 \leq 0$. Therefore both inequalities must be equalities. \square

B.3. The Aubry set in TM .

Proposition B.11. *Let $u : M \rightarrow \mathbb{R}$ be a Lipschitz function. Then u is a viscosity subsolution of $H(x, d_x u) = c$ if and only if $\partial^+ u(x, v) \leq L(x, v) + c$, for every $(x, v) \in TM$.*

Proof. Suppose the function u is a viscosity subsolution of $H(x, d_x u) = c$, then $H(x, d_x u) \leq c$ at every point where the derivative $d_x u$ exists, therefore by Fenchel's inequality

$$d_x u(v) \leq L(x, v) + H(x, p) \leq L(x, v) + c.$$

This implies that for every $p \in \partial^* u(x)$, and every $v \in T_x M$, we have $p(v) \leq L(x, v) + c$. Since $\partial^c u(x)$ is the convex envelop of $\partial^* u(x)$, the inequality $p(v) \leq L(x, v) + c$ remains true for any $p \in \partial^c u(x)$. Taking the sup over such p , yields $\partial^+ u(x, v) \leq L(x, v) + c$.

Conversely, if $\partial^+ u(x, v) \leq L(x, v) + c$, at a point x of differentiability of u we get $d_x u(v) \leq \partial^+ u(x, v) \leq L(x, v) + c$. From this, we obtain $H(x, d_x u) = \sup_{v \in T_x M} d_x u - L(x, v) \leq c$. Since H is convex in p , we infer that u is a viscosity subsolution of $H(x, d_x u) = c$. \square

For a critical subsolution $u : M \rightarrow \mathbb{R}$, we define

$$\mathcal{L}(u) = \{(x, v) \mid \partial^+ u(x, v) = L(x, v) + c(H)\},$$

where $c(H)$ is the critical value of H .

Lemma B.12. *For every critical subsolution, the set $\mathcal{L}(u) \subset TM$ is compact and $\pi(\mathcal{L}(u)) \supset \mathcal{A}$. In fact, if $\gamma : \mathbb{R} \rightarrow M$ is a static curve, then for almost every $s \in \mathbb{R}$, we have $(\gamma(s), \dot{\gamma}(s)) \in \mathcal{L}(u)$.*

Proof. Denote by κ a Lipschitz constant for u . Then $\|d_x u\|_x \leq \kappa$ at every point where $d_x u$ exists. Therefore $\partial^+ u(x, v) \leq \kappa \|v\|_x$. In particular, for every $(x, v) \in \mathcal{L}(u)$, we have $L(x, v) + c(H) \leq \kappa \|v\|_x$. Since L is superlinear, we can find a finite constant A_κ such that $L(x, v) \geq (\kappa + 1) \|v\|_x - A_\kappa$, for every $(x, v) \in TM$. Therefore $\kappa \|v\|_x \geq (\kappa + 1) \|v\|_x - A_\kappa + c(H)$, and $\|v\|_x \leq A_\kappa - c(H)$, for $(x, v) \in \mathcal{L}(u)$. This proves that $\mathcal{L}(u)$ is relatively compact in TM . We now prove that it is closed. Suppose $(x_n, v_n) \rightarrow (x, v)$ with

$$\partial^+ u(x_n, v_n) = L(x_n, v_n) + c(H).$$

By continuity of L , the right hand side tends to $L(x, v) + c(H)$. By the upper semi-continuity of $\partial^+ u$, we obtain $L(x, v) + c(H) \leq \partial^+ u(x, v)$. But the reverse inequality is true by Proposition B.11.

Suppose now that γ is a static curve. From Proposition B.11, we get

$$\partial^+ u(\gamma(s), \dot{\gamma}(s)) \leq L(\gamma(s), \dot{\gamma}(s)) + c(H),$$

for almost every $s \in \mathbb{R}$. Integrating, between a and b , with $a \leq b$, we obtain

$$\int_a^b \partial^+ u(\gamma(s), \dot{\gamma}(s)) ds \leq \int_a^b L(\gamma(s), \dot{\gamma}(s)) + c(H) ds.$$

But, by Corollary A.6, we have $u(\gamma(b)) - u(\gamma(a)) \leq \int_a^b \partial^+ u(\gamma(s), \dot{\gamma}(s)) ds$, and by Proposition B.10, we also have $u(\gamma(b)) - u(\gamma(a)) = \int_a^b L(\gamma(s), \dot{\gamma}(s)) + c(H) ds$. It follows that the integrated equality above is an equality, therefore the inequality is an almost everywhere equality. Hence, we obtain $(\gamma(s), \dot{\gamma}(s)) \in \mathcal{L}(u)$, for almost every $s \in \mathbb{R}$. Therefore $\gamma(s)$ is in the compact set $\pi(\mathcal{L}(u))$ for almost every s . By

continuity of γ , we conclude that $\gamma(t) \in \pi(\mathcal{L}(u))$, for every $t \in \mathbb{R}$. It follows from Theorem B.9 that $\mathcal{A} \subset \pi(\mathcal{L}(u))$. \square

Definition B.13 (Aubry set). *We define the Aubry set $\tilde{\mathcal{A}}$ as*

$$\tilde{\mathcal{A}} = \bigcap_u \mathcal{L}(u),$$

where the intersection is taken over all critical subsolutions $u : M \rightarrow \mathbb{R}$.

Theorem B.14. *The Aubry set $\tilde{\mathcal{A}}$ is compact non-empty, and $\pi(\tilde{\mathcal{A}}) = \mathcal{A}$. Moreover, there exists a critical subsolution $u : M \rightarrow \mathbb{R}$ with*

$$\tilde{\mathcal{A}} = \mathcal{L}(u).$$

Proof. We first prove that $\pi(\tilde{\mathcal{A}}) \subset \mathcal{A}$. By Theorem B.5, we can find a critical subsolution $u : M \rightarrow \mathbb{R}$ which is C^∞ outside of \mathcal{A} , and such that $H(x, d_x u) < c(H)$ for every $x \notin \mathcal{A}$. Since u is smooth on the open set $M \setminus \mathcal{A}$, for $(x, v) \in TM$, with $x \notin \mathcal{A}$, we have $\partial^+(x, v) = d_x u(v)$. By Fenchel inequality, we obtain $\partial^+(x, v) = d_x u(v) \leq L(x, v) + H(x, d_x u)$. Therefore $\partial^+ u(x, v) < L(x, v) + c(H)$, and no (x, v) , with $x \notin \mathcal{A}$, can be in $\mathcal{L}(u)$. Hence $\mathcal{A} \supset \pi(\mathcal{L}(u)) \supset \pi(\tilde{\mathcal{A}})$.

To finish the proof of the theorem, since $\tilde{\mathcal{A}} \subset \mathcal{L}(u)$ for every critical subsolution $u : M \rightarrow \mathbb{R}$, it suffices to find a critical solution $u : M \rightarrow \mathbb{R}$ such that $\mathcal{L}(u) \subset \tilde{\mathcal{A}}$. We recall the following fact from general topology:

If X is a separable metric space, and $(F_i)_{i \in I}$ is a family of closed sets, then we can find a sequence $i_n, n \geq 1$ with $\bigcap_{i \in I} F_i = \bigcap_{n \geq 1} F_{i_n}$.

In fact, the open set $U = X \setminus \bigcap_{i \in I} F_i$ is covered by the family of open sets $U_i = X \setminus F_i, i \in I$. Since U itself is separable metric, we can extract a countable subcover $U_{i_n}, n \geq 1$ with $U = \bigcup_{n \in \mathbb{N}} U_{i_n}$. Therefore, we have $X \setminus \bigcap_{i \in I} F_i = X \setminus \bigcap_{n \geq 1} F_{i_n}$, which implies $\bigcup_{i \in I} F_i = \bigcap_{n \geq 1} F_{i_n}$.

Therefore, we can find a sequence $u_n : M \rightarrow \mathbb{R}, n \geq 1$, of critical subsolutions such that $\tilde{\mathcal{A}} = \bigcap_{n \geq 1} \mathcal{L}(u_n)$. Fix $x_0 \in M$. If we replace u_n , by $v_n = u_n - u_n(x_0)$, we obtain another critical subsolution v_n with $\mathcal{L}(v_n) = \mathcal{L}(u_n)$. Therefore we can assume without loss of generality that $u_n(x_0) = 0$, for every $n \geq 1$. The sequence u_n of critical subsolution is equi-Lipschitz. Call κ such a constant. Since $u_n(x_0) = 0$, we get $\|u_n\|_\infty \leq \kappa \text{diam}(M)$, where $\text{diam}(M) < +\infty$ is the diameter of the compact manifold M . Not also that $|\partial^+ u(x, v)| \leq \kappa \|v\|_x$, for every $(x, v) \in TM$. Therefore if for $A \geq 0$, we denote by $\bar{B}_A = \{(x, v) \mid \|v\|_x \leq A\}$, we obtain $\|\partial^+ u_n|_{\bar{B}_A}\|_\infty \leq \kappa A$. It follows that the series $2^{-n} u_n, n \geq 1$, and $2^{-n} \partial^+ u_n, n \geq 1$ converge uniformly respectively on M and on compact subsets of TM . If we set $u = \sum_{n \geq 1} 2^{-n} u_n$, then by convexity of H in p and the fact that $\sum_{n \geq 1} 2^{-n} = 1$, we infer that the function u is also a critical subsolution. Moreover, we have

$$\forall (x, v) \in TM, \partial^+ u(x, v) \leq \sum_{n \geq 1} 2^{-n} \partial^+ u_n(x, v). \quad (\text{B.6})$$

This is a well known fact in non-smooth analysis. For the convenience of the reader, we provide the proof below.

We have

$$\partial^+ u_n(x, v) \leq L(x, v) + c(H), \text{ for every } n \geq 1 \text{ and every } (x, v) \in TM. \quad (\text{B.7})$$

By averaging these inequalities, and using the inequality (B.6) above, we get

$$\partial^+ u(x, v) \leq \sum_{n \geq 1} 2^{-n} \partial^+ u_n(x, v) \leq L(x, v) + c(H), \quad (\text{B.8})$$

for every $(x, v) \in TM$. If $(x, v) \in \mathcal{L}(u)$, then $\partial^+ u(x, v) = L(x, v) + c(H)$, and therefore all inequalities in (B.8) must be equalities. Since the right hand side inequality in (B.8) was obtained by averaging the inequalities in (B.7), we must have $\partial^+ u_n(x, v) = L(x, v) + c(H)$ for every $(x, v) \in \mathcal{L}(u)$, and every $n \geq 1$. This implies $\mathcal{L}(u) \subset \bigcap_{n \geq 1} \mathcal{L}(u_n) = \tilde{\mathcal{A}}$. \square

Lemma B.15. *Suppose that for each $n \geq 1$, the function $u_n : M \rightarrow \mathbb{R}$ is Lipschitz, with Lipschitz constant κ_n . Assume that $\sum_{n \geq 1} \|u_n\|_\infty < +\infty$, and $\sum_{n \geq 1} \kappa_n < +\infty$, then $u = \sum_{n \geq 1} u_n$ is a Lipschitz function on M , and $\sum_{n \geq 1} \partial^+ u_n$ is an upper semi-continuous function on TM . Moreover, for every $(x, v) \in TM$, we have*

$$\partial^+ u(x, v) \leq \sum_{n \geq 1} \partial^+ u_n(x, v).$$

Proof. It is obvious that the series $\sum_{n \geq 1} u_n$ converges uniformly to a function u that is Lipschitz with Lipschitz constant $\sum_{n \geq 1} \kappa_n < +\infty$. Moreover since $\partial^+ u_n(x, v) \leq \kappa_n \|v\|_x$, the series $\sum_{n \geq 1} \partial^+ u_n$ converges uniformly on any compact subset of TM . Using this last fact and the upper semi-continuity of each function $\partial^+ u_n$ we obtain that the sum $\theta = \sum_{n \geq 1} \partial^+ u_n$ is also upper semi-continuous. If we consider a C^1 path $\gamma : [a, b] \rightarrow M$, since its speed $\dot{\gamma}(t)$ is bounded, we concluded that the series $\sum_{n \geq 1} \partial^+ u_n(\gamma(t), \dot{\gamma}(t))$ converges uniformly to $\theta(\gamma(t), \dot{\gamma}(t))$. In particular, we obtain

$$\int_a^b \theta(\gamma(t), \dot{\gamma}(t)) dt = \sum_{n \geq 1} \int_a^b \partial^+ u_n(\gamma(t), \dot{\gamma}(t)) dt.$$

By Corollary A.6, we have

$$u_n(\gamma(b)) - u_n(\gamma(a)) \leq \int_a^b \partial^+ u_n(\gamma(t), \dot{\gamma}(t)) dt.$$

Since $u = \sum_{n \geq 1} u_n$, we obtain

$$u(\gamma(b)) - u(\gamma(a)) \leq \int_a^b \theta(\gamma(t), \dot{\gamma}(t)) dt.$$

Assume now $d_x u$ exists, and $v \in T_x M$. Pick a C^1 path $\gamma : [0, 1] \rightarrow M$, with $\gamma(0) = x, \dot{\gamma}(0) = v$. For very $\epsilon > 0$, we have

$$u(\gamma(\epsilon)) - u(\gamma(0)) \leq \int_0^\epsilon \theta(\gamma(t), \dot{\gamma}(t)) dt.$$

If we divide this inequality by ϵ , and let $\epsilon \rightarrow 0$, the limit of the right hand side is $d_x u(v)$, and the lim sup of the left hand side is $\leq \theta(x, v)$, since $t \mapsto \theta(\gamma(t), \dot{\gamma}(t))$ is upper semi-continuous. Therefore, we obtained $d_x u(v) \leq \theta(x, v)$ at every differentiability point x of u and every $v \in T_x M$. From this inequality, using that θ is upper semi-continuous, we get $p(v) \leq \theta(x, v)$ for every $(x, v) \in TM$, and every $p \in \partial^* u(x)$. From which it follows easily that $\partial^+ u(x, v) \leq \theta(x, v)$ on TM . \square

B.4. Mather measures and Mather set. In this work, we will deal with probability measures defined either on the compact manifold M or on its tangent bundle TM . A measure on TM will be denoted by $\tilde{\mu}$, where the tilde on the top is to keep track of the fact that the measure is on the space TM . If $\tilde{\mu}$ is a probability measure on TM , we will denote by μ its projection $\pi_{\#}\tilde{\mu}$ on M , i.e. the probability measure on M defined as

$$\pi_{\#}\tilde{\mu}(B) := \tilde{\mu}(\pi^{-1}(B)) \quad \text{for every } B \in \mathcal{B}(M).$$

Note that

$$\int_M f(x) \pi_{\#}\tilde{\mu}(x) = \int_{TM} (f \circ \pi)(x, v) d\tilde{\mu}(x, v) \quad \text{for every } f \in C(M, \mathbb{R}).$$

For a Borel measure μ on a metric separable space X , there is a largest open subset $U \subset X$ with $\mu(U) = 0$. The complement $X \setminus U$ is called the support of μ and is denoted by $\text{supp}(\mu)$. This set $\text{supp}(\mu)$ is the smallest closed subset of full μ -measure in X . In this section we generalize Mather theory to the non-smooth case, that is when the Hamiltonian is a continuous function satisfying (H1)–(H2'). We will assume that the connected manifold M is endowed with an auxiliary Riemannian metric, and we will denote by d the associated Riemannian distance.

The first step consists in showing that the constant $-c(H)$, where $c(H)$ is the critical value, can be also obtained by minimizing the integral of the Lagrangian over TM with respect to a suitable family of probability measure on TM . In the case of a Tonelli Hamiltonian, it is customary to choose this family as the one made up by probability measures on TM that are invariant by the Euler–Lagrange flow, see [22]. This approach is not feasible here due to the lack of regularity of H . It was shown that this minimization problem yields the same result if it is done on the set of closed measures [1, 16, 17, 21]. This is a set that does not depend on the Hamiltonian, and therefore this is the approach that can be adapted in a more general setting. The definition of closed measure is the following:

Definition B.16. *A probability measure $\tilde{\mu} \in \mathcal{P}(TM)$ will be called closed if it satisfies the following properties:*

- (a) $\int_{TM} \|v\|_x d\tilde{\mu}(x, v) < +\infty;$
- (b) $\int_{TM} d_x \varphi(v) d\tilde{\mu}(x, v) = 0$, for every $\varphi \in C^1(M)$.

We will denote by $\tilde{\mathfrak{M}}$ the set of closed probability measures on TM .

The following lemma is a particular case of Corollary A.7.

Lemma B.17. *Let $\tilde{\mu}$ be a closed measure on the tangent bundle TM of the compact manifold M . Then for every Lipschitz function $u : M \rightarrow \mathbb{R}$, we have*

$$\int_{TM} \partial^+ u(x, v) d\tilde{\mu}(x, v) \geq 0.$$

Let us recall that a first way to construct closed measures. If $\gamma : [a, b] \rightarrow M$ is an absolutely continuous curve, we define the probability measure $\tilde{\mu}_\gamma$ on TM , by

$$\int_{TM} \psi d\tilde{\mu}_\gamma = \frac{1}{b-a} \int_a^b \psi(\gamma(t), \dot{\gamma}(t)) dt,$$

for every bounded Borel measurable function $\psi : TM \rightarrow \mathbb{R}$. This measure is nothing but the image of the normalized Lebesgue $(b-a)^{-1}dt$ on the interval $[a, b]$ by the speed curve map $t \mapsto (\gamma(t), \dot{\gamma}(t))$ defined almost everywhere on $[a, b]$. Note that

$$\begin{aligned} \int_{TM} \|v\|_x d\tilde{\mu}_\gamma(x, v) &= \frac{1}{b-a} \int_a^b \|\dot{\gamma}(t)\|_{\gamma(t)} dt \\ &= \frac{\ell_g(\gamma)}{b-a} < +\infty, \end{aligned}$$

where $\ell_g(\gamma)$ is the Riemannian length of γ , which is finite because γ is absolutely continuous on the compact interval $[a, b]$. Note also that for $f : M \rightarrow \mathbb{R}$ of class C^1 , we have

$$\int_{TM} d_x f(v) d\tilde{\mu}_\gamma(x, v) = \frac{1}{b-a} \int_a^b d_{\gamma(t)} f(\dot{\gamma}(t)) dt = \frac{f(\gamma(b)) - f(\gamma(a))}{b-a}.$$

In particular, if γ is a loop, then $\tilde{\mu}_\gamma$ is closed. We can also use unbounded curves to define closed measures. We sum up this fact in the following lemma.

Lemma B.18. *Let $\gamma : [0, +\infty[\rightarrow M$ be a (globally) Lipschitz curve. For $t > 0$, define the measure $\tilde{\mu}_t = \tilde{\mu}_{\gamma|_{[0,t]}}$. These probability measures on TM have all support in the compact closure $\overline{\{(\gamma(s), \dot{\gamma}(s)) \mid s \in S\}}$, where S is a subset of full measure in \mathbb{R} on which $\dot{\gamma}$ is defined. In particular, the set of measures $\tilde{\mu}_t, t > 0$ is compact in $\tilde{\mathfrak{M}}$ for the weak topology. Therefore, we can find accumulation points for $\tilde{\mu}_t$, as $t \rightarrow \infty$. Any such accumulation point is a closed measure.*

Proof. Call $\tilde{\mu}$ such an accumulation point, and suppose that $\tilde{\mu}_{t_i} \rightarrow \tilde{\mu}$ in the weak topology, with $t_i \rightarrow +\infty$. Note that support of $\tilde{\mu}$ is also contained $\overline{\{(\gamma(s), \dot{\gamma}(s)) \mid s \in S\}}$. It is therefore compact, and any continuous function on TM is $\tilde{\mu}$ -integrable. In particular condition (a) in the Definition B.16 is satisfied. If $f : M \rightarrow \mathbb{R}$ is C^1 , we have

$$\begin{aligned} \int_{TM} d_x f(v) d\tilde{\mu}(x, v) &= \lim_{i \rightarrow +\infty} \int_{TM} d_x f(v) d\tilde{\mu}_{t_i}(x, v) \\ &= \frac{f(\gamma(t_i)) - f(\gamma(0))}{t_i} \\ &= 0. \end{aligned}$$

The last equality follows from the facts that the continuous function f is bounded on the compact set M , and that $t_i \rightarrow +\infty$. \square

We now clarify the relation between the critical value and minimizing measure.

Proposition B.19. *For every closed measure $\tilde{\mu}$ on TM , we have*

$$\int_{TM} L(x, v) d\tilde{\mu}(x, v) \geq -c(H),$$

where $c(H)$ is the critical value of L .

Proof. Let us consider $u : M \rightarrow \mathbb{R}$ a critical subsolution for H . We have $H(x, p) \leq c(H)$ for every $p \in \partial^c u(x)$. Therefore, for every $(x, v) \in TM$, and every $p \in \partial^c u(x)$, by Fenchel's inequality, we have

$$p(v) \leq L(x, v) + H(x, p) \leq L(x, v) + c(H).$$

Taking the sup over all $p \in \partial^c u(x)$ yields

$$\forall(x, v) \in TM, \partial^+ u(x, v) \leq L(x, v) + c(H).$$

Since $\int_{TM} \partial^+ u(x, v) d\tilde{\mu}(x, v) \geq 0$, by Lemma B.17, we obtain $\int_{TM} L(x, v) d\tilde{\mu}(x, v) + c(H) \geq 0$. \square

The relation linking closed probability measures to the critical value is clarified by the next theorem.

Theorem B.20. *The following holds:*

$$\inf_{\tilde{\mu} \in \tilde{\mathfrak{M}}} \int_{TM} L(x, v) d\tilde{\mu}(x, v) = -c(H) \quad (\text{B.9})$$

where $c(H)$ is the critical value for H . Moreover the inf is achieved by a closed measure. More precisely, for every $y \in \mathcal{A}$ there exists a minimizing measure $\tilde{\mu} \in \tilde{\mathfrak{M}}$ such that $\pi_{\#}\tilde{\mu}$ is supported in the Mather class of y .

Proof. Fix $y \in \mathcal{A}$. By Theorem B.9, there exists a static curve $\gamma : \mathbb{R} \rightarrow M$, with $\gamma(0) = y$. Since this curve γ is Lipschitz, by Lemma B.18, we can find $t_i \rightarrow +\infty$ such that $\tilde{\mu}_{t_i} = \tilde{\mu}_{\gamma|_{[0, t_i]}}$ converges weakly to a closed probability measure $\tilde{\mu}$, and have all their support contained in the compact subset $\overline{\{(\gamma(s), \dot{\gamma}(s)) \mid s \in S\}}$, where S is a subset of full measure in \mathbb{R} on which $\dot{\gamma}$ is defined. Therefore, the projection $\pi_{\#}\tilde{\mu}$ is supported in the closure $\overline{\{\gamma(s) \mid s \in \mathbb{R}\}}$. Since γ is static with $\gamma(0) = y$, the entire curve γ is contained in the Mather class of y , which is closed. This proves the last claim of the theorem. It remains to show that $\int_{TM} L d\tilde{\mu}$ is equal to $-c(H)$. Since L is continuous and the supports of both $\tilde{\mu}$ and the $\tilde{\mu}_{t_i}$ are contained in the same compact subset of TM , we have

$$\begin{aligned} \int_{TM} L(x, v) d\tilde{\mu}(x, v) &= \lim_{i \rightarrow +\infty} \int_{TM} L(x, v) d\tilde{\mu}_{t_i}(x, v) \\ &= \lim_{i \rightarrow +\infty} \frac{1}{t_i} \int_0^{t_i} L(\gamma(s), \dot{\gamma}(s)) ds = \lim_{i \rightarrow +\infty} \frac{h(\gamma(0), \gamma(t_i))}{t_i} = 0, \end{aligned}$$

where in the last two equalities, we have used that γ is a static curve and that h is bounded on $M \times M$. \square

Definition B.21. *A Mather measure for the Lagrangian L is a closed measure $\tilde{\mu} \in \tilde{\mathfrak{M}}$ such that $\int_{TM} L(x, v) d\tilde{\mu}(x, v) = -c(H)$. The set of Mather measures will be denoted by $\tilde{\mathfrak{M}}_0(L)$.*

A projected Mather measure is a Borel probability measure in μ on M of the form $\mu = \pi_{\#}\tilde{\mu}$, where $\tilde{\mu} \in \tilde{\mathfrak{M}}_0(L)$. The set of projected Mather measures is denoted by $\mathfrak{M}_0(L)$.

Some time the terminology Mather minimizing measure, rather than Mather measure, is used to emphasize that a Mather measure is solving the minimization problem (B.9).

We now show that the support of a Mather measure is always contained in the Aubry set $\tilde{\mathcal{A}}$. In particular, the support of a Mather measure is compact, and any projected Mather measure has its support contained in \mathcal{A} .

Theorem B.22. *For every Mather measure $\tilde{\mu}$, we have $\text{supp } \tilde{\mu} \subset \tilde{\mathcal{A}}$. Therefore $\tilde{\mu}$ has compact support, and the support of the projected Mather measure $\pi_{\#}\tilde{\mu}$ is contained in \mathcal{A} . It follows that the set of Mather measures, and the set of projected Mather measures are both convex and compact in the weak topology.*

Proof. By the definition of the Aubry set B.13, we have to show that $\text{supp } \tilde{\mu} \subset \mathcal{L}(u)$, for every critical subsolution $u : M \rightarrow \mathbb{R}$. By Proposition B.11, for every $(x, v) \in TM$, we have

$$\partial^+ u(x, v) \leq L(x, v) + c(H), \quad (\text{B.10})$$

where $c(H)$ is the critical value for H . If we integrate this inequality for a Mather measure, we get

$$\int_{TM} \partial^+ u d\tilde{\mu} \leq \int_{TM} L + c(H) d\tilde{\mu} = 0.$$

But the left hand side is ≥ 0 , since $\tilde{\mu}$ is closed, see Lemma B.17. Therefore the inequality (B.10) is an equality $\tilde{\mu}$ -almost everywhere. In other words, the set $\mathcal{L}(u)$ is of full $\tilde{\mu}$ -measure, but as we have shown that this set $\mathcal{L}(u)$ is closed, we get $\text{supp } \tilde{\mu} \subset \mathcal{L}(u)$, as was required.

Now that we know that all minimizing measure have support in the compact set $\tilde{\mathcal{A}}$, we can characterize the set of minimizing measures as the set of probability measures $\tilde{\mu}$ on $\tilde{\mathcal{A}}$ such that $\int_{\tilde{\mathcal{A}}} L(x, v) d\tilde{\mu} = -c(H)$, and $\int_{\tilde{\mathcal{A}}} dx f(v) d\tilde{\mu} = 0$, for every C^1 function $f : M \rightarrow \mathbb{R}$. Each one of these constraints is a closed and convex condition on $\tilde{\mu}$. This finishes the proof. \square

We end this section by extending to the current setting the notion of Mather set and by proving that it is a uniqueness set for the critical equation.

Definition B.23. *The projected Mather set is the subset of M defined as*

$$\mathcal{M} := \overline{\bigcup_{\tilde{\mu} \in \tilde{\mathfrak{M}}_0(L)} \text{supp}(\pi_{\#} \tilde{\mu})}.$$

The following holds:

Theorem B.24. *The projected Mather set \mathcal{M} is a closed subset of the projected Aubry set \mathcal{A} . Moreover, it is a uniqueness set for the critical equation (3.4), i.e. two critical solutions that coincide on \mathcal{M} coincide on the whole manifold M .*

Proof. The fact that \mathcal{M} is a closed subset of \mathcal{A} follows from its definition and from Theorem B.22. If two critical solutions coincide on \mathcal{M} , then they coincide on every Mather class of \mathcal{A} in view of Theorem B.20 and Proposition B.8, and hence on \mathcal{A} . The conclusion follows since \mathcal{A} is a uniqueness set for the critical equation by Theorem B.6. \square

REFERENCES

- [1] V. BANGERT, *Minimal measures and minimizing closed normal one-currents*, Geom. Funct. Anal., 9 (1999), pp. 413–427.
- [2] M. BARDI AND I. CAPUZZO-DOLCETTA, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*, Systems & Control: Foundations & Applications, Birkhäuser Boston Inc., Boston, MA, 1997. With appendices by Maurizio Falcone and Pierpaolo Soravia.
- [3] G. BARLES, *Solutions de viscosité des équations de Hamilton-Jacobi*, vol. 17 of Mathématiques & Applications (Berlin) [Mathematics & Applications], Springer-Verlag, Paris, 1994.
- [4] P. BERNARD, *Existence of $C^{1,1}$ critical sub-solutions of the Hamilton-Jacobi equation on compact manifolds*, Ann. Sci. École Norm. Sup. (4), 40 (2007), pp. 445–452.
- [5] ———, *The Lax-Oleinik semi-group: a Hamiltonian point of view*, Proc. Roy. Soc. Edinburgh Sect. A, 142 (2012), pp. 1131–1177.
- [6] G. BUTTAZZO, M. GIAQUINTA, AND S. HILDEBRANDT, *One-dimensional variational problems*, vol. 15 of Oxford Lecture Series in Mathematics and its Applications, The Clarendon Press Oxford University Press, New York, 1998. An introduction.

- [7] P. CANNARSA AND C. SINISTRARI, *Semiconcave functions, Hamilton-Jacobi equations, and optimal control*, Progress in Nonlinear Differential Equations and their Applications, 58, Birkhäuser Boston Inc., Boston, MA, 2004.
- [8] F. H. CLARKE, *Optimization and nonsmooth analysis*, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons Inc., New York, 1983. A Wiley-Interscience Publication.
- [9] G. CONTRERAS AND R. ITURRIAGA, *Global minimizers of autonomous Lagrangians*, 22^o Colóquio Brasileiro de Matemática. [22nd Brazilian Mathematics Colloquium], Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 1999.
- [10] G. CONTRERAS, R. ITURRIAGA, G. P. PATERNAIN, AND M. PATERNAIN, *Lagrangian graphs, minimizing measures and Mañé's critical values*, Geom. Funct. Anal., 8 (1998), pp. 788–809.
- [11] A. DAVINI AND A. SICONOLFI, *A generalized dynamical approach to the large time behavior of solutions of Hamilton-Jacobi equations*, SIAM J. Math. Anal., 38 (2006), pp. 478–502 (electronic).
- [12] A. DAVINI AND M. ZAVIDOVIQUE, *Weak KAM theory for nonregular commuting Hamiltonians*, Discrete Contin. Dyn. Syst. Ser. B, 18 (2013), pp. 57–94.
- [13] A. FATHI, *Weak KAM from a PDE point of view: viscosity solutions of the Hamilton-Jacobi equation and Aubry set*, Proc. Roy. Soc. Edinburgh Sect. A, 142 (2012), pp. 1193–1236.
- [14] ———, *Weak KAM Theorem in Lagrangian Dynamics, preliminary version 10*, Lyon. unpublished, June 15 2008.
- [15] A. FATHI AND E. MADERNA, *Weak KAM theorem on non compact manifolds*, NoDEA Nonlinear Differential Equations Appl., 14 (2007), pp. 1–27.
- [16] A. FATHI AND A. SICONOLFI, *Existence of C^1 critical subsolutions of the Hamilton-Jacobi equation*, Invent. Math., 155 (2004), pp. 363–388.
- [17] ———, *PDE aspects of Aubry-Mather theory for quasiconvex Hamiltonians*, Calc. Var. Partial Differential Equations, 22 (2005), pp. 185–228.
- [18] D. A. GOMES, *Generalized Mather problem and selection principles for viscosity solutions and Mather measures*, Adv. Calc. Var., 1 (2008), pp. 291–307.
- [19] R. ITURRIAGA AND H. SÁNCHEZ-MORGADO, *Limit of the infinite horizon discounted Hamilton-Jacobi equation*, Discrete Contin. Dyn. Syst. Ser. B, 15 (2011), pp. 623–635.
- [20] P.-L. LIONS, G. PAPANICOLAOU, AND S. VARADHAN, *Homogenization of Hamilton-Jacobi equation*. unpublished preprint, 1987.
- [21] R. MAÑÉ, *Generic properties and problems of minimizing measures of Lagrangian systems*, Nonlinearity, 9 (1996), pp. 273–310.
- [22] J. N. MATHER, *Action minimizing invariant measures for positive definite Lagrangian systems*, Math. Z., 207 (1991), pp. 169–207.
- [23] R. T. ROCKAFELLAR, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970.
- [24] A. SICONOLFI, *Hamilton-jacobi equations and weak kam theory*, in Encyclopedia of Complexity and Systems Science, 2009, pp. 4540–4561.

DIP. DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, P.LE ALDO MORO 2, 00185 ROMA, ITALY

E-mail address: `davini@mat.uniroma1.it`

UMPA, ENS-LYON, 46 ALLÉE D'ITALIE, 69364 LYON CEDEX 7, FRANCE

E-mail address: `albert.fathi@ens-lyon.fr`

CIMAT, VALENCIANA GUANAJUATO, MÉXICO 36000

E-mail address: `renato@cimat.mx`

IMJ-PRG (PROJET ANALYSE ALGÈBRIQUE), UPMC, 4, PLACE JUSSIEU, CASE 247, 75252 PARIS CEDEX 5, FRANCE

E-mail address: `zavidovique@math.jussieu.fr`