



HAL
open science

Layer potential operators for transmission problems on extension domains

Gabriel Claret, Michael Hinz, Anna Rozanova-Pierrat, Alexander Teplyaev

► **To cite this version:**

Gabriel Claret, Michael Hinz, Anna Rozanova-Pierrat, Alexander Teplyaev. Layer potential operators for transmission problems on extension domains. 2023. hal-04505158v1

HAL Id: hal-04505158

<https://hal.science/hal-04505158v1>

Preprint submitted on 14 Mar 2024 (v1), last revised 7 Jun 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Layer potential operators for transmission problems on extension domains

Gabriel Claret* Michael Hinz† Anna Rozanova-Pierrat‡ Alexander Teplyaev§

March 14, 2024

Abstract

We use the well-posedness of transmission problems on classes of two-sided Sobolev extension domains to give variational definitions for (boundary) layer potential operators and Neumann-Poincaré operators. These classes of domains contain Lipschitz domains, but also domains with fractal boundaries. Our formulation does not involve any measures on the boundary. We discuss basic properties of these operators and use them to generalize basic results in imaging beyond the Lipschitz case.

Keywords: Fractal boundaries; Layer potentials; Poincaré-Steklov operator; Neumann-Poincaré operator; Calderón projector; elliptic transmission problems; inverse problems.

Contents

1	Introduction	2
2	Boundary value problems on admissible domains	4
2.1	Admissible domains and traces	4
2.2	Orthogonality and isometries	6
2.3	Harmonic extensions	7
2.4	Neumann solutions	8
2.5	Normal derivatives	9
3	Poincaré-Steklov operators for admissible domains	10
4	Transmission problems for admissible domains	11
4.1	Two-sided admissible domains and jumps	12
4.2	Subspaces and orthogonality	13
4.3	Double layer potentials	15
4.4	Single layer potentials	16
4.5	Superposition	17

*CentraleSupélec, Université Paris-Saclay, France. Research included in the “Parcours Recherche” teaching program at CentraleSupélec.

†University of Bielefeld, Germany (mhinz@math.uni-bielefeld.de) Research supported in part by the DFG IRTG 2235: “Searching for the regular in the irregular: Analysis of singular and random systems” and by the DFG CRC 1283: “Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications”.

‡CentraleSupélec, Université Paris-Saclay, France (anna.rozanova-pierrat@centralesupelec.fr) Research supported in part by CNRS INSMI IEA (International Emerging Actions 2022) “Functional and applied analysis with fractal or non-Lipschitz boundaries”

§University of Connecticut, USA (teplyaev@uconn.edu) Research supported in part by the NSF DMS 1613025, 1950543, and the Simons and Fulbright Foundations. The project was partially conducted during a stay at CentraleSupélec, Université Paris-Saclay, whose kind hospitality is gratefully acknowledged.

5	Resolvent representations	18
5.1	Representations of single layer potentials	18
5.2	Representations of double layer potentials	20
6	Neumann-Poincaré operators for admissible domains	21
6.1	Neumann-Poincaré operators	21
6.2	Boundary layer potentials and Calderón projectors	22
6.3	Invertibility and isometries	24
7	Applications to imaging	26
7.1	Representation formula	26
7.2	Subdomain identification	27
A	Comments and proofs	28
A.1	Comments and proofs for Section 2	28

1 Introduction

In this note we study layer potential operators, Neumann-Poincaré operators, boundary layer operators and transmission problems for a fairly general class of rough (two-sided) domains, possibly with non-Lipschitz or even fractal boundaries.

In the classical theory of partial differential equations on smooth domains (boundary) layer potentials and Neumann-Poincaré operators are defined as convolutions with Green’s functions and their first derivatives, see for instance [39, 60, 67, 82, 87] or [5]. They provide explicit representation formulas for solutions of boundary value problems, and they are fundamental tools in inverse methods, numerical analysis and certain areas of spectral theory.

A systematic study of the classical approach to layer potentials in the context of Lipschitz domains was provided in [86], following seminal studies on boundary value problems, [50], and singular integrals, [19, 25]. Since then the Lipschitz case has become a standard level of generality for many applications, see for instance [3, 4, 27, 28, 31, 32, 48, 69, 73, 80]. A different approach was taken in [11], where the weak well-posedness of transmission problems was used to actually define layer potentials. In view of the well established use of Hilbert space methods in potential theory — see [26, 29, 37, 89] for classical sources and [16, 24, 41, 64] for later developments — these variational definitions in [11] are very natural.

Boundary layer operators on piecewise smooth, not necessarily Lipschitz domains were already studied in [67, Chapter 5]. Results for layer potentials in the context of Riemannian manifolds were obtained in [72], results for layer potentials on half-spaces with boundary data in Besov spaces in [12]. The research on the boundedness of singular integral operators in [19, 25, 86] culminated in the comprehensive study [71] of such operators on uniformly rectifiable sets, [34, 35]. Uniformly rectifiable sets in \mathbb{R}^n are Ahlfors $(n-1)$ -regular closed subsets having “big pieces of Lipschitz images”, see [70, Definition 5.10.1 and (5.10.2)], and the class of such sets is basically characterized by the L^2 -boundedness of singular integral operators, [70, Theorem 5.10.2]. A different major stream of research focused on the behaviour of harmonic functions on a domain Ω and harmonic measures on its boundary $\partial\Omega$, see [30, 49, 50, 53] and the later references [2, 13, 20, 33]. A main notion is that of an NTA domain, [50, Section 3]. The class of NTA domains includes all Lipschitz domains, but also certain domains with fractal boundaries such as quasidisks, [8, 50, 51, 75], and amongst them examples where different parts of the boundary have different Hausdorff dimensions. A short introduction to harmonic measures and NTA domains is given in [81]. Another question about a domain Ω is whether it is a Sobolev extension domain, i.e., whether there is a bounded linear extension operator taking Sobolev functions on Ω into Sobolev functions on \mathbb{R}^n . In [52] it was shown that uniform domains, [66, 84], and, more generally, (ε, δ) -domains, [52], are Sobolev extension domains. Any NTA domain is uniform, see for instance [8, Theorem 2.15], and any uniform domain with uniformly rectifiable boundary is NTA, [8, Theorem 1.1].

We are interested in boundary value problems involving rough boundaries that may be fractal and may even have different parts of different Hausdorff dimensions. Well-known more specific references on linear elliptic partial differential equations on fractal domains are [62, 63, 65, 75, 85, 88], some further articles close to the present note are [10, 17, 18, 22, 43]. The highly active research around domains with rough boundaries motivates an extension of the variational approach in [11] beyond the Lipschitz case, and potential follow-up questions about convergence and optimization make it desirable to have a generalization which is flexible and rather easy to handle.

The purpose of this note is to propose a lightweight and streamlined generalization. The domains we consider are “first order” Sobolev extension domains Ω in \mathbb{R}^n , [44, 52], having a boundary $\partial\Omega$ of positive capacity, [68]. Basic examples in the planar case are quasidisks or complements of large enough Cantor sets. Domains having sharp inward or outward cusps or “collapsing” boundaries are not covered, in particular, certain fractal trees, [1], do not fall within the scope of this note. We use established results, [15], to introduce trace and extension operators from the domain Ω to its boundary $\partial\Omega$ in the context of Sobolev spaces $H^1(\Omega)$ respectively homogeneous Sobolev spaces $\dot{H}^1(\Omega)$. On the boundary $\partial\Omega$ we use corresponding abstract trace spaces $\mathcal{B}(\partial\Omega)$ respectively $\dot{\mathcal{B}}(\partial\Omega)$, endowed with natural trace norms. This approach is *measure free* in the sense that we do not specify any “surface measure” on $\partial\Omega$ and, in contrast to [86], do not consider L^2 -spaces on $\partial\Omega$. Instead, we make a systematic use of the trace spaces $\mathcal{B}(\partial\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega)$ and their duals $(\mathcal{B}(\partial\Omega))'$ respectively $(\dot{\mathcal{B}}(\partial\Omega))'$. As a consequence, we observe a variety of *natural isometries*, and the preservation of these isometries might be considered a guiding theme of this note. In the special case of a Lipschitz domain Ω the spaces $\mathcal{B}(\partial\Omega)$ and $(\mathcal{B}(\partial\Omega))'$ are the familiar spaces $H^{1/2}(\partial\Omega)$ and $H^{-1/2}(\partial\Omega)$, up to equivalent norms; the “dotted” versions correspond to their homogeneous counterparts. In cases where the boundary $\partial\Omega$ carries a measure satisfying certain scaling conditions, the trace spaces $\mathcal{B}(\partial\Omega)$ admit explicit descriptions in terms of equivalent Besov space norms, see [54, 55] and the comments in [46, Section 5]. In general the present measure free formulation does not give any explicit norm representations for the trace spaces $\mathcal{B}(\partial\Omega)$ and may therefore not be sufficient to discuss regularity features. But it works under minimal assumptions on the boundary, and this seems particularly useful in view of possible later studies of convergence and compactness properties as in [45–47].

To fix spaces, operators and notations, we briefly survey Dirichlet and Neumann problems in the weak (variational) sense for one-sided domains in Section 2. They are formulated for the operator $\Delta - 1$ in the context of the spaces $H^1(\Omega)$ and for the operator Δ in the context of the spaces $\dot{H}^1(\Omega)$. The use of the spaces $\dot{H}^1(\Omega)$ may have the disadvantage that constants are factored out, but it has the advantage that arguments can proceed very much in parallel to the inhomogeneous case. In a short Section 3 we discuss basic features of related Poincaré-Steklov operators. Suitable two-sided domains are then introduced in Section 4. We roughly speaking require both the inner domain Ω and the outer domain $\mathbb{R}^n \setminus \bar{\Omega}$ to be non-empty and Sobolev extension domains and we assume that the separating boundary $\partial\Omega$ has zero Lebesgue measure. In the homogeneous case we additionally assume Ω to be bounded. Similarly as in [11], the weak well-posedness of transmission problems is used to give variational definitions of double and single layer potential operators for these two-sided domains, now measure free, see Subsections 4.3 and 4.4. Also their basic properties are discussed in these subsections. In Section 5 resolvent representations of double and single layer potential operators are surveyed. Neumann-Poincaré operators are well studied for smooth or Lipschitz domains, [5, 56, 60], their spectral properties are known to reflect regularity properties of $\partial\Omega$, such as its compactness in the C^1 -case, [39], or the presence of continuous non-real spectrum if $\partial\Omega$ has a corner [5, 21]. In Section 6 we define Neumann-Poincaré operators \mathcal{K} and $\dot{\mathcal{K}}$ on the trace spaces $\mathcal{B}(\partial\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega)$, respectively. As in the Lipschitz case, [80], they are bounded operators and satisfy the well-known jump relations, see Theorems 6.1 and 6.2. Following [80], we define boundary layer potential operators associated with $(\Delta - 1)$ and Δ in (6.1) and (6.5), respectively, and using these operators, obtain generalizations of the classical Calderón projectors, see Theorems 6.4 and 6.6. As

in [80] the boundary layer potentials give rise to equivalent Hilbert space norms on the trace spaces and their duals. These norms make the single and double layer potentials isometries, see Theorem 6.9 in Subsection 6.3. Again similarly as in [80], we observe that $(\pm\frac{1}{2}I + \mathcal{K})$ are coercive contractions and in particular, bijective, Theorem 6.9 (iii). The contractivity of $(\pm\frac{1}{2}I + \mathcal{K})$ ensures the convergence of the associated Neumann series, [74]: Recovering the jump in trace $f \in \mathcal{B}(\partial\Omega)$ of a transmission solution for $(\Delta - 1)$ with zero jump in normal derivative from its exterior trace $\varphi \in \mathcal{B}(\partial\Omega)$ amounts to solving the boundary integral equation of the second kind $-\frac{1}{2}f - \mathcal{K}f = \varphi$ in $\mathcal{B}(\partial\Omega)$, and its unique solution f is given by the Neumann series $f = \sum_{\ell=0}^{+\infty} (\frac{1}{2}I + \mathcal{K}_{\partial\Omega})^\ell \varphi$ which converges in $\mathcal{B}(\partial\Omega)$. Similarly one can invert the operator $-\frac{1}{2}I + \mathcal{K}_{\partial\Omega}^*$ on $(\mathcal{B}(\partial\Omega))'$, it recovers the jump in normal derivative of a solution with no jump in trace from its exterior normal derivative, see [80, Theorem 3.2, p.741]. As in the classical case [38, 39, 86] Theorem 6.9 (iii) can be used to see that the real spectra of $\mathcal{K}_{\partial\Omega}$ and $\mathcal{K}_{\partial\Omega}^*$ are included in $(-\frac{1}{2}, \frac{1}{2})$, Theorem 6.11. These results have homogeneous counterparts, see Theorems 6.10 and 6.12.

As an application we generalize two results from [3]. There they were shown for Lipschitz domains, here we establish them for two-sided Sobolev extension domains. Theorem 7.1 is a boundary representation formula for the unique weak solution of a specific transmission problem for the Laplacian, it generalizes [3, Theorem 2.17], see also [57, 58]. Theorem 7.2 is a uniqueness results for subdomain identification through a single boundary measurement in the monotone case, it generalizes a part of [3, Theorem A.7, p.220], see also [14].

Acknowledgements

The authors are grateful to David Hewett and Simon N. Chandler-Wilde for their helpful advice and discussions related to the observations in this note.

2 Boundary value problems on admissible domains

2.1 Admissible domains and traces

Let Ω be a nonempty open subset of \mathbb{R}^n . As usual, we write $H^1(\Omega)$ for the Hilbert space of all $u \in L^2(\Omega)$ such that $\nabla u \in L^2(\Omega, \mathbb{R}^n)$ and having the scalar product

$$\langle u, v \rangle_{H^1(\Omega)} = \int_{\Omega} \nabla u \nabla v \, dx + \int_{\Omega} uv \, dx; \quad (2.1)$$

here ∇u is interpreted in distributional sense. We write $\dot{H}^1(\Omega)$ for the Hilbert space formed by the vector space of all $u \in L^2_{\text{loc}}(\Omega)$ with $\nabla u \in L^2(\Omega, \mathbb{R}^n)$ modulo locally constant functions, endowed with the scalar product

$$\langle u, v \rangle_{\dot{H}^1(\Omega)} = \int_{\Omega} \nabla u \nabla v \, dx. \quad (2.2)$$

As usual, we call Ω a domain if it is connected. Details on $\dot{H}^1(\Omega)$ in the case of a domain Ω can be found in [24, Section 2.2.4], [37] or [68, Section 1.1.13]; the generalization to open subsets with multiple connected components is straightforward. We agree to use the notations (2.1) and (2.2) whenever the right-hand side makes sense.

We use the notions of capacity, quasi continuous representatives and quasi everywhere (q.e.) valid statements exclusively with respect to the space $H^1(\mathbb{R}^n)$; background may be found in [41, Section 2.1] or [68, Section 7.2]. Each $u \in H^1(\mathbb{R}^n)$ has a quasi continuous representative \tilde{u} . See for instance [24, Theorem 2.3.4] or [68, Section 7.2.4]. For each class $u \in \dot{H}^1(\mathbb{R}^n)$ we can find a quasi continuous representative of a representative modulo constants. This can for instance be seen from [24, Theorem 2.3.4] together with [24, Theorems 2.2.12

and 2.2.13], see (A.2) and (A.3) in the Appendix. For brevity we will again call it a quasi continuous representative of u .

Recall that Ω is called an H^1 -extension domain, [44, 52], if there is a bounded linear extension operator $E_\Omega : H^1(\Omega) \rightarrow H^1(\mathbb{R}^n)$. We say that a domain $\Omega \subset \mathbb{R}^n$ is H^1 -admissible if it is an H^1 -extension domain and its boundary $\partial\Omega$ has positive capacity. In this case we write $\mathcal{B}(\partial\Omega)$ for the vector space of all q.e. equivalence classes of pointwise restrictions $\tilde{u}|_{\partial\Omega}$ of quasi continuous representatives \tilde{u} of classes $u \in H^1(\mathbb{R}^n)$. The following result on the existence and uniqueness of natural trace operators from $H^1(\Omega)$ onto $\mathcal{B}(\partial\Omega)$ is immediate from [15, Theorem 6.1, Remark 6.2 and Corollary 6.3].

Proposition 2.1. *Let Ω be H^1 -admissible. Given $u \in H^1(\Omega)$, let $\text{Tr}_i u := (E_\Omega u)^\sim|_{\partial\Omega}$ be the restriction of any quasi continuous representative $(E_\Omega u)^\sim$ of $E_\Omega u$. Then $\text{Tr}_i : H^1(\Omega) \rightarrow \mathcal{B}(\partial\Omega)$ is a well-defined linear surjection.*

We use the subscript i in Tr_i because we will later discuss two-sided domains and Tr_i as defined here will play the role of an *inner* trace operator with respect to Ω .

We say that a domain $\Omega \subset \mathbb{R}^n$ is an \dot{H}^1 -extension domain if there is a bounded linear extension operator $\dot{E}_\Omega : \dot{H}^1(\Omega) \rightarrow \dot{H}^1(\mathbb{R}^n)$, and we call it \dot{H}^1 -admissible if it is an \dot{H}^1 -extension domain and $\partial\Omega$ is compact and of positive capacity. If so, we write $\dot{\mathcal{B}}(\partial\Omega)$ for the vector space modulo constants of all q.e. equivalence classes of pointwise restrictions $\tilde{w}|_{\partial\Omega}$ of quasi continuous representatives \tilde{w} of classes $u \in \dot{H}^1(\mathbb{R}^n)$.

A variant of Proposition 2.1 gives natural trace operators from $\dot{H}^1(\Omega)$ onto $\dot{\mathcal{B}}(\partial\Omega)$.

Proposition 2.2. *Let Ω be \dot{H}^1 -admissible. Given $u \in \dot{H}^1(\Omega)$, let \tilde{w} be a quasi continuous representative of $\dot{E}_\Omega u$. Defining $\dot{\text{Tr}}_i u$ to be the $\dot{\mathcal{B}}(\partial\Omega)$ -equivalence class of $\tilde{w}|_{\partial\Omega}$, we obtain a well-defined linear surjection $\dot{\text{Tr}}_i : \dot{H}^1(\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$.*

Proof. Suppose that $u \in \dot{H}^1(\Omega)$ and let $w \in \dot{E}_\Omega u$ be a representative of $\dot{E}_\Omega u$ modulo constants. If Ω is bounded, let B be an open ball containing $\overline{\Omega}$ and let $\chi \in C_c^\infty(B)$ be such that $0 \leq \chi \leq 1$ and $\chi \equiv 1$ on $\overline{\Omega}$. Then $v := \chi w$ is an element of $H^1(\mathbb{R}^n)$, and by [15, Theorem 6.1, Remark 6.2 and Corollary 6.3] the restriction $\tilde{w}|_{\partial\Omega} = \tilde{v}|_{\partial\Omega}$ is uniquely determined in the q.e. sense. Clearly its class modulo constants in $\dot{\mathcal{B}}(\partial\Omega)$ does not depend on the choice of w . If instead the set $\mathbb{R}^n \setminus \Omega$ is bounded, let B be an open ball containing it. Then $\Omega \cap B$ is a bounded \dot{H}^1 -extension domain, the elementary arguments are spelled out in Lemma A.1. Since $u|_{\Omega \cap B} \in \dot{H}^1(\Omega \cap B)$, we can proceed as before. \square

Remark 2.3.

- (i) *Suppose that Ω is a bounded H^1 -extension domain. Then the vector space $\dot{H}^1(\Omega)$ is isomorphic to the space of all $u \in H^1(\Omega)$ with $\int_\Omega u(x) dx = 0$ and by Poincaré's inequality (2.1) and (2.2) are equivalent scalar products on this space. In particular, the vector spaces $H^1(\Omega)$ and $\dot{H}^1(\Omega) \oplus \mathbb{R}$ are isomorphic.*
- (ii) *For a bounded domain Ω which is both H^1 - and \dot{H}^1 -admissible the vector spaces $\mathcal{B}(\partial\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega) \oplus \mathbb{R}$ are isomorphic.*

Examples 2.4.

- (i) *For $n \geq 2$ any (ε, δ) -domain $\Omega \subset \mathbb{R}^n$ is an H^1 -extension domain, [52, Theorem 1], and any (ε, ∞) -domain $\Omega \subset \mathbb{R}^n$ is an \dot{H}^1 -extension domain, [52, Theorem 2].*
- (ii) *For $n \geq 2$ any (ε, ∞) -domain $\Omega \subset \mathbb{R}^n$ with $\mathbb{R}^n \setminus \overline{\Omega}$ nonempty is H^1 -admissible, and if one of the two open sets is bounded, it is also \dot{H}^1 -admissible.*
- (iii) *For $n = 1$ any interval (a, b) with a or b finite is H^1 -admissible, and if both are finite, also \dot{H}^1 -admissible. For $n \geq 2$ the domain $\Omega = \mathbb{R}^n \setminus \{0\}$ is not H^1 -admissible.*

Remark 2.5. *Classical references on the spaces $\dot{H}^1(\Omega)$ are [37] and [68, Sections 1.1.2 and 1.1.13], a more recent discussion may be found in [42, Section II.6]. In these references different symbols are used to denote these spaces. For the full space case $\Omega = \mathbb{R}^n$ the notation $\dot{H}^1(\mathbb{R}^n)$ is established, see for instance [9, Section 1.3] and [83, Chapter 5]. Since the domains we consider are H^1 - respectively \dot{H}^1 -extension domains, we decided to follow this notation for the full space case and to use $\dot{H}^1(\Omega)$.*

2.2 Orthogonality and isometries

We center our discussion of boundary value problems around restrictions, duals and inverses of trace operators.

Let $\Omega \subset \mathbb{R}^n$ be an H^1 -admissible domain. We write $H_0^1(\Omega)$ for the closure in $H^1(\Omega)$ of the set $C_c^\infty(\Omega)$ of infinitely differentiable functions with compact support in Ω . Let $V_1(\Omega)$ denote the orthogonal complement of $H_0^1(\Omega)$ in $H^1(\Omega)$,

$$H^1(\Omega) = H_0^1(\Omega) \oplus V_1(\Omega). \quad (2.3)$$

The functions in $V_1(\Omega)$ are called *1-harmonic* in Ω . We write $\text{tr}_i := \text{Tr}_i|_{V_1(\Omega)}$ for the restriction of Tr_i defined in Proposition 2.1 to $V_1(\Omega)$. The following is well known.

Lemma 2.6. *Let $\Omega \subset \mathbb{R}^n$ be an H^1 -admissible domain.*

- (i) *The space $H_0^1(\Omega)$ is the kernel of Tr_i , that is, $H_0^1(\Omega) = \ker \text{Tr}_i$.*
- (ii) *Endowed with the norm*

$$\|f\|_{\mathcal{B}(\partial\Omega)} := \min\{\|v\|_{H^1(\Omega)} \mid v \in H^1(\Omega) \text{ and } \text{Tr}_i v = f\}, \quad (2.4)$$

the space $\mathcal{B}(\partial\Omega)$ is a Hilbert space.

- (iii) *With respect to $\|\cdot\|_{\mathcal{B}(\partial\Omega)}$ the trace operator trace Tr_i is bounded with operator norm one. Its restriction $\text{tr}_i : V_1(\Omega) \rightarrow \mathcal{B}(\partial\Omega)$ to $V_1(\Omega)$ is an isometry and onto.*

Details on statement (i) can be found in Appendix A.1, statements (ii) and (iii) are direct consequences.

We write $(\mathcal{B}(\partial\Omega))'$ and $(H^1(\Omega))'$ for the dual spaces of $\mathcal{B}(\partial\Omega)$ and $H^1(\Omega)$, and we use the notation

$$\langle \cdot, \cdot \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \quad \text{and} \quad \langle \cdot, \cdot \rangle_{(H^1(\Omega))', H^1(\Omega)}$$

for the corresponding dual pairings.

Remark 2.7. *For any $f \in \mathcal{B}(\partial\Omega)$ the assignment $\iota(f)(h) := \langle f, h \rangle_{\mathcal{B}(\partial\Omega)}$, $h \in \mathcal{B}(\partial\Omega)$, defines an isometric isomorphism ι from $\mathcal{B}(\partial\Omega)$ onto $(\mathcal{B}(\partial\Omega))'$. The dual pairing is defined by $\langle g, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} := \langle \iota^{-1}(g), f \rangle_{\mathcal{B}(\partial\Omega)} = \langle g, \iota(f) \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$, $f \in \mathcal{B}(\partial\Omega)$, $g \in (\mathcal{B}(\partial\Omega))'$. We may identify $\mathcal{B}(\partial\Omega)$ with its image $\iota(\mathcal{B}(\partial\Omega)) \subset (\mathcal{B}(\partial\Omega))'$ under ι .*

Remark 2.8. *If $(V_1(\Omega))'$ denotes the dual of the closed subspace $V_1(\Omega)$ of $H^1(\Omega)$, then $(H^1(\Omega))' \subset (V_1(\Omega))'$ by restriction. However, by the Riesz representation theorem any $w \in (V_1(\Omega))'$ is represented as $w = \langle v, \cdot \rangle_{H^1(\Omega)}$ with some suitable $v \in V_1(\Omega)$. The orthogonal decomposition (2.3) then implies that w automatically extends to a unique bounded linear functional $w' \in (H^1(\Omega))'$ on all of $H^1(\Omega)$ and zero on $H_0^1(\Omega)$, and this extension is an isometry, $\|w'\|_{(H^1(\Omega))'} = \|w\|_{(V_1(\Omega))'}$. We agree to make silent use of this extension: We write w to denote w' and use $\|\cdot\|_{(H^1(\Omega))'}$ in place of $\|\cdot\|_{(V_1(\Omega))'}$ on $(V_1(\Omega))'$.*

By Remark 2.8 the dual $\text{tr}_i^* : (\mathcal{B}(\partial\Omega))' \rightarrow (V_1(\Omega))'$ of tr_i is seen to be characterized by

$$\langle g, \text{Tr}_i v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle \text{tr}_i^* g, v \rangle_{(H^1(\Omega))', H^1(\Omega)}, \quad v \in H^1(\Omega), \quad g \in (\mathcal{B}(\partial\Omega))'. \quad (2.5)$$

Corollary 2.9. *Let $\Omega \subset \mathbb{R}^n$ be H^1 -admissible. Then the operator $\text{tr}_i^* : (\mathcal{B}(\partial\Omega))' \rightarrow (V_1(\Omega))'$ is an isometry, $\|\text{tr}_i^* g\|_{(H^1(\Omega))'} = \|g\|_{(\mathcal{B}(\partial\Omega))'}$, $g \in (\mathcal{B}(\partial\Omega))'$, and onto.*

Now let $\Omega \subset \mathbb{R}^n$ be an \dot{H}^1 -admissible domain. The space

$$\dot{V}_0(\Omega) := \left\{ u \in \dot{H}^1(\Omega) \mid \int_{\Omega} \nabla u \nabla v \, dx = 0 \text{ for all } v \in C_c^\infty(\Omega) \right\} \quad (2.6)$$

is a closed subspace of $\dot{H}^1(\Omega)$. We write $\dot{\text{tr}}_i := \dot{\text{Tr}}_i|_{\dot{V}_0(\Omega)}$ for the restriction of $\dot{\text{Tr}}_i$ to $\dot{V}_0(\Omega)$. The following counterpart of Lemma 2.6 holds; a proof of (i) is given in Appendix A.1.

Lemma 2.10. *Let $\Omega \subset \mathbb{R}^n$ be an \dot{H}^1 -admissible domain.*

- (i) *The kernel $\ker \dot{\text{Tr}}_i|_{\dot{H}^1(\Omega)}$ is the orthogonal complement $(\dot{V}_0(\Omega))^\perp$ of $\dot{V}_0(\Omega)$ in $\dot{H}^1(\Omega)$.*
- (ii) *Endowed with the norm $\|f\|_{\dot{\mathcal{B}}(\partial\Omega)} := \min\{\|v\|_{\dot{H}^1(\Omega)} \mid v \in \dot{H}^1(\Omega) \text{ and } \dot{\text{Tr}}_i v = f\}$, the space $\dot{\mathcal{B}}(\partial\Omega)$ is a Hilbert space.*
- (iii) *With respect to $\|\cdot\|_{\dot{\mathcal{B}}(\partial\Omega)}$ the operator $\dot{\text{Tr}}_i$ is bounded with operator norm one. Its restriction $\dot{\text{tr}}_i : \dot{V}_0(\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$ to $\dot{V}_0(\Omega)$ is an isometry and onto.*

We write $(\dot{H}^1(\Omega))'$, $(\dot{V}_0(\Omega))'$ and $(\dot{\mathcal{B}}(\partial\Omega))'$ for the dual spaces of $\dot{H}^1(\Omega)$, $\dot{V}_0(\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega)$. With a similar agreement as in Remark 2.8 the dual $\dot{\text{tr}}_i^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow (\dot{V}_0(\Omega))'$ of $\dot{\text{tr}}_i$ is now seen to be characterized by

$$\langle g, \dot{\text{Tr}}_i v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} = \langle \dot{\text{tr}}_i^* g, v \rangle_{(\dot{H}^1(\Omega))', \dot{H}^1(\Omega)}, \quad v \in \dot{H}^1(\Omega), \quad g \in (\dot{\mathcal{B}}(\partial\Omega))'.$$

Corollary 2.11. *Let $\Omega \subset \mathbb{R}^n$ be \dot{H}^1 -admissible. Then the operator $\dot{\text{tr}}_i^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow (\dot{V}_0(\Omega))'$ is an isometry, $\|\dot{\text{tr}}_i^* g\|_{(\dot{H}^1(\Omega))'} = \|g\|_{(\dot{\mathcal{B}}(\partial\Omega))'}$, $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, and onto.*

2.3 Harmonic extensions

Suppose that $\Omega \subset \mathbb{R}^n$ is an H^1 -admissible domain. Given $f \in \mathcal{B}(\partial\Omega)$, an element u of $H^1(\Omega)$ is called a *weak solution* of the Dirichlet problem

$$\begin{cases} -\Delta u + u = 0 & \text{in } \Omega \\ u|_{\partial\Omega} = f \end{cases} \quad (2.7)$$

if $\text{Tr}_i u = f$ and $\langle u, v \rangle_{H^1(\Omega)} = 0$ for all $v \in C_c^\infty(\Omega)$. The symbol $u|_{\partial\Omega}$ in the formal problem (2.7) stands for the restriction of u to the boundary; in the context of weak solutions it is made rigorous through the trace condition. For any $f \in \mathcal{B}(\partial\Omega)$ the Dirichlet problem (2.7) has a unique weak solution u^f , namely $u^f = \text{tr}_i^{-1} f \in V_1(\Omega)$; this is well known and follows using (2.3). By Lemma 2.6 the operator $\text{tr}_i^{-1} : \mathcal{B}(\partial\Omega) \rightarrow V_1(\Omega)$ is an isometry.

Suppose that Ω is an \dot{H}^1 -admissible domain. Given $f \in \dot{\mathcal{B}}(\partial\Omega)$, we call an element u of $\dot{H}^1(\Omega)$ a *weak solution in the \dot{H}^1 -sense* of the Dirichlet problem

$$\begin{cases} -\Delta u = 0 & \text{in } \Omega \\ u|_{\partial\Omega} = f \end{cases} \quad (2.8)$$

if $\dot{\text{Tr}}_i u = f$ and $\langle u, v \rangle_{\dot{H}^1(\Omega)} = 0$ for all $v \in C_c^\infty(\Omega)$. For any $f \in \dot{\mathcal{B}}(\partial\Omega)$ the Dirichlet problem (2.8) has the unique weak solution $u^f = \dot{\text{tr}}_i^{-1} f \in \dot{V}_0(\Omega)$ in the \dot{H}^1 -sense. By Lemma 2.10 the operator $\dot{\text{tr}}_i^{-1}$ is an isometry.

Remark 2.12.

- (i) If Ω is bounded and $f \in \mathcal{B}(\partial\Omega)$, then $w \in H^1(\Omega)$ is called a weak solution of (2.8) if $\text{Tr}_i w = f$ and $\langle w, v \rangle_{\dot{H}^1(\Omega)} = 0$ for all $v \in C_c^\infty(\Omega)$. It is well known that for any $f \in \mathcal{B}(\partial\Omega)$ there is a unique weak solution w^f in this sense of (2.8) and that w^f is an element of the space $V_0(\Omega)$ of $u \in H^1(\Omega)$ with $\langle u, v \rangle_{\dot{H}^1(\Omega)} = 0$ for all $v \in H_0^1(\Omega)$.
- (ii) If Ω is bounded and $f \in \mathcal{B}(\partial\Omega)$, then the equivalence class $[f]$ of f modulo constants is in $\dot{\mathcal{B}}(\partial\Omega)$, cf. Remark 2.3 (ii), and the unique weak solution $u^{[f]} \in \dot{H}^1(\Omega)$ in the \dot{H}^1 -sense of (2.8) with this class $[f]$ in place of f contains exactly one representative $w \in u^{[f]}$ modulo constants such that $\tilde{w}|_{\partial\Omega} = f$ q.e. This particular representative w is exactly the w^f from (i).
- (iii) In [79] a different type of function spaces based on [37] was used in order to handle Dirichlet boundary conditions for unbounded domains without losing constants. However, to avoid technicalities in later sections we decided to accept a loss of constants.

2.4 Neumann solutions

Suppose $\Omega \subset \mathbb{R}^n$ be an H^1 -admissible domain. Given $g \in (\mathcal{B}(\partial\Omega))'$, we call $u \in H^1(\Omega)$ a weak solution of the Neumann problem

$$\begin{cases} -\Delta u + u &= 0 & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} &= g \end{cases} \quad (2.9)$$

if for all $v \in H^1(\Omega)$ we have

$$\langle u, v \rangle_{H^1(\Omega)} = \langle g, \text{Tr}_i v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}. \quad (2.10)$$

The symbol $\frac{\partial u}{\partial \nu}$ in the formal problem (2.9) stands for the (interior) normal derivative; in the context of weak solutions it is implicitly made rigorous by (2.10). It is well known and easily seen from the Riesz representation theorem that for any $g \in (\mathcal{B}(\partial\Omega))'$ the Neumann problem (2.9) has a unique weak solution u_g , it is an element of $V_1(\Omega)$. We write $\mathcal{N}_1 : (\mathcal{B}(\partial\Omega))' \rightarrow V_1(\Omega)$, $\mathcal{N}_1 g := u_g$ for the linear operator taking a given element g of $(\mathcal{B}(\partial\Omega))'$ into the unique weak solution of (2.9). By (2.5) we have

$$\langle \mathcal{N}_1 g, v \rangle_{H^1(\Omega)} = \langle \text{tr}_i^* g, v \rangle_{(H^1(\Omega))', H^1(\Omega)}, \quad v \in H^1(\Omega),$$

hence $\|\mathcal{N}_1 g\|_{H^1(\Omega)} = \|\text{tr}_i^* g\|_{(H^1(\Omega))'} = \|g\|_{(\mathcal{B}(\partial\Omega))'}$, $g \in (\mathcal{B}(\partial\Omega))'$, that is, \mathcal{N}_1 is an isometry.

Corollary 2.13. *Let Ω be H^1 -admissible. The linear operator $\text{tr}_i \circ \mathcal{N}_1 : (\mathcal{B}(\partial\Omega))' \rightarrow \mathcal{B}(\partial\Omega)$ is an isometry and onto. It satisfies*

$$\langle g, \text{tr}_i \circ \mathcal{N}_1 h \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle g, h \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle h, \text{tr}_i \circ \mathcal{N}_1 g \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}, \quad g, h \in (\mathcal{B}(\partial\Omega))'.$$

Proof. The first statement is clear from (2.10), (2.5) and the preceding. The second follows because for all $g, h \in (\mathcal{B}(\partial\Omega))'$ we have

$$\begin{aligned} \langle g, \text{tr}_i \circ \mathcal{N}_1 h \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} &= \langle g, \text{tr}_i u_h \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \\ &= \langle \text{tr}_i^* g, u_h \rangle_{(H^1(\Omega))', H^1(\Omega)} = \langle u_g, u_h \rangle_{H^1(\Omega)}. \end{aligned}$$

□

Now suppose that Ω is \dot{H}^1 -admissible. Given $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, we call $u \in \dot{H}^1(\Omega)$ a weak solution in the \dot{H}^1 -sense of the Neumann problem

$$\begin{cases} -\Delta u &= 0 & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} \Big|_{\partial\Omega} &= g \end{cases} \quad (2.11)$$

if for all $v \in \dot{H}^1(\Omega)$ we have

$$\langle u, v \rangle_{\dot{H}^1(\Omega)} = \langle g, \dot{\text{Tr}}_i v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}. \quad (2.12)$$

For any $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ the Neumann problem (2.11) has a unique weak solution u_g in the \dot{H}^1 -sense, and u_g is an element of $\dot{V}_0(\Omega)$ defined in (2.6). We write $\dot{\mathcal{N}}_0 : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{V}_0(\Omega)$, $\dot{\mathcal{N}}_0 g := u_g$, for the linear operator taking a given element of $(\dot{\mathcal{B}}(\partial\Omega))'$ into u_g . Similarly as before we see that $\dot{\mathcal{N}}_0$ is an isometry, $\|\dot{\mathcal{N}}_0 g\|_{\dot{H}^1(\Omega)} = \|\dot{\text{tr}}_i^* g\|_{(\dot{H}^1(\Omega))'} = \|g\|_{(\dot{\mathcal{B}}(\partial\Omega))'}$, $g \in (\dot{\mathcal{B}}(\partial\Omega))'$.

Corollary 2.14. *Let Ω be \dot{H}^1 -admissible. The linear operator $\text{tr}_i \circ \dot{\mathcal{N}}_0 : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{\mathcal{B}}(\partial\Omega)$ is an isometry and onto. It satisfies*

$$\langle g, \dot{\text{tr}}_i \circ \dot{\mathcal{N}}_0 h \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} = \langle g, h \rangle_{(\dot{\mathcal{B}}(\partial\Omega))'} = \langle h, \dot{\text{tr}}_i \circ \dot{\mathcal{N}}_0 g \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}, \quad g, h \in (\dot{\mathcal{B}}(\partial\Omega))'.$$

2.5 Normal derivatives

Abstract normal derivatives have been defined and studied by various authors in different contexts, see for instance [26, p. 218], [64, Section 3.2] and [61]. We formulate a variant of these definitions which suits our purposes.

Suppose $\Omega \subset \mathbb{R}^n$ is H^1 -admissible. Let

$$H_\Delta^1(\Omega) := \{u \in H^1(\Omega) \mid \Delta u \in L^2(\Omega)\} \quad (2.13)$$

with Δu understood in distributional sense. Clearly $V_1(\Omega) \subset H_\Delta^1(\Omega)$. Given $u \in H_\Delta^1(\Omega)$, there is a unique element g of $(\mathcal{B}(\partial\Omega))'$ such that

$$\langle g, \text{Tr}_i v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \int_\Omega (\Delta u)v \, dx + \int_\Omega \nabla u \nabla v \, dx, \quad v \in H^1(\Omega). \quad (2.14)$$

We call this element g the *weak interior normal derivative* of u (with respect to Ω) and denote it by $\frac{\partial_i u}{\partial \nu} := g$. The operator $\frac{\partial_i}{\partial \nu} : H_\Delta^1(\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$ is linear and bounded in the sense that

$$\left\| \frac{\partial_i u}{\partial \nu} \right\|_{(\mathcal{B}(\partial\Omega))'} \leq \|u\|_{H^1(\Omega)} + \|\Delta u\|_{L^2(\Omega)}. \quad (2.15)$$

Corollary 2.15. *Let Ω be H^1 -admissible.*

(i) *Both \mathcal{N}_1 and the operator $\frac{\partial_i}{\partial \nu} : V_1(\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$ are isometries and onto, and we have $\frac{\partial_i}{\partial \nu} = \mathcal{N}_1^{-1}$ on $V_1(\Omega)$.*

(ii) *For $u, v \in V_1(\Omega)$ we have*

$$\left\langle \frac{\partial_i u}{\partial \nu}, \text{tr}_i v \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle u, v \rangle_{H^1(\Omega)} = \left\langle \frac{\partial_i v}{\partial \nu}, \text{tr}_i u \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}. \quad (2.16)$$

(iii) *The dual $(\frac{\partial_i}{\partial \nu})^* : \mathcal{B}(\partial\Omega) \rightarrow (V_1(\Omega))'$ of $\frac{\partial_i}{\partial \nu}$ on $V_1(\Omega)$ is an isometry and onto.*

Proof. Statement (i) follows using the surjectivity of tr observed in Lemma 2.6 (iii) and a comparison of (2.10) and (2.14). Statement (ii) is a special case of (2.14), and (iii) is a consequence of (i). \square

Now suppose $\Omega \subset \mathbb{R}^n$ is \dot{H}^1 -admissible. We then consider the space

$$\dot{H}_\Delta^1(\Omega) := \{u \in \dot{H}^1(\Omega) \mid \Delta u \in L^2(\Omega)\},$$

it contains $\dot{V}_0(\Omega)$. Given $u \in \dot{H}_\Delta^1(\Omega)$, there is a unique element g of $(\dot{\mathcal{B}}(\partial\Omega))'$ such that

$$\langle g, \dot{\text{Tr}}_i v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} = \int_\Omega (\Delta u)v \, dx + \int_\Omega \nabla u \nabla v \, dx, \quad v \in \dot{H}^1(\Omega),$$

and we call also $\frac{\partial_i u}{\partial \nu} := g$ the *weak interior normal derivative of u in the \dot{H}^1 -sense* (with respect to Ω). The operator $\frac{\partial_i}{\partial \nu} : \dot{H}_\Delta^1(\Omega) \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$ is linear and

$$\left\| \frac{\partial_i u}{\partial \nu} \right\|_{(\mathcal{B}(\partial\Omega))'} \leq \|u\|_{\dot{H}^1(\Omega)} + \|\Delta u\|_{L^2(\Omega)}. \quad (2.17)$$

Remark 2.16. *Suppose that Ω is \dot{H}^1 -admissible and bounded. Then by Remark 2.3 (ii) the spaces $\mathcal{B}(\partial\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega) \oplus \mathbb{R}$ can be identified. Each bounded linear functional $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ on $\dot{\mathcal{B}}(\partial\Omega)$ induces a linear functional on $\mathcal{B}(\partial\Omega)$ through extension by zero on \mathbb{R} . In this sense an equality of two elements of $(\dot{\mathcal{B}}(\partial\Omega))'$, such as the Neumann boundary condition $\frac{\partial_i \dot{N}_0 g}{\partial \nu} = g$ for $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, may be seen as an equality of two linear functionals on $\mathcal{B}(\partial\Omega)$.*

Corollary 2.17. *Let Ω be \dot{H}^1 -admissible.*

(i) *Both \dot{N}_0 and the operator $\frac{\partial_i}{\partial \nu} : \dot{V}_0(\Omega) \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$ are isometries and onto, and we have $\frac{\partial_i}{\partial \nu} = \dot{N}_0^{-1}$ on $\dot{V}_0(\Omega)$.*

(ii) *For $u, v \in \dot{V}_0(\Omega)$ we have*

$$\left\langle \frac{\partial_i u}{\partial \nu}, \text{tr}_i v \right\rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} = \langle u, v \rangle_{\dot{H}^1(\Omega)} = \left\langle \frac{\partial_i v}{\partial \nu}, \text{tr}_i u \right\rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}. \quad (2.18)$$

(iii) *The dual $(\frac{\partial_i}{\partial \nu})^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow (\dot{V}_0(\Omega))'$ of $\frac{\partial_i}{\partial \nu}$ on $\dot{V}_0(\Omega)$ is an isometry and onto.*

Remark 2.18. *Weak normal derivatives can also be used to characterize weak solutions of (2.7): An element u of $H^1(\Omega)$ is a weak solution to (2.7) if and only if $\langle u, v \rangle_{H^1(\Omega)} = 0$ for all $v \in C_c^\infty(\Omega)$ and $\langle v, u \rangle_{H^1(\Omega)} = \langle \frac{\partial v}{\partial \nu}, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ for all $v \in V_1(\Omega)$. Similarly for (2.8).*

3 Poincaré-Steklov operators for admissible domains

We provide some comments on Poincaré-Steklov operators for bounded admissible domains.

Let $\Omega \subset \mathbb{R}^n$ be a bounded H^1 -admissible domain. We write Δ_D for the self-adjoint Dirichlet Laplacian on $L^2(\Omega)$ and denote its spectrum by $\sigma(\Delta_D)$. Recall that $\sigma(\Delta_D) \subset (-\infty, 0)$ is pure point with eigenvalues accumulating at minus infinity.

Given $k \in \mathbb{R}$ and $f \in \mathcal{B}(\partial\Omega)$, we call $u \in H^1(\Omega)$ a weak solution of the Dirichlet problem

$$\begin{cases} -\Delta u + ku = 0 & \text{in } \Omega \\ u|_{\partial\Omega} = f \end{cases} \quad (3.1)$$

if $\text{Tr}_i u = f$ and $\langle u, v \rangle_{\dot{H}^1(\Omega)} + k \langle u, v \rangle_{L^2(\Omega)} = 0$ for all $v \in C_c^\infty(\Omega)$. This corresponds to (2.7) with $-\Delta u + ku$ in place of $-\Delta u + u$; problem (2.7) is the special case for $k = 1$.

If $k \in \mathbb{R} \setminus \sigma(\Delta_D)$, then for any $f \in \mathcal{B}(\partial\Omega)$ there is a unique weak solution $u_f \in H_\Delta(\Omega)$ of (3.1). For such k one can define a linear operator $\mathcal{A}_k : \mathcal{B}(\partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$ by

$$\mathcal{A}_k f := \frac{\partial_i u_f}{\partial \nu}.$$

this operator is called the *Poincaré-Steklov* (or *Dirichlet-to-Neumann*) *operator* associated with $(\Delta - k)$ on Ω . See for instance [6, 7, 40, 77] for studies of Poincaré-Steklov operators under more restrictive assumptions on Ω .

Lemma 3.1. *Let Ω be a bounded H^1 -admissible domain.*

(i) *For any $k \in \mathbb{R} \setminus \sigma(\Delta_D)$ the Poincaré-Steklov operator $\mathcal{A}_k : \mathcal{B}(\partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$ is a bounded linear operator and coincides with its adjoint. It is injective if and only if k is not an eigenvalue of the self-adjoint Neumann Laplacian for Ω .*

(ii) The Poincaré-Steklov operator $\mathcal{A}_1 : \mathcal{B}(\partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$ is an isometry and

$$\mathcal{A}_1 = \frac{\partial_i}{\partial\nu} \circ (\text{tr}_i)^{-1}.$$

Proof. By (2.15) and (3.1) we have

$$\|\mathcal{A}_k f\|_{(\mathcal{B}(\partial\Omega))'} \leq (1+k) \|u_f\|_{H^1(\Omega)} = (1+k) \|f\|_{\mathcal{B}(\partial\Omega)}, \quad f \in \mathcal{B}(\partial\Omega).$$

From (2.14) it is easily seen that

$$\langle \mathcal{A}_k f_1, f_2 \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle \mathcal{A}_k f_2, f_1 \rangle_{\mathcal{B}(\partial\Omega), (\mathcal{B}(\partial\Omega))'},$$

the special case for $k = 1$ was stated in (2.16). The statement on injectivity is clear. Item (ii) follows using Lemma 2.6 (iii) and Corollary 2.15 (i). \square

Now suppose that $\Omega \subset \mathbb{R}^n$ is bounded and \dot{H}^1 -admissible. Then we call the linear operator $\dot{\mathcal{A}}_0 : \dot{\mathcal{B}}(\partial\Omega) \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$, defined as

$$\dot{\mathcal{A}}_0 := \frac{\dot{\partial}_i}{\dot{\partial}\nu} \circ (\dot{\text{tr}}_i)^{-1},$$

the Poincaré-Steklov (or Dirichlet-to-Neumann) operator in the \dot{H}^1 -sense associated with Δ on Ω . Lemma 2.10 (iii) and Corollary 2.17 (i) give the following analog of Lemma 3.1.

Lemma 3.2. *Let Ω be a bounded \dot{H}^1 -admissible domain. Then $\dot{\mathcal{A}}_0 : \dot{\mathcal{B}}(\partial\Omega) \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$ is a linear isometry, and it coincides with its adjoint.*

Writing ι to denote the Riesz isomorphism from a Hilbert space to its dual (regardless of that space), we obtain the following commutative diagram (3.2):

$$\begin{array}{ccc} V_1(\Omega) = H_0^1(\Omega)^\perp & \begin{array}{c} \xrightarrow{\text{tr}_i} \\ \xleftarrow{\text{tr}_i^{-1}} \end{array} & \text{Tr}_i(H^1(\Omega)) = \mathcal{B}(\partial\Omega) \\ \begin{array}{c} \uparrow \iota^{-1} \\ \downarrow \iota \end{array} & \begin{array}{c} \searrow \frac{\partial_i}{\partial\nu} \\ \swarrow (\frac{\partial_i}{\partial\nu})^{-1} \end{array} & \begin{array}{c} \downarrow \mathcal{A}_1^{-1} = \iota^{-1} \\ \uparrow \mathcal{A}_1 = \iota \end{array} \\ V_1'(\Omega) = (H_0^1(\Omega)^\perp)' & \begin{array}{c} \xrightarrow{(\text{tr}_i^*)^{-1}} \\ \xleftarrow{\text{tr}_i^*} \end{array} & (\text{Tr}_i(H^1(\Omega)))' = (\mathcal{B}(\partial\Omega))' \end{array} \quad (3.2)$$

Note that indeed for any $f_1, f_2 \in \mathcal{B}(\partial\Omega)$ we have

$$\begin{aligned} \langle \iota(f_1), f_2 \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} &= \langle f_1, f_2 \rangle_{\mathcal{B}(\partial\Omega)} \\ &= \langle \text{tr}_i^{-1} f_1, \text{tr}_i^{-1} f_2 \rangle_{H^1(\Omega)} = \left\langle \frac{\partial_i}{\partial\nu} \circ (\text{tr}_i)^{-1} f_1, f_2 \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}. \end{aligned}$$

Obvious adjustments give an analogous commutative diagram involving the Poincaré-Steklov operator in the \dot{H}^1 -sense.

4 Transmission problems for admissible domains

We introduce layer potential operators for transmission problems on two-sided rough domains through well-posedness results for transmission problems of form

$$\begin{cases} (-\Delta + k)u & = 0 \quad \text{on } \mathbb{R}^n \setminus \partial\Omega \\ u_i|_{\partial\Omega} - u_e|_{\partial\Omega} & = -f \\ \frac{\partial_i u_i}{\partial\nu}|_{\partial\Omega} - \frac{\partial_e u_e}{\partial\nu}|_{\partial\Omega} & = g; \end{cases} \quad (4.1)$$

here $\Omega \subset \mathbb{R}^n$ is a bounded domain, u_i and u_e are the restrictions of the prospective solution to Ω and $\mathbb{R}^n \setminus \overline{\Omega}$, f and g are given data on $\partial\Omega$ and $k = 0, 1$. We first describe our setup for (4.1) and then proceed to well-posedness results, the definition of layer potentials and statements on some of their properties.

4.1 Two-sided admissible domains and jumps

We call $\Omega \subset \mathbb{R}^n$ a *two-sided H^1 -admissible domain* if both Ω and $\mathbb{R}^n \setminus \overline{\Omega}$ are nonempty and H^1 -extension domains and the Lebesgue measure of $\partial\Omega$ is zero. This implies in particular that the Hausdorff dimension of $\partial\Omega$ is at least $n - 1$, hence its capacity is positive.

Suppose that $\Omega \subset \mathbb{R}^n$ is a two-sided H^1 -admissible domain. We discuss (4.1) with $k = 1$ in terms of the space $H^1(\mathbb{R}^n \setminus \partial\Omega)$. It admits the orthogonal decomposition $H^1(\mathbb{R}^n \setminus \partial\Omega) = H^1(\Omega) \oplus H^1(\mathbb{R}^n \setminus \overline{\Omega})$, so every $u \in H^1(\mathbb{R}^n \setminus \partial\Omega)$ can be written $u = u_i + u_e$ with uniquely determined $u_i \in H^1(\Omega)$ and $u_e \in H^1(\mathbb{R}^n \setminus \overline{\Omega})$. We denote the closure of $C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$ in $H^1(\mathbb{R}^n \setminus \partial\Omega)$ by $H_0^1(\mathbb{R}^n \setminus \partial\Omega)$ and write $V_1(\mathbb{R}^n \setminus \partial\Omega)$ for its orthogonal complement, which is the space of functions that are 1-harmonic in $\mathbb{R}^n \setminus \partial\Omega$.

To distinguish between the two operators, we now write $\text{Tr}_i : H^1(\Omega) \rightarrow \mathcal{B}(\partial\Omega)$ to denote the boundary trace operator for Ω and $\text{Tr}_e : H^1(\mathbb{R}^n \setminus \overline{\Omega}) \rightarrow \mathcal{B}(\partial\Omega)$ to denote the boundary trace operator for $\mathbb{R}^n \setminus \overline{\Omega}$; by construction both map onto $\mathcal{B}(\partial\Omega)$. We refer to Tr_i as the *interior* trace operator with respect to Ω and to Tr_e as the *exterior*. In the context of transmission problems we endow $\mathcal{B}(\partial\Omega)$ with the Hilbert space norm

$$\|f\|_{\mathcal{B}(\partial\Omega),t} := (\|f\|_{\mathcal{B}(\partial\Omega),i}^2 + \|f\|_{\mathcal{B}(\partial\Omega),e}^2)^{1/2}, \quad (4.2)$$

where $\|\cdot\|_{\mathcal{B}(\partial\Omega),i}^2$ denotes the norm defined in (2.4) with respect to Ω and $\|\cdot\|_{\mathcal{B}(\partial\Omega),e}^2$ denotes the norm defined similarly but with $\mathbb{R}^n \setminus \overline{\Omega}$ in place of Ω . Since both Ω and $\mathbb{R}^n \setminus \overline{\Omega}$ are H^1 -extension domains, the norms $\|\cdot\|_{\mathcal{B}(\partial\Omega),i}$, $\|\cdot\|_{\mathcal{B}(\partial\Omega),e}$ and $\|\cdot\|_{\mathcal{B}(\partial\Omega),t}$ are all equivalent. As a consequence, we may view Tr_i and Tr_e as bounded linear operators from $H^1(\mathbb{R}^n \setminus \partial\Omega)$ onto $\mathcal{B}(\partial\Omega)$, no matter which norm is used. Since only $\|\cdot\|_{\mathcal{B}(\partial\Omega),t}$ will be used in the sequel, we agree to denote it again by $\|\cdot\|_{\mathcal{B}(\partial\Omega)}$.

Given $u \in H^1(\mathbb{R}^n \setminus \partial\Omega)$, we write

$$\llbracket \text{Tr } u \rrbracket := \text{Tr}_i u - \text{Tr}_e u$$

for its *jump in trace across $\partial\Omega$* . The map $u \mapsto \llbracket \text{Tr } u \rrbracket$ defines a bounded linear operator $\llbracket \text{Tr } u \rrbracket : H^1(\mathbb{R}^n \setminus \partial\Omega) \rightarrow \mathcal{B}(\partial\Omega)$, it is onto.

If $\Delta u \in L^2(\Omega)$, then the *interior* normal derivative $\frac{\partial_i u}{\partial \nu}$ of u with respect to Ω is as defined in (2.14). If $\Delta u \in L^2(\mathbb{R}^n \setminus \overline{\Omega})$, then we define the *exterior* normal derivative $\frac{\partial_e u}{\partial \nu}$ of u with respect to Ω as minus the interior normal derivative of u with respect to $\mathbb{R}^n \setminus \overline{\Omega}$, that is, the unique element $g \in (\mathcal{B}(\partial\Omega))'$ such that

$$\langle g, \text{Tr}_e v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = - \int_{\mathbb{R}^n \setminus \overline{\Omega}} (\Delta u) v \, dx - \int_{\mathbb{R}^n \setminus \overline{\Omega}} \nabla u \cdot \nabla v \, dx, \quad v \in H^1(\mathbb{R}^n \setminus \overline{\Omega}). \quad (4.3)$$

Now let

$$H_\Delta^1(\mathbb{R}^n \setminus \partial\Omega) := \{u \in H^1(\mathbb{R}^n \setminus \partial\Omega) \mid \Delta u \in L^2(\mathbb{R}^n \setminus \partial\Omega)\}.$$

Clearly this space contains $V_1(\mathbb{R}^n \setminus \partial\Omega)$. For $u \in H_\Delta^1(\mathbb{R}^n \setminus \partial\Omega)$ we write

$$\llbracket \frac{\partial u}{\partial \nu} \rrbracket := \frac{\partial_i u}{\partial \nu} - \frac{\partial_e u}{\partial \nu}$$

for the *jump of its normal derivative across $\partial\Omega$* . This defines a linear operator $u \mapsto \llbracket \frac{\partial u}{\partial \nu} \rrbracket$ from $H_\Delta^1(\mathbb{R}^n \setminus \partial\Omega)$ onto $(\mathcal{B}(\partial\Omega))'$, bounded in the sense that

$$\left\| \llbracket \frac{\partial u}{\partial \nu} \rrbracket \right\|_{(\mathcal{B}(\partial\Omega))'} \leq \|u\|_{H^1(\mathbb{R}^n \setminus \partial\Omega)} + \|\Delta u\|_{L^2(\mathbb{R}^n \setminus \partial\Omega)}.$$

We call $\Omega \subset \mathbb{R}^n$ a *two-sided \dot{H}^1 -admissible domain* if both Ω and $\mathbb{R}^n \setminus \overline{\Omega}$ are nonempty and \dot{H}^1 -extension domains, Ω is bounded and the Lebesgue measure of $\partial\Omega$ is zero.

Suppose that $\Omega \subset \mathbb{R}^n$ is a two-sided \dot{H}^1 -admissible domain. We discuss (4.1) with $k = 0$ using the space $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$, which decomposes $\dot{H}^1(\Omega) \oplus \dot{H}^1(\mathbb{R}^n \setminus \overline{\Omega})$. By $\dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega)$ we denote the closure of $C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$ in $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ and by $\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$ its orthogonal complement.

By $\dot{\text{Tr}}_i : \dot{H}^1(\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$ and $\dot{\text{Tr}}_e : \dot{H}^1(\mathbb{R}^n \setminus \overline{\Omega}) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$ we denote the interior and exterior trace with respect to Ω , defined in an analogous manner. We endow $\dot{\mathcal{B}}(\partial\Omega)$ with the Hilbert space norm

$$\|f\|_{\dot{\mathcal{B}}(\partial\Omega),t} := (\|f\|_{\dot{\mathcal{B}}(\partial\Omega),i}^2 + \|f\|_{\dot{\mathcal{B}}(\partial\Omega),e}^2)^{1/2}$$

with the summands defined similarly as before. Since only $\|\cdot\|_{\dot{\mathcal{B}}(\partial\Omega),t}$ will be used in the sequel, we agree to denote it again by $\|\cdot\|_{\dot{\mathcal{B}}(\partial\Omega)}$.

Given $u \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$, we set

$$[[\dot{\text{Tr}} u]] := \dot{\text{Tr}}_i u - \dot{\text{Tr}}_e u$$

for its jump in trace across Ω in the \dot{H}^1 -sense. The map $[[\dot{\text{Tr}}]] : \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$, $u \mapsto [[\dot{\text{Tr}} u]]$, defines a bounded linear operator which is onto.

Let $\dot{H}_\Delta^1(\mathbb{R}^n \setminus \partial\Omega)$ denote the space of all $u \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ with $\Delta u \in L^2(\mathbb{R}^n \setminus \partial\Omega)$; it contains $\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$. Given an element $u \in \dot{H}_\Delta^1(\mathbb{R}^n \setminus \partial\Omega)$, we write $\frac{\partial_i u}{\partial \nu}$ for its interior normal derivative in the \dot{H}^1 -sense with respect to Ω and $\frac{\partial_e u}{\partial \nu}$ for its exterior, again defined as minus the interior with respect to $\mathbb{R}^n \setminus \overline{\Omega}$. We then write

$$\left[\left[\frac{\partial u}{\partial \nu} \right] \right] := \frac{\partial_i u}{\partial \nu} - \frac{\partial_e u}{\partial \nu}$$

for the jump of its normal derivative across $\partial\Omega$ in the \dot{H}^1 -sense. This defines a linear operator $u \mapsto \left[\left[\frac{\partial u}{\partial \nu} \right] \right]$ from $\dot{H}_\Delta^1(\mathbb{R}^n \setminus \partial\Omega)$ onto $(\dot{\mathcal{B}}(\partial\Omega))'$, bounded in the sense that

$$\left\| \left[\left[\frac{\partial u}{\partial \nu} \right] \right] \right\|_{(\dot{\mathcal{B}}(\partial\Omega))'} \leq \|u\|_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} + \|\Delta u\|_{L^2(\mathbb{R}^n \setminus \partial\Omega)}.$$

Remark 4.1. *By elliptic regularity any element of $V_1(\mathbb{R}^n \setminus \partial\Omega)$ and any representative modulo (two) constants of any element $\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$ is in $C^\infty(\mathbb{R}^n \setminus \partial\Omega)$.*

4.2 Subspaces and orthogonality

Suppose that $\Omega \subset \mathbb{R}^n$ is a two-sided H^1 -admissible domain. We write

$$V_{1,s}(\mathbb{R}^n \setminus \partial\Omega) := \{u \in V_1(\mathbb{R}^n \setminus \partial\Omega) \mid [[\text{Tr} u]] = 0\}$$

and

$$V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega) := \{u \in V_1(\mathbb{R}^n \setminus \partial\Omega) \mid \left[\left[\frac{\partial u}{\partial \nu} \right] \right] = 0\}.$$

Lemma 4.2. *Let $\Omega \subset \mathbb{R}^n$ be two-sided H^1 -admissible.*

(i) *We have $\ker [[\text{Tr}]] = H^1(\mathbb{R}^n)$. The linear operator $\text{Tr} : H^1(\mathbb{R}^n) \rightarrow \mathcal{B}(\partial\Omega)$, defined as $\text{Tr} := \text{Tr}_i|_{H^1(\mathbb{R}^n)} = \text{Tr}_e|_{H^1(\mathbb{R}^n)}$, is bounded with operator norm one. Its kernel agrees with $H_0^1(\mathbb{R}^n \setminus \partial\Omega)$, which is the orthogonal complement of $V_{1,s}(\mathbb{R}^n \setminus \partial\Omega)$ in $H^1(\mathbb{R}^n)$.*

(ii) *The space $V_{1,s}(\mathbb{R}^n \setminus \partial\Omega)$ is a closed subspace of $H^1(\mathbb{R}^n \setminus \partial\Omega)$. The operator*

$$\text{tr} : V_{1,s}(\mathbb{R}^n \setminus \partial\Omega) \rightarrow \mathcal{B}(\partial\Omega), \tag{4.4}$$

defined as the restriction $\text{tr} := \text{Tr}|_{V_{1,s}(\mathbb{R}^n \setminus \partial\Omega)}$, is a linear isometry and onto.

(iii) The space $V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ is a closed subspace of $H^1(\mathbb{R}^n \setminus \partial\Omega)$. The operator

$$\frac{\partial}{\partial\nu} : V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))', \quad (4.5)$$

defined as $\frac{\partial}{\partial\nu} := \frac{\partial_i}{\partial\nu} = \frac{\partial_e}{\partial\nu}$, is a linear isometry and onto.

(iv) The space $V_1(\mathbb{R}^n \setminus \partial\Omega)$ admits the orthogonal decomposition

$$V_1(\mathbb{R}^n \setminus \partial\Omega) = V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega).$$

Proof. For (i), note that $u \in V_1(\mathbb{R}^n \setminus \Omega) \cap \ker \llbracket \text{Tr} \rrbracket$ agrees with the unique element w of $H^1(\mathbb{R}^n)$ of minimal norm such that $w|_{\partial\Omega} = \text{Tr}_i u = \text{Tr}_e u$. The remaining part of (i) now follows using (4.2) and (A.1). Statements (ii) and (iii) follow from Lemma 2.6 (iii) and Corollary 2.15 (i). To see (iv), note that given $u \in V_1(\mathbb{R}^n \setminus \partial\Omega)$, we find that for any $v \in V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ we have

$$\left\langle \left[\frac{\partial u}{\partial\nu} \right], \text{tr } v \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \int_{\mathbb{R}^n \setminus \partial\Omega} (\Delta u)v \, dx + \int_{\mathbb{R}^n \setminus \partial\Omega} \nabla u \nabla v \, dx = \langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)}. \quad (4.6)$$

Since tr in (4.4) is surjective, it follows that $\left[\frac{\partial u}{\partial\nu} \right] = 0$ in $(\mathcal{B}(\partial\Omega))'$ if and only if (4.6) is zero for all $v \in V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$, and this is the case if and only if u belongs to the orthogonal complement of $V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ in $V_1(\mathbb{R}^n \setminus \partial\Omega)$. \square

Remark 4.3. Alternatively, we could observe that given $v \in V_1(\mathbb{R}^n \setminus \partial\Omega)$, we have

$$\left\langle \frac{\partial u}{\partial\nu}, \llbracket \text{Tr } v \rrbracket \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)}. \quad (4.7)$$

for all $u \in V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$. Since $\frac{\partial}{\partial\nu}$ in (4.5) is surjective and $\mathcal{B}(\partial\Omega)$ is a Hilbert space, it follows that $\llbracket \text{Tr } v \rrbracket = 0$ in $\mathcal{B}(\partial\Omega)$ if and only if v belongs to the orthogonal complement of $V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ in $V_1(\mathbb{R}^n \setminus \partial\Omega)$.

By $P_{1,\mathcal{S}}$ and $P_{1,\mathcal{D}}$ we denote the orthogonal projections from $H^1(\mathbb{R}^n \setminus \partial\Omega)$ onto the closed subspaces $V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ and $V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$, respectively. The following is a straightforward consequence of Lemma 4.2 (iv).

Corollary 4.4. Let Ω be two-sided H^1 -admissible. If $u, v \in V_1(\mathbb{R}^n \setminus \partial\Omega)$, then

$$\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = \left\langle \left[\frac{\partial u}{\partial\nu} \right], \text{tr} \circ P_{1,\mathcal{S}} v \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} + \left\langle \frac{\partial}{\partial\nu} \circ P_{1,\mathcal{D}} v, \llbracket \text{Tr } u \rrbracket \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}.$$

Now suppose that $\Omega \subset \mathbb{R}^n$ is a two-sided H^1 -admissible domain. We write

$$\dot{V}_{0,\dot{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega) := \{u \in \dot{V}_0(\mathbb{R}^n \setminus \partial\Omega) \mid \llbracket \dot{\text{Tr}} u \rrbracket = 0\}$$

and

$$\dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega) := \{u \in \dot{V}_0(\mathbb{R}^n \setminus \partial\Omega) \mid \left[\frac{\partial u}{\partial\nu} \right] = 0\}.$$

The following is seen similarly as before.

Lemma 4.5. Let $\Omega \subset \mathbb{R}^n$ be two-sided \dot{H}^1 -admissible.

(i) The linear operator $\dot{\text{Tr}} : \ker \llbracket \dot{\text{Tr}} \rrbracket \rightarrow \dot{\mathcal{B}}(\partial\Omega)$, defined as $\dot{\text{Tr}} := \dot{\text{Tr}}_i = \dot{\text{Tr}}_e$, is bounded with operator norm one. Its kernel is the orthogonal complement of $\dot{V}_{0,\dot{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega)$ in $\ker \llbracket \dot{\text{Tr}} \rrbracket$, this complement is the closure of $C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$.

(ii) The space $\dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega)$ is a closed subspace of $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$. The operator

$$\dot{\text{tr}} : \dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega),$$

defined as the restriction $\dot{\text{tr}} := \dot{\text{Tr}}|_{\dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega)}$, is a linear isometry and onto.

(iii) The space $\dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$ is a closed subspace of $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$. The operator

$$\frac{\dot{\partial}}{\partial\nu} : \dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))',$$

defined as $\frac{\dot{\partial}}{\partial\nu} := \frac{\dot{\partial}_i}{\partial\nu} = \frac{\dot{\partial}_e}{\partial\nu}$, is a linear isometry and onto.

(iv) The space $\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$ admits the orthogonal decomposition

$$\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega) = \dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega) \oplus \dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega).$$

By $P_{0,\dot{s}}$ and $P_{0,\dot{\mathcal{D}}}$ we denote the orthogonal projections from $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ onto the closed subspaces $\dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$, respectively. Lemma 4.2 (iv) now gives the following.

Corollary 4.6. *Let Ω be two-sided \dot{H}^1 -admissible. If $u, v \in \dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$, then*

$$\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = \left\langle \left[\frac{\dot{\partial}}{\partial\nu} \right], \dot{\text{tr}} \circ P_{0,\dot{s}} v \right\rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} + \left\langle \frac{\dot{\partial}}{\partial\nu} \circ P_{0,\dot{\mathcal{D}}} v, \left[\dot{\text{Tr}} u \right] \right\rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}.$$

Remark 4.7. *Suppose that Ω is two-sided \dot{H}^1 -admissible and $u \in \ker \left[\dot{\text{Tr}} \right]$. Let $w_i \in u_i$ and $w_e \in u_e$ be representatives modulo constants of u_i and u_e . By the arguments used to show Proposition 2.2 both $\text{Tr}_i w_i$ and $\text{Tr}_e w_e$ are well-defined elements of $\mathcal{B}(\partial\Omega)$, and by construction there is a constant $c \in \mathbb{R}$ such that $\text{Tr}_i w_i - \text{Tr}_e w_e = c$. Setting $\bar{w} := w_i + w_e + c \mathbf{1}_{\mathbb{R}^n \setminus \bar{\Omega}}$ we obtain a representative \bar{w} of u that satisfies $\left[\text{Tr} \bar{w} \right] = 0$ in $\mathcal{B}(\partial\Omega)$. The equivalence class \bar{u} modulo single constants is a uniquely determined element of $\dot{H}^1(\mathbb{R}^n)$, it could be called the zero trace jump readjusted representative of u .*

The zero trace jump readjusted representative \bar{u} of $u \in \dot{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega)$ is an element of the orthogonal complement of $C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$ in $\dot{H}^1(\mathbb{R}^n)$; we denote it by $\bar{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega)$. Setting $\bar{\text{tr}} \bar{u} := \dot{\text{tr}} u$, we obtain a linear bijection

$$\bar{\text{tr}} : \bar{V}_{0,\dot{s}}(\mathbb{R}^n \setminus \partial\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega). \quad (4.8)$$

4.3 Double layer potentials

Let $\Omega \subset \mathbb{R}^n$ be two-sided H^1 -admissible. Given $f \in \mathcal{B}(\partial\Omega)$, we call $u \in H^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution* of (4.1) with $k = 1$ and $g = 0$, that is, a weak solution of

$$\begin{cases} -\Delta u + u & = 0 & \text{on } \mathbb{R}^n \setminus \partial\Omega \\ u_i|_{\partial\Omega} - u_e|_{\partial\Omega} & = -f \\ \frac{\partial_i u_i}{\partial\nu}|_{\partial\Omega} - \frac{\partial_e u_e}{\partial\nu}|_{\partial\Omega} & = 0, \end{cases} \quad (4.9)$$

if $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega) \cup V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle v, u \rangle_{H^1(\mathbb{R}^n \setminus \Omega)} = \langle \frac{\partial v}{\partial\nu}, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ for all $v \in V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$.

Lemma 4.8. *Let $\Omega \subset \mathbb{R}^n$ be two-sided H^1 -admissible. For any $f \in \mathcal{B}(\partial\Omega)$ there is a unique weak solution u^f of (4.9). It is an element of $V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u^f\|_{H^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|f\|_{\mathcal{B}(\partial\Omega)}$.*

Proof. Since by Lemma 4.2 (iii) the linear functional $v \mapsto \langle \frac{\partial v}{\partial \nu}, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ is bounded on the closed subspace $V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ of $H^1(\mathbb{R}^n \setminus \partial\Omega)$, the result follows from the Riesz representation theorem. \square

To the bounded linear operator $\mathcal{D} : \mathcal{B}(\partial\Omega) \rightarrow V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ defined by

$$\mathcal{D}f := u^f$$

we refer as the *double layer potential operator associated with the transmission problem for $1 - \Delta$ and Ω* .

Corollary 4.9. *Let Ω be two-sided H^1 -admissible. The operator \mathcal{D} is bijective, and its inverse is $\mathcal{D}^{-1} = -\llbracket \text{Tr} \rrbracket$.*

Now let $\Omega \subset \mathbb{R}^n$ be two-sided \dot{H}^1 -admissible. Given $f \in \dot{\mathcal{B}}(\partial\Omega)$, we call $u \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution in the \dot{H}^1 -sense* of (4.1) with $k = 0$ and $g = 0$, that is,

$$\begin{cases} -\Delta u & = 0 & \text{on } \mathbb{R}^n \setminus \partial\Omega \\ u_i|_{\partial\Omega} - u_e|_{\partial\Omega} & = -f \\ \frac{\partial_i u_i}{\partial \nu}|_{\partial\Omega} - \frac{\partial_e u_e}{\partial \nu}|_{\partial\Omega} & = 0, \end{cases} \quad (4.10)$$

if $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega) \cup \dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle v, u \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \Omega)} = \langle \frac{\partial v}{\partial \nu}, f \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}$ for all $v \in \dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$.

Lemma 4.10. *Let $\Omega \subset \mathbb{R}^n$ be two-sided \dot{H}^1 -admissible. For any $f \in \dot{\mathcal{B}}(\partial\Omega)$ there is a unique weak solution u^f of (4.10) in the \dot{H}^1 -sense. It is an element of $\dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u^f\|_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|f\|_{\dot{\mathcal{B}}(\partial\Omega)}$.*

To the bounded linear operator $\dot{\mathcal{D}} : \dot{\mathcal{B}}(\partial\Omega) \rightarrow \dot{V}_{0,\dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$ defined by $\dot{\mathcal{D}}f := u^f$ we refer as the *double layer potential operator associated with the transmission problem for $-\Delta$ and Ω in the \dot{H}^1 -sense*.

Corollary 4.11. *Let Ω be two-sided \dot{H}^1 -admissible. The operator $\dot{\mathcal{D}}$ is bijective, and its inverse is $\dot{\mathcal{D}}^{-1} = -\llbracket \dot{\text{Tr}} \rrbracket$.*

Remark 4.12. *A priori the trace jump condition in (4.10) is only an equality modulo additive constants. However, by arguments similar to Remarks 2.12 and 4.7 the situation is auto-improving: If Ω is two-sided \dot{H}^1 -admissible and $f \in \mathcal{B}(\partial\Omega)$ is given, let $u^{[f]}$ be the unique weak solution in the \dot{H}^1 -sense of (4.10) with the equivalence class $[f] \in (\dot{\mathcal{B}}(\partial\Omega))'$ of f modulo constants in place of f . We can find representatives $w_i \in u_i^{[f]}$ and $w_e \in u_e^{[f]}$ modulo constants and some $c \in \mathbb{R}$ such that $\text{Tr}_i w_i - \text{Tr}_e w_e = -f + c$ in $\mathcal{B}(\partial\Omega)$. We set $\bar{w} := w_i + w_e + c\mathbf{1}_{\mathbb{R}^n \setminus \bar{\Omega}}$. The equivalence class $\bar{u}^{[f]}$ of \bar{w} modulo a single constant is uniquely determined. It satisfies $\Delta \bar{u}^{[f]} = 0$ on $\mathbb{R}^n \setminus \partial\Omega$, $\llbracket \text{Tr} \bar{u}^{[f]} \rrbracket = -f$ as an equality in $\mathcal{B}(\partial\Omega)$ and $\llbracket \frac{\partial \bar{u}^{[f]}}{\partial \nu} \rrbracket = 0$, which by Remark 2.16 may be interpreted as an equality of linear functionals on $\mathcal{B}(\partial\Omega)$. One could call $\bar{u}^{[f]}$ the trace jump readjusted solution of (4.10).*

4.4 Single layer potentials

Let Ω be two-sided H^1 -admissible. Given $g \in (\mathcal{B}(\partial\Omega))'$, we call $u \in H^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution* of (4.1) with $k = 1$ and $f = 0$, that is,

$$\begin{cases} -\Delta u + u & = 0 & \text{on } \mathbb{R}^n \setminus \partial\Omega \\ u_i|_{\partial\Omega} - u_e|_{\partial\Omega} & = 0 \\ \frac{\partial_i u_i}{\partial \nu}|_{\partial\Omega} - \frac{\partial_e u_e}{\partial \nu}|_{\partial\Omega} & = g, \end{cases} \quad (4.11)$$

if $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle g, \text{Tr} v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ for all $v \in H^1(\mathbb{R}^n)$.

Lemma 4.13. *Let Ω be two-sided H^1 -admissible. For any $g \in (\mathcal{B}(\partial\Omega))'$ there is a unique weak solution u_g of (4.11). It is in $V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u_g\|_{H^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|g\|_{(\mathcal{B}(\partial\Omega))'}$.*

Proof. By Lemma 4.2 (ii) the linear functional $v \mapsto \langle g, \text{Tr } v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ is bounded on the closed subspace $V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ of $H^1(\mathbb{R}^n \setminus \partial\Omega)$, so the result follows again from the Riesz representation theorem. \square

To the bounded linear operator $\mathcal{S} : (\mathcal{B}(\partial\Omega))' \rightarrow V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$, defined by

$$\mathcal{S}g := u_g$$

we refer as the *single layer potential operator associated with the transmission problem for $1 - \Delta$ and Ω* .

Corollary 4.14. *Let Ω be two-sided H^1 -admissible. The operator \mathcal{S} is bijective, and its inverse is $\mathcal{S}^{-1} = \llbracket \frac{\partial}{\partial\nu} \rrbracket$.*

Suppose that Ω is two-sided \dot{H}^1 -admissible. Given $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, we call $u \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution in the \dot{H}^1 -sense* of (4.1) with $k = 0$ and $f = 0$, that is,

$$\begin{cases} -\Delta u & = 0 & \text{on } \mathbb{R}^n \setminus \partial\Omega, \\ u_i|_{\partial\Omega} - u_e|_{\partial\Omega} & = 0 \\ \frac{\partial_i u_i}{\partial\nu}|_{\partial\Omega} - \frac{\partial_e u_e}{\partial\nu}|_{\partial\Omega} & = g \end{cases} \quad (4.12)$$

if $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in \dot{V}_{0,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle g, \dot{\text{Tr}} v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}$ for all $v \in \dot{H}^1(\mathbb{R}^n)$.

Lemma 4.15. *Let Ω be two-sided \dot{H}^1 -admissible. For any $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ there is a unique weak solution u_g of (4.12) in the \dot{H}^1 -sense. It is an element of $\dot{V}_{0,\dot{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u_g\|_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|g\|_{(\dot{\mathcal{B}}(\partial\Omega))'}$.*

To the bounded linear operator $\dot{\mathcal{S}} : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{V}_{0,\dot{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega)$ defined by $\dot{\mathcal{S}}g := u_g$ we refer as the *single layer potential operator associated with the transmission problem for $-\Delta$ and Ω in the \dot{H}^1 -sense*.

Corollary 4.16. *Let Ω be two-sided \dot{H}^1 -admissible. The operator $\dot{\mathcal{S}}$ is bijective, and its inverse is $\dot{\mathcal{S}}^{-1} = \llbracket \frac{\partial}{\partial\nu} \rrbracket$.*

Remark 4.17. *Let Ω be two-sided \dot{H}^1 -admissible, $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ and let u_g be as above. Let $\bar{u}_g \in \bar{V}_{0,\bar{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega)$ be the zero trace jump readjusted representative of u_g as in Remark 4.7. Since it is uniquely determined, $\bar{\mathcal{S}}g := \bar{u}_g$ defines a bounded linear map*

$$\bar{\mathcal{S}} : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \bar{V}_{0,\bar{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega); \quad (4.13)$$

it is a zero trace jump readjusted variant of $\dot{\mathcal{S}}$.

4.5 Superposition

Superposition gives well-posedness for (4.1). Suppose Ω is H^1 -admissible. Given $f \in \mathcal{B}(\partial\Omega)$ and $g \in (\mathcal{B}(\partial\Omega))'$ we call $u \in H^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution* for (4.1) with $k = 1$ if $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$, $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle g, \text{Tr } v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ for all $v \in V_{1,\mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = - \langle \llbracket \frac{\partial u}{\partial\nu} \rrbracket, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$ for all $v \in V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$.

Corollary 4.18. *Let Ω be H^1 -admissible. For any $f \in \mathcal{B}(\partial\Omega)$ and $g \in (\mathcal{B}(\partial\Omega))'$ the unique weak solution u of (4.1) with $k = 1$ is $u = \mathcal{D}f + \mathcal{S}g$. It is an element of $V_1(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u\|_{H^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|f\|_{\mathcal{B}(\partial\Omega)} + \|g\|_{(\mathcal{B}(\partial\Omega))'}$.*

If Ω is \dot{H}^1 -admissible, $f \in \dot{\mathcal{B}}(\partial\Omega)$ and $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, then we call $u \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ a *weak solution* for (4.1) with $k = 0$ in the \dot{H}^1 -sense if $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = 0$ for all $v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$, $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle g, \text{Tr } v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}$ for all $v \in \dot{V}_{0, \dot{\mathcal{S}}}(\mathbb{R}^n \setminus \partial\Omega)$ and $\langle u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = - \left\langle \left[\frac{\partial v}{\partial \nu} \right], f \right\rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}$ for all $v \in \dot{V}_{0, \dot{\mathcal{D}}}(\mathbb{R}^n \setminus \partial\Omega)$.

Corollary 4.19. *Let Ω be \dot{H}^1 -admissible. For any $f \in \dot{\mathcal{B}}(\partial\Omega)$ and $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ the unique weak solution u in the \dot{H}^1 -sense of (4.1) with $k = 0$ is $u = \dot{\mathcal{D}}f + \dot{\mathcal{S}}g$. It is an element of $\dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$ and satisfies $\|u\|_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} \leq \|f\|_{\dot{\mathcal{B}}(\partial\Omega)} + \|g\|_{(\dot{\mathcal{B}}(\partial\Omega))'}$.*

5 Resolvent representations

5.1 Representations of single layer potentials

Let \mathcal{G} be the Bessel potential operator $u \mapsto \mathcal{G}u = ((1 + |\xi|^2)^{-1}\hat{u})^\vee$ of order 2, where $u \mapsto \hat{u}$ denotes the Fourier transform and $u \mapsto \check{u}$ its inverse, a priori defined on Schwartz functions. It is well known that $\mathcal{G} = (I - \Delta)^{-1}$ on Schwartz functions and that \mathcal{G} extends to a bounded linear operator of $L^2(\mathbb{R}^2)$.

If $\Omega \subset \mathbb{R}^n$ is two-sided H^1 -admissible, then this operator \mathcal{G} is the resolvent operator uniquely associated with the symmetric bilinear form (2.1) when endowed with the domain $H^1(\mathbb{R}^n) = H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1, \mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega)$.

The operator \mathcal{G} extends further to an isometric isomorphism $H^{-1}(\mathbb{R}^n) \rightarrow H^1(\mathbb{R}^n)$. Using a similar agreement as in Remark 2.8 together with Lemma 4.2 (ii) the adjoint $\text{tr}^* : (\mathcal{B}(\partial\Omega))' \rightarrow (V_{1, \mathcal{S}}(\mathbb{R}^n \setminus \partial\Omega))'$ of the restricted trace operator tr as in Lemma 4.2 (iii) can be viewed as a bounded linear operator $\text{tr}^* : (\mathcal{B}(\partial\Omega))' \rightarrow H^{-1}(\mathbb{R}^n)$. We obtain the following representation for the single layer potential operator.

Lemma 5.1. *Let Ω be two-sided H^1 -admissible. Then $\mathcal{S} = \mathcal{G} \circ \text{tr}^*$.*

Proof. Let $g \in (\mathcal{B}(\partial\Omega))'$. Then $\text{tr}^* g \in H^{-1}(\mathbb{R}^n)$ and consequently $\mathcal{G} \circ \text{tr}^* g \in H^1(\mathbb{R}^n)$. By general theory, (2.5) and the definition of \mathcal{S} we find that

$$\langle \mathcal{G} \circ \text{tr}^* g, v \rangle_{H^1(\mathbb{R}^n)} = \langle \text{tr}^* g, v \rangle_{H^{-1}(\mathbb{R}^n), H^1(\mathbb{R}^n)} = \langle g, \text{Tr } v \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \langle \mathcal{S}g, v \rangle_{H^1(\mathbb{R}^n)} \quad (5.1)$$

for all $v \in H^1(\mathbb{R}^n)$. \square

Let G be the Bessel kernel of order two, that is, the fundamental solution for $\Delta - 1$ on \mathbb{R}^n . It is well known that for any $w \in L^2(\mathbb{R}^n)$ we have $\mathcal{G}w = G * w$. For a nonnegative Radon measure ν on $\partial\Omega$ one defines

$$G * \nu(x) := \int_{\partial\Omega} G(x - y)\nu(dy), \quad x \in \mathbb{R}^n;$$

this is a lower semicontinuous function taking values in $[0, +\infty]$, finite at all points $x \in \mathbb{R}^n \setminus \partial\Omega$. For a finite signed Borel measure ν on $\partial\Omega$ we set

$$G * \nu(x) := G * \nu^+(x) - G * \nu^-(x), \quad x \in \mathbb{R} \setminus \partial\Omega;$$

this gives a well-defined integrable function $G * \nu$ on \mathbb{R}^n .

We say that a nonnegative Radon measure ν on $\partial\Omega$ has *finite energy* if there is a constant $c > 0$ such that

$$\int_{\partial\Omega} |v| d\nu \leq c \|v\|_{H^1(\mathbb{R}^n)}, \quad v \in H^1(\mathbb{R}^n) \cap C_c(\mathbb{R}^n). \quad (5.2)$$

Given a finite signed Borel measure ν on $\partial\Omega$, we say that it has finite energy if its total variation measure $\nu^+ + \nu^-$ has.

Remark 5.2. It is well known that by the Riesz representation theorem for measures the cone of nonnegative elements of $H^{-1}(\mathbb{R}^n)$ is in one-to-one correspondence with the cone of nonnegative Radon measures of finite energy on \mathbb{R}^n , see [78, Chapter 6, Exercise 4]; a variant of the argument can be found in [16, Proposition 9.2.1].

Proposition 5.3. Let Ω be two-sided H^1 -admissible and let ν be a finite signed Borel measure on $\partial\Omega$ of finite energy. Then sets of zero capacity have zero ν -measure, and ν defines an element of $(\mathcal{B}(\partial\Omega))'$ by

$$\langle \nu, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} := \int_{\partial\Omega} f d\nu, \quad f \in \mathcal{B}(\partial\Omega). \quad (5.3)$$

Moreover, $\mathcal{S}\nu(x) = G * \nu(x)$, $x \in \mathbb{R}^n \setminus \partial\Omega$.

Proof. It suffices to prove the result for a nonnegative Radon measure ν of finite energy. The first claim is proved in [41, Lemma 2.2.3]. One can follow [41, Theorem 2.2.2] to see that

$$\int_{\partial\Omega} \tilde{v} d\nu = \langle G * \nu, v \rangle_{H^1(\mathbb{R}^n)}, \quad v \in H^1(\mathbb{R}^n). \quad (5.4)$$

Estimate (5.2) extends to all $v \in H^1(\mathbb{R}^n)$ and gives $|\int_{\partial\Omega} \tilde{v} d\nu| \leq c\|v\|_{H^1(\mathbb{R}^n)}$; optimizing over v gives $|\langle \nu, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}| \leq c\|f\|_{\mathcal{B}(\partial\Omega), t}$, $f \in \mathcal{B}(\partial\Omega)$, with a (different) constant $c > 0$. Consequently $\nu \in (\mathcal{B}(\partial\Omega))'$. The rest now follows from (5.1), (5.3) and (5.4). \square

Corollary 5.4. Let Ω be two-sided H^1 -admissible and let μ be a nonnegative Radon measure on $\partial\Omega$ such that $\mathcal{B}(\partial\Omega) \subset L^2(\partial\Omega, \mu)$ is a continuous embedding. Then $L^2(\partial\Omega, \mu) \subset (\mathcal{B}(\partial\Omega))'$, for any $g \in L^2(\partial\Omega, \mu)$ the finite signed measure $g \cdot \mu$ is of finite energy and

$$\mathcal{S}g(x) = \int_{\partial\Omega} G(x-y)g(y)\mu(dy), \quad x \in \mathbb{R}^n \setminus \partial\Omega. \quad (5.5)$$

Remark 5.5.

- (i) Suppose that $\partial\Omega$ is compact, $\text{supp } \nu = \partial\Omega$, $n-2 < d < n$, $c > 0$ and $\nu^\pm(B(x, r)) \leq cr^d$ for all $x \in \partial\Omega$, $0 < r < 1$. Then $x \mapsto G * \nu(x)$ is bounded and continuous on \mathbb{R}^n .
- (ii) If Ω is a bounded Lipschitz domain and μ is the surface measure σ on $\partial\Omega$, then $(\mathcal{B}(\partial\Omega))'$ equals $H^{-1/2}(\partial\Omega)$, and for any $g \in L^2(\partial\Omega) \subset H^{-1/2}(\partial\Omega)$ the right-hand side of (5.5) is a weak solution of (4.11). In this case (5.5) can alternatively be concluded from the uniqueness part of Lemma 4.13.

Now assume that $n \geq 2$. Then one can, with some care, introduce the Newton potential operator $u \mapsto \mathcal{J}u = (|\xi|^{-2}\hat{u})^\vee$. It is well known that \mathcal{J} extends to an isometric isomorphism $\dot{H}^{-1}(\mathbb{R}^n) \rightarrow \dot{H}^1(\mathbb{R}^n)$.

Suppose that Ω is two-sided \dot{H}^1 -admissible and recall the operator (4.8). Similarly as before we may view its dual $\overline{\text{tr}}^*$ as a bounded linear operator $\overline{\text{tr}}^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{H}^{-1}(\mathbb{R}^n)$.

Lemma 5.6. Let $n \geq 2$ and let Ω be a two-sided \dot{H}^1 -admissible domain in \mathbb{R}^n . Then $\overline{\mathcal{S}} = \mathcal{J} \circ \overline{\text{tr}}^*$.

Proof. Given $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, we have $\mathcal{J} \circ \overline{\text{tr}}^* g \in \dot{H}^1(\mathbb{R}^n)$, and for all $v \in \dot{H}^1(\mathbb{R}^n)$ then

$$\begin{aligned} \langle \mathcal{J} \circ \overline{\text{tr}}^* g, v \rangle_{\dot{H}^1(\mathbb{R}^n)} &= \langle \overline{\text{tr}}^* g, v \rangle_{\dot{H}^{-1}(\mathbb{R}^n), \dot{H}^1(\mathbb{R}^n)} = \langle g, \hat{\text{Tr}} v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} \\ &= \langle \dot{\mathcal{S}}g, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle \overline{\mathcal{S}}g, v \rangle_{\dot{H}^1(\mathbb{R}^n)}. \end{aligned} \quad (5.6)$$

\square

Let K denote the Newton kernel, that is, the fundamental solution (Green's function) for Δ on \mathbb{R}^n . It is well known that $\mathcal{J}w = K * w$ for all $w \in L^2(\mathbb{R}^n)$. For a nonnegative Radon measure ν on $\partial\Omega$ the function

$$K * \nu(x) := \int_{\mathbb{R}^n} K(x-y)\nu(dy), \quad x \in \mathbb{R}^n$$

is lower semicontinuous. In the case $n \geq 3$ it takes values in $[0, +\infty]$, in the case $n = 2$ in $(-\infty, +\infty]$, and in either case it is finite on $\mathbb{R}^n \setminus \partial\Omega$. Given a finite signed Borel measure ν , we consider $K * \nu := K * \nu^+ - K * \nu^-$ on $\mathbb{R}^n \setminus \partial\Omega$. We call ν *centered* if $\nu(\mathbb{R}^n) = 0$.

Proposition 5.7. *Let $n \geq 2$, let Ω be \dot{H}^1 -admissible and let ν be a centered finite signed Borel measure on $\partial\Omega$ of finite energy. Then sets of zero capacity have zero ν -measure, and ν defines an element of $(\dot{\mathcal{B}}(\partial\Omega))'$ by*

$$\langle \nu, f \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} := \int_{\partial\Omega} f \, d\nu, \quad f \in \dot{\mathcal{B}}(\partial\Omega). \quad (5.7)$$

Moreover, $K * \nu$ is a representative modulo constants of $\overline{\mathcal{S}}\nu = \mathcal{J} \circ \overline{\text{tr}}^* \nu$.

Proof. By Proposition 5.3 the measure ν charges no sets of zero capacity, and since ν is centered, the right hand side of (5.7) is linear in $f \in \dot{\mathcal{B}}(\partial\Omega)$. Let B be a ball containing $\partial\Omega$, let $\chi \in C_c^\infty(B)$ be such that $0 \leq \chi \leq 1$ and $\chi \equiv 1$ on $\overline{\Omega}$. Given $f \in (\dot{\mathcal{B}}(\partial\Omega))'$, let $v \in \dot{H}^1(\mathbb{R}^n)$ be such that $\text{Tr } v = f$ and let $w \in v$ be such that $\int_B \chi w \, dx = 0$. By Poincaré's inequality for B we find that $\|\chi w\|_{H^1(\mathbb{R}^n)} \leq c \|\chi\|_{C^1(B)} \|v\|_{\dot{H}^1(\Omega)}$ with $c > 0$ independent of v . By (5.2) the function $\tilde{w}|_{\partial\Omega} = (\chi w)^\sim|_{\partial\Omega}$ is integrable with respect to ν^+ and ν^- . Similarly as before we find that

$$\int_{\partial\Omega} \tilde{w} \, d\nu = \langle K * \nu, v \rangle_{\dot{H}^1(\mathbb{R}^n)}, \quad v \in \dot{H}^1(\mathbb{R}^n), \quad (5.8)$$

and this implies that $|\langle \nu, f \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}| \leq c \|f\|_{\dot{\mathcal{B}}(\partial\Omega)}$, $f \in \dot{\mathcal{B}}(\partial\Omega)$, where $c > 0$ is another constant. The rest now follows from (5.6), (5.7) and (5.8). \square

Corollary 5.8. *Let $n \geq 2$, let Ω be two-sided \dot{H}^1 -admissible and let μ be a nonnegative Radon measure on $\partial\Omega$ such that $\mathcal{B}(\partial\Omega) \subset L^2(\partial\Omega, \mu)$ is a continuous embedding. Then $L_0^2(\partial\Omega, \mu) = \{f \in L_0^2(\partial\Omega, \mu) : \int_{\partial\Omega} f \, d\mu = 0\}$ is a subspace of $(\dot{\mathcal{B}}(\partial\Omega))'$, for any $g \in L^2(\partial\Omega, \mu)$ the finite signed measure $g \cdot \mu$ is centered and of finite energy, and $K * (g \cdot \mu)(x) = \int_{\partial\Omega} K(x-y)g(y)\mu(dy)$, $x \in \mathbb{R}^n \setminus \partial\Omega$, is a representative modulo constants of $\overline{\mathcal{S}}g$.*

Remark 5.9. *One can formulate a corresponding analog of Remark 5.5.*

5.2 Representations of double layer potentials

Given a two-sided H^1 -admissible domain $\Omega \subset \mathbb{R}^n$, let \mathcal{R} denote the unique bounded linear operator \mathcal{R} from $L^2(\mathbb{R}^n \setminus \partial\Omega)$ into $H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ such that

$$\langle \mathcal{R}u, v \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle u, v \rangle_{L^2(\mathbb{R}^n \setminus \partial\Omega)}$$

for all $u \in L^2(\mathbb{R}^n \setminus \partial\Omega)$ and $v \in H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$. This operator is symmetric on $L^2(\mathbb{R}^n \setminus \partial\Omega)$ and symmetric with respect to $\langle \cdot, \cdot \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)}$. It extends to an isometric isomorphism from $(H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega))'$ onto $H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$. The adjoint $\left(\frac{\partial}{\partial\nu}\right)^*$ of $\frac{\partial}{\partial\nu}$ as in Lemma 4.2 (iii) maps from $\mathcal{B}(\partial\Omega)$ into $(V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega))'$. Following the philosophy of Remark 2.8, it can be viewed as a bounded linear operator from $\mathcal{B}(\partial\Omega)$ into $(H^1(\mathbb{R}^n \setminus \partial\Omega))'$ if for each $f \in \mathcal{B}(\partial\Omega)$ the functional $\left(\frac{\partial}{\partial\nu}\right)^* f$ is silently extended by zero to a bounded linear functional on all of $H^1(\mathbb{R}^n \setminus \partial\Omega)$. The operator \mathcal{D} can be represented in terms of these operators.

Lemma 5.10. *Let Ω be two-sided H^1 -admissible. Then $\mathcal{D} = -\mathcal{R} \circ \left(\frac{\partial}{\partial \nu}\right)^*$.*

Proof. Given $f \in \mathcal{B}(\partial\Omega)$ and $\varphi \in H_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus V_{1,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ Corollary 4.4 gives

$$\begin{aligned} & -\langle \mathcal{D}f, \varphi \rangle_{(H^1(\mathbb{R}^n \setminus \partial\Omega))', H^1(\mathbb{R}^n \setminus \partial\Omega)} = -\langle \mathcal{R} \circ \mathcal{D}f, \varphi \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = -\langle \mathcal{D}f, \mathcal{R}\varphi \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} \\ & = \left\langle \frac{\partial}{\partial \nu} \circ P_{1,\mathcal{D}} \circ \mathcal{R}\varphi, f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \left\langle \left(\frac{\partial}{\partial \nu}\right)^* f, P_{1,\mathcal{D}} \circ \mathcal{R}\varphi \right\rangle_{(H^1(\mathbb{R}^n \setminus \partial\Omega))', H^1(\mathbb{R}^n \setminus \partial\Omega)} \\ & = \left\langle \left(\frac{\partial}{\partial \nu}\right)^* f, \mathcal{R}\varphi \right\rangle_{(H^1(\mathbb{R}^n \setminus \partial\Omega))', H^1(\mathbb{R}^n \setminus \partial\Omega)} = \left\langle \mathcal{R} \circ \left(\frac{\partial}{\partial \nu}\right)^* f, \varphi \right\rangle_{(H^1(\mathbb{R}^n \setminus \partial\Omega))', H^1(\mathbb{R}^n \setminus \partial\Omega)}. \end{aligned}$$

□

Proposition 5.11. *Let Ω be a bounded Lipschitz domain. Then $\mathcal{B}(\partial\Omega)$ equals $H^{1/2}(\partial\Omega)$, and for any and $f \in H^{1/2}(\partial\Omega)$ we have*

$$\mathcal{D}f(x) = - \int_{\partial\Omega} \frac{\partial}{\partial \nu_y} G(x-y) f(y) \sigma(dy), \quad x \in \mathbb{R}^n \setminus \partial\Omega; \quad (5.9)$$

here $\frac{\partial}{\partial \nu_y}$ denotes the classical interior normal derivative and σ the surface measure on $\partial\Omega$.

Proof. The classical double layer potential on the right hand side of (5.9) is known to solve (4.9) in the weak sense, so the uniqueness part of Lemma 4.8 implies (5.9). □

Given a two-sided \dot{H}^1 -admissible domain $\Omega \subset \mathbb{R}^n$. Let $\dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega)$ denote the closure of $C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$ in $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$. Let $\dot{\mathcal{R}}$ denote the unique bounded linear operator from $(\dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus \dot{V}_{0,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega))'$ into $\dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus \dot{V}_{0,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$ such that

$$\langle \dot{\mathcal{R}}u, v \rangle_{\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)} = \langle u, v \rangle_{(\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega))', \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)}$$

for all $u \in (\dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus \dot{V}_{0,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega))'$ and $v \in \dot{H}_0^1(\mathbb{R}^n \setminus \partial\Omega) \oplus \dot{V}_{0,\mathcal{D}}(\mathbb{R}^n \setminus \partial\Omega)$. With a similar interpretation as before, we obtain the following.

Lemma 5.12. *Let Ω be two-sided \dot{H}^1 -admissible. Then $\dot{\mathcal{D}} = -\dot{\mathcal{R}} \circ \left(\frac{\partial}{\partial \nu}\right)^*$.*

Proposition 5.13. *Let $n \geq 2$, let Ω be a bounded Lipschitz domain. Then $\dot{\mathcal{B}}(\partial\Omega)$ equals $\dot{H}^{1/2}(\partial\Omega)$, and for any $f \in \dot{H}^{1/2}(\partial\Omega)$ the trace jump readjusted version of $\dot{\mathcal{D}}f$ differs only by a constant from $-\int_{\partial\Omega} \frac{\partial}{\partial \nu_y} K(\cdot - y) f(y) \mu(dy)$ on $\mathbb{R}^n \setminus \partial\Omega$.*

6 Neumann-Poincaré operators for admissible domains

6.1 Neumann-Poincaré operators

Suppose that $\Omega \subset \mathbb{R}^n$ is a two-sided H^1 -admissible domain. To the bounded linear operator $\mathcal{K} : \mathcal{B}(\partial\Omega) \rightarrow \mathcal{B}(\partial\Omega)$, defined by

$$\mathcal{K} := \frac{1}{2}(\text{Tr}_i + \text{Tr}_e) \circ \mathcal{D},$$

we refer as the *Neumann-Poincaré operator* for (4.9). By $\mathcal{K}^* : (\mathcal{B}(\partial\Omega))' \rightarrow (\mathcal{B}(\partial\Omega))'$ we denote its dual.

In the sequel we use the symbol I to denote the identity operator, the space on which it acts will be clear from the context.

The following identities generalize results well-known in the Lipschitz case, see [3, Theorem 2.4] or [86].

Theorem 6.1. *Let Ω be two-sided H^1 -admissible.*

- (i) We have $\text{Tr}_i \circ \mathcal{D} = -\frac{1}{2}I + \mathcal{K}$ and $\text{Tr}_e \circ \mathcal{D} = \frac{1}{2}I + \mathcal{K}$.
- (ii) We have $\frac{\partial_i}{\partial \nu} \circ \mathcal{S} = \frac{1}{2}I + \mathcal{K}^*$ and $\frac{\partial_e}{\partial \nu} \circ \mathcal{S} = -\frac{1}{2}I + \mathcal{K}^*$. In particular,

$$\mathcal{K}^* = \frac{1}{2} \left(\frac{\partial_i}{\partial \nu} + \frac{\partial_e}{\partial \nu} \right) \circ \mathcal{S}.$$

Proof. Statement (i) and the first part of (ii) follow from the definitions of \mathcal{D} and \mathcal{S} . To see the second part of (ii), note that for any $f \in \mathcal{B}(\partial\Omega)$ and $g \in (\mathcal{B}(\partial\Omega))'$ we have

$$0 = \langle \mathcal{D}f, \mathcal{S}g \rangle_{H^1(\mathbb{R}^n \setminus \partial\Omega)} = \left\langle \frac{\partial_i \mathcal{S}g}{\partial \nu}, \text{Tr}_i \mathcal{D}f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} - \left\langle \frac{\partial_e \mathcal{S}g}{\partial \nu}, \text{Tr}_e \mathcal{D}f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$$

by Lemma 4.2 (iv), (2.14) and (4.3), and that adding this zero to

$$\left\langle \frac{\partial_i \mathcal{S}g}{\partial \nu}, \text{Tr}_e \mathcal{D}f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} - \left\langle \frac{\partial_e \mathcal{S}g}{\partial \nu}, \text{Tr}_i \mathcal{D}f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$$

respectively subtracting it gives

$$\begin{aligned} 2 \langle g, \mathcal{K}f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} &= \left\langle \left[\frac{\partial \mathcal{S}g}{\partial \nu} \right], \text{Tr}_i \mathcal{D}f + \text{Tr}_e \mathcal{D}f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \\ &= \left\langle \frac{\partial_i \mathcal{S}g}{\partial \nu} + \frac{\partial_e \mathcal{S}g}{\partial \nu}, -\llbracket \text{Tr} \mathcal{D}f \rrbracket \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \left\langle \left(\frac{\partial_i}{\partial \nu} + \frac{\partial_e}{\partial \nu} \right) \circ \mathcal{S}g, f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}. \end{aligned}$$

□

For a two-sided \dot{H}^1 -admissible domain $\Omega \subset \mathbb{R}^n$ similar observations can be made. To the bounded linear operator $\dot{\mathcal{K}} : \dot{\mathcal{B}}(\partial\Omega) \rightarrow \dot{\mathcal{B}}(\partial\Omega)$, defined by

$$\dot{\mathcal{K}} := \frac{1}{2}(\dot{\text{Tr}}_i + \dot{\text{Tr}}_e) \circ \dot{\mathcal{D}},$$

we refer as the *Neumann-Poincaré operator* for (4.10). By $\dot{\mathcal{K}}^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$ we denote its dual.

Theorem 6.2. *Let Ω be two-sided \dot{H}^1 -admissible.*

- (i) We have $\dot{\text{Tr}}_i \circ \dot{\mathcal{D}} = -\frac{1}{2}I + \dot{\mathcal{K}}$ and $\dot{\text{Tr}}_e \circ \dot{\mathcal{D}} = \frac{1}{2}I + \dot{\mathcal{K}}$.
- (ii) We have $\frac{\dot{\partial}_i}{\partial \nu} \circ \dot{\mathcal{S}} = \frac{1}{2}I + \dot{\mathcal{K}}^*$ and $\frac{\dot{\partial}_e}{\partial \nu} \circ \dot{\mathcal{S}} = -\frac{1}{2}I + \dot{\mathcal{K}}^*$. In particular,

$$\dot{\mathcal{K}}^* = \frac{1}{2} \left(\frac{\dot{\partial}_i}{\partial \nu} + \frac{\dot{\partial}_e}{\partial \nu} \right) \circ \dot{\mathcal{S}} = \frac{1}{2} \left(\frac{\dot{\partial}_i}{\partial \nu} + \frac{\dot{\partial}_e}{\partial \nu} \right) \circ \bar{\mathcal{S}},$$

where $\bar{\mathcal{S}}$ is as in Remark 4.13.

6.2 Boundary layer potentials and Calderón projectors

Let Ω be two-sided H^1 -admissible. By Lemma 4.2, Corollary 4.9 and Corollary 4.14 the *boundary single layer potential operator* $\mathcal{V} : (\mathcal{B}(\partial\Omega))' \rightarrow \mathcal{B}(\partial\Omega)$ and the *boundary double layer potential operator* $\mathcal{W} : \mathcal{B}(\partial\Omega) \rightarrow (\mathcal{B}(\partial\Omega))'$, defined by

$$\mathcal{V} := \text{tr} \circ \mathcal{S} \quad \text{respectively} \quad \mathcal{W} := -\frac{\partial}{\partial \nu} \circ \mathcal{D}, \quad (6.1)$$

are bounded linear bijections with bounded inverses.

Remark 6.3. In a classical context the trace $\mathcal{V}g$ on $\partial\Omega$ of the single layer potential $\mathcal{S}g$ can be defined as a pointwise limit, while the (co-)normal derivative $\mathcal{W}f$ of the double layer potential $\mathcal{D}f$ is a hypersingular integral which for smooth enough $\partial\Omega$ and f can be expressed as a principal value. See for instance [80, Section 2] for details; note that our notation differs slightly from the one used there.

Now let $\mathcal{M} : \mathcal{B}(\partial\Omega) \times (\mathcal{B}(\partial\Omega))' \rightarrow \mathcal{B}(\partial\Omega) \times (\mathcal{B}(\partial\Omega))'$ be the linear operator defined by

$$\mathcal{M} := \begin{pmatrix} -\mathcal{K} & \mathcal{V} \\ \mathcal{W} & \mathcal{K}^* \end{pmatrix}$$

and set

$$\mathcal{C}_i := \frac{1}{2}I + \mathcal{M} \quad \text{and} \quad \mathcal{C}_e := \frac{1}{2}I - \mathcal{M}.$$

To the linear operators \mathcal{C}_i and \mathcal{C}_e one refers as the *interior* respectively *exterior Calderón projector* for Ω . We obtain generalizations of known symmetrization formulas referred to as *Calderón relations*, see [48, Lemma 1.2.4], [73, Theorem 3.1.3] or [80, Proposition 5.1].

Theorem 6.4. Let Ω be two-sided H^1 -admissible. The operators \mathcal{C}_i and \mathcal{C}_e are continuous projectors and satisfy $\mathcal{C}_i + \mathcal{C}_e = I$. Moreover, we have $\mathcal{M}^2 = \frac{1}{4}I$, that is,

$$\begin{cases} \mathcal{K}\mathcal{V} = \mathcal{V}\mathcal{K}^*, \\ \mathcal{W}\mathcal{K} = \mathcal{K}^*\mathcal{W}, \end{cases} \quad \text{and} \quad \begin{cases} \mathcal{K}^2 + \mathcal{V}\mathcal{W} = \frac{1}{4}I, \\ (\mathcal{K}^*)^2 + \mathcal{W}\mathcal{V} = \frac{1}{4}I. \end{cases} \quad (6.2)$$

Proof. Continuity and the first identity are clear. To see that $\mathcal{C}_i^2 = \mathcal{C}_i$, suppose that $f \in \mathcal{B}(\partial\Omega)$ and $g \in (\mathcal{B}(\partial\Omega))'$ are given and that $u = \mathcal{S}g - \mathcal{D}f$ is the unique weak solution of (4.1) with $k = 1$, with $\llbracket \frac{\partial u}{\partial \nu} \rrbracket = g$ and with $\llbracket \text{tr } u \rrbracket = f$ in place of $\llbracket \text{tr } u \rrbracket = -f$. Then (6.1) and Theorem 6.1 give

$$\mathcal{C}_i \begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} \mathcal{V}g - \left(-\frac{1}{2}I + \mathcal{K}\right)f \\ \left(\frac{1}{2}I + \mathcal{K}^*\right)g + \mathcal{W}f \end{pmatrix} = \begin{pmatrix} \text{tr } \mathcal{S}g - \text{tr}_i \mathcal{D}f \\ \frac{\partial_i u}{\partial \nu} \mathcal{S}g - \frac{\partial}{\partial \nu} \mathcal{D}f \end{pmatrix} = \begin{pmatrix} \text{tr}_i u \\ \frac{\partial_i u}{\partial \nu} \end{pmatrix} \quad (6.3)$$

Now let v be the unique weak solution of problem (4.1) with $k = 1$, with $\llbracket \text{tr } u \rrbracket = \text{tr}_i u$ and $\llbracket \frac{\partial u}{\partial \nu} \rrbracket = \frac{\partial_i u}{\partial \nu}$. Since also $u\mathbf{1}_\Omega$ is a weak solution of this problem, it follows that $v = u\mathbf{1}_\Omega$ and consequently

$$\mathcal{C}_i^2 \begin{pmatrix} f \\ g \end{pmatrix} = \mathcal{C}_i \begin{pmatrix} \text{tr}_i u \\ \frac{\partial_i u}{\partial \nu} \end{pmatrix} = \begin{pmatrix} \text{tr}_i v \\ \frac{\partial_i v}{\partial \nu} \end{pmatrix} = \begin{pmatrix} \text{tr}_i u \\ \frac{\partial_i u}{\partial \nu} \end{pmatrix} = \mathcal{C}_i \begin{pmatrix} f \\ g \end{pmatrix}.$$

In the same way, we prove $\mathcal{C}_e^2 = \mathcal{C}_e$. The remaining identities then follow. \square

Remark 6.5. The unique weak solution $u = \mathcal{S}g - \mathcal{D}f$ of (4.1) with $k = 1$ and boundary data $f \in \mathcal{B}(\partial\Omega)$ and $g \in (\mathcal{B}(\partial\Omega))'$ is zero on $\mathbb{R}^n \setminus \bar{\Omega}$ if and only if $\text{tr}_e u = 0$ and $\frac{\partial_e u}{\partial \nu} = 0$. By (6.3) this is the case if and only if $\mathcal{C}_i \begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} f \\ g \end{pmatrix}$ and this happens if and only if (f, g) is an element of the graph of the Poincaré-Steklov operator \mathcal{A}_1 for Ω . As in [80, Section 5], an evaluation of (6.3) gives

$$\mathcal{A}_1 = \mathcal{V}^{-1} \left(\frac{1}{2}I + \mathcal{K} \right) = \mathcal{W} + \left(\frac{1}{2}I + \mathcal{K}^* \right) \mathcal{V}^{-1} \left(\frac{1}{2}I + \mathcal{K} \right). \quad (6.4)$$

Now suppose that Ω is two-sided \dot{H}^1 -admissible. By Lemma 4.5, Corollary 4.11 and Corollary 4.16 the operators $\dot{\mathcal{V}} : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{\mathcal{B}}(\partial\Omega)$ and $\dot{\mathcal{W}} : \dot{\mathcal{B}}(\partial\Omega) \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$, defined by

$$\dot{\mathcal{V}} := \text{tr} \circ \dot{\mathcal{S}} \quad \text{and} \quad \dot{\mathcal{W}} := -\frac{\dot{\partial}}{\partial \nu} \circ \dot{\mathcal{D}}, \quad (6.5)$$

are bounded linear bijections with bounded inverses. Let $\dot{\mathcal{M}} : \dot{\mathcal{B}}(\partial\Omega) \times (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow \dot{\mathcal{B}}(\partial\Omega) \times (\dot{\mathcal{B}}(\partial\Omega))'$ be defined by

$$\dot{\mathcal{M}} := \begin{pmatrix} -\dot{\mathcal{K}} & \dot{\mathcal{V}} \\ \dot{\mathcal{W}} & \dot{\mathcal{K}}^* \end{pmatrix}$$

and set

$$\dot{\mathcal{C}}_i := \frac{1}{2}I + \dot{\mathcal{M}} \quad \text{and} \quad \dot{\mathcal{C}}_e := \frac{1}{2}I - \dot{\mathcal{M}},$$

The same arguments as before give the following counterpart of Theorem 6.4.

Theorem 6.6. *Let Ω be two-sided \dot{H}^1 -admissible. The operators $\dot{\mathcal{C}}_i$ and $\dot{\mathcal{C}}_e$ are continuous projectors and satisfy $\dot{\mathcal{C}}_i + \dot{\mathcal{C}}_e = I$. We have $\dot{\mathcal{M}}^2 = \frac{1}{4}I$, that is,*

$$\begin{cases} \dot{\mathcal{K}}\dot{\mathcal{V}} = \dot{\mathcal{V}}\dot{\mathcal{K}}^*, \\ \dot{\mathcal{W}}\dot{\mathcal{K}} = \dot{\mathcal{K}}^*\dot{\mathcal{W}}, \end{cases} \quad \text{and} \quad \begin{cases} \dot{\mathcal{K}}^2 + \dot{\mathcal{V}}\dot{\mathcal{W}} = \frac{1}{4}I, \\ (\dot{\mathcal{K}}^*)^2 + \dot{\mathcal{W}}\dot{\mathcal{V}} = \frac{1}{4}I. \end{cases} \quad (6.6)$$

6.3 Invertibility and isometries

As in the classical case one can observe invertibility properties depending on a spectral parameter $\lambda \in \mathbb{R}$. The following observation is similar to [3, Lemma 2.4].

Lemma 6.7. *Let Ω be two-sided H^1 -admissible and $|\lambda| \geq \frac{1}{2}$. Then $\lambda I + \mathcal{K}^* : (\mathcal{B}(\partial\Omega))' \rightarrow (\mathcal{B}(\partial\Omega))'$ is injective.*

Proof. Assume that there is some nonzero $g \in \ker(\lambda I + \mathcal{K}^*)$. Then $\mathcal{S}g \neq 0$ by Corollary 4.14, and since $\mathcal{S}g$ is 1-harmonic in $\mathbb{R}^n \setminus \partial\Omega$, both $\mathcal{S}g|_\Omega$ and $\mathcal{S}g|_{\mathbb{R}^n \setminus \overline{\Omega}}$ must be nonzero. By Theorem 6.1, together with (2.14) and (4.3), this implies that

$$A := \left(\frac{1}{2} - \lambda\right) \langle g, \text{Tr } \mathcal{S}g \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = \left\langle \left(\frac{1}{2}I + \mathcal{K}^*\right)g, \text{Tr } \mathcal{S}g \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} > 0$$

and

$$B := \left(\frac{1}{2} + \lambda\right) \langle g, \text{Tr } \mathcal{S}g \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} = -\left\langle \left(-\frac{1}{2}I + \mathcal{K}^*\right)g, \text{Tr } \mathcal{S}g \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} > 0.$$

For $|\lambda| = \frac{1}{2}$ this is impossible. For $|\lambda| > \frac{1}{2}$ we can use that $\langle g, \text{Tr } \mathcal{S}g \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} > 0$ by Corollary 4.4. This gives $\lambda = \frac{1}{2} \frac{B-A}{B+A} \in \left[-\frac{1}{2}, \frac{1}{2}\right]$, which again is impossible. \square

The corresponding result for two-sided \dot{H}^1 -admissible domains is seen analogously.

Lemma 6.8. *Let Ω be two-sided H^1 -admissible and $|\lambda| \geq \frac{1}{2}$. Then $\lambda I + \dot{\mathcal{K}}^* : (\dot{\mathcal{B}}(\partial\Omega))' \rightarrow (\dot{\mathcal{B}}(\partial\Omega))'$ is injective.*

Using the operators in (6.1) as metrics, we can introduce alternative Hilbert space norms on the trace spaces and their duals by

$$\|\cdot\|_{\mathcal{B}(\partial\Omega), \mathcal{W}}^2 := \langle \mathcal{W}\cdot, \cdot \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \quad \text{and} \quad \|\cdot\|_{(\mathcal{B}(\partial\Omega))', \mathcal{V}}^2 := \langle \cdot, \mathcal{V}\cdot \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)}$$

in the case of a two-sided H^1 -admissible domain Ω and by

$$\|\cdot\|_{\dot{\mathcal{B}}(\partial\Omega), \dot{\mathcal{W}}}^2 := \langle \dot{\mathcal{W}}\cdot, \cdot \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)} \quad \text{and} \quad \|\cdot\|_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{V}}}^2 := \langle \cdot, \dot{\mathcal{V}}\cdot \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}$$

in the case of a two-sided \dot{H}^1 -admissible domain Ω . The positive definiteness follows from the boundedness of inverses or, alternatively, from Corollaries 4.4 and 4.6 together with (6.1) and (6.5), respectively. On $\mathcal{B}(\partial\Omega)$ and $\dot{\mathcal{B}}(\partial\Omega)$ we also use the Hilbert space norms

$$\|\cdot\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}}^2 := \langle \mathcal{V}^{-1}\cdot, \cdot \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \quad \text{and} \quad \|\cdot\|_{\dot{\mathcal{B}}(\partial\Omega), \dot{\mathcal{V}}^{-1}}^2 := \langle \dot{\mathcal{V}}^{-1}\cdot, \cdot \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)},$$

where \mathcal{V}^{-1} and $\dot{\mathcal{V}}^{-1}$ denote the inverses of \mathcal{V} and $\dot{\mathcal{V}}$, respectively.

Theorem 6.9. *Let Ω be two-sided H^1 -admissible.*

(i) *There are constants $\alpha, \beta > 0$ such that*

$$\frac{1}{\alpha} \|g\|_{(\mathcal{B}(\partial\Omega))'}^2 \leq \|g\|_{(\mathcal{B}(\partial\Omega))', \mathcal{V}}^2 \leq \alpha \|g\|_{(\mathcal{B}(\partial\Omega))'}^2, \quad g \in (\mathcal{B}(\partial\Omega))',$$

and

$$\frac{1}{\beta} \|f\|_{\mathcal{B}(\partial\Omega)}^2 \leq \|f\|_{\mathcal{B}(\partial\Omega), \mathcal{W}}^2 \leq \beta \|f\|_{\mathcal{B}(\partial\Omega)}^2, \quad f \in \mathcal{B}(\partial\Omega).$$

(ii) *The operator \mathcal{S} is an isometry with respect to $\|\cdot\|_{(\mathcal{B}(\partial\Omega))', \mathcal{V}}$, and \mathcal{D} is an isometry with respect to $\|\cdot\|_{\mathcal{B}(\partial\Omega), \mathcal{W}}$.*

(iii) *For any $f \in \mathcal{B}(\partial\Omega)$ we have*

$$(1-c) \|f\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}} \leq \left\| \left(\pm \frac{1}{2} I + \mathcal{K} \right) f \right\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}} \leq c \|f\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}},$$

where

$$c = \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{1}{\alpha\beta}} < 1$$

with constants α and β as in (i) chosen large enough so that $\alpha\beta > 4$. In particular, the operators $\pm \frac{1}{2} I + \mathcal{K} : \mathcal{B}(\partial\Omega) \rightarrow \mathcal{B}(\partial\Omega)$ are isomorphisms.

Proof. Item (i) follows from the boundedness of the operators \mathcal{V} and \mathcal{W} and their inverses. Using Corollaries 4.4, 4.9 and Corollary 4.14 also (ii) follows. Item (iii) can be shown similarly as in [80, p. 744]; note that by (6.4) we have

$$\begin{aligned} \left\| \left(\frac{1}{2} + \mathcal{K} \right) f \right\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}}^2 &= \left\langle \mathcal{V}^{-1} \left(\frac{1}{2} + \mathcal{K} \right) f, \left(\frac{1}{2} + \mathcal{K} \right) f \right\rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} \\ &= \langle \mathcal{A}_1 f, f \rangle_{(\mathcal{B}(\partial\Omega))', \mathcal{B}(\partial\Omega)} - \langle \mathcal{W} f, f \rangle_{\mathcal{B}(\partial\Omega), (\mathcal{B}(\partial\Omega))'} \end{aligned}$$

for any $f \in \mathcal{B}(\partial\Omega)$, the first summand on the right-hand side is

$$\langle \mathcal{V} \mathcal{A}_1 f, f \rangle_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}} \leq \left\| \left(\frac{1}{2} + \mathcal{K} \right) f \right\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}} \|f\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}}$$

and by (i) the second is bounded below by $\frac{1}{\alpha\beta} \|f\|_{\mathcal{B}(\partial\Omega), \mathcal{V}^{-1}}^2$. \square

Theorem 6.10. *Let Ω be two-sided \dot{H}^1 -admissible. Then counterparts of Theorem 6.9 (i), (ii) and (iii) hold for $\dot{\mathcal{V}}, \dot{\mathcal{W}}, \dot{\mathcal{S}}, \dot{\mathcal{D}}$ and $\dot{\mathcal{K}}$.*

We use a result from perturbation theory to establish spectral properties of the Neumann-Poincaré operators, generalizing well-known results in the Lipschitz case, [23, 76].

Theorem 6.11. *Let Ω be two-sided H^1 -admissible. If $|\lambda| \geq \frac{1}{2}$, then the operators $\lambda I + \mathcal{K}$ and $\lambda I + \mathcal{K}^*$ are invertible on $\mathcal{B}(\partial\Omega)$ and $(\mathcal{B}(\partial\Omega))'$ respectively.*

Proof. For any $\lambda \neq -\frac{1}{2}$ we have $\lambda I + \mathcal{K} = (\lambda + \frac{1}{2}) I + (-\frac{1}{2} I + \mathcal{K})$. By Theorem 6.9, $\|-\frac{1}{2} I + \mathcal{K}\| < 1$. Hence, by [36, Theorem 1.2.9], $\lambda I + \mathcal{K}$ is invertible anytime $|\lambda + \frac{1}{2}| \geq 1$, which is valid when $\lambda \geq \frac{1}{2}$. Proceeding in the same way with $\frac{1}{2} I + \mathcal{K}$ instead, we find that $\lambda I + \mathcal{K}$ is invertible for $\lambda \leq -\frac{1}{2}$. The real spectrum of \mathcal{K} is included in $(-\frac{1}{2}, \frac{1}{2})$. By Banach's closed range theorem and Lemma 6.8 the same holds for \mathcal{K}^* , see [36, Theorem 8.1.5]. \square

Analogous arguments give a parallel result in the homogeneous case.

Theorem 6.12. *Let Ω be two-sided \dot{H}^1 -admissible. If $|\lambda| \geq \frac{1}{2}$, then the operators $\lambda I + \dot{\mathcal{K}}$ and $\lambda I + \dot{\mathcal{K}}^*$ are invertible on $\dot{\mathcal{B}}(\partial\Omega)$ and $(\dot{\mathcal{B}}(\partial\Omega))'$ respectively.*

7 Applications to imaging

We generalize two basic results on imaging results that are well known in the case of bounded Lipschitz domains, [3, 59].

7.1 Representation formula

Let Ω and D be two two-sided \dot{H}^1 -admissible domains in \mathbb{R}^n such that $D \subset\subset \Omega$. We write $\dot{\text{Tr}}_{\partial\Omega,i}$ and $\dot{\text{Tr}}_{\partial\Omega,e}$ for the interior and exterior trace and $[[\dot{\text{Tr}}]]_{\partial\Omega}$ for the jump in trace with respect to Ω . Abusing notation slightly, we write $\frac{\dot{\partial}_i}{\partial\nu}|_{\partial\Omega}$ and $\frac{\dot{\partial}_e}{\partial\nu}|_{\partial\Omega}$ to denote the interior and exterior normal derivative in the \dot{H}^1 -sense and $[[\frac{\dot{\partial}}{\partial\nu}]]_{\partial\Omega}$ for the corresponding jump in normal derivatives with respect to $\partial\Omega$. Similarly for ∂D in place of $\partial\Omega$.

Given $k \in (0, 1) \cup (1, +\infty)$ and $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, we call $u \in \dot{H}^1(\Omega)$ a *weak solution* of the Neumann problem

$$\begin{cases} \nabla \cdot \left((1 + (k-1)\mathbb{1}_D) \nabla u \right) = 0 & \text{on } \Omega, \\ \frac{\dot{\partial}_i u}{\partial\nu} |_{\partial\Omega} = g \end{cases} \quad (7.1)$$

in the \dot{H}^1 -sense if for any $v \in \dot{H}^1(\Omega)$ we have

$$\int_{\Omega} (1 + (k-1)\mathbb{1}_D) \nabla u \cdot \nabla v \, dx = \langle g, \dot{\text{Tr}}_{\partial\Omega,i} v \rangle_{(\dot{\mathcal{B}}(\partial\Omega))', \dot{\mathcal{B}}(\partial\Omega)}.$$

Since $\min(1, k) \leq 1 + (k-1)\mathbb{1}_D \leq \max(1, k)$, the existence of a unique weak solution u is clear from the Riesz representation theorem.

We write $\dot{\mathcal{S}}_{\partial\Omega}$ and $\dot{\mathcal{S}}_{\partial D}$ for the single layer potential operators with respect to Ω and D , $\dot{\mathcal{D}}_{\partial\Omega}$ for the double layer operator with respect to $\partial\Omega$ and $\dot{\mathcal{K}}_{\partial D}$ for the Neumann-Poincaré operator with respect to ∂D , all in the \dot{H}^1 -sense. The following representation formula for u is a generalization of [3, Theorem 2.17].

Theorem 7.1. *Let Ω and D two two-sided \dot{H}^1 -admissible such that $D \subset\subset \Omega$ and let $k \in (0, 1) \cup (1, +\infty)$. Given $g \in (\dot{\mathcal{B}}(\partial\Omega))'$, let $u \in \dot{H}^1(\Omega)$ be the unique weak solution of (7.1) in the \dot{H}^1 -sense, let $f := \dot{\text{Tr}}_{\partial\Omega,i} u$ and $H = \dot{\mathcal{S}}_{\partial\Omega} g - \dot{\mathcal{D}}_{\partial\Omega} f$. Then we have*

$$u = H - \dot{\mathcal{S}}_{\partial D} \varphi, \quad (7.2)$$

seen as an equality in $\dot{H}^1(\Omega)$, where φ is the unique element of $(\dot{\mathcal{B}}(\partial D))'$ such that

$$\left(\frac{k+1}{2(k-1)} I + \dot{\mathcal{K}}_{\partial D}^* \right) \varphi = \frac{\dot{\partial} H}{\partial\nu} |_{\partial D}. \quad (7.3)$$

The function $H - \dot{\mathcal{S}}_{\partial D} \varphi$ is constant on $\mathbb{R}^n \setminus \overline{\Omega}$.

The function $H|_{\Omega}$ is the harmonic part of u in Ω , while $(-\dot{\mathcal{S}}_{\partial D} \varphi)|_{\Omega}$ is its refraction part. The existence and uniqueness of $\varphi \in (\dot{\mathcal{B}}(\partial D))'$ solving (7.3) follows from Theorem 6.12.

Proof. Let $w \in \dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ be the unique weak solution in the \dot{H}^1 -sense of the transmission problem

$$\begin{cases} \nabla \cdot \left((1 + (k-1)\mathbb{1}_D) \nabla w \right) = 0 & \text{on } \mathbb{R}^n \setminus \partial\Omega, \\ w_i |_{\partial\Omega} - w_e |_{\partial\Omega} = f \\ \frac{\dot{\partial}_i w_i}{\partial\nu} |_{\partial\Omega} - \frac{\dot{\partial}_e w_e}{\partial\nu} |_{\partial\Omega} = g, \end{cases} \quad (7.4)$$

the definition is completely analogous to that in Subsection 4.5. Similarly as for (7.1), a unique weak solution w in the \dot{H}^1 -sense exists by comparability of quadratic forms. The function w , defined by $w := u$ on Ω and $w := 0$ on $\mathbb{R}^n \setminus \overline{\Omega}$ is a weak solution of (7.4) in

the \dot{H}^1 -sense. We now show that also $\tilde{w} := H - \dot{S}_{\partial D}\varphi$ is a weak solution of (7.4) in the \dot{H}^1 -sense. Using Corollary 4.16 and Theorem 6.2 (ii) the identity (7.3) can be rewritten as

$$k \frac{\dot{\partial}_i(\dot{S}_{\partial D}\varphi)}{\partial\nu} \Big|_{\partial D} - \frac{(\dot{\partial}_e \dot{S}_{\partial D}\varphi)}{\partial\nu} \Big|_{\partial D} = (k-1) \frac{\dot{\partial}H}{\partial\nu} \Big|_{\partial D},$$

and since $H \in \dot{V}_0(\mathbb{R}^n \setminus \partial\Omega)$, it follows that $k \frac{\dot{\partial}_i \tilde{w}}{\partial\nu} \Big|_{\partial D} - \frac{\dot{\partial}_e \tilde{w}}{\partial\nu} \Big|_{\partial D} = 0$. Given arbitrary $v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$, we find that

$$k \int_D \nabla \tilde{w} \nabla v \, dx + \int_{\Omega \setminus \overline{D}} \nabla \tilde{w} \nabla v \, dx = \left\langle k \frac{\dot{\partial}_i \tilde{w}}{\partial\nu} \Big|_{\partial D} - \frac{\dot{\partial}_e \tilde{w}}{\partial\nu} \Big|_{\partial D}, \dot{\text{Tr}}_{\partial D} v \right\rangle_{(\mathcal{B}(\partial D))', \mathcal{B}(\partial D)} = 0.$$

By definition H satisfies the boundary conditions in (7.4) and since $\dot{S}_{\partial D}\varphi$ is harmonic on $\mathbb{R}^n \setminus \overline{D} \supset \partial\Omega$, it does not alter these boundary conditions. Consequently \tilde{w} is a weak solution of (7.4) in the \dot{H}^1 -sense, and $u = \tilde{w}$ in $\dot{H}^1(\mathbb{R}^n \setminus \partial\Omega)$ by uniqueness.

If (7.2) holds with $\tilde{H} \in \dot{V}_0(\Omega)$ and $\tilde{\varphi} \in (\dot{\mathcal{B}}(\partial\Omega))'$ in place of H and φ , then $H - \dot{S}_{\partial D}\varphi = \tilde{H} - \dot{S}_{\partial D}\tilde{\varphi}$, consequently $\dot{S}_{\partial D}(\varphi - \tilde{\varphi}) \in \dot{V}_0(\Omega)$ and

$$\left[\frac{\dot{\partial}}{\partial\nu} \dot{S}_{\partial D}(\varphi - \tilde{\varphi}) \right]_{\partial D} = 0.$$

But this implies that $\tilde{\varphi} = \varphi$ and therefore $\tilde{H} = H$. □

7.2 Subdomain identification

We give a generalization of a theorem on the identification of “monotone” inclusions through a single boundary measurement. For Lipschitz domains it can be found as a part of [3, Theorem A.7], the formulation below works for two-sided \dot{H}^1 -admissible domains.

Theorem 7.2. *Let Ω , D_1 and D_2 be two-sided \dot{H}^1 -admissible and such that $D_1 \subset D_2 \subset\subset \Omega$ and let $k \in (0, 1) \cup (1, +\infty)$. Suppose that $g \in (\dot{\mathcal{B}}(\partial\Omega))'$ is nonzero and that u_1 and u_2 are the unique weak solutions in the \dot{H}^1 -sense of (7.1) with D_1 and D_2 in place of D , respectively. Then $\dot{\text{Tr}}_{\partial\Omega, i} u_1 = \dot{\text{Tr}}_{\partial\Omega, i} u_2$ implies $D_1 = D_2$.*

Proof. Since

$$\int_{\Omega} (1 + (k-1)\mathbb{1}_{D_1}) \nabla u_1 \nabla v \, dx = \int_{\Omega} (1 + (k-1)\mathbb{1}_{D_2}) \nabla u_2 \nabla v \, dx$$

for all $v \in \dot{H}^1(\Omega)$, it follows that

$$\int_{\Omega} (1 + (k-1)\mathbb{1}_{D_1}) \nabla(u_1 - u_2) \nabla v \, dx = (k-1) \int_{D_2 \setminus D_1} \nabla u_2 \nabla v \, dx.$$

Testing with $v = u_1 - u_2$ gives

$$\begin{aligned} \int_{\Omega} (1 + (k-1)\mathbb{1}_{D_1}) |\nabla(u_1 - u_2)|^2 \, dx + (k-1) \int_{D_2 \setminus D_1} |\nabla u_2|^2 \, dx \\ = (k-1) \int_{D_2 \setminus D_1} \nabla u_2 \nabla u_1 \, dx, \end{aligned} \quad (7.5)$$

and testing with $v = u_1$ shows that the right-hand side equals

$$\int_{\Omega} (1 + (k-1)\mathbb{1}_{D_1}) \nabla(u_1 - u_2) \nabla u_1 \, dx. \quad (7.6)$$

Since u_1 is harmonic with respect to the equivalent scalar product

$$(u, w) \mapsto \int_{\Omega} (1 + (k-1)\mathbb{1}_{D_1}) \nabla u \nabla w \, dx$$

on $\dot{H}^1(\Omega) \times \dot{H}^1(\Omega)$ and $u_1 - u_2 \in \ker \dot{\text{Tr}}_{\partial\Omega, i}$, an analog of Lemma 2.10 (i) shows that (7.6) is zero and therefore also (7.5).

In the case $k > 1$ then necessarily $u_1 = u_2$ in $\dot{H}^1(\Omega)$. For $0 < k < 1$ the result follows basically the same arguments but with D_1 and D_2 interchanged. \square

A Comments and proofs

A.1 Comments and proofs for Section 2

Lemma A.1. *If $\Omega \subset \mathbb{R}^n$ is an H^1 -extension domain (resp. \dot{H}^1 -extension domain) and $\mathbb{R}^n \setminus \Omega$ is contained in an open ball B , then $\Omega \cap B$ is an H^1 -extension domain (resp. \dot{H}^1 -extension domain).*

Proof. Let V be an open set containing $\mathbb{R}^n \setminus \Omega$ and such that $\bar{V} \subset B$. Let $\chi \in C_c^\infty(B)$ be such that $0 \leq \chi \leq 1$ and $\chi \equiv 1$ on \bar{V} . Given $u \in H^1(\Omega \cap B)$, the continuation by zero of χu to all of Ω is in $H^1(\Omega)$ and $\|\chi u\|_{H^1(\Omega)} \leq \|\chi\|_{C^1(B)} \|u\|_{H^1(\Omega \cap B)}$. Similarly, the continuation by zero of $(1 - \chi)u$ to all of B is in $H^1(B)$ and $\|\chi u\|_{H^1(B)} \leq \|\chi\|_{C^1(B)} \|u\|_{H^1(\Omega \cap B)}$. The sum $E_{\Omega \cap B} u := E_\Omega(\chi u) + E_B((1 - \chi)u)$ is in $H^1(\mathbb{R}^n)$ and is linear in u , satisfies $E_{\Omega \cap B} u|_{\Omega \cap B} = u$ a.e. for all $u \in H^1(\Omega \cap B)$ and

$$\|E_{\Omega \cap B} u\|_{H^1(\mathbb{R}^n)} \leq \|E_\Omega(\chi u)\|_{H^1(\mathbb{R}^n)} + \|E_B((1 - \chi)u)\|_{H^1(\mathbb{R}^n)} \leq c \|\chi\|_{C^1(B)} \|u\|_{H^1(\Omega \cap B)}.$$

The arguments for an \dot{H}^1 -extension domain are similar. \square

We provide arguments and references for Lemma 2.6 (i).

Proof of Lemma 2.6 (i). We consider the quadratic form defined by (2.2) with $\Omega = \mathbb{R}^n$ and the domain $H^1(\mathbb{R}^n)$. For (i) it suffices to note that by [41, Corollary 2.3.1 and Example 2.3.1] we have

$$H_0^1(\mathbb{R}^n \setminus \partial\Omega) = \{w \in H^1(\mathbb{R}^n) \mid \tilde{w} = 0 \text{ q.e. on } \partial\Omega\}, \quad (\text{A.1})$$

where \tilde{w} is a quasi continuous version of w . Now $u \in H^1(\Omega)$ has an extension to an element of $H_0^1(\mathbb{R}^n \setminus \partial\Omega)$ if and only if $u \in H_0^1(\Omega)$, and it has an extension to an element of the right-hand side of (A.1) if and only if $\text{Tr } u = 0$. \square

A proof of Lemma 2.10 (i) is given in [41, Example 2.3.2], just in a slightly different language. We sketch how to adapt it to our formulation. For any open set $\Omega \subset \mathbb{R}^n$ the bilinear form (2.2), endowed with the domain $H^1(\Omega)$, is a Dirichlet form. By $H_e^1(\Omega)$ we denote its extended Dirichlet space. For $\Omega = \mathbb{R}^n$ it is well-known that

$$\text{if } n \geq 3, \text{ then } (H_e^1(\mathbb{R}^n), \langle \cdot, \cdot \rangle_{\dot{H}^1(\mathbb{R}^n)}) \cong (\dot{H}^1(\mathbb{R}^n), \langle \cdot, \cdot \rangle_{\dot{H}^1(\mathbb{R}^n)}) \quad (\text{A.2})$$

in the sense that the vector spaces are isomorphic and the isomorphism is a Hilbert space isometry, and

$$\text{if } n \leq 2, \text{ then } (H_e^1(\mathbb{R}^n), \langle \cdot, \cdot \rangle_{\dot{H}^1(\mathbb{R}^n)}) = (\dot{H}^1(\mathbb{R}^n) \oplus \mathbb{R}, \langle \cdot, \cdot \rangle_{\dot{H}^1(\mathbb{R}^n)}), \quad (\text{A.3})$$

seen as an equality of vector spaces endowed with bilinear forms. Proofs can for instance be found in [24, Theorems 2.2.12 and 2.2.13].

Proof of Lemma 2.10 (i). An element $u \in \dot{H}^1(\Omega)$ is in $\ker \dot{\text{Tr}}$ if and only if it is the restriction to Ω of an element of

$$\{w \in \dot{H}^1(\mathbb{R}^n) \mid \tilde{w} = 0 \text{ q.e. on } \partial\Omega\}, \quad (\text{A.4})$$

where $\tilde{w} = 0$ is understood representative wise and modulo constants. For $n \geq 3$ the space (A.4) may be identified with

$$\{v \in H_e^1(\mathbb{R}^n) \mid \tilde{v} = 0 \text{ q.e. on } \partial\Omega\} \quad (\text{A.5})$$

under the isometry in (A.2). By [41, Theorem 2.3.3 and Example 2.3.2] the space (A.5) coincides with $H_{0,e}^1(\mathbb{R}^n \setminus \partial\Omega)$, the extended Dirichlet space of (2.2), endowed with the smaller domain $H_0^1(\mathbb{R}^n \setminus \partial\Omega)$. The complement of $H_{0,e}^1(\mathbb{R}^n \setminus \partial\Omega)$ in $H_e^1(\mathbb{R}^n)$ is isometric to

$$\{w \in \dot{H}^1(\mathbb{R}^n) \mid \int_{\mathbb{R}^n} \nabla w \nabla v \, dx = 0 \text{ for all } v \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)\}, \quad (\text{A.6})$$

and since $\mathbb{R}^n \setminus \partial\Omega$ is the disjoint union of Ω and $\mathbb{R}^n \setminus \overline{\Omega}$, the restriction of this space to Ω is $\dot{V}_0(\Omega)$. For $n = 2$ it is shown in [41, Example 2.3.2] that the space (A.5) still coincides with $H_{0,e}^1(\mathbb{R}^n \setminus \partial\Omega)$, and (simplifications of) the same arguments give this coincidence also for $n = 1$. By (A.3) the space (A.5) may then be regarded as a closed subspace of $\dot{H}^1(\mathbb{R}^n)$, and the complement of this closed subspace is (A.6). \square

References

- [1] Y. ACHDOU AND N. TCHOU, *Trace results on domains with self-similar fractal boundaries*, J. Math. Pures Appl., 89 (2008), pp. 596–623.
- [2] H. AIKAWA, *Equivalence between the boundary Harnack principle and the Carleson estimate*, Math. Scand., 103 (2008), pp. 61–76.
- [3] H. AMMARI AND H. KANG, *Reconstruction of Small Inhomogeneities from Boundary Measurements*, Lecture Notes in Mathematics, Springer Berlin Heidelberg New York, 2004.
- [4] H. AMMARI, H. KANG., AND H. LEE, *Layer Potential Techniques in Spectral Analysis*, vol. 153, Mathematical Survey and Monographs, 2009.
- [5] K. ANDO, H. KANG, Y. MIYANISHI, AND M. PUTINAR, *Spectral analysis of Neumann-Poincaré operator*, Rev. Roumaine Math. Pures Appl., 66 (2021), pp. 545–575.
- [6] W. ARENDT AND R. MAZZEO, *Friedlander’s eigenvalue inequalities and the Dirichlet-to-Neumann semigroup*, Communications on Pure and Applied Analysis, 11 (2012), pp. 2201–2212.
- [7] K. ARFI AND A. ROZANOVA-PIERRAT, *Dirichlet-to-Neumann or Poincaré-Steklov operator on fractals described by d -sets*, Discrete & Continuous Dynamical Systems - S, 12 (2019), pp. 1–26.
- [8] J. AZZAM, S. HOFMANN, J. M. MARTELL, K. NYSTRÖM, AND T. TORO, *A new characterization of chord-arc domains*, Journal of the European Mathematical Society, 19 (2017), pp. 967–981.
- [9] H. BAHOURI, J.-Y. CHEMIN, AND R. DANCHIN, *Fourier Analysis and Nonlinear Partial Differential Equations*, vol. 343 of Grundlehren math. Wiss., Springer, Berlin, 2011.
- [10] J. BANNISTER, A. GIBBS, AND D. P. HEWETT, *Acoustic scattering by impedance screens/cracks with fractal boundary: well-posedness analysis and boundary element approximation*, Math. Models Methods Appl. Sci., 32 (2022), pp. 291–319.
- [11] A. BARTON, *Layer potentials for general linear elliptic systems*, Electronic Journal of Differential Equations, 2017 (2017), pp. 1–23.
- [12] A. BARTON AND S. MAYBORODA, *Layer potentials and boundary-value problems for second order elliptic operators with data in Besov spaces*, Mem. Amer. Math. Soc., 243 (2016).

- [13] R. F. BASS AND K. BURDZY, *A boundary Harnack principle in twisted Hölder domains*, Ann. of Math. (2), 134 (1991), pp. 253–276.
- [14] H. BELLOUT AND A. FRIEDMAN, *Identification problems in potential theory*, Archive for Rational Mechanics and Analysis, 101 (1988), pp. 143–160.
- [15] M. BIEGERT, *On traces of Sobolev functions on the boundary of extension domains*, Proceedings of the American Mathematical Society, 137 (2009), pp. 4169–4176.
- [16] N. BOULEAU AND F. HIRSCH, *Dirichlet Forms and Analysis on Wiener Space*, de Gruyter, Berlin, 1991.
- [17] A. M. CAETANO, S. N. CHANDLER-WILDE, A. GIBBS, D. P. HEWETT, AND A. MOIOLA, *A Hausdorff-measure boundary element method for acoustic scattering by fractal screens*, (2022).
- [18] A. M. CAETANO, D. P. HEWETT, AND A. MOIOLA, *Density results for Sobolev, Besov and Triebel-Lizorkin spaces on rough sets*, J. Funct. Anal., 281 (2021), pp. Paper No. 109019, 43.
- [19] A.-P. CALDERON, *Cauchy integrals on Lipschitz curves and related operators*, Proc. Natl. Acad. Sci. USA, 74 (1977), pp. 33–49.
- [20] L. CAPOGNA, C. E. KENIG, AND L. LANZANI, *Harmonic measure. Geometric and analytic points of view*, vol. 35 of University Lecture Series, American Mathematical Society, Providence, RI, 2005.
- [21] T. CARLEMAN, *Über das Neumann–Poincarésche Problem für ein Gebiet mit Ecken*, Almqvist and Wiksells, Uppsala, 1916.
- [22] S. N. CHANDLER-WILDE, D. P. HEWETT, AND A. MOIOLA, *Sobolev spaces on non-Lipschitz subsets of \mathbb{R}^n with application to boundary integral equations on fractal screens*, Integral Equations and Operator Theory, 87 (2017), pp. 179–224.
- [23] T. CHANG AND K. LEE, *Spectral properties of the layer potentials on Lipschitz domains*, Illinois Journal of Mathematics, 52 (2008), pp. 463–472.
- [24] Z.-Q. CHEN AND M. FUKUSHIMA, *Symmetric Markov Processes, Time Change, and Boundary Theory*, London Mathematical Society Monographs, Princeton University Press, Princeton, 2012.
- [25] R. COIFMAN, A. MCINTOSH, AND Y. MEYER, *L 'intégrale de Cauchy définit un opérateur borné sur l^2 pour les courbes Lipschitziennes*, Ann. of Math., 116 (1982), pp. 361–388.
- [26] C. CONSTANTINESCU AND A. CORNEA, *Ideale Ränder Riemannscher Flächen*, vol. 32 of Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, Berlin, 1963.
- [27] M. COSTABEL, *Boundary integral operators on Lipschitz domains: Elementary results*, SIAM Journal Math. Anal., 19 (1988), pp. 613–626.
- [28] M. COSTABEL AND W. WENDLAND, *Strong ellipticity of boundary integral operators*, J. Reine Angew. Math., 372 (1988), pp. 34–63.
- [29] R. COURANT AND D. HILBERT, *Methods of Mathematical Physics, vol. I*, Springer, New York, 1937.
- [30] B. DAHLBERG, *On estimates for harmonic measure*, Arch. Rat. Mech. Analysis, 65 (1977), pp. 272–288.

- [31] B. DAHLBERG, C. KENIG, AND G. VERCHOTA, *Boundary value problems for the systems of elastostatics in Lipschitz domains*, Duke Math. J, 57 (1988), pp. 795–818.
- [32] B. DAHLBERG AND G. VERCHOTA, *Galerkin methods for the boundary integral equations of elliptic equations in nonsmooth domains*, Contemp. Math., 107 (1990), pp. 39–60.
- [33] G. DAVID AND D. JERISON, *Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals*, Indiana Univ. Math. J., 39 (1990), pp. 831–845.
- [34] G. DAVID AND S. SEMMES, *Singular integrals and rectifiable sets in \mathbb{R}^n : Beyond Lipschitz graphs*, Astérisque, 193 (1991).
- [35] ———, *Analysis of and on Uniformly Rectifiable Sets*, AMS Series, Amer. Math. Soc., Providence, 1993.
- [36] E. B. DAVIES, *Linear Operators and their Spectra*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2007.
- [37] J. DENY AND J.-L. LIONS, *Les espaces du type de Beppo Levi*, Annales de l’institut Fourier, 5 (1954), pp. 305–370.
- [38] L. ESCAURIAZA, E. B. FABES, AND G. VERCHOTA, *On a regularity theorem for weak solutions to transmission problems with internal Lipschitz boundaries*, Proceedings of the American Mathematical Society, 115 (1992), pp. 1069–1076.
- [39] E. B. FABES, M. JODEIT, AND N. M. RIVIÈRE, *Potential techniques for boundary value problems on C^1 -domains*, Acta Mathematica, 141 (1978), pp. 165–186.
- [40] L. FRIEDLANDER, *Some inequalities between Dirichlet and Neumann eigenvalues*, Arch. Rational Mech. Anal., 11 (1991), pp. 153–160.
- [41] M. FUKUSHIMA, Y. OSHIMA, AND M. TAKEDA, *Dirichlet Forms and Symmetric Markov Processes*, deGruyter, Berlin, New York, 1994.
- [42] G. GALDI, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations. Steady State Problems*, Springer, New York, 2011.
- [43] A. GIBBS, D. HEWETT, AND A. MOIOLA, *Numerical quadrature for singular integrals on fractals*, Numer. Algorithms, 92 (2023), pp. 2071–2124.
- [44] P. HAJŁASZ, P. KOSKELA, AND H. TUOMINEN, *Sobolev embeddings, extensions and measure density condition*, Journal of Functional Analysis, 254 (2008), pp. 1217–1234.
- [45] M. HINZ, F. MAGOULÈS, A. ROZANOVA-PIERRAT, M. RYNKOVSKAYA, AND A. TEPLYAEV, *On the existence of optimal shapes in architecture*, Applied Mathematical Modelling, 94 (2021), pp. 676–687.
- [46] M. HINZ, A. ROZANOVA-PIERRAT, AND A. TEPLYAEV, *Non-Lipschitz uniform domain shape optimization in linear acoustics*, SIAM Journal on Control and Optimization, 59 (2021), pp. 1007–1032.
- [47] ———, *Boundary value problems on non-Lipschitz uniform domains: stability, compactness and the existence of optimal shapes*, Asymptotic Analysis, (2023), pp. 1–37.
- [48] G. C. HSIAO AND W. L. WENDLAND, *Boundary Integral Equations*, Applied Mathematical Sciences 164, Springer-Verlag Berlin Heidelberg, 2008.
- [49] D. S. JERISON AND C. E. KENIG, *The Dirichlet problem in nonsmooth domains*, Ann. of Math. (2), 113 (1981), pp. 367–382.

- [50] D. S. JERISON AND C. E. KENIG, *Boundary behavior of harmonic functions in nontangentially accessible domains*, Advances in Mathematics, 46 (1982), pp. 80–147.
- [51] P. JONES, *Extension theorems for BMO*, Indiana University Mathematics Journal, 29 (1980), p. 41.
- [52] P. W. JONES, *Quasiconformal mappings and extendability of functions in Sobolev spaces*, Acta Mathematica, 147 (1981), pp. 71–88.
- [53] P. W. JONES, *A geometric localization theorem*, Adv. in Math., 46 (1982), pp. 71–79.
- [54] A. JONSSON, *Besov spaces on closed subsets of \mathbb{R}^n* , Transactions of the American Mathematical Society, 341 (1994), pp. 355–370.
- [55] A. JONSSON AND H. WALLIN, *Function spaces on subsets of \mathbb{R}^n* , Math. Reports 2, Part 1, Harwood Acad. Publ. London, 1984.
- [56] H. KANG, *Spectral Geometry and Analysis of the Neumann-Poincaré Operator, a Review*, in Recent progress in mathematics, N.-G. Kang, J. Choe, K. Choi, and S.-h. Kim, eds., vol. 1 of KIAS Springer Ser. Math., Springer Nature Singapore, Singapore, 2022, pp. 119–153.
- [57] H. KANG AND J. K. SEO, *The layer potential technique for the inverse conductivity problem*, Inverse Problems, 12 (1996), pp. 267–278.
- [58] ———, *Identification of domains with near-extreme conductivity: global stability and error estimates*, Inverse Problems, 15 (1999), pp. 851–867.
- [59] H. KANG AND J. K. SEO, *Recent Progress in the Inverse Conductivity Problem with Single Measurement*, in Inverse problems and related topics, Chapman and Hall/CRC, 2000. Num Pages: 12.
- [60] D. KHAVINSON, M. PUTINAR, AND H. S. SHAPIRO, *Poincaré’s variational problem in potential theory*, Archive for Rational Mechanics and Analysis, 185 (2006), pp. 143–184.
- [61] M. R. LANCIA, *A transmission problem with a fractal interface*, Zeitschrift für Analysis und ihre Anwendungen, 21 (2002), pp. 113–133.
- [62] M. LAPIDUS, *Fractal drum, inverse spectral problem for elliptic operators and a partial resolution of the Weyl- Berry conjecture*, Trans. Amer. Math. Soc., 325 (1991), pp. 465–529.
- [63] M. L. LAPIDUS AND M. M. H. PANG, *Eigenfunctions of the Koch snowflake domain*, Comm. Math. Phys., 172 (1995), pp. 359–376.
- [64] Y. LEJAN, *Mesures associées à une forme de Dirichlet. Applications*, Bull. S.M.F., 106 (1978), pp. 61–112.
- [65] M. LEVITIN AND D. VASSILIEV, *Spectral asymptotics, renewal theorem, and the Berry conjecture for a class of fractals*, Proceedings of the London Mathematical Society, s3-72 (1996), pp. 188–214.
- [66] O. MARTIO AND J. SARVAS, *Injectivity theorems in plane and space*, Annales Academiae Scientiarum Fennicae Series A I Mathematica, 4 (1979), pp. 383–401.
- [67] V. G. MAZ’YA, *Boundary integral equations*, in Analysis IV. Linear and Boundary Integral Equations, V. G. Maz’ya and S. M. Nikolskij, eds., vol. 27 of Encycl. Math. Sci., Springer, Berlin, 1991, pp. 127–222.
- [68] V. G. MAZ’JA, *Sobolev Spaces*, Springer-Verlag, Berlin, 1985.

- [69] W. MCLEAN, *Strongly elliptic systems and boundary integral equations*, Cambridge University Press, 2000.
- [70] D. MITREA, I. MITREA, AND M. MITREA, *Geometric Harmonic Analysis I: A Sharp Divergence Theorem with Nontangential Pointwise Traces*, vol. 75 of *Developments in Mathematics*, Springer Nature, Switzerland, 2023.
- [71] ———, *Geometric Harmonic Analysis IV: Boundary Layer Potentials in Uniformly Rectifiable Domains, and Applications to Complex Analysis*, vol. 75 of *Developments in Mathematics*, Springer Nature, Switzerland, 2023.
- [72] M. MITREA AND M. TAYLOR, *Boundary layer methods for Lipschitz domains in Riemannian manifolds*, *J. Funct. Anal.*, 163 (1999), pp. 181–251.
- [73] J.-C. NÉDÉLEC, *Acoustic and Electromagnetic Equations. Integral Representations for Harmonic Problems*, vol. 144 of *Applied Mathematical Sciences*, Springer, New York, 2001.
- [74] C. NEUMANN, *Zur Theorie des logarithmischen und des Newtonschen Potentials*, *Ber. Verh. Math.-Phys. Classe Königl. Sächs. Akad. Wiss. Leipzig*, 32 (1870), pp. 49–56, 264–321.
- [75] K. NYSTRÖM, *Integrability of Green potentials in fractal domains*, *Arkiv för Matematik*, 34 (1996), pp. 335–381.
- [76] K.-M. PERFEKT AND M. PUTINAR, *The essential spectrum of the Neumann–Poincaré operator on a domain with corners*, *Archive for Rational Mechanics and Analysis*, 223 (2017), pp. 1019–1033.
- [77] A. ROZANOVA-PIERRAT, *Generalization of Rellich-Kondrachov theorem and trace compactness in the framework of irregular and fractal boundaries*, M.R. Lancia, A. Rozanova-Pierrat (Eds.), *Fractals in engineering: Theoretical aspects and Numerical approximations*, 8, *ICIAM 2019 SEMA SIMAI Springer Series Springer Intl. Publ.*, 2021.
- [78] W. RUDIN, *Functional Analysis*, McGraw-Hill, Inc., Singapore, 1991.
- [79] C. SIMADER AND H. SOHR, *The Dirichlet Problem for the Laplacian in Bounded and Unbounded Domains*, Chapman and Hall, London, 1996.
- [80] O. STEINBACH AND W. WENDLAND, *On C. Neumann’s method for second-order elliptic systems in domains with non-smooth boundaries*, *Journal of Mathematical Analysis and Applications*, 262 (2001), pp. 733–748.
- [81] T. TORO, *Analysis and geometry on non-smooth domains*, *Rev. Acad. Colombiana Cienc. Exact. Fís. Natur.*, 41 (2017), pp. 521–527.
- [82] H. TRIEBEL, *Higher Analysis*, J.A. Barth, Leipzig, 1992.
- [83] H. TRIEBEL, *Theory of Function Spaces*, Springer, Basel, 2010.
- [84] J. VÄISÄLÄ, *Uniform domains*, *Tohoku Math. J.*, 40 (1988), pp. 101–118.
- [85] M. VAN DEN BERG, *Heat equation on the arithmetic von Koch snowflake*, *Probability Theory and Related Fields*, 118 (2000), pp. 17–36.
- [86] G. VERCHOTA, *Layer potentials and regularity for the Dirichlet problem for Laplace’s equation in Lipschitz domains*, *Journal of Functional Analysis*, 59 (1984), pp. 572–611.
- [87] V. VLADIMIROV, *Equations of Mathematical Physics*, Pure and Applied Mathematics, 1971.

- [88] H. WALLIN, *The trace to the boundary of Sobolev spaces on a snowflake*, *Manuscripta Math*, 73 (1991), pp. 117–125.
- [89] H. WEYL, *The method of orthogonal projection in potential theory*, *Duke Math. J*, 7 (1940), pp. 411–444.