



**HAL**  
open science

# Singular limit of a chemotaxis model with indirect signal production and phenotype switching

Philippe Laurençot, Christian Stinner

► **To cite this version:**

Philippe Laurençot, Christian Stinner. Singular limit of a chemotaxis model with indirect signal production and phenotype switching. 2024. hal-04511472

**HAL Id: hal-04511472**

**<https://hal.science/hal-04511472>**

Preprint submitted on 19 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SINGULAR LIMIT OF A CHEMOTAXIS MODEL WITH INDIRECT SIGNAL PRODUCTION AND PHENOTYPE SWITCHING

PHILIPPE LAURENÇOT AND CHRISTIAN STINNER

ABSTRACT. Convergence of solutions to a partially diffusive chemotaxis system with indirect signal production and phenotype switching is shown in a two-dimensional setting when the switching rate increases to infinity, thereby providing a rigorous justification of formal computations performed in the literature. The expected limit system being the classical parabolic-parabolic Keller-Segel system, the obtained convergence is restricted to a finite time interval for general initial conditions but valid for arbitrary bounded time intervals when the mass of the initial condition is appropriately small. Furthermore, if the solution to the limit system blows up in finite time, then neither of the two phenotypes in the partially diffusive system can be uniformly bounded with respect to the  $L_2$ -norm beyond that time.

## 1. INTRODUCTION

The singular limit  $\gamma \rightarrow \infty$  of a chemotaxis model with indirect signal production and phenotype switching

$$\begin{aligned} \partial_t u_\gamma &= \operatorname{div}(D_u \nabla u_\gamma - u_\gamma \nabla w_\gamma) + \gamma(\theta v_\gamma - u_\gamma) & \text{in } (0, \infty) \times \Omega, \\ \partial_t v_\gamma &= D_v \Delta v_\gamma + \gamma(u_\gamma - \theta v_\gamma) & \text{in } (0, \infty) \times \Omega, \\ \partial_t w_\gamma &= D \Delta w_\gamma - \alpha w_\gamma + v_\gamma & \text{in } (0, \infty) \times \Omega, \end{aligned} \tag{1.1}$$

supplemented with no-flux boundary conditions and initial conditions  $(u_\gamma, v_\gamma, w_\gamma)(0) = (u_0, v_0, w_0)$ , is studied in [19] in a bounded domain  $\Omega \subset \mathbb{R}^n$  with  $n \leq 3$  and positive constants  $D_u$ ,  $D_v$ ,  $D$ ,  $\alpha$ , and  $\theta$ , which were all set to 1 in [19]. The above model describes the dynamics of a population of cells divided into two phenotypes with respective densities  $u_\gamma$  and  $v_\gamma$  and a chemoattractant with concentration  $w_\gamma$ . The model is designed to account for phenotype-dependent responses of cells to the presence of a chemoattractant [16]. While both phenotypes are diffusing with possibly different diffusion rates  $D_u$  and  $D_v$ , only the motion of the phenotype with density  $u_\gamma$  features a chemotactic bias, the chemoattractant being only secreted by the other phenotype. For more details concerning the modeling we refer to [16, 19] and the references therein.

---

*Date:* March 19, 2024.

*2010 Mathematics Subject Classification.* 35B25, 35K57, 35M33, 35Q92.

*Key words and phrases.* chemotaxis system, species with two subpopulations, fast reaction limit, partially diffusive system.

As pointed out in [19, Section 2], letting  $\gamma \rightarrow \infty$  formally, one finds  $u_\gamma \sim \theta v_\gamma$ , so that, introducing the total density  $n_\gamma := u_\gamma + v_\gamma$ , we expect that  $u_\gamma \sim \theta n_\gamma / (1 + \theta)$  and  $v_\gamma \sim n_\gamma / (1 + \theta)$ . Since  $n_\gamma$  solves

$$\partial_t n_\gamma = \operatorname{div}(D_u \nabla u_\gamma - u_\gamma \nabla w_\gamma) + D_v \Delta v_\gamma \quad \text{in } (0, \infty) \times \Omega,$$

cluster points of  $(n_\gamma, w_\gamma)_\gamma$  as  $\gamma \rightarrow \infty$  (if any) are bound to solve the following parabolic-parabolic Keller-Segel system

$$\begin{aligned} \partial_t n &= \frac{D_u \theta + D_v}{1 + \theta} \Delta n - \frac{\theta}{1 + \theta} \operatorname{div}(n \nabla w) && \text{in } (0, \infty) \times \Omega, \\ \partial_t w &= D \Delta w - \alpha w + \frac{n}{1 + \theta} && \text{in } (0, \infty) \times \Omega, \end{aligned}$$

supplemented with no-flux boundary conditions and initial conditions  $(n, w)(0) = (n_0, w_0)$  with  $n_0 = u_0 + v_0$ . A challenging issue in establishing the connection between these two systems is that, while the former is globally well-posed in space dimension 2 and 3 [7, 19], finite time blowup may occur for the latter [9, 22].

As the diffusion rates of the two phenotypes may be of different orders, we focus here on the limit case where the phenotype which produces the chemoattractant is not diffusing and static ( $D_v = 0$ ) and set  $D_u = 1$  for simplicity. For the corresponding system, global well-posedness and the existence of a critical mass in a bounded domain of  $\mathbb{R}^2$  are established in [14]. It is a particular case of the model from [20] for foraging ants and also is a variant of chemotaxis models for building termites nests, mountain pine beetles, and cancer cells. For more details we refer to the introduction of [14] and the references therein.

We therefore assume that  $\Omega \subset \mathbb{R}^2$  is a bounded domain with smooth boundary and investigate the singular limit  $\gamma \rightarrow \infty$  of the system

$$\partial_t u_\gamma = \operatorname{div}(\nabla u_\gamma - u_\gamma \nabla w_\gamma) + \gamma(\theta v_\gamma - u_\gamma) \quad \text{in } (0, \infty) \times \Omega, \quad (1.2a)$$

$$\partial_t v_\gamma = \gamma(u_\gamma - \theta v_\gamma) \quad \text{in } (0, \infty) \times \Omega, \quad (1.2b)$$

$$\partial_t w_\gamma = D \Delta w_\gamma - \alpha w_\gamma + v_\gamma \quad \text{in } (0, \infty) \times \Omega, \quad (1.2c)$$

supplemented with no-flux boundary conditions

$$(\nabla u_\gamma - u_\gamma \nabla w_\gamma) \cdot \mathbf{n} = \nabla w_\gamma \cdot \mathbf{n} = 0 \quad \text{on } (0, \infty) \times \partial\Omega \quad (1.2d)$$

and initial conditions

$$(u_\gamma, v_\gamma, w_\gamma)(0) = (u_0, v_0, w_0) \quad \text{in } \Omega, \quad (1.3)$$

where the parameters  $D$ ,  $\alpha$ , and  $\theta$  are positive.

According to the above formal calculation, we expect that in the limit  $\gamma \rightarrow \infty$  solutions to (1.2)–(1.3) converge to that of

$$\partial_t n = \frac{\theta}{1 + \theta} \operatorname{div}(\nabla n - n \nabla w) \quad \text{in } (0, \infty) \times \Omega, \quad (1.4a)$$

$$\partial_t w = D \Delta w - \alpha w + \frac{n}{1 + \theta} \quad \text{in } (0, \infty) \times \Omega, \quad (1.4b)$$

supplemented with no-flux boundary conditions

$$(\nabla n - n\nabla w) \cdot \mathbf{n} = \nabla w \cdot \mathbf{n} = 0 \quad \text{on } (0, \infty) \times \partial\Omega, \quad (1.4c)$$

and initial conditions

$$(n, w)(0) = (n_0, w_0) \quad \text{in } \Omega, \quad (1.5)$$

with  $n_0 = u_0 + v_0$ , and the purpose of this work is to provide a proof of this fact. Such a result supplements nicely the analysis performed in [19, Lemmas 4.9 and 4.10] for the fully diffusive system (1.1), where only a conditional convergence result is provided. As already mentioned, the first difficulty to be faced is that, while solutions to (1.2)–(1.3) are always global in time, see [14, Theorem 1.1] and Proposition 2.1 below, solutions to (1.4)–(1.5) may blow up in finite time when  $\|n_0\|_1 = \|u_0 + v_0\|_1$  exceeds the threshold value  $4\pi(1 + \theta)D$ , see [10–12] and Appendix B below. This feature excludes convergence on arbitrary time intervals and we shall thus prove only convergence on a small time interval for arbitrary initial data. It is actually proved in [19, Theorem 1.4] that there are initial conditions such that, for the fully diffusive system (1.1), the family  $(u_\gamma, v_\gamma)_{\gamma>0}$  is not bounded in  $L_\infty((0, T) \times \Omega)$  for some  $T > 0$  sufficiently large and we provide a corresponding result for (1.2)–(1.3) in Theorem 1.2 below. When  $\|n_0\|_1 < 4\pi(1 + \theta)D$ , convergence on any finite time interval is established. The second difficulty is of a more technical nature and is due to the assumption that one of the species does not diffuse in space [20]. As a consequence, no compactness with respect to the space variable is available for  $(v_\gamma)_{\gamma>0}$ .

**Theorem 1.1.** *Let  $M > 0$  and consider*

$$(u_0, v_0, w_0) \in W_{3,+}^1(\Omega) \times W_{3,+}^1(\Omega) \times W_{2,+}^2(\Omega) \quad (1.6)$$

*satisfying*

$$\|u_0 + v_0\|_1 = M. \quad (1.7)$$

*For  $\gamma > 0$ , we denote the solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$  and set  $n_\gamma := u_\gamma + v_\gamma$ . Then there is  $T_\infty \in (0, \infty]$  such that, for all  $T \in (0, T_\infty)$ ,*

$$\lim_{\gamma \rightarrow \infty} \int_0^T \|n_\gamma(t) - n(t)\|_2^2 dt = 0, \quad (1.8a)$$

$$\lim_{\gamma \rightarrow \infty} \sup_{t \in [0, T]} \|w_\gamma(t) - w(t)\|_{W_2^1} = 0, \quad (1.8b)$$

*where  $(n, w)$  is the unique weak-strong solution to (1.4)–(1.5) with  $n_0 := u_0 + v_0$  in the sense of Definition C.1.*

*Moreover,  $T_\infty = \infty$  if  $M = \|u_0 + v_0\|_1 \in (0, 4\pi(1 + \theta)D)$ .*

The final statement of Theorem 1.1 is valid for all masses below the critical mass and this property can be established thanks to the construction of a Liapunov functional [14]. The latter does not seem to be available for the fully diffusive system (1.1) and it is yet unclear whether Theorem 1.1 extends to that case.

We next establish the analogue of [19, Theorem 1.4] for the partially diffusive system (1.2)–(1.3).

**Theorem 1.2.** Consider  $(u_0, v_0, w_0)$  satisfying (1.6) and set  $n_0 := u_0 + w_0$ . Assume that the corresponding classical solution  $(n, w)$  to (1.4)–(1.5) blows up at some finite time  $T_{bu} \in (0, \infty)$ ; that is,

$$\lim_{t \rightarrow T_{bu}} \|n(t)\|_\infty = \infty. \quad (1.9)$$

Then, for all  $T > T_{bu}$ ,

$$\sup_{\gamma \geq 1} \{ \|u_\gamma\|_{L_\infty((0,T),L_2(\Omega))} \} = \infty \quad \text{and} \quad \sup_{\gamma \geq 1} \{ \|v_\gamma\|_{L_\infty((0,T),L_2(\Omega))} \} = \infty, \quad (1.10)$$

where  $(u_\gamma, v_\gamma, w_\gamma)$  denotes the solution to (1.2)–(1.3) with initial condition  $(u_0, v_0, w_0)$  given by Proposition 2.1.

Before we start to prove the above results, we recall in Section 2 the global well-posedness for (1.2)–(1.3) which is established in [14]. The proof of Theorem 1.1 involves two steps: the derivation of estimates on  $(u_\gamma, v_\gamma, w_\gamma)_\gamma$  which do not depend on  $\gamma$ , eventually leading to compactness, and afterwards the convergence of  $(u_\gamma, v_\gamma, w_\gamma)_\gamma$ . The proofs of the former in Section 4 bear some similarities with computations performed to establish the local or global existence of solutions to the Keller-Segel system (1.4)–(1.5) and are divided into small time estimates for arbitrary masses  $M > 0$  and global estimates for  $M \in (0, 4\pi(1 + \theta)D)$ . The convergence proof is more involved and we shall deal with it in Section 3. One important step in the proof is the strong convergence of  $(v_\gamma)_\gamma$  which does not follow from classical compact embeddings due to the non-diffusive character of (1.2b). The proof of Theorem 1.1 is completed in Section 5. The final Section 6 is devoted to the proof of Theorem 1.2, where we show that uniform bounds on the  $L_2$ -norm of either  $v_\gamma$  or  $u_\gamma$  imply the validity of the estimates which are required for the convergence proof of Theorem 1.1 and contradict the assumed blowup in view of the uniqueness of the weak-strong solution to (1.4)–(1.5). We collect in three appendices some results used in the proofs. In Appendix A we recall the well-known Moser-Trudinger inequality, while the rescaling performed in Appendix B shows the optimality of the mass constraint from Theorem 1.1. In Appendix C we give the definition of the weak-strong solution to (1.4)–(1.5) and prove its uniqueness, adapting an argument from [13].

## 2. WELL-POSEDNESS

In this section we recall the global well-posedness for (1.2)–(1.3) established in [14, Theorem 1.1] in an appropriate functional setting and give the notation of the involved function spaces. More precisely, for  $r \in (1, \infty)$ , we set

$$\begin{aligned} W_{r,\mathcal{B}}^m(\Omega) &:= \{z \in W_r^m(\Omega) : \nabla z \cdot \mathbf{n} = 0 \text{ on } \partial\Omega\} \quad \text{if } 1 + \frac{1}{r} < m \leq 2, \\ W_{r,\mathcal{B}}^m(\Omega) &:= W_r^m(\Omega) \quad \text{if } -1 + \frac{1}{r} < m < 1 + \frac{1}{r}, \\ W_{r,\mathcal{B}}^m(\Omega) &:= W_{r/(r-1)}^{-m}(\Omega)' \quad \text{if } -2 + \frac{1}{r} < m \leq -1 + \frac{1}{r}, \end{aligned} \quad (2.1)$$

and

$$W_{r,\mathcal{B},+}^m(\Omega) := \{z \in W_{r,\mathcal{B}}^m(\Omega) : z \geq 0 \text{ in } \Omega\}, \quad (2.2)$$

where  $W_r^m(\Omega)$ ,  $m \in [0, \infty)$ ,  $r \in [1, \infty)$ , denote the usual Sobolev spaces, see [3, Section 5].

**Proposition 2.1.** *Let  $M > 0$ ,  $\gamma > 0$ , and  $(u_0, v_0, w_0) \in W_{3,+}^1(\Omega; \mathbb{R}^3)$  satisfying (1.7). Then the system (1.2)–(1.3) has a unique non-negative weak solution  $(u_\gamma, v_\gamma, w_\gamma)$  in  $W_3^1(\Omega)$  defined on  $[0, \infty)$  satisfying*

$$\begin{aligned} u_\gamma &\in C([0, \infty), W_{3,+}^1(\Omega)) \cap C^1([0, \infty), W_{3/2}^1(\Omega, \mathbb{R}^2)'), \\ v_\gamma &\in C^1([0, \infty), W_{3,+}^1(\Omega)), \\ w_\gamma &\in C([0, \infty), W_{3,+}^1(\Omega)) \cap C^1([0, \infty), W_{3/2}^1(\Omega, \mathbb{R}^2)'), \end{aligned}$$

and

$$\|(u_\gamma + v_\gamma)(t)\|_1 = M, \quad t \geq 0. \quad (2.3)$$

Moreover,

$$\begin{aligned} u_\gamma &\in C((0, \infty), W_{3,\mathcal{B}}^2(\Omega)) \cap C^1((0, \infty), L_3(\Omega)), \\ w_\gamma &\in C((0, \infty), W_{3,\mathcal{B}}^2(\Omega)) \cap C^1((0, \infty), L_3(\Omega)). \end{aligned}$$

The local well-posedness of (1.2)–(1.3) relies on the abstract theory developed in [1, 2] for partially diffusive parabolic systems. It is however likely that a larger set of initial data may be handled as well, in light of the recent result in [17, Section 5.2]. Global existence is a consequence from several estimates and makes in particular use of the following Gagliardo-Nirenberg inequality

$$\|z\|_4 \leq c_0 \|z\|_{W_2^1}^{1/2} \|z\|_2^{1/2}, \quad z \in W_2^1(\Omega), \quad (2.4)$$

which is also an important tool in the forthcoming analysis.

### 3. CONVERGENCE

We begin with the convergence issue and assume that we have already obtained several bounds on the family  $(u_\gamma, v_\gamma, w_\gamma)_{\gamma \geq 1}$ , which do not depend on  $\gamma \geq 1$  and are valid on a time interval  $[0, T]$ . We shall devote the next sections to the derivation of these bounds.

**Proposition 3.1.** *Let  $M > 0$  and consider  $(u_0, v_0, w_0) \in W_{3,+}^1(\Omega; \mathbb{R}^3)$  satisfying (1.7). For  $\gamma > 0$ , we denote the solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$  and set*

$$n_\gamma := u_\gamma + v_\gamma. \quad (3.1)$$

Assume that there are  $T \in (0, \infty)$  and  $c_1 > 0$  such that, for any  $\gamma \geq 1$ ,

$$\sup_{t \in [0, T]} \{ \|u_\gamma(t)\|_2 + \|v_\gamma(t)\|_2 + \|w_\gamma(t)\|_{W_2^1} \} \leq c_1, \quad (3.2a)$$

$$\int_0^T [\|\nabla u_\gamma(t)\|_2^2 + \|\partial_t w_\gamma(t)\|_2^2 + \gamma \|e_\gamma(t)\|_2^2] dt \leq c_1^2, \quad (3.2b)$$

with

$$e_\gamma := \theta v_\gamma - u_\gamma. \quad (3.3)$$

Then there are a sequence  $(\gamma_j)_{j \geq 1}$ ,  $\gamma_j \rightarrow \infty$ , and non-negative functions

$$\begin{aligned} n &\in W_2^1((0, T), W_2^1(\Omega)') \cap L_\infty((0, T), L_2(\Omega)) \cap L_2((0, T), W_2^1(\Omega)), \\ w &\in W_2^1((0, T), L_2(\Omega)) \cap L_\infty((0, T), W_2^1(\Omega)) \cap L_2((0, T), W_2^2(\Omega)), \end{aligned}$$

such that

$$\lim_{j \rightarrow \infty} \sup_{t \in [0, T]} \{ \|(n_{\gamma_j} - n)(t)\|_{(W_2^1)'} + \|(w_{\gamma_j} - w)(t)\|_2 \} = 0, \quad (3.4)$$

$$\lim_{j \rightarrow \infty} \int_0^T [ \|(w_{\gamma_j} - w)(t)\|_{W_2^1}^2 + \|(n_{\gamma_j} - n)(t)\|_2^2 + \|e_{\gamma_j}(t)\|_2^2 ] dt = 0. \quad (3.5)$$

In addition,

$$\lim_{j \rightarrow \infty} \int_0^T \left[ \left\| \left( u_{\gamma_j} - \frac{\theta n}{1 + \theta} \right) (t) \right\|_2^2 + \left\| \left( v_{\gamma_j} - \frac{n}{1 + \theta} \right) (t) \right\|_2^2 \right] dt = 0 \quad (3.6)$$

and

$$\lim_{j \rightarrow \infty} \int_0^T \left\| \left( u_{\gamma_j} \nabla w_{\gamma_j} - \frac{\theta}{1 + \theta} n \nabla w \right) (t) \right\|_2^{4/3} dt = 0. \quad (3.7)$$

In the following,  $c$  and  $(c_i)_{i \geq 2}$  denote positive constants depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $(u_0, v_0, w_0)$ , and  $c_1$ . The dependence upon additional parameters will be indicated explicitly.

Several additional estimates can be derived from (3.2).

**Lemma 3.2.** *Assume (3.2). There is  $c_2(T) > 0$  such that, for all  $\gamma \geq 1$ ,*

$$\int_0^T [ \|w_\gamma(t)\|_{W_2^2}^2 + \|u_\gamma(t)\|_4^4 + \|\nabla w_\gamma(t)\|_4^4 + \|(u_\gamma \nabla w_\gamma)(t)\|_2^2 ] dt \leq c_2^2(T), \quad (3.8)$$

$$\int_0^T \|\partial_t n_\gamma(t)\|_{(W_2^1)'}^2 dt \leq c_2^2(T), \quad (3.9)$$

recalling that  $n_\gamma = u_\gamma + v_\gamma$ , see (3.1).

*Proof.* It first readily follows from (1.2c), (3.2), and parabolic regularity that

$$\int_0^T \|w_\gamma(t)\|_{W_2^2}^2 dt \leq c(T). \quad (3.10)$$

We next combine (2.4) with (3.2a) and (3.10) to find

$$\begin{aligned} \int_0^T \|\nabla w_\gamma(t)\|_4^4 dt &\leq c_0^4 \int_0^T \|\nabla w_\gamma(t)\|_{W_2^1}^2 \|\nabla w_\gamma(t)\|_2^2 dt \\ &\leq c_0^4 c_1^2 \int_0^T \|w_\gamma(t)\|_{W_2^2}^2 dt \leq c(T). \end{aligned} \quad (3.11)$$

Similarly, we infer from (2.4) and (3.2) that

$$\begin{aligned} \int_0^T \|u_\gamma(t)\|_4^4 dt &\leq c_0^4 \int_0^T \|u_\gamma(t)\|_{W_2^1}^2 \|u_\gamma(t)\|_2^2 dt \\ &\leq c_0^4 c_1^2 \left( c_1^2 T + \int_0^T \|\nabla u_\gamma(t)\|_2^2 dt \right) \leq c(T). \end{aligned} \quad (3.12)$$

Gathering (3.10), (3.11), and (3.12) and using Hölder's inequality lead us to (3.8).

We next consider  $\varphi \in W_2^1(\Omega)$  and deduce from (1.2a), (1.2b), (1.2d), and Hölder's inequality that

$$|\langle \partial_t n_\gamma, \varphi \rangle_{(W_2^1)', W_2^1}| = \left| \int_\Omega \nabla \varphi \cdot (\nabla u_\gamma - u_\gamma \nabla w_\gamma) dx \right| \leq \|\nabla \varphi\|_2 (\|\nabla u_\gamma\|_2 + \|u_\gamma \nabla w_\gamma\|_2),$$

so that

$$\|\partial_t n_\gamma\|_{(W_2^1)'} \leq \|\nabla u_\gamma\|_2 + \|u_\gamma \nabla w_\gamma\|_2$$

by a duality argument. Combining the above inequality with (3.2b) and (3.8) readily gives (3.9) and completes the proof.  $\square$

We are now ready to investigate the compactness properties of the family  $(u_\gamma, v_\gamma, w_\gamma)_\gamma$ . As a preliminary step, let us recall the following lemma [15, Chapitre I, Lemme 5.2], see also [21, Lemma 8], which is a consequence of the compactness of the embedding of  $W_2^1(\Omega)$  in  $L_2(\Omega)$  and the continuity of that of  $L_2(\Omega)$  in  $W_2^1(\Omega)'$ .

**Lemma 3.3.** *Given  $\eta > 0$ , there is  $c_3(\eta) > 0$  such that*

$$\|z\|_2 \leq \eta \|z\|_{W_2^1} + c_3(\eta) \|z\|_{(W_2^1)'}, \quad z \in W_2^1(\Omega).$$

**Proposition 3.4.** *Assume (3.2). There are a sequence  $(\gamma_j)_{j \geq 1}$ ,  $\gamma_j \rightarrow \infty$ , and non-negative functions*

$$n \in W_2^1((0, T), W_2^1(\Omega)') \cap L_\infty((0, T), L_2(\Omega)) \cap L_2((0, T), W_2^1(\Omega)),$$

$$w \in W_2^1((0, T), L_2(\Omega)) \cap L_\infty((0, T), W_2^1(\Omega)) \cap L_2((0, T), W_2^2(\Omega)),$$

such that the convergences (3.4), (3.5), and (3.6) hold true; that is,

$$\lim_{j \rightarrow \infty} \sup_{t \in [0, T]} \{ \|n_{\gamma_j}(t) - n(t)\|_{(W_2^1)'} + \|w_{\gamma_j}(t) - w(t)\|_2 \} = 0,$$

$$\lim_{j \rightarrow \infty} \int_0^T [\|w_{\gamma_j}(t) - w(t)\|_{W_2^1}^2 + \|n_{\gamma_j}(t) - n(t)\|_2^2 + \|e_{\gamma_j}(t)\|_2^2] dt = 0,$$

and

$$\lim_{j \rightarrow \infty} \int_0^T \left[ \left\| \left( u_{\gamma_j} - \frac{\theta n}{1 + \theta} \right) (t) \right\|_2^2 + \left\| \left( v_{\gamma_j} - \frac{n}{1 + \theta} \right) (t) \right\|_2^2 \right] dt = 0.$$

*Proof.* We first observe that a straightforward consequence of (3.2b) is that

$$\lim_{\gamma \rightarrow \infty} \int_0^T \|e_\gamma(t)\|_2^2 dt = 0. \quad (3.13)$$



Next, since  $(n_\gamma)_\gamma$  is bounded in  $L_\infty((0, T), L_2(\Omega))$  by (3.1) and (3.2a) and  $L_2(\Omega)$  is compactly embedded in  $W_2^1(\Omega)'$ , we deduce from (3.9) and [21, Corollary 4] that

$$(n_\gamma)_\gamma \text{ is relatively compact in } C([0, T], W_2^1(\Omega)').$$

A similar argument based on (3.2), (3.8), and the compactness of the embeddings of  $W_2^1(\Omega)$  in  $L_2(\Omega)$  and of  $W_2^2(\Omega)$  in  $W_2^1(\Omega)$  entails that

$$(w_\gamma)_\gamma \text{ is relatively compact in } C([0, T], L_2(\Omega)) \text{ and in } L_2((0, T), W_2^1(\Omega)).$$

Consequently, there are a sequence  $(\gamma_j)_{j \geq 1}$ ,  $\gamma_j \rightarrow \infty$ , and non-negative functions

$$\begin{aligned} n &\in W_2^1((0, T), W_2^1(\Omega)') \cap L_\infty((0, T), L_2(\Omega)), \\ w &\in W_2^1((0, T), L_2(\Omega)) \cap L_\infty((0, T), W_2^1(\Omega)) \cap L_2((0, T), W_2^2(\Omega)) \end{aligned}$$

such that the convergences (3.4) hold true, along with

$$\lim_{j \rightarrow \infty} \int_0^T \|w_{\gamma_j}(t) - w(t)\|_{W_2^1}^2 dt = 0. \quad (3.14)$$

Moreover, it readily follows from (3.2) and (3.8) that there are non-negative functions

$$u \in L_\infty((0, T), L_2(\Omega)) \cap L_2((0, T), W_2^1(\Omega)) \text{ and } v \in L_\infty((0, T), L_2(\Omega)) \quad (3.15)$$

such that, up to the extraction of a further subsequence,

$$(u_{\gamma_j}, v_{\gamma_j})_j \xrightarrow{*} (u, v) \text{ in } L_\infty((0, T), L_2(\Omega; \mathbb{R}^2)), \quad (3.16a)$$

$$(u_{\gamma_j})_j \rightharpoonup u \text{ in } L_2((0, T), W_2^1(\Omega)) \cap L^4((0, T) \times \Omega), \quad (3.16b)$$

$$(w_{\gamma_j})_j \rightharpoonup w \text{ in } L_2((0, T), W_2^2(\Omega)). \quad (3.16c)$$

Since

$$u_{\gamma_j} = \frac{\theta n_{\gamma_j} - e_{\gamma_j}}{1 + \theta} \text{ and } v_{\gamma_j} = \frac{n_{\gamma_j} + e_{\gamma_j}}{1 + \theta}$$

by (3.1) and (3.3), a first consequence of (3.4) and (3.13) is that

$$u = \frac{\theta n}{1 + \theta} \text{ and } v = \frac{n}{1 + \theta} \text{ a.e. in } (0, T) \times \Omega. \quad (3.17)$$

In particular, it follows from (3.15), (3.17), and the positivity of  $\theta$  that  $n \in L_2((0, T), W_2^1(\Omega))$  and we have thus established the regularity properties of  $(n, w)$  listed in Proposition 3.4.

We next turn to the proof of (3.5) and first recall that the convergences of  $(w_{\gamma_j})_{j \geq 1}$  and  $(e_{\gamma_j})_{j \geq 1}$  stated therein are already shown in (3.13) and (3.14). We are thus left with the strong convergence of  $(n_{\gamma_j})_{j \geq 1}$  which will follow from Lemma 3.3. Indeed, for  $\eta > 0$ , we infer from (3.1), (3.3), and Lemma 3.3 that

$$\begin{aligned} \|n_{\gamma_j} - n\|_2 &= \|u_{\gamma_j} + v_{\gamma_j} - n\|_2 = \left\| \frac{1 + \theta}{\theta} u_{\gamma_j} - n + \frac{e_{\gamma_j}}{\theta} \right\|_2 \\ &\leq \frac{1 + \theta}{\theta} \left\| u_{\gamma_j} - \frac{\theta n}{1 + \theta} \right\|_2 + \frac{\|e_{\gamma_j}\|_2}{\theta} \end{aligned}$$

$$\leq \frac{(1+\theta)\eta}{\theta} \left\| u_{\gamma_j} - \frac{\theta n}{1+\theta} \right\|_{W_2^1} + \frac{(1+\theta)c_3(\eta)}{\theta} \left\| u_{\gamma_j} - \frac{\theta n}{1+\theta} \right\|_{(W_2^1)'} + \frac{\|e_{\gamma_j}\|_2}{\theta}.$$

After integration with respect to time, we deduce from (3.1), (3.2), and (3.3) that

$$\begin{aligned} \|n_{\gamma_j} - n\|_{L_2((0,T)\times\Omega)} &\leq \frac{(1+\theta)\eta}{\theta} \left( \|u_{\gamma_j}\|_{L_2((0,T),W_2^1(\Omega))} + \frac{\theta}{1+\theta} \|n\|_{L_2((0,T),W_2^1(\Omega))} \right) \\ &\quad + c_3(\eta) \|n_{\gamma_j} - n\|_{L_2((0,T),W_2^1(\Omega)')} + \frac{c_3(\eta)}{\theta} \|e_{\gamma_j}\|_{L_2((0,T),W_2^1(\Omega)')} \\ &\quad + \frac{\|e_{\gamma_j}\|_{L_2((0,T)\times\Omega)}}{\theta} \\ &\leq \frac{(1+\theta)\eta}{\theta} \left( c_1\sqrt{T} + c_1 + \frac{\theta}{1+\theta} \|n\|_{L_2((0,T),W_2^1(\Omega))} \right) \\ &\quad + c_3(\eta) \|n_{\gamma_j} - n\|_{L_2((0,T),W_2^1(\Omega)')} + \frac{c_3(\eta)}{\theta} \|e_{\gamma_j}\|_{L_2((0,T),W_2^1(\Omega)')} \\ &\quad + \frac{\|e_{\gamma_j}\|_{L_2((0,T)\times\Omega)}}{\theta}. \end{aligned}$$

Owing to (3.4), (3.13), and the continuous embedding of  $L_2(\Omega)$  in  $W_2^1(\Omega)'$ , we may take the limit  $j \rightarrow \infty$  in the above inequality to obtain

$$\limsup_{j \rightarrow \infty} \|n_{\gamma_j} - n\|_{L_2((0,T)\times\Omega)} \leq \frac{(1+\theta)\eta}{\theta} \left( c_1\sqrt{T} + c_1 + \frac{\theta}{1+\theta} \|n\|_{L_2((0,T),W_2^1(\Omega))} \right).$$

The above inequality being valid for any  $\eta > 0$ , we conclude that

$$\lim_{j \rightarrow \infty} \|n_{\gamma_j} - n\|_{L_2((0,T)\times\Omega)} = 0,$$

which completes the proof of (3.5). Finally, the convergences (3.6) are immediate consequences of (3.1), (3.3), (3.5), and (3.13).  $\square$

**Corollary 3.5.** *Assume (3.2). There holds*

$$\lim_{j \rightarrow \infty} \int_0^T \left\| \left( u_{\gamma_j} \nabla w_{\gamma_j} - \frac{\theta}{1+\theta} n \nabla w \right) (t) \right\|_2^{4/3} dt = 0.$$

*Proof.* We first recall that  $\theta n/(1+\theta) = u$  by (3.17).

On the one hand, thanks to (2.4), (3.2), the regularity of  $u$ , and Hölder's inequality, we obtain

$$\begin{aligned} \int_0^T \|(u_{\gamma_j} - u)(t)\|_4^2 dt &\leq c_0^2 \int_0^T \|(u_{\gamma_j} - u)(t)\|_{W_2^1} \|(u_{\gamma_j} - u)(t)\|_2 dt \\ &\leq c_0^2 (\|u_{\gamma_j}\|_{L_2((0,T),W_2^1(\Omega))} + \|u\|_{L_2((0,T),W_2^1(\Omega))}) \|u_{\gamma_j} - u\|_{L_2((0,T)\times\Omega)} \\ &\leq c_0^2 \left( c_1 + c_1\sqrt{T} + \|u\|_{L_2((0,T),W_2^1(\Omega))} \right) \|u_{\gamma_j} - u\|_{L_2((0,T)\times\Omega)}. \end{aligned}$$

Hence, by (3.6),

$$\lim_{j \rightarrow \infty} \int_0^T \|(u_{\gamma_j} - u)(t)\|_4^2 dt = 0. \quad (3.18)$$

On the other hand, we deduce from (2.4), (3.10), and Hölder's inequality that

$$\begin{aligned} & \int_0^T \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_4^2 dt \\ & \leq c_0^2 \int_0^T \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_{W_2^1} \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_2 dt \\ & \leq c_0^2 \left( c(T) + \|w\|_{L_2((0,T),W_2^2(\Omega))} \right) \|w_{\gamma_j} - w\|_{L_2((0,T),W_2^1(\Omega))}, \end{aligned}$$

and we use (3.5) to conclude that

$$\lim_{j \rightarrow \infty} \int_0^T \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_4^2 dt = 0. \quad (3.19)$$

Now, using Hölder's inequality, along with (3.8) and (3.16b), give

$$\begin{aligned} & \int_0^T \|(u_{\gamma_j} \nabla w_{\gamma_j} - u \nabla w)(t)\|_2^{4/3} dt \\ & \leq c \int_0^T \|[(u_{\gamma_j} - u) \nabla w_{\gamma_j}](t)\|_2^{4/3} dt + c \int_0^T \| [u(\nabla w_{\gamma_j} - \nabla w)](t) \|_2^{4/3} dt \\ & \leq c \int_0^T \|(u_{\gamma_j} - u)(t)\|_4^{4/3} \|\nabla w_{\gamma_j}(t)\|_4^{4/3} dt + c \int_0^T \|u(t)\|_4^{4/3} \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_4^{4/3} dt \\ & \leq c \left( \int_0^T \|(u_{\gamma_j} - u)(t)\|_4^2 dt \right)^{2/3} \left( \int_0^T \|\nabla w_{\gamma_j}(t)\|_4^4 dt \right)^{1/3} \\ & \quad + c \left( \int_0^T \|u(t)\|_4^4 dt \right)^{1/3} \left( \int_0^T \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_4^2 dt \right)^{2/3} \\ & \leq c(T) \left[ \left( \int_0^T \|(u_{\gamma_j} - u)(t)\|_4^2 dt \right)^{2/3} + \left( \int_0^T \|(\nabla w_{\gamma_j} - \nabla w)(t)\|_4^2 dt \right)^{2/3} \right] \end{aligned}$$

and we may pass to the limit  $j \rightarrow \infty$  in the above inequality with the help of (3.18) and (3.19) to obtain Corollary 3.5.  $\square$

*Proof of Proposition 3.1.* Gathering the outcome of Proposition 3.4 and Corollary 3.5 gives Proposition 3.1.  $\square$

## 4. UNIFORM ESTIMATES

This section is devoted to the derivation of estimates on solutions to (1.2)–(1.3) which do not depend on  $\gamma$  and allow us to apply Proposition 3.1 to prove Theorem 1.1 in Section 5. In the following,  $C$  and  $(C_i)_{i \geq 0}$  denote positive constants depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ , and  $(u_0, v_0, w_0)$ . The dependence upon additional parameters will be indicated explicitly.

Two different cases are handled: first, no constraint is placed on  $M = \|u_0 + v_0\|_1$  and the estimates we derive are only valid on a finite time interval, a feature which complies with the possible finite time blowup of solutions to (1.4)–(1.5). We next assume that  $M = \|u_0 + v_0\|_1 < 4\pi D(1 + \theta)$  and show that estimates are available on any finite time interval, which is again consistent with the properties of (1.4)–(1.5), as global solutions to (1.4)–(1.5) exist in that case. Specifically, we prove the following result.

**Proposition 4.1.** *Let  $M > 0$  and consider  $(u_0, v_0, w_0)$  satisfying (1.6) and (1.7). For  $\gamma \geq 1$ , we denote the corresponding solution to (1.2)–(1.3) provided by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$ .*

- (a) *There is  $T_\infty \in (0, \infty)$  depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ , and  $(u_0, v_0, w_0)$  with the following property: for any  $T \in (0, T_\infty)$ , there is  $C(T) > 0$  depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $(u_0, v_0, w_0)$ , and  $T$  such that, for any  $\gamma \geq 1$ ,*

$$\|u_\gamma(t)\|_2 + \|v_\gamma(t)\|_2 + \|w_\gamma(t)\|_{W_2^1} \leq C(T), \quad t \in [0, T], \quad (4.1a)$$

$$\int_0^T [\|\nabla u_\gamma(t)\|_2^2 + \|\partial_t w_\gamma(t)\|_2^2 + \gamma \|e_\gamma(t)\|_2^2] dt \leq C(T), \quad (4.1b)$$

recalling that  $e_\gamma$  is defined in (3.3).

- (b) *Assume further that  $M \in (0, 4\pi D(1 + \theta))$ . Then, for all  $T > 0$ , there is  $C(T) > 0$  depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $(u_0, v_0, w_0)$ , and  $T$  such that the estimates (4.1) are valid for all  $\gamma \geq 1$ .*

**4.1. Small time estimates.** We begin with the following estimate which is valid whatever the value of  $M$  but only on a finite time interval.

**Lemma 4.2.** *There is  $T_\infty \in (0, \infty)$  (defined in (4.9) below) depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ , and  $(u_0, v_0, w_0)$  with the following property: for any  $T \in (0, T_\infty)$ , there is  $C_0(T) > 0$  depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $(u_0, v_0, w_0)$ , and  $T$  such that, for any  $\gamma \geq 1$ ,*

$$\|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1 + \|w_\gamma(t)\|_{W_2^1} \leq C_0(T), \quad t \in (0, T), \quad (4.2)$$

$$\int_0^T \|\partial_t w_\gamma(t)\|_2^2 dt \leq C_0(T), \quad (4.3)$$

where

$$L(r) := r \ln r - r + 1 \geq 0, \quad L_\theta(r) := \frac{L(\theta r)}{\theta} = r \ln(\theta r) - r + \frac{1}{\theta} \geq 0, \quad r \geq 0. \quad (4.4)$$

*Proof.* On the one hand, it follows from (1.2a), (1.2c), and (1.2d) that

$$\frac{d}{dt} \int_\Omega L(u_\gamma) dx = - \int_\Omega \nabla \ln u_\gamma \cdot (\nabla u_\gamma - u_\gamma \nabla w_\gamma) dx + \gamma \int_\Omega \ln u_\gamma (\theta v_\gamma - u_\gamma) dx$$

$$\begin{aligned}
&= -4\|\nabla\sqrt{u_\gamma}\|_2^2 - \int_\Omega u_\gamma\Delta w_\gamma \, dx + \gamma \int_\Omega \ln u_\gamma(\theta v_\gamma - u_\gamma) \, dx \\
&= -4\|\nabla\sqrt{u_\gamma}\|_2^2 + \frac{1}{D} \int_\Omega u_\gamma(v_\gamma - \alpha w_\gamma - \partial_t w_\gamma) \, dx + \gamma \int_\Omega \ln u_\gamma(\theta v_\gamma - u_\gamma) \, dx.
\end{aligned}$$

On the other hand, we infer from (1.2b), (1.2c), and (1.2d) that

$$\begin{aligned}
\frac{d}{dt} \int_\Omega L_\theta(v_\gamma) \, dx &= -\gamma \int_\Omega \ln(\theta v_\gamma)(\theta v_\gamma - u_\gamma) \, dx, \\
\frac{1}{2\gamma} \frac{d}{dt} \|v_\gamma\|_2^2 &= \int_\Omega u_\gamma v_\gamma \, dx - \theta \|v_\gamma\|_2^2,
\end{aligned}$$

and

$$\begin{aligned}
\frac{\theta}{2} \frac{d}{dt} \left( \|\nabla w_\gamma\|_2^2 + \frac{\alpha}{D} \|w_\gamma\|_2^2 \right) &= \frac{\theta}{D} \int_\Omega v_\gamma \partial_t w_\gamma \, dx - \frac{\theta \|\partial_t w_\gamma\|_2^2}{D} \\
\frac{1}{2\gamma D} \frac{d}{dt} \|\partial_t w_\gamma\|_2^2 &= \frac{1}{\gamma D} \int_\Omega \partial_t v_\gamma \partial_t w_\gamma \, dx - \frac{\|\nabla \partial_t w_\gamma\|_2^2}{\gamma} - \frac{\alpha \|\partial_t w_\gamma\|_2^2}{\gamma D}.
\end{aligned}$$

Introducing

$$P_\gamma(u_\gamma, v_\gamma, w_\gamma) := \|L(u_\gamma)\|_1 + \|L_\theta(v_\gamma)\|_1 + \frac{\theta}{2} \|\nabla w_\gamma\|_2^2 + \frac{\alpha\theta}{2D} \|w_\gamma\|_2^2 + \frac{\|\partial_t w_\gamma\|_2^2}{2\gamma D} + \frac{\|v_\gamma\|_2^2}{2\gamma},$$

we sum up the above five identities and use (1.2b) to obtain

$$\begin{aligned}
\frac{d}{dt} P_\gamma(u_\gamma, v_\gamma, w_\gamma) &= -4\|\nabla\sqrt{u_\gamma}\|_2^2 + \frac{D+1}{D} \int_\Omega u_\gamma v_\gamma \, dx - \frac{\alpha}{D} \int_\Omega u_\gamma w_\gamma \, dx \\
&\quad - \left( \frac{\theta}{D} + \frac{\alpha}{\gamma D} \right) \|\partial_t w_\gamma\|_2^2 - \frac{\|\nabla \partial_t w_\gamma\|_2^2}{\gamma} - \theta \|v_\gamma\|_2^2 \\
&\quad - \gamma \int_\Omega (\theta v_\gamma - u_\gamma) \ln \left( \frac{\theta v_\gamma}{u_\gamma} \right) \, dx + \frac{1}{D} \int_\Omega \left( \frac{1}{\gamma} \partial_t v_\gamma + \theta v_\gamma - u_\gamma \right) \partial_t w_\gamma \, dx \\
&= -4\|\nabla\sqrt{u_\gamma}\|_2^2 + \frac{D+1}{D} \int_\Omega u_\gamma v_\gamma \, dx - \frac{\alpha}{D} \int_\Omega u_\gamma w_\gamma \, dx - \left( \frac{\theta}{D} + \frac{\alpha}{\gamma D} \right) \|\partial_t w_\gamma\|_2^2 \\
&\quad - \frac{\|\nabla \partial_t w_\gamma\|_2^2}{\gamma} - \theta \|v_\gamma\|_2^2 - \gamma \int_\Omega (\theta v_\gamma - u_\gamma) \ln \left( \frac{\theta v_\gamma}{u_\gamma} \right) \, dx.
\end{aligned}$$

Since both  $u_\gamma$  and  $w_\gamma$  are non-negative and the logarithm is an increasing function on  $(0, \infty)$ , we infer from Young's inequality that

$$\frac{d}{dt} P_\gamma(u_\gamma, v_\gamma, w_\gamma) \leq -4\|\nabla\sqrt{u_\gamma}\|_2^2 + \frac{(D+1)^2}{2\theta D^2} \|u_\gamma\|_2^2 + \frac{\theta}{2} \|v_\gamma\|_2^2 - \theta \|v_\gamma\|_2^2 - \frac{\theta}{D} \|\partial_t w_\gamma\|_2^2.$$

Hence,

$$\frac{d}{dt} P_\gamma(u_\gamma, v_\gamma, w_\gamma) \leq -4\|\nabla\sqrt{u_\gamma}\|_2^2 + \frac{(D+1)^2}{2\theta D^2} \|u_\gamma\|_2^2 - \frac{\theta}{2} \|v_\gamma\|_2^2 - \frac{\theta}{D} \|\partial_t w_\gamma\|_2^2. \quad (4.5)$$

Now, let  $K > 1$  to be determined later. By the Gagliardo-Nirenberg inequality (2.4),

$$\begin{aligned}
\|u_\gamma\|_2^2 &= \|\sqrt{u_\gamma}\|_4^4 = \int_\Omega (\sqrt{u_\gamma} - K + K)^4 \mathbf{1}_{(K^2, \infty)}(u_\gamma) \, dx + \int_\Omega u_\gamma^2 \mathbf{1}_{[0, K^2]}(u_\gamma) \, dx \\
&\leq \int_\Omega ((\sqrt{u_\gamma} - K)_+ + K)^4 \mathbf{1}_{(K^2, \infty)}(u_\gamma) \, dx + K^2 \int_\Omega u_\gamma \mathbf{1}_{[0, K^2]}(u_\gamma) \, dx \\
&\leq 8 \int_\Omega (\sqrt{u_\gamma} - K)_+^4 \, dx + 8K^4 \int_\Omega \mathbf{1}_{(K^2, \infty)}(u_\gamma) \, dx + K^2 \int_\Omega u_\gamma \mathbf{1}_{[0, K^2]}(u_\gamma) \, dx \\
&\leq 8c_0^4 \left( \|\nabla(\sqrt{u_\gamma} - K)_+\|_2^2 \|\sqrt{u_\gamma} - K)_+\|_2^2 + \|(\sqrt{u_\gamma} - K)_+\|_2^4 \right) \\
&\quad + 8K^2 \int_\Omega u_\gamma \mathbf{1}_{(K^2, \infty)}(u_\gamma) \, dx + K^2 \int_\Omega u_\gamma \mathbf{1}_{[0, K^2]}(u_\gamma) \, dx \\
&\leq 8c_0^4 \|u_\gamma \mathbf{1}_{(K^2, \infty)}(u_\gamma)\|_1 \|\nabla \sqrt{u_\gamma}\|_2^2 + 8c_0^4 \|u_\gamma\|_1^2 + 8K^2 \|u_\gamma\|_1.
\end{aligned}$$

Since

$$\|u_\gamma \mathbf{1}_{(K^2, \infty)}(u_\gamma)\|_1 \leq \frac{1}{2 \ln K} \int_\Omega u_\gamma \ln u_\gamma \mathbf{1}_{(K^2, \infty)}(u_\gamma) \, dx \leq \frac{\|L(u_\gamma)\|_1 + \|u_\gamma\|_1}{2 \ln K},$$

we combine the above two inequalities and (2.3) to obtain

$$\|u_\gamma\|_2^2 \leq \frac{4c_0^4 (\|L(u_\gamma)\|_1 + M)}{\ln K} \|\nabla \sqrt{u_\gamma}\|_2^2 + 8M (c_0^4 M + K^2).$$

At this point, we choose

$$K = \exp \left\{ \frac{c_0^4 (D+1)^2}{\theta D^2} (\|L(u_\gamma)\|_1 + M) \right\}$$

in the above inequality and end up with

$$\|u_\gamma\|_2^2 \leq \frac{4\theta D^2}{(D+1)^2} \|\nabla \sqrt{u_\gamma}\|_2^2 + 8M \left( c_0^4 M + \exp \left\{ \frac{2c_0^4 (D+1)^2}{\theta D^2} (\|L(u_\gamma)\|_1 + M) \right\} \right). \quad (4.6)$$

Since  $\|L(u_\gamma)\|_1 \leq P_\gamma(u_\gamma, v_\gamma, w_\gamma)$ , it follows from (4.5) and (4.6) that there is  $C_1 > 0$  such that

$$\frac{d}{dt} P_\gamma(u_\gamma, v_\gamma, w_\gamma) \leq -2 \|\nabla \sqrt{u_\gamma}\|_2^2 - \frac{\theta}{2} \|v_\gamma\|_2^2 - \frac{\theta}{D} \|\partial_t w_\gamma\|_2^2 + C_1 e^{C_1 P_\gamma(u_\gamma, v_\gamma, w_\gamma)}. \quad (4.7)$$

We first infer from (4.7) that

$$\frac{d}{dt} P_\gamma(u_\gamma, v_\gamma, w_\gamma) \leq C_1 e^{C_1 P_\gamma(u_\gamma, v_\gamma, w_\gamma)}, \quad t > 0;$$

hence, after integration with respect to time,

$$P_\gamma(u_\gamma(t), v_\gamma(t), w_\gamma(t)) \leq P_\gamma(u_0, v_0, w_0) - \frac{1}{C_1} \ln \left( 1 - \frac{t}{T_\gamma} \right), \quad t \in [0, T_\gamma), \quad (4.8)$$

where

$$T_\gamma := \frac{e^{-C_1 P_\gamma(u_0, v_0, w_0)}}{C_1^2} \in (0, \infty).$$

Now, in view of  $\gamma \geq 1$ ,

$$\begin{aligned} P_\gamma(u_0, v_0, w_0) &\leq P_1(u_0, v_0, w_0) := \|L(u_0)\|_1 + \|L_\theta(v_0)\|_1 + \frac{\theta}{2} \|\nabla w_0\|_2^2 + \frac{\alpha\theta}{2D} \|w_0\|_2^2 \\ &\quad + \frac{1}{2D} \|D\Delta w_0 - \alpha w_0 + v_0\|_2^2 + \frac{1}{2} \|v_0\|_2^2, \end{aligned}$$

so that

$$T_\gamma \geq T_\infty := \frac{e^{-C_1 P_1(u_0, v_0, w_0)}}{C_1^2}, \quad (4.9)$$

and the estimates (4.2) readily follow from (4.8), while (4.3) is a consequence of (4.7) and (4.8) after integrating the former with respect to time.  $\square$

**4.2. Global estimates.** We next show that the outcome of Lemma 4.2 is valid for any  $T > 0$  provided  $M = \|u_0\|_1 + \|v_0\|_1$  is appropriately small. The building block of the proof, which is actually one of the main contributions of [14], is the construction of a Liapunov function for (1.2)–(1.3), which we recall now. We set

$$\begin{aligned} \mathcal{L}_\gamma(u, v, w) &:= \int_\Omega (L(u) + L_\theta(v) - (u + v)w) \, dx + \frac{1 + \theta}{2} (D\|\nabla w\|_2^2 + \alpha\|w\|_2^2) \\ &\quad + \frac{1}{2\gamma} \|D\Delta w - \alpha w + v\|_2^2, \end{aligned} \quad (4.10)$$

where  $L$  and  $L_\theta$  are defined in (4.4). Adapting the computation performed in [14, Lemma 2.1] to (1.2) leads to the following differential inequality.

**Proposition 4.3.** *Let  $M > 0$  and  $\gamma > 0$ . Consider  $(u_0, v_0, w_0)$  satisfying (1.6) and (1.7) and denote the corresponding solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$ . Then*

$$\frac{d}{dt} \mathcal{L}_\gamma(u_\gamma, v_\gamma, w_\gamma) + \mathcal{D}_\gamma(u_\gamma, v_\gamma, w_\gamma) = 0, \quad t > 0, \quad (4.11)$$

where  $\mathcal{D}_\gamma$  is non-negative and given by

$$\begin{aligned} \mathcal{D}_\gamma(u, v, w) &:= \int_\Omega u |\nabla(\ln u - w)|^2 \, dx + \gamma \int_\Omega (\theta v - u)(\ln(\theta v) - \ln u) \, dx \\ &\quad + \frac{D}{\gamma} \|\nabla(D\Delta w - \alpha w + v)\|_2^2 + \left(1 + \theta + \frac{\alpha}{\gamma}\right) \|D\Delta w - \alpha w + v\|_2^2. \end{aligned} \quad (4.12)$$

From now on, we fix

$$M \in (0, 4\pi(1 + \theta)D) \quad (4.13)$$

and consider  $(u_0, v_0, w_0) \in W_{3,+}^1(\Omega; \mathbb{R}^3)$  with  $w_0 \in W_2^2(\Omega)$  satisfying (1.7). For  $\gamma > 0$ , we denote the corresponding solution to (1.2)–(1.3) provided by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$  and set  $n_\gamma := u_\gamma + v_\gamma$ . By (2.3), there holds

$$\|n_\gamma(t)\|_1 = M, \quad t \geq 0. \quad (4.14)$$

We begin with a lower bound on  $\mathcal{L}_\gamma(u_\gamma, v_\gamma, w_\gamma)$  which is obtained as in [14, Lemma 4.3] with the help of the Moser-Trudinger inequality recalled in Proposition A.1.

**Lemma 4.4.** For  $\gamma > 0$  and  $t \geq 0$ ,

$$\begin{aligned} \mathcal{L}_\gamma(u_\gamma(t), v_\gamma(t), w_\gamma(t)) &\geq \frac{4\pi(1+\theta)D - M}{8\pi} \|\nabla w_\gamma(t)\|_2^2 + \frac{\alpha(1+\theta)}{2} \|w_\gamma(t)\|_2^2 \\ &\quad + \frac{\|\partial_t w_\gamma(t)\|_2^2}{2\gamma} - C_2. \end{aligned}$$

From Lemma 4.4, we derive the same estimates as in Lemma 4.2 but valid for arbitrary positive times.

**Lemma 4.5.** For  $\gamma \geq 1$  and  $t \geq 0$ ,

$$\begin{aligned} \|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1 + \|w_\gamma(t)\|_{W_2^1} &\leq C_3, \\ \int_0^\infty \|\partial_t w_\gamma(s)\|_2^2 ds &\leq C_3. \end{aligned}$$

*Proof.* We argue as in [14, Lemma 4.4]. Let  $t > 0$ . On the one hand, it follows from Proposition 4.3 that

$$\mathcal{L}_\gamma(u_\gamma(t), v_\gamma(t), w_\gamma(t)) + \int_0^t \mathcal{D}_\gamma(u_\gamma(s), v_\gamma(s), w_\gamma(s)) ds \leq \mathcal{L}_\gamma(u_0, v_0, w_0) \leq \mathcal{L}_1(u_0, v_0, w_0).$$

On the other hand, we infer from Lemma 4.4 that

$$\mathcal{L}_\gamma(u_\gamma(t), v_\gamma(t), w_\gamma(t)) \geq \frac{4\pi(1+\theta)D - M}{8\pi} \|\nabla w_\gamma(t)\|_2^2 + \frac{\alpha(1+\theta)}{2} \|w_\gamma(t)\|_2^2 - C_2$$

and

$$\mathcal{D}_\gamma(u(s), v(s), w(s)) \geq (1+\theta) \|(D\Delta w_\gamma - \alpha w_\gamma + v_\gamma)(s)\|_2^2 = (1+\theta) \|\partial_t w_\gamma(s)\|_2^2, \quad s \geq 0.$$

Combining the above inequalities gives

$$\|w_\gamma(t)\|_{W_2^1}^2 + \int_0^t \|\partial_t w_\gamma(s)\|_2^2 ds \leq C. \quad (4.15)$$

Next, by (4.10), Proposition 4.3, and Young's inequality  $ab \leq L(a) + e^b - 1$ ,

$$\begin{aligned} &\|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1 \\ &\leq \mathcal{L}_\gamma(u_\gamma(t), v_\gamma(t), w_\gamma(t)) + \int_\Omega (u_\gamma + v_\gamma)(t) w_\gamma(t) dx \\ &\leq \mathcal{L}_\gamma(u_0, v_0, w_0) + \int_\Omega \left[ L\left(\frac{u_\gamma(t)}{2}\right) + \frac{1}{\theta} L\left(\frac{\theta v_\gamma(t)}{2}\right) + \frac{1+\theta}{\theta} (e^{2w_\gamma(t)} - 1) \right] dx \\ &\leq \mathcal{L}_1(u_0, v_0, w_0) + \int_\Omega \left[ \frac{L(u_\gamma(t)) + 1}{2} + \frac{L(\theta v_\gamma(t)) + 1}{2\theta} + \frac{1+\theta}{\theta} e^{2w_\gamma(t)} \right] dx \\ &\leq C + \frac{1}{2} [\|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1] + \frac{1+\theta}{\theta} \|e^{2w_\gamma(t)}\|_1. \end{aligned}$$



We now use the Moser-Trudinger inequality recalled in Proposition A.1, along with (4.15), to estimate the last term on the right-hand side of the above inequality and conclude that

$$\|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1 \leq C,$$

thereby completing the proof of Lemma 4.5.  $\square$

### 4.3. Proof of Proposition 4.1.

*Proof of Proposition 4.1.* Setting  $T_\infty = \infty$  when  $M \in (0, 4\pi D(1 + \theta))$ , we consider  $T \in (0, T_\infty)$  and recall that Lemma 4.2 and Lemma 4.5 imply that, for  $\gamma \geq 1$ ,

$$\|L(u_\gamma(t))\|_1 + \|L_\theta(v_\gamma(t))\|_1 + \|w_\gamma(t)\|_{W_2^1} \leq C_4(T), \quad t \in (0, T), \quad (4.16)$$

$$\int_0^T \|\partial_t w_\gamma(t)\|_2^2 dt \leq C_4(T). \quad (4.17)$$

Let  $t \in (0, T]$ . On the one hand, it follows from (1.2) that

$$\begin{aligned} \frac{d}{dt} \left( \|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2 \right) &= -2 \|\nabla u_\gamma\|_2^2 + 2 \int_\Omega u_\gamma \nabla u_\gamma \cdot \nabla w_\gamma dx - 2\gamma \|e_\gamma\|_2^2 \\ &= -2 \|\nabla u_\gamma\|_2^2 - 2\gamma \|e_\gamma\|_2^2 - \int_\Omega u_\gamma^2 \Delta w_\gamma dx \\ &= -2 \|\nabla u_\gamma\|_2^2 - 2\gamma \|e_\gamma\|_2^2 + \frac{1}{D} \int_\Omega u_\gamma^2 (v_\gamma - \alpha w_\gamma - \partial_t w_\gamma) dx. \end{aligned}$$

Hence, by Hölder's inequality,

$$\frac{d}{dt} \left( \|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2 \right) \leq -2 \|\nabla u_\gamma\|_2^2 - 2\gamma \|e_\gamma\|_2^2 + \frac{1}{D} \int_\Omega u_\gamma^2 v_\gamma dx + \frac{1}{D} \|u_\gamma\|_4^2 \|\partial_t w_\gamma\|_2.$$

On the other hand, we deduce from (1.2b) that

$$\frac{1}{\gamma} \frac{d}{dt} \|v_\gamma\|_3^3 = 3 \int_\Omega u_\gamma v_\gamma^2 dx - 3\theta \|v_\gamma\|_3^3.$$

Combining the previous two inequalities and introducing

$$Y_\gamma := 1 + \|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2 + \frac{1}{\gamma} \|v_\gamma\|_3^3$$

lead us to the differential inequality

$$\begin{aligned} \frac{dY_\gamma}{dt} &\leq -2 \|\nabla u_\gamma\|_2^2 - 2\gamma \|e_\gamma\|_2^2 - 3\theta \|v_\gamma\|_3^3 \\ &\quad + \frac{1}{D} \int_\Omega u_\gamma^2 v_\gamma dx + 3 \int_\Omega u_\gamma v_\gamma^2 dx + \frac{1}{D} \|u_\gamma\|_4^2 \|\partial_t w_\gamma\|_2. \end{aligned} \quad (4.18)$$

We now infer from Young's inequality that

$$\frac{1}{D} \int_\Omega u_\gamma^2 v_\gamma dx + 3 \int_\Omega u_\gamma v_\gamma^2 dx \leq \frac{\theta}{3} \|v_\gamma\|_3^3 + \frac{2}{3} \frac{\|u_\gamma\|_3^3}{\sqrt{\theta D^3}} + \frac{2\theta}{3} \|v_\gamma\|_3^3 + \frac{9}{\theta^2} \|u_\gamma\|_3^3$$

$$\leq \theta \|v_\gamma\|_3^3 + C \|u_\gamma\|_3^3,$$

while the Gagliardo-Nirenberg inequality (2.4), along with Young's inequality, entails that

$$\begin{aligned} \frac{1}{D} \|u_\gamma\|_4^2 \|\partial_t w_\gamma\|_2 &\leq \frac{c_0^2}{D} \|u_\gamma\|_{W_2^1} \|u_\gamma\|_2 \|\partial_t w_\gamma\|_2 \leq \frac{c_0^2}{D} \left( \|\nabla u_\gamma\|_2 \|u_\gamma\|_2 + \|u_\gamma\|_2^2 \right) \|\partial_t w_\gamma\|_2 \\ &\leq \frac{1}{2} \|\nabla u_\gamma\|_2^2 + C \|u_\gamma\|_2^2 \|\partial_t w_\gamma\|_2^2 + C \|u_\gamma\|_2^2 \|\partial_t w_\gamma\|_2 \\ &\leq \frac{1}{2} \|\nabla u_\gamma\|_2^2 + C \|u_\gamma\|_2^2 (1 + \|\partial_t w_\gamma\|_2^2). \end{aligned}$$

Collecting the above estimates gives

$$\frac{dY_\gamma}{dt} + \frac{3}{2} \|\nabla u_\gamma\|_2^2 + 2\gamma \|e_\gamma\|_2^2 + 2\theta \|v_\gamma\|_3^3 \leq C_5 \|u_\gamma\|_3^3 + C_5 \|u_\gamma\|_2^2 (1 + \|\partial_t w_\gamma\|_2^2). \quad (4.19)$$

Finally, let  $\eta > 0$ . By [5, Equation (22)], there exists  $C_6(\eta) > 0$  (depending actually only on  $\Omega$  and  $\eta$ ) such that

$$\|z\|_3^3 \leq \eta \|z \ln |z|\|_1 \|z\|_{W_2^1}^2 + C_6(\eta) \|z\|_1, \quad z \in W_2^1(\Omega).$$

Since  $r |\ln r| \leq L(r) + r$  for  $r \geq 0$ , we deduce from (2.3), (4.16), and the above functional inequality that

$$\begin{aligned} C_5 \|u_\gamma\|_3^3 &\leq \eta C_5 \|L(u_\gamma) + u_\gamma\|_1 \|u_\gamma\|_{W_2^1}^2 + C_5 C_6(\eta) \|u_\gamma\|_1 \\ &\leq \eta C_5 (C_4(T) + M) \|\nabla u_\gamma\|_2^2 + \eta C_5 (C_4(T) + M) \|u_\gamma\|_2^2 + M C_5 C_6(\eta). \end{aligned}$$

Choosing  $\eta = 1/[2C_5(C_4(T) + M)]$  in the above inequality and inserting the outcome in (4.19), we end up with

$$\frac{dY_\gamma}{dt} + \|\nabla u_\gamma\|_2^2 + 2\gamma \|e_\gamma\|_2^2 + 2\theta \|v_\gamma\|_3^3 \leq C_7(T) (1 + \|u_\gamma\|_2^2) (1 + \|\partial_t w_\gamma\|_2^2). \quad (4.20)$$

A first consequence of (4.20) is that

$$\frac{dY_\gamma}{dt} \leq C_7(T) (1 + \|\partial_t w_\gamma\|_2^2) Y_\gamma, \quad t \in (0, T].$$

Hence, after integrating with respect to time and using (4.17), we obtain

$$Y_\gamma(t) \leq Y_\gamma(0) \exp \left\{ C_7(T) \left( t + \int_0^t \|\partial_t w_\gamma(s)\|_2^2 ds \right) \right\} \leq C e^{C_7(T)t + C_7(T)C_4(T)} \leq C_8(T) \quad (4.21)$$

for  $t \in [0, T]$ , from which (4.1a) follows.

We next integrate (4.20) with respect to time and use the non-negativity of  $Y_\gamma$ , along with (4.17) and (4.21), to find

$$\begin{aligned} \int_0^t \left[ \|\nabla u_\gamma(s)\|_2^2 + \gamma \|e_\gamma(s)\|_2^2 \right] ds &\leq Y_\gamma(0) + C_7(T) \sup_{s \in [0, t]} \{1 + \|u_\gamma(s)\|_2^2\} \left( t + \int_0^t \|\partial_t w_\gamma(s)\|_2^2 ds \right) \\ &\leq C + C_7(T) C_8(T) (t + C_4(T)), \end{aligned}$$

and thus complete the proof.  $\square$

## 5. CONVERGENCE: PROOF OF THEOREM 1.1

*Proof of Theorem 1.1.* Let  $M > 0$  and initial conditions  $(u_0, v_0, w_0)$  satisfying (1.6) and (1.7). For  $\gamma > 0$ , we denote the solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$  and set  $n_\gamma = u_\gamma + v_\gamma$  and  $e_\gamma = \theta v_\gamma - u_\gamma$ . According to Proposition 4.1, there is  $T_\infty \in (0, \infty]$  such that the estimates (4.1) hold true for all  $T \in (0, T_\infty)$ , having set  $T_\infty = \infty$  in Case (b). We are thus in a position to apply Proposition 3.1 for any  $T \in (0, T_\infty)$  and use a diagonal process to construct a sequence  $(\gamma_j)_{j \geq 1}$ ,  $\gamma_j \rightarrow \infty$ , and non-negative functions

$$\begin{aligned} n &\in W_{2,\text{loc}}^1([0, T_\infty), W_2^1(\Omega)') \cap L_{\infty,\text{loc}}([0, T_\infty), L_2(\Omega)) \cap L_{2,\text{loc}}([0, T_\infty), W_2^1(\Omega)), \\ w &\in W_{2,\text{loc}}^1([0, T_\infty), L_2(\Omega)) \cap L_{\infty,\text{loc}}([0, T_\infty), W_2^1(\Omega)) \cap L_{2,\text{loc}}([0, T_\infty), W_2^2(\Omega)), \end{aligned}$$

such that the convergences (3.4), (3.5), (3.6), and (3.7) hold true for any  $T \in (0, T_\infty)$ .

Now, it readily follows from (1.2a), (1.2b), and (1.2d) that, for  $j \geq 1$ ,  $t \in (0, T_\infty)$ , and  $\varphi \in W_2^1(\Omega)$ ,

$$\int_0^t \langle \partial_t n_{\gamma_j}(s), \varphi \rangle_{(W_2^1)', W_2^1} ds + \int_0^t \int_\Omega \nabla \varphi \cdot (\nabla u_{\gamma_j} - u_{\gamma_j} \nabla w_{\gamma_j})(s) \, dx ds = 0.$$

Owing to (3.4), (3.6), and (3.7), it is straightforward to pass to the limit  $j \rightarrow \infty$  in the above identity and conclude that  $n$  solves

$$\int_0^t \langle \partial_t n(s), \varphi \rangle_{(W_2^1)', W_2^1} ds + \frac{\theta}{1+\theta} \int_0^t \int_\Omega \nabla \varphi \cdot (\nabla n - n \nabla w)(s) \, dx ds = 0 \quad (5.1)$$

for all  $t \in (0, T_\infty)$  and  $\varphi \in W_2^1(\Omega)$ . Similarly, we infer from (1.2c), (3.4), (3.5), and (3.6) that

$$\frac{d}{dt} \int_\Omega w \varphi \, dx + D \int_\Omega \nabla w \cdot \nabla \varphi \, dx + \alpha \int_\Omega w \varphi \, dx = \frac{1}{1+\theta} \int_\Omega n \varphi \, dx \quad (5.2)$$

for all  $t \in (0, T_\infty)$  and  $\varphi \in W_2^1(\Omega)$ . Actually, the regularity of  $w$  and  $n$ , together with (5.2), implies that  $w$  is a strong solution to

$$\begin{aligned} \partial_t w - D \Delta w + \alpha w &= \frac{n}{1+\theta} && \text{in } (0, T_\infty) \times \Omega, \\ \nabla w \cdot \mathbf{n} &= 0 && \text{on } (0, T_\infty) \times \partial\Omega, \\ w(0) &= w_0 && \text{in } \Omega. \end{aligned} \quad (5.3)$$

In other words,  $(n, w)$  is a weak-strong solution to (1.4)–(1.5) on  $(0, T)$  in the sense of Definition C.1 below for all  $T \in (0, T_\infty)$ , which is unique according to Proposition C.2 below. A classical argument then implies that the convergences (3.4), (3.5), (3.6), and (3.7) hold true for all  $\gamma \geq 1$  and  $T \in (0, T_\infty)$ .

We are left with improving the convergence of  $(w_\gamma)_{\gamma \geq 1}$  from the  $L_2$ -convergence derived in Proposition 3.4 to the  $W_2^1$ -convergence (1.8b) claimed in Theorem 1.1. To this end, we infer from (1.2c), (1.2d), (5.3), and Hölder's and Young's inequalities that

$$\begin{aligned} \|\partial_t(w_\gamma - w)\|_2^2 + \frac{1}{2} \frac{d}{dt} [D \|\nabla(w_\gamma - w)\|_2^2 + \alpha \|w_\gamma - w\|_2^2] \\ = \int_\Omega \left( v_\gamma - \frac{n}{1+\theta} \right) \partial_t(w_\gamma - w) \, dx \end{aligned}$$

$$\leq \frac{1}{2} \|\partial_t(w_\gamma - w)\|_2^2 + \frac{1}{2} \left\| v_\gamma - \frac{n}{1+\theta} \right\|_2^2.$$

Hence,

$$\frac{d}{dt} [D \|\nabla(w_\gamma - w)\|_2^2 + \alpha \|w_\gamma - w\|_2^2] \leq \left\| v_\gamma - \frac{n}{1+\theta} \right\|_2^2,$$

from which we deduce, after integrating with respect to time, that

$$D \|\nabla(w_\gamma - w)(t)\|_2^2 + \alpha \|(w_\gamma - w)(t)\|_2^2 \leq \int_0^t \left\| \left( v_\gamma - \frac{n}{1+\theta} \right) (s) \right\|_2^2 ds$$

for  $t \in [0, T_\infty)$ . Consequently, for  $T \in (0, T_\infty)$ ,

$$\sup_{t \in [0, T]} [D \|\nabla(w_\gamma - w)(t)\|_2^2 + \alpha \|(w_\gamma - w)(t)\|_2^2] \leq \int_0^T \left\| \left( v_\gamma - \frac{n}{1+\theta} \right) (s) \right\|_2^2 ds.$$

Since the right-hand side of the above inequality converges to zero as  $\gamma \rightarrow \infty$  by (3.6), we readily conclude that

$$\lim_{\gamma \rightarrow \infty} \sup_{t \in [0, T]} [D \|\nabla(w_\gamma - w)(t)\|_2^2 + \alpha \|(w_\gamma - w)(t)\|_2^2] = 0.$$

The convergence (1.8b) then follows due to the positivity of  $D$  and  $\alpha$ , and the proof of Theorem 1.1 is complete.  $\square$

## 6. UNBOUNDEDNESS: PROOF OF THEOREM 1.2

The main step of the proof of Theorem 1.2 is the following result asserting that the estimates (3.2) can be derived from a bound on the  $L_2$ -norm of  $v_\gamma$ , whenever available.

**Lemma 6.1.** *Consider  $(u_0, v_0, w_0) \in W_{3,+}^1(\Omega; \mathbb{R}^3)$  and, for  $\gamma > 0$ , denote the solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$ . We assume that there are  $T \in (0, \infty)$  and  $K > 0$  such that*

$$\|v_\gamma(t)\|_2 \leq K \quad \text{for all } t \in (0, T), \quad \gamma \geq 1. \quad (6.1)$$

*Then  $(u_\gamma, v_\gamma, w_\gamma)$  satisfies (3.2) with  $e_\gamma = \theta v_\gamma - u_\gamma$ .*

*Proof.* In the following,  $C$  denotes positive constants depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $u_0$ ,  $v_0$ ,  $w_0$ ,  $T$ , and  $K$ .

It readily follows from (1.2c), (1.2d), (6.1), and Young's inequality that

$$\begin{aligned} \|\partial_t w_\gamma\|_2^2 + \frac{1}{2} \frac{d}{dt} [D \|\nabla w_\gamma\|_2^2 + \alpha \|w_\gamma\|_2^2] &= \int_\Omega v_\gamma \partial_t w_\gamma \, dx \\ &\leq \frac{1}{2} \|v_\gamma\|_2^2 + \frac{1}{2} \|\partial_t w_\gamma\|_2^2 \leq \frac{K^2}{2} + \frac{1}{2} \|\partial_t w_\gamma\|_2^2. \end{aligned}$$

Hence,

$$\|\partial_t w_\gamma\|_2^2 + \frac{d}{dt} [D \|\nabla w_\gamma\|_2^2 + \alpha \|w_\gamma\|_2^2] \leq K^2,$$

from which we deduce, after integration with respect to time,

$$\sup_{t \in [0, T]} \|w_\gamma(t)\|_{W_2^1}^2 + \int_0^T \|\partial_t w_\gamma(t)\|_2^2 dt \leq C. \quad (6.2)$$

We next infer from (1.2a), (1.2b), (1.2d), the Gagliardo-Nirenberg inequality (2.4), and Hölder's inequality that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} [\|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2] &= - \int_\Omega \nabla u_\gamma \cdot (\nabla u_\gamma - u_\gamma \nabla w_\gamma) dx - \gamma \|e_\gamma\|_2^2 \\ &= - \|\nabla u_\gamma\|_2^2 - \frac{1}{2} \int_\Omega u_\gamma^2 \Delta w_\gamma dx - \gamma \|e_\gamma\|_2^2 \\ &\leq - \|u_\gamma\|_{W_2^1}^2 + \|u_\gamma\|_2^2 + \frac{1}{2} \|u_\gamma\|_4^2 \|\Delta w_\gamma\|_2 - \gamma \|e_\gamma\|_2^2 \\ &\leq - \|u_\gamma\|_{W_2^1}^2 + \|u_\gamma\|_2^2 + \frac{c_0^2}{2} \|u_\gamma\|_{W_2^1} \|u_\gamma\|_2 \|\Delta w_\gamma\|_2 - \gamma \|e_\gamma\|_2^2. \end{aligned}$$

Using now (1.2c), Young's inequality, and (6.1), we further obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} [\|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2] &\leq - \|u_\gamma\|_{W_2^1}^2 + \|u_\gamma\|_2^2 + \frac{1}{2} \|u_\gamma\|_{W_2^1}^2 \\ &\quad + \frac{c_0^4}{8D^2} \|u_\gamma\|_2^2 \|\partial_t w_\gamma + \alpha w_\gamma - v_\gamma\|_2^2 - \gamma \|e_\gamma\|_2^2 \\ &\leq - \frac{1}{2} \|u_\gamma\|_{W_2^1}^2 + C \|u_\gamma\|_2^2 (1 + \|\partial_t w_\gamma\|_2^2 + \|w_\gamma\|_2^2 + \|v_\gamma\|_2^2) - \gamma \|e_\gamma\|_2^2 \\ &\leq - \frac{1}{2} \|u_\gamma\|_{W_2^1}^2 - \gamma \|e_\gamma\|_2^2 + C \|u_\gamma\|_2^2 (1 + \|\partial_t w_\gamma\|_2^2 + \|w_\gamma\|_2^2). \end{aligned} \quad (6.3)$$

As this yields

$$\frac{d}{dt} [\|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2] \leq C (\|u_\gamma\|_2^2 + \theta \|v_\gamma\|_2^2) (1 + \|\partial_t w_\gamma\|_2^2 + \|w_\gamma\|_2^2),$$

Gronwall's lemma in conjunction with (6.2) implies that, for all  $t \in [0, T]$ ,

$$\|u_\gamma(t)\|_2^2 + \theta \|v_\gamma(t)\|_2^2 \leq C \exp \left\{ C \int_0^T (1 + \|\partial_t w_\gamma(s)\|_2^2 + \|w_\gamma(s)\|_2^2) ds \right\} \leq C. \quad (6.4)$$

Integrating (6.3) with respect to time, we deduce from (6.2) and (6.4) that

$$\int_0^T [\|u_\gamma(t)\|_{W_2^1}^2 + \gamma \|e_\gamma(t)\|_2^2] dt \leq C. \quad (6.5)$$

Collecting (6.1), (6.2), (6.4), and (6.5) shows that  $(u_\gamma, v_\gamma, w_\gamma)$  satisfies (3.2) on  $(0, T)$  for all  $\gamma \geq 1$ , as claimed.  $\square$

We further show that a corresponding bound on  $u_\gamma$  implies the bound (6.1) on  $v_\gamma$ . Here we adapt the idea from [19, Lemma 4.11].

**Lemma 6.2.** Consider  $(u_0, v_0, w_0) \in W_{3,+}^1(\Omega; \mathbb{R}^3)$  and, for  $\gamma > 0$ , denote the solution to (1.2)–(1.3) given by Proposition 2.1 by  $(u_\gamma, v_\gamma, w_\gamma)$ . Assume that there are  $T \in (0, \infty)$  and  $\tilde{K} > 0$  such that

$$\|u_\gamma(t)\|_2 \leq \tilde{K} \quad \text{for all } t \in (0, T), \quad \gamma \geq 1. \quad (6.6)$$

Then there is  $K > 0$  such that (6.1) is fulfilled.

*Proof.* Using (1.2b), Hölder's inequality, and (6.6), we have for  $t \in (0, T)$

$$\frac{d}{dt} \|v_\gamma\|_2^2 = 2\gamma \left( \int_{\Omega} u_\gamma v_\gamma \, dx - \theta \|v_\gamma\|_2^2 \right) \leq 2\gamma (\|u_\gamma\|_2 \|v_\gamma\|_2 - \theta \|v_\gamma\|_2^2) \leq 2\gamma (\tilde{K} - \theta \|v_\gamma\|_2) \|v_\gamma\|_2.$$

In view of  $v_\gamma(0) = v_0$ , this implies (6.1) with  $K := \max \left\{ \frac{\tilde{K}}{\theta}, \|v_0\|_2 \right\}$  by the comparison principle.  $\square$

*Proof of Theorem 1.2.* It first follows from (1.9) and [12, Theorem 3.6] that

$$\lim_{t \rightarrow T_{bu}} \|n(t)\|_2 = \infty. \quad (6.7)$$

Next, let  $T > T_{bu}$  and assume for contradiction that there is  $K > 0$  such that

$$\|v_\gamma(t)\|_2 \leq K \quad \text{for all } t \in (0, T), \quad \gamma \geq 1. \quad (6.8)$$

Then (3.2) is fulfilled by Lemma 6.1. Hence, by Proposition 3.1 and the proof of Theorem 1.1,  $(n_\gamma, w_\gamma)$  converges as  $\gamma \rightarrow \infty$  to a weak-strong solution  $(\tilde{n}, \tilde{w})$  to (1.4)–(1.5) on  $(0, T)$ . As the classical solution  $(n, w)$  to (1.4)–(1.5) is a weak-strong solution and the latter is unique by Proposition C.2, we have  $(n, w) = (\tilde{n}, \tilde{w})$ . Consequently,  $n \in L_\infty((0, T), L_2(\Omega))$  which contradicts (6.7) in view of  $T > T_{bu}$ . Hence, (6.8) cannot be valid. In view of Lemma 6.2, this further implies that

$$\|u_\gamma(t)\|_2 \leq \tilde{K} \quad \text{for all } t \in (0, T), \quad \gamma \geq 1$$

cannot be satisfied for some  $\tilde{K} > 0$ , and the proof of Theorem 1.2 is complete.  $\square$

## APPENDIX A. THE MOSER-TRUDINGER INEQUALITY

We recall here the version of the Moser-Trudinger inequality derived in [8, Corollary 2.7] and in [18, Section 2] from [6, Proposition 2.3] and which is used in the proof of Lemma 4.5.

**Proposition A.1.** *There is  $K_0 > 0$  depending only on  $\Omega$  such that, for all  $z \in W_2^1(\Omega)$ ,*

$$\int_{\Omega} e^{|z|} \, dx \leq K_0 \exp \left( \frac{\|\nabla z\|_2^2}{8\pi} + \frac{\|z\|_1}{|\Omega|} \right).$$

## APPENDIX B. RESCALING (1.4)

Let  $(n, w)$  be a solution to (1.4) with initial condition  $(n_0, w_0)$ . Performing a simple scaling transforms (1.4) to a version of the parabolic-parabolic chemotaxis Keller-Segel system which is usually used in the literature. Indeed, introducing

$$U(s, x) := \frac{1}{\theta} n \left( \frac{1+\theta}{\theta} s, x \right), \quad V(s, x) := w \left( \frac{1+\theta}{\theta} s, x \right), \quad (s, x) \in (0, \infty) \times \Omega, \quad (\text{B.1})$$

we readily deduce from (1.4) that  $(U, V)$  solves the Keller-Segel system

$$\partial_s U = \operatorname{div}(\nabla U - U \nabla V) \quad \text{in } (0, \infty) \times \Omega, \quad (\text{B.2a})$$

$$\frac{\theta}{(1+\theta)D} \partial_s V = \Delta w - \frac{\alpha}{D} V + \frac{\theta}{(1+\theta)D} U \quad \text{in } (0, \infty) \times \Omega, \quad (\text{B.2b})$$

$$(\nabla U - U \nabla V) \cdot \mathbf{n} = \nabla V \cdot \mathbf{n} = 0 \quad \text{on } (0, \infty) \times \partial\Omega, \quad (\text{B.2c})$$

$$(U, V)(0) = \left( \frac{n_0}{\theta}, w_0 \right) \quad \text{in } \Omega. \quad (\text{B.2d})$$

According to [8, 18], global existence holds true for (B.2) if

$$\|n_0\|_1 = \theta \frac{\|n_0\|_1}{\theta} < \theta \frac{4\pi(1+\theta)D}{\theta} = 4\pi(1+\theta)D,$$

while finite time blowup may occur when  $\|n_0\|_1$  exceeds the above threshold [10–12].

## APPENDIX C. UNIQUENESS OF WEAK-STRONG SOLUTIONS TO (1.4)–(1.5)

We provide here a proof of the uniqueness result of weak-strong solutions to (1.4)–(1.5) which is used in the proof of Theorem 1.1 to obtain the convergence of the whole family  $(n_\gamma)_{\gamma \geq 1}$ . We first make precise the meaning of weak-strong solutions to (1.4)–(1.5).

**Definition C.1.** *Let  $T \in (0, \infty)$  and  $(n_0, w_0) \in L_{2,+}(\Omega) \times W_{2,+}^1(\Omega)$ . A weak-strong solution to (1.4)–(1.5) on  $[0, T]$  is a pair of functions*

$$n \in W_2^1((0, T), W_2^1(\Omega)') \cap L_\infty((0, T), L_{2,+}(\Omega)) \cap L_2((0, T), W_2^1(\Omega)), \quad n(0) = n_0, \quad (\text{C.1a})$$

$$w \in W_2^1((0, T), L_2(\Omega)) \cap L_\infty((0, T), W_{2,+}^1(\Omega)) \cap L_2((0, T), W_2^2(\Omega)), \quad (\text{C.1b})$$

such that

$$\int_0^t \langle \partial_t n(s), \varphi \rangle_{(W_2^1)', W_2^1} ds + \frac{\theta}{1+\theta} \int_0^t \int_\Omega \nabla \varphi \cdot (\nabla n - n \nabla w)(s) \, dx ds = 0 \quad (\text{C.1c})$$

for all  $t \in (0, T)$  and  $\varphi \in W_2^1(\Omega)$ , while  $w$  is a strong solution to

$$\begin{aligned} \partial_t w - D \Delta w + \alpha w &= \frac{n}{1+\theta} && \text{in } (0, T) \times \Omega, \\ \nabla w \cdot \mathbf{n} &= 0 && \text{on } (0, T) \times \partial\Omega, \\ w(0) &= w_0 && \text{in } \Omega. \end{aligned} \quad (\text{C.1d})$$

Clearly, classical solutions to (1.4) are weak-strong solutions to (1.4), see [13, Theorem 3.1] for instance for the existence of the former. Also, the weak solutions to (1.4) constructed in [4, Theorem 1] for  $(n_0, w_0) \in L_{q,+}(\Omega) \times W_{q,+}^1(\Omega)$ ,  $q > 2$ , in [8, Theorem 3.3] for  $(n_0, w_0) \in L_{\infty,+}(\Omega) \times W_{q,+}^1(\Omega)$ ,  $q > 2$ , in [23] for  $(n_0, w_0) \in W_{2,+}^s(\Omega; \mathbb{R}^2)$ ,  $s > 1$ , and in [24, Chapter 12] for  $(n_0, w_0) \in L_{2,+}(\Omega) \times W_{2,\mathcal{B},+}^2(\Omega)$  are weak-strong solutions to (1.4). Uniqueness results are also provided in [8, 13, 23, 24] but require more regularity than that stated in Definition C.1. Let us also mention that well-posedness in  $L_{p,+}(\Omega) \times W_{q,+}^1(\Omega)$  with  $p > 1$  and  $q > 2$  is obtained in [4]. We shall show now that the assumptions in Definition C.1 are sufficient to guarantee uniqueness. To this end, we adapt the uniqueness argument used in the proof of [13, Theorem 3.1] and rely heavily on the two-dimensional setting and the Gagliardo-Nirenberg inequality (2.4).

**Proposition C.2.** *Let  $T \in (0, \infty)$  and  $(n_0, w_0) \in L_{2,+}(\Omega) \times W_{2,+}^1(\Omega)$ , where  $\Omega \subset \mathbb{R}^2$  is a bounded domain with smooth boundary. There is at most one weak-strong solution to (1.4)–(1.5) on  $[0, T]$ .*

*Proof.* Let  $T > 0$ ,  $(n_0, w_0) \in L_{2,+}(\Omega) \times W_{2,+}^1(\Omega)$ , and consider two weak-strong solutions  $(n_1, w_1)$  and  $(n_2, w_2)$  to (1.4)–(1.5) on  $(0, T)$ . We set  $N := n_1 - n_2$ ,  $W := w_1 - w_2$ ,  $\psi := \|n_2\|_{W_2^1}^2 + \|w_1\|_{W_2^1}^2$  and infer from (C.1a) and (C.1b) that

$$\kappa := \sup_{t \in (0, T)} \left[ \|n_2(t)\|_2 + \|w_1(t)\|_{W_2^1} \right] < \infty \quad \text{and} \quad \psi \in L_1((0, T)). \quad (\text{C.2})$$

In the following,  $C$  denotes positive constants depending only on  $\Omega$ ,  $\theta$ ,  $D$ ,  $\alpha$ ,  $n_0$ ,  $w_0$ ,  $T$ , and  $\kappa$ .

On the one hand, it follows from (C.1a), (C.1c), and Hölder's inequality that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|N\|_2^2 &= \langle \partial_t N, N \rangle_{(W_2^1)', W_2^1} = -\frac{\theta}{1+\theta} \int_{\Omega} \nabla N \cdot (\nabla N - n_1 \nabla w_1 + n_2 \nabla w_2) \, dx \\ &= -\frac{\theta}{1+\theta} \|\nabla N\|_2^2 + \frac{\theta}{1+\theta} \int_{\Omega} N \nabla N \cdot \nabla w_1 \, dx + \frac{\theta}{1+\theta} \int_{\Omega} n_2 \nabla N \cdot \nabla W \, dx \\ &\leq -\frac{\theta}{1+\theta} \|\nabla N\|_2^2 + \frac{\theta}{1+\theta} \|N\|_4 \|\nabla N\|_2 \|\nabla w_1\|_4 + \frac{\theta}{1+\theta} \|n_2\|_4 \|\nabla N\|_2 \|\nabla W\|_4. \end{aligned}$$

We now use the Gagliardo-Nirenberg inequality (2.4) to estimate the  $L_4$ -norms, along with Young's inequality, to obtain

$$\begin{aligned} \frac{1+\theta}{2\theta} \frac{d}{dt} \|N\|_2^2 + \|N\|_{W_2^1}^2 &\leq \|N\|_2^2 + c_0^2 \|N\|_{W_2^1}^{1/2} \|N\|_2^{1/2} \|\nabla N\|_2 \|\nabla w_1\|_{W_2^1}^{1/2} \|\nabla w_1\|_2^{1/2} \\ &\quad + c_0^2 \|n_2\|_{W_2^1}^{1/2} \|n_2\|_2^{1/2} \|\nabla N\|_2 \|\nabla W\|_{W_2^1}^{1/2} \|\nabla W\|_2^{1/2} \\ &\leq \|N\|_2^2 + c_0^2 \kappa^{1/2} \|N\|_2^{1/2} \|N\|_{W_2^1}^{3/2} \|w_1\|_{W_2^2}^{1/2} \\ &\quad + c_0^2 \kappa^{1/2} \|n_2\|_{W_2^1}^{1/2} \|N\|_{W_2^1} \|W\|_{W_2^2}^{1/2} \|W\|_{W_2^1}^{1/2} \\ &\leq C(1 + \|w_1\|_{W_2^2}^2) \|N\|_2^2 + \frac{1}{4} \|N\|_{W_2^1}^2 + C \|n_2\|_{W_2^1} \|W\|_{W_2^2} \|W\|_{W_2^1}. \end{aligned}$$



Hence,

$$\frac{1+\theta}{\theta} \frac{d}{dt} \|N\|_2^2 + \frac{3}{2} \|N\|_{W_2^1}^2 \leq C(1+\psi) \|N\|_2^2 + C \|n_2\|_{W_2^1} \|W\|_{W_2^2} \|W\|_{W_2^1}. \quad (\text{C.3})$$

On the other hand, by (C.1b), (C.1d), and Hölder's and Young's inequalities,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|W\|_{W_2^1}^2 + D(\|\nabla W\|_2^2 + \|\Delta W\|_2^2) + \alpha \|W\|_{W_2^1}^2 &= \frac{1}{1+\theta} \int_{\Omega} (NW + \nabla N \cdot \nabla W) \, dx \\ &\leq \frac{1}{2(1+\theta)} (\|N\|_2^2 + \|W\|_2^2) + \frac{1}{4} \|\nabla N\|_2^2 + \frac{1}{(1+\theta)^2} \|\nabla W\|_2^2. \end{aligned}$$

Since there is  $\delta > 0$  depending only on  $\Omega$ ,  $D$ , and  $\alpha$  such that

$$\delta \|z\|_{W_2^2}^2 \leq D(\|\nabla z\|_2^2 + \|\Delta z\|_2^2) + \alpha \|z\|_{W_2^1}^2, \quad z \in W_2^2(\Omega),$$

we further obtain

$$\frac{d}{dt} \|W\|_{W_2^1}^2 + 2\delta \|W\|_{W_2^2}^2 \leq \frac{1}{2} \|N\|_{W_2^1}^2 + C(\|N\|_2^2 + \|W\|_{W_2^1}^2). \quad (\text{C.4})$$

Combining (C.3) and (C.4) and using once more Young's inequality give

$$\begin{aligned} \frac{d}{dt} \left[ \frac{1+\theta}{\theta} \|N\|_2^2 + \|W\|_{W_2^1}^2 \right] + \|N\|_{W_2^1}^2 + 2\delta \|W\|_{W_2^2}^2 \\ \leq C(1+\psi) \|N\|_2^2 + C \|n_2\|_{W_2^1} \|W\|_{W_2^2} \|W\|_{W_2^1} + C(\|N\|_2^2 + \|W\|_{W_2^1}^2) \\ \leq \delta \|W\|_{W_2^2}^2 + C(1+\psi) (\|N\|_2^2 + \|W\|_{W_2^1}^2). \end{aligned}$$

Consequently,

$$\frac{d}{dt} \left[ \frac{1+\theta}{\theta} \|N\|_2^2 + \|W\|_{W_2^1}^2 \right] \leq C(1+\psi) \left( \frac{1+\theta}{\theta} \|N\|_2^2 + \|W\|_{W_2^1}^2 \right),$$

and the integrability (C.2) of  $\psi$ , along with Gronwall's lemma, entails that  $N = W \equiv 0$  on  $(0, T)$  and completes the proof.  $\square$

#### ACKNOWLEDGMENTS

Part of this work was done while Ph. L. enjoyed the hospitality and support from the Mathematical Institute, Tohoku University, Sendai.

#### REFERENCES

- [1] H. AMANN, *Global existence for a class of highly degenerate parabolic systems*, Japan J. Indust. Appl. Math., 8 (1991), pp. 143–151.
- [2] ———, *Highly degenerate quasilinear parabolic systems*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 18 (1991), pp. 135–166.
- [3] ———, *Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems*, in Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992), vol. 133 of Teubner-Texte Math., Teubner, Stuttgart, 1993, pp. 9–126.

- [4] P. BILER, *Local and global solvability of some parabolic systems modelling chemotaxis*, Adv. Math. Sci. Appl., 8 (1998), pp. 715–743.
- [5] P. BILER, W. HEBISCH, AND T. NADZIEJA, *The Debye system: existence and large time behavior of solutions*, Nonlinear Anal., 23 (1994), pp. 1189–1209.
- [6] S.-Y. A. CHANG AND P. C. YANG, *Conformal deformation of metrics on  $S^2$* , J. Differential Geom., 27 (1988), pp. 259–296.
- [7] K. FUJIE AND T. SENBA, *Application of an Adams type inequality to a two-chemical substances chemotaxis system*, J. Differential Equations, 263 (2017), pp. 88–148.
- [8] H. GAJEWSKI AND K. ZACHARIAS, *Global behaviour of a reaction-diffusion system modelling chemotaxis*, Math. Nachr., 195 (1998), pp. 77–114.
- [9] M. A. HERRERO AND J. J. L. VELÁZQUEZ, *A blow-up mechanism for a chemotaxis model*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 24 (1997), pp. 633–683.
- [10] D. HORSTMANN, *The nonsymmetric case of the Keller-Segel model in chemotaxis: some recent results*, NoDEA Nonlinear Differential Equations Appl., 8 (2001), pp. 399–423.
- [11] ———, *On the existence of radially symmetric blow-up solutions for the Keller-Segel model*, J. Math. Biol., 44 (2002), pp. 463–478.
- [12] D. HORSTMANN AND G. WANG, *Blow-up in a chemotaxis model without symmetry assumptions*, European J. Appl. Math., 12 (2001), pp. 159–177.
- [13] D. HORSTMANN AND M. WINKLER, *Boundedness vs. blow-up in a chemotaxis system*, J. Differential Equations, 215 (2005), pp. 52–107.
- [14] PH. LAURENÇOT AND C. STINNER, *Mass threshold for infinite-time blowup in a chemotaxis model with split population*, SIAM J. Math. Anal., 53 (2021), pp. 3385–3419.
- [15] J. L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Études mathématiques. Paris: Dunod; Paris: Gauthier-Villars. xx, 554 p., 1969.
- [16] F. R. MACFARLANE, T. LORENZI, AND K. J. PAINTER, *The impact of phenotypic heterogeneity on chemotactic self-organisation*, Bull. Math. Biol., 84 (2022), p. 35. Id/No 143.
- [17] B.-V. MATIOC AND CH. WALKER, *Well-posedness of quasilinear parabolic equations in time-weighted spaces*. arXiv: 2312.07974.
- [18] T. NAGAI, T. SENBA, AND K. YOSHIDA, *Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis*, Funkcial. Ekvac., 40 (1997), pp. 411–433.
- [19] K. J. PAINTER AND M. WINKLER, *Phenotype switching in chemotaxis aggregation models controls the spontaneous emergence of large densities*, SIAM J. Appl. Math., 83 (2023), pp. 2096–2117.
- [20] S. RAMAKRISHNAN, T. LAURENT, M. KUMAR, AND A. L. BERTOZZI, *Spatiotemporal chemotactic model for ant foraging*, Modern Phys. Lett. B, 28 (2014), p. 1450238.
- [21] J. SIMON, *Compact sets in the space  $L^p(0, T; B)$* , Ann. Mat. Pura Appl. (4), 146 (1987), pp. 65–96.
- [22] M. WINKLER, *Finite-time blow-up in the higher-dimensional parabolic-parabolic Keller-Segel system*, J. Math. Pures Appl. (9), 100 (2013), pp. 748–767.
- [23] A. YAGI, *Norm behavior of solutions to a parabolic system of chemotaxis*, Math. Japon., 45 (1997), pp. 241–265.
- [24] ———, *Abstract parabolic evolution equations and their applications*, Springer Monogr. Math., Berlin: Springer, 2010.

LABORATOIRE DE MATHÉMATIQUES (LAMA), UMR 5127, UNIVERSITÉ SAVOIE MONT BLANC, CNRS, F-73000 CHAMBÉRY, FRANCE

*Email address:* philippe.laurencot@univ-smb.fr

TECHNISCHE UNIVERSITÄT DARMSTADT, FACHBEREICH MATHEMATIK, SCHLOSSGARTENSTR. 7, D-64289 DARMSTADT, GERMANY

*Email address:* stinner@mathematik.tu-darmstadt.de