



HAL
open science

Emerging Functional Connections Between Metabolism and Epigenetic Remodeling in Neural Differentiation

Edgar Sánchez-Ramírez, Thi Phuong Lien Ung, Chiara Stringari, Lorena Aguilar-Arnal

► **To cite this version:**

Edgar Sánchez-Ramírez, Thi Phuong Lien Ung, Chiara Stringari, Lorena Aguilar-Arnal. Emerging Functional Connections Between Metabolism and Epigenetic Remodeling in Neural Differentiation. *Molecular Neurobiology*, In press, 10.1007/s12035-024-04006-w . hal-04514297

HAL Id: hal-04514297

<https://hal.science/hal-04514297>

Submitted on 21 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Emerging Functional Connections Between Metabolism and Epigenetic Remodeling in Neural Differentiation

Edgar Sánchez-Ramírez¹ · Thi Phuong Lien Ung² · Chiara Stringari² · Lorena Aguilar-Arnal¹

Received: 13 September 2023 / Accepted: 30 January 2024
© The Author(s) 2024

Abstract

Stem cells possess extraordinary capacities for self-renewal and differentiation, making them highly valuable in regenerative medicine. Among these, neural stem cells (NSCs) play a fundamental role in neural development and repair processes. NSC characteristics and fate are intricately regulated by the microenvironment and intracellular signaling. Interestingly, metabolism plays a pivotal role in orchestrating the epigenome dynamics during neural differentiation, facilitating the transition from undifferentiated NSC to specialized neuronal and glial cell types. This intricate interplay between metabolism and the epigenome is essential for precisely regulating gene expression patterns and ensuring proper neural development. This review highlights the mechanisms behind metabolic regulation of NSC fate and their connections with epigenetic regulation to shape transcriptional programs of stemness and neural differentiation. A comprehensive understanding of these molecular gears appears fundamental for translational applications in regenerative medicine and personalized therapies for neurological conditions.

Keywords Neural stem cells · Neurogenesis · Energy metabolism · Chromatin · Epigenetics · Transcriptional regulation

Spotlights

- Neural stem cells rely on anaerobic glycolysis for energy. They maintain low reactive oxygen species levels and have immature or non-functional mitochondria.
- As neural stem cells differentiate, specific metabolic shifts determine cell fate. Terminally differentiated neurons rely on oxidative phosphorylation for energy.
- Oxygen levels play a crucial role in neural stem cell proliferation and differentiation, with hypoxia influencing the balance between self-renewal and differentiation.
- Specific modifications on chromatin conform to an epigenetic regulatory landscape shaping transcriptional transitions in neural development.
- Fluctuating metabolites during neural differentiation drive enzymatic reactions for epigenetic modifications and activate transcription factors, crucial for proper neuronal differentiation.

✉ Lorena Aguilar-Arnal
loreaguilararnal@iibiomedicas.unam.mx

¹ Departamento de Biología Celular y Fisiología, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Mexico City, Mexico

² Laboratory for Optics and Biosciences, Ecole Polytechnique, CNRS, INSERM, Institut Polytechnique de Paris, Palaiseau, France

Introduction

Stem cells (SCs) are undifferentiated cells that possess two unique features: self-renewal, which refers to the ability to divide and proliferate, generating identical cells while maintaining the characteristics of stemness, and potentiality, which is the ability to differentiate into different types of functional cells from daughter cells that have lost their stemness during the differentiation process [1]. Stem cells can be classified according to their differentiation capacity into totipotent, pluripotent, multipotent, and unipotent stem cells [2]. Pluripotent stem cells have the capacity to differentiate into all three germ layers, multipotent stem cells differentiate into a range of cell types within a specific tissue; and unipotent stem cells have the potential to produce only one cell type. Additionally, totipotent stem cells can be obtained in the early stages of embryonic development when the zygote divides to form the morula. These cells can differentiate into any type of embryonic and extra-embryonic tissue, and even generate a complete organism [2]. As embryonic development progresses, the self-renewal and differentiation potential of stem cells changes [3]. For example, embryonic stem cells (ESCs) are isolated from the inner mass of the blastocyst, which

forms between days 5–6 of human embryonic development. These cells have pluripotent characteristics and can generate the three embryonic germ layers (mesoderm, ectoderm, and endoderm) but not extra-embryonic tissues [4]. Yet, stem cells are not limited to the embryonic stage and can also be found in adult tissues such as bone marrow, brain, and skin, which are referred to as adult stem cells. Neural stem cells (NSCs) are an example of adult stem cells, which reside in neurogenic niches within the adult brain [5].

NSCs are derived from the ectoderm during gastrulation and are a type of multipotent stem cell that generates different lineages of neurons, astrocytes, and oligodendrocytes during embryonic development. The central nervous system is formed from NSC lining the neural tube [6]. Subsequently, NSCs are “inactivated” at the end of the nervous system development, giving rise to quiescent NSC (NSCq). Thus, NSCq will be in a relatively inactive state until specific signals trigger their activation to participate in various physiological or pathophysiological functions in the brain, for example, neuronal plasticity, or repair of brain lesions [7]. The behavior of neural stem cells (NSCs), including their equilibrium between quiescent and active states, proliferation, and differentiation, is governed by a multitude of stimuli present in the NSC microenvironment, both during embryonic development and in the adult organism. For instance, growth factors such as fibroblast growth factor 2 (FGF-2), epidermal growth factor (EGF), or bone morphogenetic protein (BMPs) [8, 9] have demonstrated pivotal roles in regulating NSC proliferation and differentiation in embryonic and adult brains. Furthermore, extracellular matrix (ECM) proteins like laminin and fibronectin play a crucial role by offering structural support and modulating cell signaling [10]. In addition to ECM proteins, neurotransmitters such as dopamine and serotonin, which are known to influence the proliferation and differentiation of neural stem cells in the adult brain, contribute to this intricate regulatory network [11, 12]. The sophisticated interplay among these diverse factors within the NSC microenvironment significantly impacts NSC behavior, ultimately shaping neural development and repair processes [13, 14].

During development, NSCs initially expand by symmetric self-renewing divisions, in which NSC divides to produce two identical daughter cells, both of which retain their stem cell characteristics. Later, NSCs can divide via asymmetric division, originating either another NSC through self-renewing or neural progenitor cells (NPC) [15]. This population of cells is defined by their ability to proliferate through symmetric self-renewal divisions in a controlled manner for a limited period before expressing a differentiated phenotype [6]. The number of neurons generated during neurogenesis is determined by the initial number of neural stem/progenitor cells (NS/PC), the duration of their proliferative period, and

their lineage. NSCs can generate all cell types of the brain, whereas NPC has a more restricted potential [16].

Differentiation of stem cells into specific cell types involves the expression of lineage-specific genes, inhibition of factors inducing differentiation into other lineages, and expression of factors promoting differentiation into a specific lineage [17]. Differentiation is regulated by extracellular signals, such as cytokine signaling, and intracellular programs such as epigenetic regulation, which involves DNA methylation, chromatin and histone modifications, functional non-coding RNAs, dynamic chromatin long-range interactions, and distinct chromatin spatial conformations, among other mechanisms [18, 19]. NS/PC must integrate cell-intrinsic programs and environmental signals in time and space to promote correct development [20], assisted by coordinated and subsequent gene expression programs regulated by transcription factors and the dynamic epigenome. Moreover, stem cell metabolism also plays a role in determining cell fate. Recent studies of stem cell metabolism have revealed coordinated fluctuations in energy metabolism and oxidative stress during NS/PC maturation and differentiation [21–23]. For example, the differentiation of NSCs into neurons shows significant changes in cell morphology and metabolism, such as the glycolytic activity which progressively decreases [24]. Interestingly, NSCs remain quiescent in a hypoxic niche with low O_2 tension and obtain their energy by anaerobic glycolysis [25]. This glycolytic metabolism is substantially less energy efficient, suggesting that anaerobic glycolysis is not merely an environmental adaptation but an intrinsic necessity for quiescent adult stem cells. However, the differentiation of NPC to neurons is accompanied by a metabolic reprogramming, where the main source of energy will switch to oxidative phosphorylation (OXPHOS) [21, 23]. Remarkably, the metabolic state of stem cells can influence their epigenetic regulation. During differentiation, transitions from a quiescent state to a more active state implicate that metabolic preferences change. This metabolic shift, in turn, affects the availability of metabolites and cofactors necessary for epigenetic modifications, including DNA methylation and histone modifications [26]. For example, the acetyl-CoA produced during glycolysis is a critical substrate for histone acetylation, a key epigenetic modification that can activate gene expression. Changes in glycolytic flux can alter the levels of acetyl-CoA and, in turn, influence the epigenetic landscape to coordinate specific gene expression patterns.

Here, we review how certain metabolites playing an important role in energy and intermediary metabolism fluctuate during neural differentiation, and coordinately promote enzymatic reactions leading to epigenetic modifications or the activation of transcription factors that regulate specific transcriptional programs necessary for proper neuronal differentiation. We structured this review in two major sections,

the first one oriented to outline metabolic transitions during NSC differentiation, and a second section describing advances on how these fluctuations in certain metabolites directly regulate specific epigenetic mechanisms contributing to the dynamic transcriptional reprogramming. Finally, we describe technological approaches which could contribute to further expanding our knowledge in this molecular interplay. By delving deeper into these mechanisms, we can gain a greater understanding of how to harness neural differentiation for cell-based therapies and potentially unlock new avenues for the treatment of debilitating conditions affecting the brain and nervous system.

Metabolic Fluxes Define Neural Stem Cell Fate

Neuronal differentiation is a tightly controlled process through multiple regulatory layers. During the past few years, metabolic transitions are emerging as direct regulators of neural differentiation [27]. Accordingly, energy-regulating pathways are highly dynamic, showing flexibility to adapt in response to environmental factors including developmental signals [28]. Glycolysis in the cytoplasm and the tricarboxylic acid (TCA) cycle inside the mitochondria are coupled by pyruvate, through the pyruvate dehydrogenase complex, and their activity constitutes a major source of metabolic precursors and energy fluxes (Fig. 1) [29]. In fact, most terminally differentiated cells rely on the coordinated action between the TCA cycle and mitochondrial OXPHOS to meet energy demands. In this regard, dehydrogenases in the TCA cycle transfer electrons to the reducing equivalents nicotinamide adenine dinucleotide, NAD(H) and flavine adenine dinucleotide, FAD(H₂), which in turn carry them into the electron transport chain to generate an electrochemical proton gradient across the inner mitochondrial membrane used by the ATP synthase to produce ATP [30]. Because the OXPHOS process consumes oxygen, in hypoxic conditions, mitochondrial function is inhibited and glucose is metabolized in the cytoplasm to lactate by lactate dehydrogenase (LDH), in a process known as anaerobic glycolysis. Surprisingly, many stem cell types rely on lactate production for their metabolic needs even in the presence of oxygen, a phenomenon known as the Warburg effect or aerobic glycolysis, first identified in cancer cells [31, 32]. Indeed, the incomplete oxidation of glucose to lactate permits a less efficient but faster rate of ATP production, but it also allows for the conservation of glycolytic intermediates which serve as substrates for a number of metabolic pathways whose fluxes are essential to meet the biosynthetic demands in highly proliferative cells, such as stem cells. For example, channeling glucose-6-phosphate, fructose-6-phosphate, and fructose-1,6-bisphosphate through the pentose phosphate pathway

(PPP) generates NADPH and ribose-5-phosphate, which is necessary to sustain high rates of nucleic acid biosynthesis in proliferating cells [33]. Also, 3-phosphoglycerate can be shunted into the serine biosynthesis pathway, leading to the generation of serine, which is a precursor for proteins and participates in phospholipid biosynthesis. Definitively, it is clear that metabolic transitions occur during neural differentiation to support specific metabolic requirements at each stage.

Metabolic Reprogramming in Neural Differentiation: From Glycolysis to OXPHOS


During neurogenesis, a switch from glycolytic to oxidative metabolic phenotype has been well characterized [27] (see Table 1). Evidence from a *Xenopus* model of retinal differentiation demonstrated that proliferating stages are highly dependent on glycolysis *in vivo*, with cells presenting higher levels of intracellular lactate and lactate dehydrogenase (LDH) and lower oxygen consumption rates than these cells at terminally differentiated stages (Fig. 1A, B) [34]. Importantly, a perturbation of this metabolic balance leads to impaired development of the retina. Similarly, in *Drosophila*, inhibiting OXPHOS in NSC impedes neurogenesis during development [35], van den [36]. Concomitantly, in human NPC, a transcriptional rewiring of metabolic genes accompanies and contributes to the metabolic switch during neurogenesis [23]. For example, downregulation of the glycolytic limiting enzymes hexokinase (HK2) and lactate dehydrogenase (LDHA) expression both at the mRNA and protein levels assists in transitioning from aerobic glycolysis in NPC to neuronal oxidative phosphorylation [23, 37, 38]. Alongside, hindering glycolysis through deletion of HK2 disrupts neural progenitor proliferation [37, 38].

As expected, increased OXPHOS during neurogenesis is tightly linked to mitochondrial function, and it has been postulated that mitochondrial dynamics regulate neural differentiation (Fig. 1). For example, during embryonic development in mice, NSC depicts elongated morphology, which becomes fragmented in committed neural progenitors, and regains an elongated shape in post-mitotic neurons. Disturbing these dynamics hinders stem cell self-renewal and neurogenesis, likely through dysfunctional metabolic fluxes [39, 40].

In human NSC, neural differentiation is accompanied by increased gene expression programs controlling mitochondrial biogenesis, including the peroxisome proliferator-activated receptor gamma coactivator 1-alpha (*PGC-1α*) and its downstream targets, *POLG*, *POLRMT*, *ERRα*, and *NRF1*, which is paralleled by decreased glycolytic flux (Fig. 1B) [41]. Accordingly, mitochondrial dysfunction as a result of a genetic or neurodegenerative disease drives cognitive decline, at least in part, through loss of adult NSC and neurogenesis, and leads to neuronal death [42–46]. Notably,

A

Neural stem /progenitor cell (NSPC)



B

Terminally differentiated neural cell




Fig. 1 A Metabolism in neural stem and progenitor cells (NS/PC). In a hypoxic niche, neural stem cells (NSCs) maintain a quiescent state and predominantly rely on anaerobic glycolysis for energy production. These cells actively regulate reactive oxygen species (ROS) levels and exhibit immature or non-functional mitochondria. Neural progenitors (NPCs) utilize distinct metabolic pathways, including the pentose phosphate pathway and the folate 1C cycle, to support their proliferation. Furthermore, NPCs facilitate the synthesis of phospholipid membranes in collaboration with cholesterol biosynthesis, accompanied by elevated levels of lipid droplets essential for sustaining stem cell properties. **B** Metabolism of terminally differentiated neural cells. A notable metabolic reprogramming is evident, characterized by increased activity of glycolytic enzymes, such as hexokinase 1 and pyruvate kinase isoform 1, alongside reduced expression of hexokinase 2 and lactate dehydrogenase A. Simultaneously, pyruvate undergoes efficient conversion to acetyl-CoA for entry into the TCA cycle, where oxidative phosphorylation becomes the primary source of ATP. The upregulation of mitochondrial biogenesis, orchestrated by PGC-1 α /ERR α , further accentuates this shift. Concurrently, lipid metabolism, encompassing eicosanoids and fatty acid oxidation, assumes a pivotal role in elevating acetyl-CoA levels, fueling the TCA cycle and augmenting NADH and FADH₂ production critical for establishing the proton gradient driving ATP synthase. Concomitantly, elevated mitochondrial activity contributes to heightened ROS levels, playing a crucial role in determining cell fate. Abbreviations: glucose 6-phosphate (G6P); fructose 6-phosphate (F6P); fructose 1,6-bisphosphate (FBP); S-adenosylmethionine (SAM); acetyl-CoA (AcCoA); fatty acid (FA), reduced nicotinamide adenine dinucleotide phosphate (NADPH); polyunsaturated fatty acids (PUFAs); peroxisome proliferator-activated receptor gamma (PPAR γ); hypoxia-inducible factor 1 (HIF-1); mammalian target of rapamycin complex 1 (mTORC1); pentose phosphate pathway (PPP); reactive oxygen species (ROS); phosphoenolpyruvic acid (PEP); nicotinamide adenine dinucleotide (NAD⁺); reduced nicotinamide adenine dinucleotide (NADH); reduced flavin adenine dinucleotide (FADH₂); fatty acid oxidation (FAO); oxidative phosphorylation (OxPhos); adenosine triphosphate (ATP); tricarboxylic acid cycle (TCA cycle); peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC-1 α); estrogen-related receptor alpha (ERR α). Enzymes: hexokinase 1 (HK1); pyruvate kinase isozymes M1 (PKM1); lactate dehydrogenase A (LDHA); pyruvate dehydrogenase (PDH); pyruvate dehydrogenase kinase (PDK), carnitine palmitoyltransferase 1 (CPT1); acetyl-CoA carboxylase (ACC). Parts of the figure were drawn by using pictures from Servier Medical Art by Servier, licensed under a Creative Commons Attribution 3.0 Unported License (<https://creativecommons.org/licenses/by/3.0/>)




enhancing mitochondrial function, for example overexpressing key proteins such as the inner membrane GTPase OPA1, or the manganese superoxide dismutase SOD2, has been proven an effective strategy opposing neuronal decline in mice [47–50]. This evidence indicates that a tight balance between metabolic fluxes and mitochondrial function and dynamics controls NSC maintenance and neurogenesis.

Increased OXPHOS to provide ATP during differentiation consumes oxygen and generates high levels of reactive oxygen species (ROS). It is well established that ROS production has a profound impact on neural differentiation. For example, low ROS levels below 1 nM H₂O₂ contribute to maintaining a quiescent state in stem cells, while increasing ROS promotes self-renewal and differentiation [51].

However, above 10 nM H₂O₂, oxidative stress increases leading to cell death and neural tissue degeneration. Physiological fluctuations in ROS levels contribute to neurogenesis, maturation, and synaptic plasticity, although the mechanisms are not fully understood [52, 53]. Notably, mitochondria serve as the primary source of ROS, influencing mitochondrial dynamics and potentially leading to the formation of granular mitochondria [54]. Concurrently, changes in mitochondrial shape can impact mitochondrial function and redox status, amplifying ROS production and oxidative stress [55]. Mitochondrially derived ROS function in redox signaling, but in excess, they contribute to cellular injury and death. Importantly, the accumulation of mitochondrial ROS affects neural cell fate by inducing oxidative stress and DNA damage and activating cell death pathways, while alterations in mitochondrial dynamics influence cellular bioenergetics, differentiation, and fate determination [56, 57]. This evidence underscores the intertwined roles of mitochondrial dynamics and ROS in shaping neural cell fate, while the implicated mechanisms need further exploration.

It is generally assumed that stem cells reside in hypoxic niches which might contribute to sustaining glycolytic metabolism; however, evidence for low oxygen tension in adult NSC niches is scarce [58]. Notably, adult NSCs in the brain, such as those in the subventricular zone (SVZ) and the dentate gyrus of the hippocampus, exist in a more oxygenated environment compared to embryonic NSCs. Local oxygen levels in the neurogenic niches where adult NSCs reside can influence their behavior. For example, normoxia can stimulate the activation of adult NSCs and their differentiation into neurons or glial cells. The glycolytic phenotype of NSC and NPC, characterized by a preference for glycolysis over oxidative phosphorylation to promote energy-efficient maintenance of their quiescent state (NSC) or for energy production (NPC), renders them highly resistant to hypoxic conditions [59]. Interestingly, numerous data support that low oxygen tension increases self-renewal and proliferation of NSC in vitro, and defines cell fate [60–65]. Several molecular mechanisms are implicated. For example, progressive relief of hypoxia during the development of the mouse cerebral cortex shapes NSC differentiation in vivo, a mechanism involving a gradual decline in hypoxia-inducible factor (HIF)-1 α signaling [66]. Importantly, the transcription factor HIF-1 α is stabilized under hypoxic conditions, at oxygen levels below approximately 6 to 2% oxygen (or 60–20 mm Hg) [67], and controls a transcriptional program of genes which ultimately promote glycolysis and repress OXPHOS [68]. HIF-1 α is essential for NSC maintenance and contributes to adult neurogenesis [69, 70], and also influences important pathways such as Wnt/ β -catenin and Notch function, which fine tune NSC proliferation, differentiation, and neuronal maturation (Fig. 1A) [71–74]. HIF-1 also promotes overexpression of its downstream target *BNIP3*,

Table 1 Metabolic transitions from neural stem cells to differentiated neural cells

Metabolic aspect	Neural Stem Cells		Differentiated cells (Neurons/ Glia)
	Proliferation	Differentiation	
Energy source	Glycolysis	OXPHOS	OXPHOS
Oxygen utilization			
Glycolysis rate			
Mitochondrial activity			
Glucose uptake	Consistently high	Variable	Variable
Biosynthesis	Nucleotides and amino acids	Lipids and proteins	Lipids and proteins
Redox state	NAD ⁺ /NADH shifted towards NADH	NAD ⁺ /NADH shifted towards NAD ⁺	NAD ⁺ /NADH shifted towards NAD ⁺
ROS production	Low	Moderate	Variable
Serine metabolism	biosynthesis of nucleotides and amino acids	lipid synthesis and redox maintenance	lipid synthesis and redox maintenance
Pentose Phosphate Pathway (PPP)	Active. Contributes to nucleotide and NADPH production	Variable activity	Variable activity
Lipogenesis	Increased fatty acids, cholesterol, sphingolipids and phospholipids for membrane synthesis		Specialized: myelin in glia or synaptic lipids in neurons

which in turn inhibits mTORC1 activity, hereby increasing autophagy which appears necessary for NSC proliferation [75]. Another advantage of hypoxia-induced glycolysis in NSC is the decreased ROS production as compared to terminally differentiated neurons relying on OXPHOS. In fact, raising ROS during neural differentiation constitutes a signaling event influencing NSC fate decision. For example, ROS increase the activity of the transcription factor NRF2, a master regulator of the antioxidant response, which also targets the expression of pro-neuronal factors such as Isl1, Nkx2.1, Lhx5, and Sim1 [40]. Along these lines, maintaining low ROS levels in NPC is also influenced by the forkhead box O (FoxO) family of transcription factors, which tightly control metabolic fluxes and sustain redox balance

in these cells [76, 77], and autophagy rates in developing and adult neurons [78]. Recent research reveals a notable divergence in energy metabolism among neurons based on subcellular structures. Specifically, somata exhibit elevated aerobic glycolysis and reduced OXPHOS compared to terminals [79]. Further investigations are needed to clarify the physiological significance of this metabolic compartmentalization and the underlying mechanisms.

In summary, cumulative evidence suggests that a shift in metabolic pathways from glycolysis to OXPHOS takes place during neural differentiation. The relationship between oxygen levels and metabolic shifts during NSC differentiation and proliferation is bidirectional: low oxygen (hypoxia) promotes glycolysis and proliferation by activating HIF-1 α

to maintain stemness and support rapid cell division, while higher oxygen levels (normoxia) are associated with the transition to OXPHOS, which supports differentiation, and the increased energy demands of terminally differentiated neural cells. These transitions are meticulously regulated and synchronize with gene expression programs and intracellular signaling. Collectively, these mechanisms sustain the self-renewal and proliferative potential of NSC and play a role in determining cell fate and shaping neurogenesis and homeostasis in adult neurons.

Fatty Acid Metabolism in Neural Stem Cells

Besides glucose fluxes, fatty acid metabolism plays a crucial role in the biology of NSC, regulating proliferation, differentiation, and self-renewal. The eicosanoid pathway and fatty acid synthesis, as well as fatty acid oxidation (FAO), contribute to neurogenesis at distinct stages [80]. In neural cells, phospholipids such as phosphatidylcholine, phosphatidylethanolamine, and phosphatidylserine are essential for membrane structure and function [81]. Glial cells require sphingomyelin and glycosphingolipids, particularly for constituting myelin sheaths. Gangliosides, a type of glycosphingolipid with sialic acid residues, are essential for neuronal development, signaling, and synaptic function [82]. Additionally, lipid droplets, the lipid-storing organelles, are highly abundant in mouse NPC and decay during differentiation [83]. Increased lipid droplets promote proliferative and differentiation capacities and fuel FAO to sustain NPC survival [83]. It is generally accepted that the brain mostly relies on glucose as a source of energy over lipids and FAO, hereby avoiding ROS production and oxidative damage [84]. However, FAO is now emerging as a critical regulator of neural development [85]. For example, quiescent mouse NPC expresses high levels of the FAO rate-limiting enzyme carnitine palmitoyltransferase I (CPT1), and FAO is more active than in proliferative NPC and mature neurons. Inhibiting CPT1 leads to defective neurogenesis and cell death, while stimulating it alters NPC proliferation (Fig. 1) [86, 87]. Additionally, FAO is a critical regulator of the transition from mouse NSC to intermediate progenitor cells during brain development, which may underlie defective cell division leading to autism spectrum disorder (ASD) [88]. In fact, fatty acids serve as energy sources, but can also be incorporated into various lipid molecules such as long-chain polyunsaturated fatty acids like docosahexaenoic acid (DHA), which is particularly important for neural development and function. DHA is an omega-3 fatty acid and is the major prevalent fatty acid in the brain membrane [89]. The brain maintains its fatty acid levels mainly via the uptake of plasma-free DHA, and its metabolization to N-docosahexaenoylethanolamide leads to neurite growth, synaptogenesis,

and synaptic function, which is relevant for learning and memory [90].

Aberrations in lipid metabolism can have significant consequences for neural development, homeostasis, and function, contributing to various neurological disorders. For example, abnormal oleic acid-enriched triglyceride accumulation within ependymal cells, the main support cells of the forebrain NSC niche, during brain development might promote Alzheimer's disease (AD) in mice [91], underscoring the relevance of lipid metabolism in neural development.

On the other hand, *de novo* lipogenesis is catalyzed by the fatty acid synthase (FASN) enzyme, which is elevated in mouse NSC when compared to mature neurons [92]. FASN activity is required for normal NPC proliferation and neurogenesis [92–94]. In mice and humans, a *FASN* R1819W mutation impairs adult hippocampal NSPC activity and cognitive defects due to lipid accumulation and lipogenic ER stress in these cells [93, 94], reinforcing the idea that lipid metabolism contributes to cognitive function [95].

Certain lipids appear to be involved in neurogenesis (Fig. 1). For example, oleic acid is an endogenous ligand for the nuclear receptor NR2E1/TLX, which controls NSPC self-renewal and proliferation. When bound to oleic acid, TLX becomes a transcriptional activator of neurogenic genes, driving hippocampal development in mice [96]. Along the same lines, omega-3 polyunsaturated fatty acids (n-3 PUFAs) are emerging as neuroprotective compounds [97], and recent research shows that in a human hippocampal progenitor cell line, they exert a transcriptional response for pathways involved in oxidative stress and inflammation [98]. Importantly, the nuclear receptor peroxisome proliferator-activated receptor alpha (PPAR α) in the brain binds a class of omega-3 hydroxy fatty acid promoting multiple dendritic responses in mouse cortical neurons, which might result in neuroprotective effects [99].

Lastly, cholesterol metabolism plays a central role in the central nervous system. It is considered that the brain contains about 25% of the whole body's cholesterol [100]. It contributes to maintaining membrane fluidity and modulating the function of membrane proteins and, importantly, is involved in myelin formation [101]. Because cholesterol is a key component of membranes in mammalian cells, neural functions are tightly linked to cholesterol bioavailability, including the formation of synapses [102]. Several diseases are associated with defects in cholesterol metabolic pathways, such as Alzheimer's disease, Huntington's disease, and Parkinson's disease [103]. Interestingly, selective ablation of cholesterol biosynthesis in NSC in the adult brain is accompanied by increased VEGF expression, which promotes angiogenesis to supply the niche with circulating cholesterol-loaded lipoproteins [104]. Yet, in the developing

brain, altering cholesterol biosynthesis leads to accelerated neurogenesis and abnormal migration and proliferation patterns of cerebellar granule precursors [105, 106].

Folate-Dependent One-Carbon Metabolism Is Critical to the Developing Brain

Mounting research highlights an essential role for folate-dependent one-carbon (1C) metabolism in the developing brain [107, 108]. 1C metabolism consists of the folate cycle, the methionine remethylation, and the transsulfuration pathways, which are essential for amino acid metabolism, purines and thymidine biosynthesis, and synthesis of S-adenosylmethionine (SAM), the universal methyl donor for methylation reactions [107–109]. A strong association between folate-1C metabolism deficiencies and defects in neural tube development has been established, which generally leads to adverse pregnancy outcomes [110]. In this regard, many GWAS studies have identified certain SNPs in genes encoding rate-limiting enzymes in 1C metabolism, such as the methylenetetrahydrofolate reductase (*MTHFR*) or betaine-homocysteine S-methyltransferase 1 (*BHMT1*), associated with increased risk of neural tube defects during development [111], Rampersaud, [112, 113]. Mounting research has emphasized the significance of folate in the self-renewal and differentiation of neural stem cells (NSCs). Inadequate levels of folate have been found to impede the growth of NSCs in the adult hippocampus in vivo and prompt NSC cell death in vitro [114, 115]. Accordingly, NSCs exhibit increased proliferation and neuronal differentiation in response to folate [116, 117].

Epigenetic Regulation and Metabolic Fluxes Intersect to Shape Neurogenesis

As related, neural fate commitment and differentiation are typically accompanied by significant transitions in metabolic activity, as specific pathways assume control over the supply of fuel. These metabolic transitions are accompanied by extensive transcriptional rewiring of gene regulatory networks to sustain the differentiation process [118, 119]. Indeed, transcription occurs in the chromatin fiber, hereby, chromatin states are key controlling neural development. As a result, distinct modifications on chromatin conform to an epigenetic regulatory landscape shaping transcriptional transitions in neural development [120].

During neural commitment, ESCs lose their pluripotency through activating epigenetic mechanisms leading to chromatin transitions in regulatory regions of multiple genes. Extrinsic signaling promotes the expression of proneural genes such as *Sox1*, *Pax6*, and *Ngn1* through chromatin remodeling at their regulatory elements, while

genes activated only in terminally differentiated cells remain poised until their cell fate is defined [121, 122]. While technological advances in high-throughput sequencing have contributed to defining these transcriptional and chromatin transitions with high precision and even to a single-cell level, the interplay between metabolic and epigenetic reprogramming and how it evolves with time during neurogenesis is emerging.

Metabolism and epigenetics intersect at the level of metabolites that are used as substrates and cofactors for enzymes catalyzing reactions to directly modify chromatin and hereby regulate gene expression [123]. Certain metabolites such as acetyl coenzyme A, SAM, NAD⁺, FAD, or even glucose and lactate are necessary for the activity of chromatin modifiers that define the epigenetic landscape [124]. Since the availability of these metabolites relies on the intermediary metabolism, the wide metabolic fluctuations observed during neural differentiation are thought to impact the epigenetic landscape.

In the next sub-sections, we describe current knowledge of this direct interplay between metabolic and epigenetic transitions to control gene expression in neurogenesis. Of note, a comprehensive view of epigenetic and epigenomic reprogramming in neurogenesis is not the focus of this review and has been extensively reviewed elsewhere [125, 126], which we recommend to the reader aiming to get a complete understanding on this specific topic.

SAM-Directed DNA Methylation Contributes to Neural Development

DNA methylation is a pivotal process that plays a crucial role in development through controlling gene expression, genomic imprinting, or chromatin structure. It involves the transfer of the methyl group from the metabolite SAM to a cytosine by DNA methyltransferase enzymes (DNMTs), forming 5-methylcytosine (5mC). Moreover, there is a distinct form of DNA methylation known as 5-hydroxymethylcytosine (5hmC) that occurs frequently in the mouse brain. During embryonic development, de novo DNMTs DNMT3A and DNMT3B establish the DNA methylation pattern, while DNMT1 maintains this pattern. DNA methylation can be removed passively by blocking methylation of newly synthesized DNA during replication or through enzymatic reactions that remove the methyl-modifications [127] such as the ten-eleven translocation (TET) family [128]. DNA methylation regulates gene expression via preventing transcription factor binding and silencing genes through recognition by transcriptional repressors with a methyl-CpG-binding domain (MBD) such as MBD1 and methyl-CpG binding protein 2 (MeCP2)[129]. Importantly, extensive methylation reconfiguration occurs during mammalian brain

development, which is cell-type specific and modulates the function of regulatory regions in the genome [130]. For example, during embryonic neural development, the DNA methylation status of genes such as *Pax6*, which codes for a transcription factor, can determine cell fate [131]. Methylation of the *Pax6* gene regulatory elements regulates its activity and promotes neurogenesis from ESC [132]. Recent literature supports the idea that metabolites such as folate and SAM can regulate DNA methylation in neural development. For example, during axonal myelination in mice, global DNA demethylation occurs, which is associated with the activation of genes such as *Srebf1*, *Hmgcr*, *Dgat1*, *Alc27a1*, and *Abca2*, which are crucial to activate lipid metabolism required for myelination process [133]. Interestingly, SAM levels appear reduced during Schwann cell myelination, and increased SAM leads to abnormal hypermethylation and peripheral nerve defects [133].

The relationship between folic acid metabolism and neural development has been largely known; however, the direct impact of 1C metabolism on the DNA methylation dynamics in neurogenesis is still a matter of debate [134, 135]. In humans, evidence of a direct impact of 1C metabolism intermediaries in controlling DNA methylation during neurogenesis is scarce. A remarkable study was conducted on the *POMC* gene which encodes several peptides involved in the appetite control in the hypothalamus [136]. *POMC* methylation status in the brain from post-mortem subjects was positively correlated with individual weight. Interestingly, lower *POMC* methylation was found in children conceived from mothers with a lower supply of 1C metabolites, while a robust negative correlation for SAH and positive correlations with betaine and SAM were found at several *POMC* promoter methylation sites [137]. Notably, the availability and dynamics of 1C metabolism intermediaries have been implicated in the development of a number of neurodegenerative conditions such as Alzheimer's disease, Huntington's disease, and autistic spectrum disorders [138–140]. In rodents, mounting evidence sustains a direct relationship, for example, in NSCs, folate treatment leads to increased proliferation while inhibition of DNMT1 counteracts it [141, 142]. This is paralleled by differential methylation of a set of genes involved in critical pathways such as neuroactive ligand–receptor interaction, Jak-STAT signaling pathway, steroid biosynthesis, fatty acid elongation, PI3K/Akt, MAPK signaling pathway, and cytokine–cytokine receptor interaction [141]. Furthermore, a low folate diet alters the 5mC and 5hmC epigenetic patterns in mESC, altering the expression of neural development genes pertaining to the hedgehog signaling pathway which is involved in the formation of the neural tube [143]. Concomitantly, severe maternal undernutrition leads to reduced proliferation and differentiation in NSC, with

decreased levels of DNMT1 and altered expression of neural markers such as *Nestin* or *Hes1* [144]. Along the same lines, increased levels of S-adenosylhomocysteine (SAH), a bypass metabolite from SAM consumption during methylation reactions, inhibit methyltransferases, blocking proliferation in NPCs (Fig. 2A) [145]. Increased SAH leads to hypomethylation at the *Sprouty2* (*Spry2*) promoter and its overexpression, which negatively regulates the fibroblast growth factor receptor (FGFR)-Erk1/2-cyclin E signaling pathway and, hereby, impairs proliferative signaling [146].


Metabolic Control of Histone Methylation During Neural Differentiation

Similar to DNA, methylation of histones consists of the transfer of a methyl group from SAM to a lysine or arginine residue of a histone protein, by histone methyltransferase (HMT) enzymes. Pioneering studies within the past decade demonstrated that SAM accumulation in mouse embryonic stem cells (mESC) selectively influences trimethylation of histone H3 lysine 4 (H3K4me3) levels, a transcriptionally activating mark involved in the maintenance of pluripotency [147]. Similarly, methionine-deprived human pluripotent stem cells (hPSC) also show decreased SAM and H3K4me3, which is accompanied by impaired self-renewal [148]. Interestingly, when induced to neural differentiation, methionine-deprived hPSC express higher levels of the neural marker *MAP2*, suggesting increased differentiation potency [148]. This evidence supports the idea that the metabolome regulates the epigenetic landscape in multipotent cells and during differentiation. In addition, at very early stages in human development, the repressive H3K27me3 and H3K9me3 marks are responsive to variations in SAM levels (Fig. 2B) [149].

Interesting research has recently demonstrated a link between one-carbon metabolism and histone methylation which determines neural differentiation. Using NSC, the authors demonstrated that activation of the ephrin receptor Eph-B induced downregulation of the transcript *Dhfr*, encoding a key enzyme in the 1C metabolic pathway [150]. Decreased DHFR expression and activity depletes the progenitor pool and promotes differentiation through an epigenetic switch consisting of decreased H3K4me3 levels on progenitor-specific genes such as *Sox2* and *Nestin* [150]. Together, these data indicate that the modulation of metabolic pathways influences stem cell fate through epigenetic modulation.

On the other hand, histone demethylation can be catalyzed by the JMJC (Jumonji-C) family of lysine demethylases (KDM). Among them, KDM5A, KDM6A, and KDM3A have been described as effective oxygen sensors [151–153]. KDM5A contributes to the maintenance of NPC by suppressing genes related to astrocyte

A
Neural stem /progenitor cell
(NSPC)



B
Terminally differentiated
neural cell




Fig. 2 The interplay between metabolism and the epigenome in neural stem/progenitor cells and terminally differentiated neural cells. **A** Metabolism plays a crucial role in establishing and maintaining the epigenetic state that underlies the neural stem/progenitor cell phenotype. Enzyme-mediated post-translational modifications of DNA or histones relying on small metabolites as cofactors contribute to this process. Repressive DNA marks, such as 5mC, catalyzed by DNMTs, and histone methylation marks H3K9me3 and H3K27me3, catalyzed by SAM-dependent KMTs, silence pluripotency genes. Conversely, the presence of H3K9ac, catalyzed by acetyl-CoA-dependent HATs, activates certain genes. The maintenance of the multipotent state is favored by FAD-dependent LSD1 and the KDM5A protein. **B** In terminally differentiated cells, metabolism, particularly oxidative metabolism, promotes an acetyl-CoA pool, which facilitates the activity of HAT, CBP, and p300, leading to an active transcriptional state for differentiation genes. TIGAR inhibits glycolysis, whereas NaBt promotes mitochondrial biogenesis. Multipotent genes are transcriptionally repressed by DNA and histone repressive marks. In addition, the NAD⁺-dependent deacetylase SIRT1 targets H3K9ac, promoting transcriptional repression at specific genes which determine cell fate. Abbreviations: NAD⁺, nicotinamide adenine dinucleotide; acetyl-CoA, acetyl coenzyme A; SAM, S-adenosylmethionine; SAH, S-adenosyl-L-homocysteine; FAD, flavin adenine dinucleotide; HAT, histone acetyltransferases; DNMTs, DNA methyltransferases; KMT, lysine methyltransferase; LSD1, lysine-specific demethylase 1; 5mC, 5-methylcytosine; Me, methyl group; SIRT1, NAD-dependent deacetylase sirtuin-1; HDAC, histone deacetylases; ACSS2, acetyl-CoA synthetase 2; ACL, ATP-citrate synthase; TIGAR, TP53-inducible glycolysis and apoptosis regulator; NaBt, sodium butyrate; CREB binding protein (CBP); p300, histone acetyltransferase p300; Ac, acetyl group. Parts of the figure were drawn by using pictures from Servier Medical Art by Servier, licensed under a Creative Commons Attribution 3.0 Unported License (<https://creativecommons.org/licenses/by/3.0/>)

differentiation [154]. KDM6A participates in resolving bivalent promoters, a particular class of chromatin enriched in both H3K4me3 and H3K27me3, during human neural differentiation [155], and activates specific neurogenic genes [156]. Also, KDM3A facilitates recruitment to chromatin of the transcription factor Neurog2, which is essential for neurodevelopment [157]. Hereby, it is tempting to speculate that the hypoxia signaling might coordinate dynamic changes in the epigenome shaping the neurogenic program. Further research is necessary to clarify this possibility.

Another kind of histone demethylases is the flavin adenine dinucleotide (FAD)-dependent lysine-specific histone demethylase (LSD) proteins. LSD1 removes mono- or di-methylation on lysine 4 of histone H3 (H3K4me1/me2) and also demethylates H3K9 or H4K20 [158–160]. In rodents, LSD1 supports the maintenance of NS/PC identity and proliferation, and a progressive decrease in LSD1 levels is required for neural differentiation [158, 161]. Interestingly, the pharmacological increase of FAD activates LSD1 in NSC, which is sufficient to induce neuronal differentiation [162]. Hereby, it is possible that histone demethylation, regulated by FAD-dependent LSD1 activation, may contribute to the regulation of neural differentiation coordinated by metabolic transitions (Fig. 2A).

Histone Acetylation Is Responsive to Metabolic Cues

Modulation of histone acetylation is a fundamental process to define the fate of neural progenitors toward a specific lineage during neural development [163, 164]. It consists of the covalent addition of an acetyl group from acetyl coenzyme A (AcCoA) to a lysine residue on a histone by a histone acetyltransferase (HAT) enzyme, hereby neutralizing the positive charge to weaken DNA-histone interactions. Histone acetylation can be reverted by histone deacetylases (HDACs), leading to transcriptional repression. Mounting research illustrates the critical role of histone acetylation in the control of the neurogenic transcriptional programs and neurodegenerative conditions [165–167]. Interestingly, this epigenetic mark is widely responsive to metabolic cues [168], and the implications of this crosstalk for neurogenesis are beginning to emerge.

Histone Acetylation in Neurogenesis Depends on AcCoA

In mammals, two main enzymes sustain AcCoA production for histone acetylation: acetate-dependent acetyl-CoA synthetase 2 (ACSS2) and citrate-dependent ATP-citrate lyase (ACL). Unexpectedly, during neural differentiation, ACSS2 locates to the nucleus and regulates the expression of neural markers such as *NeuN*, *Map2*, or *Snap25* [169]. ACSS2 is recruited to chromatin sites proximal to genes linked to neural differentiation, which are also enriched for acetylation of lysines 5 and 12 of histone H4 (H4K5/K12Ac) and lysine 9 of histone H3 (H3K9Ac). This is consistent with the idea that ACSS2 locally provides a constant AcCoA fuel for histone acetylation by HATs CBP and p300 during neurodevelopment at neuron-specific genes (Fig. 2). Consistently, altering ACSS2 function in the hippocampus impairs long-term memory consolidation in mice [169]. Interestingly, oxygen and serum limitation increases nuclear localization of ACSS2 which contributes to sustaining histone acetylation levels and cell survival [170, 171]. Along these lines, recent research in mice demonstrates that alcohol intake is metabolized in the liver to acetate, which impacts the brain metabolism leading to increased histone acetylation and transcriptional activity in the dorsal hippocampus in an ACSS2-dependent manner [172]. This evidence illustrates important connections between metabolism and histone acetylation in neural development, with potential implications for cognitive conditions such as addictive behavior or age-associated memory decline.

We have previously discussed that during neural development, a switch from glycolytic to oxidative phenotype occurs. This metabolic rewiring might be connected to the wide epigenomic transitions driving neurogenesis. For example, TIGAR is a bisphosphatase which reduces intracellular fructose-2,6-bisphosphate levels, leading

to decreased glycolytic flux [173]. Accordingly, TIGAR expression increases during embryonic cortex development and along the course of differentiation of NSC, which is accompanied by increased expression of neural markers such as MAP2, GFAP, *Tuj1*, *NeuroD1*, and *Ngn1* [174]. Interestingly, overexpression of TIGAR in NSC promotes differentiation favoring the glycolytic-to-oxidative metabolic switch, which is accompanied by increased AcCoA, which ultimately contributes to increased H3K9Ac at the promoters of critical neural genes including *Ngn1*, *Neurod1*, and *Gfap* (Fig. 2B) [174].

Histone Deacetylation in Neurogenesis and Its Interaction with Metabolism

HDAC activity manipulation has been linked to the metabolic reprogramming driving neural differentiation. For example, pan-HDAC inhibitors such as sodium butyrate (NaBt) or valproic acid (VPA) induce neuronal differentiation of NPC [175, 176]. NaBt promotes mitochondrial biogenesis and enhances the oxidative metabolism in NPCs [164]. This is accompanied by increased CBP HAT activity and relocation of the H3K27Ac mark at regulatory regions of genes involved in mitochondrial metabolism and OXPHOS, providing a mechanistic link between epigenetic and metabolic reprogramming in neurogenesis [164]. Similarly, VPA suppresses the proliferation of NPCs and promotes the expression of neural markers including *Tuj1* and CaMKII. VPA also increases the glycolytic flux in NPCs and the H3K9Ac mark at the promoters of *Ngn2* and *Mash1*, and H4Ac within the *Ngn1*, *Math1*, and *p15* promoters [56, 57, 176].

Another deacetylase that is influenced by metabolism is the class III HDAC SIRT1. The deacetylase activity of SIRT1 is coupled to the hydrolysis of the metabolite NAD⁺, and has been characterized as a redox sensor, as SIRT1 activation is dependent on the NAD⁺/NADH ratio. The glycolytic phenotype of NSC tends to reduce the NAD⁺ pool, a process that intensifies with aging [177]. Interestingly, the pharmacological increase of NAD⁺ in aged mice promotes proliferation and neurogenesis in the hippocampus [177, 178], partially mediated by SIRT1 [177, 179]. Furthermore, the redox state is a critical regulator of self-renewal and differentiation of NPCs through controlling SIRT1, which when activated downregulates the H3K9Ac epigenetic mark, hereby dictating cell fate decisions (Fig. 2B) [180, 181]. Accordingly, mitochondrial impairment through increased oxidative stress and accumulation of mtDNA damage in NSC activates SIRT1 to epigenetically support *Mash1*-mediated astrogliosis [182], or ultimately drive cellular senescence and mitophagy [94]. Together, these findings indicate that SIRT1 is an

important regulator of NS/PC self-renewal and differentiation potential, and an epigenetic effector of metabolic transitions in redox states [183].

Novel Methods to Study Metabolic Transitions in Live Cells: Fluorescence Lifetime Imaging Microscopy of Endogenous Biomarkers

We have shown that small molecule metabolites play much underappreciated roles in cell differentiation and homeostasis by controlling the activity of several epigenetic remodelers. Generally, metabolic states are determined from whole cell lysates. Yet, local concentrations of certain metabolites could alter enzymatic properties in discrete microenvironments as shown in the case of ACSS2 locally supplying AcCoA. A redox switch from glycolytic to oxidative metabolism and wide epigenetic reprogramming are hallmarks for NSC maintenance and cell fate. However, the molecular links between these major processes during differentiation remain largely unknown. Hereby, novel approaches to measure metabolism across scales are very much needed in this field. To approach this challenge, it is necessary to consider cellular heterogeneity, which is an intrinsic mechanism of determination of cell fate. Deciphering heterogeneity has proven crucial to understanding differentiation processes, and single-cell transcriptomics has become the gold standard to untangle heterogeneity at the transcriptional level. While mounting research is seeding light into transcriptional and epigenetic regulation of neurogenesis in single cells, disentangling metabolism at the cellular level has fallen behind due to the limited availability of technological approaches. Currently, single-cell metabolomics is a rapidly advancing field. The main techniques used for single-cell analysis include mass spectrometry (MS) and MS imaging (MSI), electrochemistry, fluorescence microscopy, vibrational spectroscopy, and capillary electrophoresis, which have been thoroughly reviewed elsewhere [184–187].

A number of these techniques have been applied to reveal the metabolic intricacies of neural development. For example, a single-cell profiling method named matrix-enhanced-secondary ion mass spectrometry (ME-SIMS) was used to investigate the lipid profiles of neuronal cells, allowing to classify single-cell populations and subpopulations using SIMS profiling of lipid and metabolite contents [188]. Additionally, MSI has been proven effective in characterizing metabolism in human brain organoids, allowing direct comparison of certain lipid signal intensities and distributions, including ceramides and phosphatidylethanolamine [189].

In the last decade, optical metabolic imaging has been extensively applied to quantify redox states at subcellular scales. Two-photon excited fluorescence (2PEF) microscopy-based techniques provide functional images of tissues and cells through intrinsic fluorophores that are naturally present in cells and tissues [190]. Importantly, the metabolic coenzymes NADH and FAD constitute biomarkers for metabolism, mitochondrial function, and oxidative stress in live cells and tissues, due to their intrinsic fluorescent properties [191, 192]. Since the pioneering work of Britton Chance [193], NADH and FAD fluorescence intensity has been extensively used to monitor changes in metabolism in different fields of biomedical research such as neuroscience and differentiation [194, 195].

Multiphoton microscopy, such as third harmonic generation (THG), has been successfully used to label-free visualization of structures with high lipid content, including myelin sheaths, cellular membranes, or ECM structures in live organisms [196–198]. Two-photon fluorescence lifetime microscopy (2P-FLIM) of the metabolic coenzymes NADH and FAD has been increasingly used in recent years to perform metabolic imaging as it reveals the richness and complexity of several metabolic processes in a label-free and non-invasive way. For example, 2P-FLIM of NADH provides very sensitive measurements of the redox states (NADH/NAD⁺) of cells as well as glycolysis, OXPHOS, and oxidative stress rates [199–204] and has the spatiotemporal resolution required to characterize dynamic physiological states and NADH intracellular subcellular compartmentalization [183, 205, 206]. In recent years, fluorescence lifetime imaging of NADH has been increasingly used to monitor metabolic shifts in neurodegenerative diseases [207] as well as during stem cell differentiation, for example, to map subcellular metabolism of stem cells in a label-free and non-invasive way measuring metabolic trajectories during differentiation [183, 202, 203, 208–210]. 2P-FLIM of NADH was used to characterize the metabolic state of NPCs during differentiation in live cells, observing that the ratio of free to protein-bound NADH strongly correlates with the differentiation state. Undifferentiated NPCs have a glycolytic phenotype characterized by high free/bound NADH, while the OXPHOS phenotype from differentiated neurons is characterized by low free/bound NADH. Hereby, the metabolic signature of NPCs correlates with their differentiation potential, showing that neuronal progenitors and glial progenitors have a different free/bound NADH ratio in line with evidence that energy metabolism and the redox state are important regulators of NSC [211, 212, 181, 213, 214]. Ultimately, multiparametric metabolic imaging has been recently implemented in an efficient way with multicolor multiphoton simultaneous excitation of NADH and FAD [215, 216] and it could bring new insights into the understanding of the spatial and temporal metabolic patterns

during neural differentiation, as NADH and FAD provide complementary information on different metabolic pathways [209]. Using fluorescently labeled epigenetic remodelers or histone modifications, or even genomic regions through for example dCas9-based techniques [217–220], a combination of FLIM-FRET techniques could be implemented, representing new avenues to uncover the intimate relationship between metabolism and epigenome regulation during differentiation in live cells and tissues.

Conclusions

In conclusion, the process of neural differentiation is a complex and tightly regulated event that involves wide metabolic and epigenetic transitions. Through the regulation of metabolic pathways, the cell generates energy and biosynthetic precursors necessary for stemness maintenance and determination of cell fate. Additionally, epigenetic modifications, such as DNA methylation, histone modifications, and non-coding RNA, play critical roles in gene expression regulation during neural differentiation.

Recent studies have highlighted the importance of the crosstalk between metabolism and epigenetic during early neural development, which have provided critical insights into the mechanisms that govern neural differentiation. These findings have potential implications for understanding the etiology of neurodevelopmental disorders and for developing new therapies. However, there is still much to be learned about the precise mechanisms that underlie metabolic and epigenetic transitions during neural differentiation. Although the intricate relationship between epigenetics and metabolism has been emerging, there are still many outstanding questions. For example, why are histone post-translational modifications and epigenetic marks at certain genomic loci more susceptible to fluctuations in metabolic cofactor availability than others? Another outstanding question is how metabolic signals shape brain DNA methylation during adulthood and how manipulations of the diet could modulate DNA methylation in the adult brain. Additionally, it is still not fully understood how metabolic pathways and epigenetic modifications interact to regulate NSC fate. Finally, it is important to understand how metabolic transitions translate into changes in redox balance, cell signaling, and epigenetics, thereby regulating stem cell activation and differentiation. For example, in light of current knowledge, it is of interest to define the precise mechanisms through which IC metabolism modulates DNA methylation and subsequently regulates neuronal fate during development. This represents a pivotal yet challenging area of investigation, necessitating the application of cutting-edge methodologies and integrative approaches to unravel the intricate interplay between these biological processes. Further research

is needed to unravel the intricacies of these processes and to determine how they are coordinated and regulated. The future prospects are undoubtedly related to advances in technologies such as simultaneous single-cell transcriptomics and metabolomics in real time which will certainly contribute to our understanding of the molecular mechanisms underlying neural differentiation.

Acknowledgements We thank all members of Aguilar-Arnal and Chiara Stringari laboratories for helpful discussions and particularly Marcia Bustamante Zepeda, MSc, at the IIBo, for critically reading the manuscript. E. Sánchez-Ramírez acknowledges the Ph.D. program Doctorado en Ciencias Bioquímicas-Universidad Nacional Autónoma de México.

Author Contribution Lorena Aguilar-Arnal and Edgar Sánchez-Ramírez conceptualized and designed the manuscript. Edgar Sánchez-Ramírez, Thi Phuong Lien Ung, Chiara Stringari, and Lorena Aguilar-Arnal wrote the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by the Human Frontier Science Program under the contract RGY0078/2017 ChroMet project and by the Chan Zuckerberg Initiative grant 2023-331942 scNeuroMET project to C. Stringari and L. Aguilar-Arnal, grants DGAPA-PAPIIT IN210619 and IN208022 from Universidad Nacional Autónoma de México, and The National Council of Science and Technology (FORDECYT-PRONACES/15758/2020) to L. Aguilar-Arnal. E. Sánchez-Ramírez acknowledges the reception of PhD fellowship from the National Council of Science and Technology (CONAHCyT).

Data Availability Not applicable.

Code Availability Not applicable.

Declarations

Ethics Approval Not applicable.

Consent to Participate Not applicable.

Consent for Publication All authors consent to publish this manuscript.

Conflict of Interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Wu J, Izpisua Belmonte JC (2016) Stem cells: a renaissance in human biology research. *Cell* 165(7):1572–1585
2. Weger M, Diotel N, Dorseman AC, Dickmeis T, Weger BD (2017) Stem cells and the circadian clock. *Dev Biol* 431(2):111–123
3. Zakrzewski W, Dobrzynski M, Szymonowicz M, Rybak Z (2019) Stem cells: past, present, and future. *Stem Cell Res Ther* 10(1):68
4. Rossant J, Tam PPL (2022) Early human embryonic development: blastocyst formation to gastrulation. *Dev Cell* 57(2):152–165
5. Bystron I, Blakemore C, Rakic P (2008) Development of the human cerebral cortex: Boulder Committee revisited. *Nat Rev Neurosci* 9(2):110–122
6. Taupin P (2006) Adult neurogenesis in mammals. *Curr Opin Mol Ther* 8(4):345–351
7. Llorente V, Velarde P, Desco M, Gomez-Gavio MV (2022) Current understanding of the neural stem cell niches. *Cells* 11(19):3002. <https://doi.org/10.3390/cells11193002>
8. Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH (1998) Neurogenesis in the adult human hippocampus. *Nat Med* 4(11):1313–1317
9. Ming GL, Song H (2005) Adult neurogenesis in the mammalian central nervous system. *Annu Rev Neurosci* 28:223–250
10. Gattazzo F, Urciuolo A, Bonaldo P (2014) Extracellular matrix: a dynamic microenvironment for stem cell niche. *Biochim Biophys Acta* 1840(8):2506–2519
11. Gage FH, Temple S (2013) Neural stem cells: generating and regenerating the brain. *Neuron* 80(3):588–601
12. Luo Y, Coskun V, Liang A, Yu J, Cheng L, Ge W, Shi Z, Zhang KC (2015) Single-cell transcriptome analyses reveal signals to activate dormant neural stem cells. *Cell* 161(5):1175–1186. <https://doi.org/10.1016/j.cell.2015.04.001>
13. Kim DY, Rhee I, Paik J (2014) Metabolic circuits in neural stem cells. *Cell Mol Life Sci* 71(21):4221–4241
14. Suh H, Deng W, Gage FH (2009) Signaling in adult neurogenesis. *Annu Rev Cell Dev Biol* 25:253–275
15. Gomez-Lopez S, Lerner RG, Petritsch C (2014) Asymmetric cell division of stem and progenitor cells during homeostasis and cancer. *Cell Mol Life Sci* 71(4):575–597
16. Homem CC, Repic M, Knoblich JA (2015) Proliferation control in neural stem and progenitor cells. *Nat Rev Neurosci* 16(11):647–659
17. Namihira M, Nakashima K (2013) Mechanisms of astrocytogenesis in the mammalian brain. *Curr Opin Neurobiol* 23(6):921–927
18. Bernstein BE, Meissner A, Lander ES (2007) The mammalian epigenome. *Cell* 128(4):669–681
19. Namihira M, Kohyama J, Abematsu M, Nakashima K (2008) Epigenetic mechanisms regulating fate specification of neural stem cells. *Philos Trans R Soc Lond B Biol Sci* 363(1500):2099–2109
20. Hu XL, Wang Y, Shen Q (2012) Epigenetic control on cell fate choice in neural stem cells. *Protein Cell* 3(4):278–290
21. Agostini M, Romeo F, Inoue S, Niklison-Chirou MV, Elia AJ, Dinsdale D, Morone N, Knight RA et al (2016) Metabolic reprogramming during neuronal differentiation. *Cell Death Differ* 23(9):1502–1514
22. French HM, Reid M, Mamontov P, Simmons RA, Grinspan JB (2009) Oxidative stress disrupts oligodendrocyte maturation. *J Neurosci Res* 87(14):3076–3087
23. Zheng X, Boyer L, Jin M, Mertens J, Kim Y, Ma L, Ma L, Hamm M et al (2016) Metabolic reprogramming during neuronal differentiation from aerobic glycolysis to neuronal oxidative phosphorylation. *Elife* 5. <https://doi.org/10.7554/eLife.13374>

24. Xu X, Duan S, Yi F, Ocampo A, Liu GH, Izpisua Belmonte JC (2013) Mitochondrial regulation in pluripotent stem cells. *Cell Metab* 18(3):325–332
25. De Filippis L, Delia D (2011) Hypoxia in the regulation of neural stem cells. *Cell Mol Life Sci* 68(17):2831–2844
26. Tsogetbaatar E, Landin C, Minter-Dykhouse K, Folmes CDL (2020) Energy metabolism regulates stem cell pluripotency. *Front Cell Dev Biol* 8:87
27. Angelopoulos I, Gakis G, Birmpas K, Kyrousi C, Habeos EE, Kaplani K, Lygerou Z, Habeos I et al (2022) Metabolic regulation of the neural stem cell fate: unraveling new connections, establishing new concepts. *Front Neurosci* 16:1009125
28. Zhang J, Zhao J, Dahan P, Lu V, Zhang C, Li H, Teitell MA (2018) Metabolism in pluripotent stem cells and early mammalian development. *Cell Metab* 27(2):332–338
29. Martinez-Reyes I, Chandel NS (2020) Mitochondrial TCA cycle metabolites control physiology and disease. *Nat Commun* 11(1):102
30. Spinelli JB, Haigis MC (2018) The multifaceted contributions of mitochondria to cellular metabolism. *Nat Cell Biol* 20(7):745–754
31. Potter M, Newport E, Morten KJ (2016) The Warburg effect: 80 years on. *Biochem Soc Trans* 44(5):1499–1505
32. Vander Heiden MG, Cantley LC, Thompson CB (2009) Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 324(5930):1029–1033
33. Lunt SY, Vander Heiden MG (2011) Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu Rev Cell Dev Biol* 27(1):441–464
34. Agathocleous M, Love NK, Randlett O, Harris JJ, Liu J, Murray AJ, Harris WA (2012) Metabolic differentiation in the embryonic retina. *Nat Cell Biol* 14(8):859–864
35. Homem CCF, Steinmann V, Burkard TR, Jais A, Esterbauer H, Knoblich JA (2014) Ecdysone and mediator change energy metabolism to terminate proliferation in *Drosophila* neural stem cells. *Cell* 158(4):874–888
36. van den Ameele J, Brand AH (2019) Neural stem cell temporal patterning and brain tumour growth rely on oxidative phosphorylation. *Elife* 8. <https://doi.org/10.7554/eLife.47887>
37. Gershon TR, Crowther AJ, Liu H, Miller CR, Deshmukh M (2013) Cerebellar granule neuron progenitors are the source of Hk2 in the postnatal cerebellum. *Cancer Metab* 1(1):15
38. Gershon TR, Crowther AJ, Tikunov A, Garcia I, Annis R, Yuan H, Miller CR, Macdonald J et al (2013) Hexokinase-2-mediated aerobic glycolysis is integral to cerebellar neurogenesis and pathogenesis of medulloblastoma. *Cancer Metab* 1(1):2
39. Beckervordersandforth R, Ebert B, Schaffner I, Moss J, Fiebig C, Shin J, Moore DL, Ghosh L et al (2017) Role of mitochondrial metabolism in the control of early lineage progression and aging phenotypes in adult hippocampal neurogenesis. *Neuron* 93(3):560–573 e566
40. Khacho M, Clark A, Svoboda DS, Azzi J, MacLaurin JG, Meghaizel C, Sesaki H, Lagace DC et al (2016) Mitochondrial dynamics impacts stem cell identity and fate decisions by regulating a nuclear transcriptional program. *Cell Stem Cell* 19(2):232–247
41. O'Brien LC, Keeney PM, Bennett JP Jr (2015) Differentiation of human neural stem cells into motor neurons stimulates mitochondrial biogenesis and decreases glycolytic flux. *Stem Cells Dev* 24(17):1984–1994
42. Chen J, Ren Y, Gui C, Zhao M, Wu X, Mao K, Li W, Zou F (2018) Phosphorylation of Parkin at serine 131 by p38 MAPK promotes mitochondrial dysfunction and neuronal death in mutant A53T alpha-synuclein model of Parkinson's disease. *Cell Death Dis* 9(6):700
43. Khacho M, Clark A, Svoboda DS, MacLaurin JG, Lagace DC, Park DS, Slack RS (2017) Mitochondrial dysfunction underlies cognitive defects as a result of neural stem cell depletion and impaired neurogenesis. *Hum Mol Genet* 26(17):3327–3341
44. Prieto J, Leon M, Ponsoda X, Sendra R, Bort R, Ferrer-Lorente R, Raya A, Lopez-Garcia C et al (2016) Early ERK1/2 activation promotes DRP1-dependent mitochondrial fission necessary for cell reprogramming. *Nat Commun* 7(1):11124
45. Xie W, Guo D, Li J, Yue L, Kang Q, Chen G, Zhou T, Wang H et al (2022) CEND1 deficiency induces mitochondrial dysfunction and cognitive impairment in Alzheimer's disease. *Cell Death Differ* 29(12):2417–2428
46. Zhao Y, Sun X, Hu D, Prosdocimo DA, Hoppel C, Jain MK, Ramachandran R, Qi X (2019) ATAD3A oligomerization causes neurodegeneration by coupling mitochondrial fragmentation and bioenergetics defects. *Nat Commun* 10(1):1371
47. Jahani-Asl A, Pilon-Larose K, Xu W, MacLaurin JG, Park DS, McBride HM, Slack RS (2011) The mitochondrial inner membrane GTPase, optic atrophy 1 (Opa1), restores mitochondrial morphology and promotes neuronal survival following excitotoxicity. *J Biol Chem* 286(6):4772–4782
48. Klivenyi P, St Clair D, Wermer M, Yen HC, Oberley T, Yang L, Flint Beal M (1998) Manganese superoxide dismutase overexpression attenuates MPTP toxicity. *Neurobiol Dis* 5(4):253–258
49. Massaad CA, Washington TM, Pautler RG, Klann E (2009) Overexpression of SOD-2 reduces hippocampal superoxide and prevents memory deficits in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 106(32):13576–13581
50. Wu W, Zhao D, Shah SZA, Zhang X, Lai M, Yang D, Wu X, Guan Z et al (2019) OPA1 overexpression ameliorates mitochondrial cristae remodeling, mitochondrial dysfunction, and neuronal apoptosis in prion diseases. *Cell Death Dis* 10(10):710
51. Sies H, Jones DP (2020) Reactive oxygen species (ROS) as pleiotropic physiological signalling agents. *Nat Rev Mol Cell Biol* 21(7):363–383
52. Oswald MCW, Garnham N, Sweeney ST, Landgraf M (2018) Regulation of neuronal development and function by ROS. *FEBS Lett* 592(5):679–691
53. Wilson C, Muñoz-Palma E, González-Billault C (2018) From birth to death: a role for reactive oxygen species in neuronal development. *Semin Cell Dev Biol* 80:43–49
54. Hung CH-L, Cheng SS-Y, Cheung Y-T, Wuwongse S, Zhang NQ, Ho Y-S, Lee SM-Y, Chang RC-C (2018) A reciprocal relationship between reactive oxygen species and mitochondrial dynamics in neurodegeneration. *Redox Biol* 14:7–19
55. Brillo V, Chieragato L, Leanza L, Muccioli S, Costa R (2021) Mitochondrial dynamics, ROS, and cell signaling: a blended overview. *Life* 11(4):332
56. Chen A, Wang M, Xu C, Zhao Y, Xian P, Li Y, Zheng W, Yi X et al (2023) Glycolysis mediates neuron specific histone acetylation in valproic acid-induced human excitatory neuron differentiation. *Front Mol Neurosci* 16:1151162
57. Chen W, Zhao H, Li Y (2023) Mitochondrial dynamics in health and disease: mechanisms and potential targets. *Signal Transduct Target Ther* 8(1):333
58. Meacham CE, DeVilbiss AW, Morrison SJ (2022) Metabolic regulation of somatic stem cells in vivo. *Nat Rev Mol Cell Biol* 23(6):428–443
59. Candelario KM, Shuttleworth CW, Cunningham LA (2013) Neural stem/progenitor cells display a low requirement for oxidative metabolism independent of hypoxia inducible factor-1alpha expression. *J Neurochem* 125(3):420–429
60. Chen HL, Pistollato F, Hoepfner DJ, Ni HT, McKay RD, Panchision DM (2007) Oxygen tension regulates survival and fate of

- mouse central nervous system precursors at multiple levels. *Stem Cells* 25(9):2291–2301
61. Francis KR, Wei L (2010) Human embryonic stem cell neural differentiation and enhanced cell survival promoted by hypoxic preconditioning. *Cell Death Dis* 1(2):e22
 62. Horie N, So K, Moriya T, Kitagawa N, Tsutsumi K, Nagata I, Shinohara K (2008) Effects of oxygen concentration on the proliferation and differentiation of mouse neural stem cells in vitro. *Cell Mol Neurobiol* 28(6):833–845
 63. Li G, Liu J, Guan Y, Ji X (2021) The role of hypoxia in stem cell regulation of the central nervous system: from embryonic development to adult proliferation. *CNS Neurosci Ther* 27(12):1446–1457
 64. Rodrigues CA, Diogo MM, da Silva CL, Cabral JM (2010) Hypoxia enhances proliferation of mouse embryonic stem cell-derived neural stem cells. *Biotechnol Bioeng* 106(2):260–270
 65. Studer L, Csete M, Lee SH, Kabbani N, Walikonis J, Wold B, McKay R (2000) Enhanced proliferation, survival, and dopaminergic differentiation of CNS precursors in lowered oxygen. *J Neurosci* 20(19):7377–7383
 66. Lange C, Turrero Garcia M, Decimo I, Bifari F, Eelen G, Quaegebeur A, Boon R, Zhao H et al (2016) Relief of hypoxia by angiogenesis promotes neural stem cell differentiation by targeting glycolysis. *EMBO J* 35(9):924–941
 67. Kumar A, Vaish M, Karuppagounder SS, Gazaryan I, Cave JW, Starkov AA, Anderson ET, Zhang S et al (2021) HIF1 α stabilization in hypoxia is not oxidant-initiated. *Elife* 10:e72873
 68. Majmundar AJ, Wong WJ, Simon MC (2010) Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell* 40(2):294–309
 69. Carrica L, Li L, Newville J, Kenton J, Gustus K, Brigman J, Cunningham LA (2019) Genetic inactivation of hypoxia inducible factor 1- α (HIF-1 α) in adult hippocampal progenitors impairs neurogenesis and pattern discrimination learning. *Neurobiol Learn Mem* 157:79–85
 70. Li L, Candelario KM, Thomas K, Wang R, Wright K, Messier A, Cunningham LA (2014) Hypoxia inducible factor-1 α (HIF-1 α) is required for neural stem cell maintenance and vascular stability in the adult mouse SVZ. *J Neurosci* 34(50):16713–16719
 71. Cui P, Zhang P, Yuan L, Wang L, Guo X, Cui G, Zhang Y, Li M et al (2021) HIF-1 α affects the neural stem cell differentiation of human induced pluripotent stem cells via MFN2-mediated Wnt/ β -catenin signaling. *Front Cell Dev Biol* 9:671704
 72. Gustafsson MV, Zheng X, Pereira T, Gradin K, Jin S, Lundkvist J, Ruas JL, Poellinger L et al (2005) Hypoxia requires notch signaling to maintain the undifferentiated cell state. *Dev Cell* 9(5):617–628
 73. Li Y, Wu L, Yu M, Yang F, Wu B, Lu S, Tu M, Xu H (2018) HIF-1 α is critical for the activation of notch signaling in neurogenesis during acute epilepsy. *Neuroscience* 394:206–219
 74. Mazumdar J, O'Brien WT, Johnson RS, Lamanna JC, Chavez JC, Klein PS, Simon MC (2010) O₂ regulates stem cells through Wnt/ β -catenin signalling. *Nat cell biol* 12(10):1007–1013
 75. Li J, Gong SH, He YL, Cao Y, Chen Y, Huang GH, Wang YF, Zhao M et al (2023) autophagy is essential for neural stem cell proliferation promoted by hypoxia. *Stem Cells* 41(1):77–92
 76. Renault VM, Rafalski VA, Morgan AA, Salih DA, Brett JO, Webb AE, Villeda SA, Thekkat PU et al (2009) FoxO3 regulates neural stem cell homeostasis. *Cell Stem Cell* 5(5):527–539
 77. Yeo H, Lyssiotis CA, Zhang Y, Ying H, Asara JM, Cantley LC, Paik JH (2013) FoxO3 coordinates metabolic pathways to maintain redox balance in neural stem cells. *EMBO J* 32(19):2589–2602
 78. Schaffner I, Minakaki G, Khan MA, Balta EA, Schlotzer-Schrehardt U, Schwarz TJ, Beckervordersandforth R et al (2018) FoxO function is essential for maintenance of autophagic flux and neuronal morphogenesis in adult neurogenesis. *Neuron* 99(6):1188–1203 e1186
 79. Wei Y, Miao Q, Zhang Q, Mao S, Li M, Xu X, Xia X, Wei K et al (2023) Aerobic glycolysis is the predominant means of glucose metabolism in neuronal somata, which protects against oxidative damage. *Nat Neurosci* 26(12):2081–2089
 80. Yanes O, Clark J, Wong DM, Patti GJ, Sanchez-Ruiz A, Benton HP, Trauger SA, Despons C et al (2010) Metabolic oxidation regulates embryonic stem cell differentiation. *Nat Chem Biol* 6(6):411–417
 81. Tracey TJ, Steyn FJ, Wolvetang EJ, Ngo ST (2018) Neuronal lipid metabolism: multiple pathways driving functional outcomes in health and disease. *Front Mol Neurosci* 11:10
 82. Sipione S, Monyror J, Galleguillos D, Steinberg N, Kadam V (2020) Gangliosides in the brain: physiology, pathophysiology and therapeutic applications. *Front Neurosci* 14:572965
 83. Ramosaj M, Madsen S, Maillard V, Scandella V, Sudria-Lopez D, Yuizumi N, Telley L, Knobloch M (2021) Lipid droplet availability affects neural stem/progenitor cell metabolism and proliferation. *Nat Commun* 12(1):7362
 84. Schonfeld P, Reiser G (2013) Why does brain metabolism not favor burning of fatty acids to provide energy? Reflections on disadvantages of the use of free fatty acids as fuel for brain. *J Cereb Blood Flow Metab* 33(10):1493–1499
 85. Madsen S, Ramosaj M, Knobloch M (2021) Lipid metabolism in focus: how the build-up and breakdown of lipids affects stem cells. *Development* 148(10). <https://doi.org/10.1242/dev.191924>
 86. Knobloch M, Pilz GA, Ghesquiere B, Kovacs WJ, Wegleiter T, Moore DL, Hruzova M, Zamboni N et al (2017) A fatty acid oxidation-dependent metabolic shift regulates adult neural stem cell activity. *Cell Rep* 20(9):2144–2155
 87. Stoll EA, Makin R, Sweet IR, Trevelyan AJ, Miwa S, Horner PJ, Turnbull DM (2015) Neural stem cells in the adult subventricular zone oxidize fatty acids to produce energy and support neurogenic activity. *Stem Cells* 33(7):2306–2319
 88. Xie Z, Jones A, Deeney JT, Hur SK, Bankaitis VA (2016) Inborn errors of long-chain fatty acid beta-oxidation link neural stem cell self-renewal to autism. *Cell Rep* 14(5):991–999
 89. Mallick R, Basak S, Duttaroy AK (2019) Docosahexaenoic acid, 22:6n-3: its roles in the structure and function of the brain. *Int J Dev Neurosci* 79(1):21–31
 90. Kim HY, Moon HS, Cao D, Lee J, Kevala K, Jun SB, Lovinger DM, Akbar M et al (2011) N-Docosahexaenoyl ethanolamide promotes development of hippocampal neurons. *Biochem J* 435(2):327–336
 91. Hamilton LK, Dufresne M, Joppe SE, Petryszyn S, Aumont A, Calon F, Barnabe-Heider F, Furtos A et al (2015) Aberrant lipid metabolism in the forebrain niche suppresses adult neural stem cell proliferation in an animal model of Alzheimer's disease. *Cell Stem Cell* 17(4):397–411
 92. Knobloch M, Braun SM, Zurkirchen L, von Scholtz C, Zamboni N, Arauzo-Bravo MJ, Kovacs WJ, Karalay O et al (2013) Metabolic control of adult neural stem cell activity by Fasn-dependent lipogenesis. *Nature* 493(7431):226–230
 93. Bowers M, Liang T, Gonzalez-Bohorquez D, Zocher S, Jaeger BN, Kovacs WJ, Rohrl C, Cramb KML et al (2020) FASN-dependent lipid metabolism links neurogenic stem/progenitor cell activity to learning and memory deficits. *Cell Stem Cell* 27(1):98–109 e111
 94. Liang KX, Kristiansen CK, Mostafavi S, Vatne GH, Zantingh GA, Kianian A, Tzoulis C, Hoyland LE et al (2020) Disease-specific phenotypes in iPSC-derived neural stem cells with POLG mutations. *EMBO Mol Med* 12(10):e12146

95. Bogie JFJ, Haidar M, Kooij G, Hendriks JJA (2020) Fatty acid metabolism in the progression and resolution of CNS disorders. *Adv Drug Deliv Rev* 159:198–213
96. Kandel P, Semerci F, Mishra R, Choi W, Bajic A, Baluya D, Ma L, Chen K et al (2022) Oleic acid is an endogenous ligand of TLX/NR2E1 that triggers hippocampal neurogenesis. *Proc Natl Acad Sci U S A* 119(13):e2023784119
97. Borsini A, Alboni S, Horowitz MA, Tojo LM, Cannazza G, Su KP, Pariante CM, Zunszain PA (2017) Rescue of IL-1beta-induced reduction of human neurogenesis by omega-3 fatty acids and antidepressants. *Brain Behav Immun* 65:230–238
98. Borsini A, Stangl D, Jeffries AR, Pariante CM, Thuret S (2020) The role of omega-3 fatty acids in preventing glucocorticoid-induced reduction in human hippocampal neurogenesis and increase in apoptosis. *Transl Psychiatry* 10(1):219
99. Liu J, Sahin C, Ahmad S, Magomedova L, Zhang M, Jia Z, Metherell AH, Orellana A et al (2022) The omega-3 hydroxy fatty acid 7(S)-HDHA is a high-affinity PPARalpha ligand that regulates brain neuronal morphology. *Sci Signal* 15(741):eabo1857
100. Hussain G, Wang J, Rasul A, Anwar H, Imran A, Qasim M, Zafar S, Kamran SKS et al (2019) Role of cholesterol and sphingolipids in brain development and neurological diseases. *Lipids Health Dis* 18(1):26
101. Li D, Zhang J, Liu Q (2022) Brain cell type-specific cholesterol metabolism and implications for learning and memory. *Trends Neurosci* 45(5):401–414
102. Mauch DH, Nagler K, Schumacher S, Goritz C, Muller EC, Otto A, Pfrieger FW (2001) CNS synaptogenesis promoted by gliaderived cholesterol. *Science* 294(5545):1354–1357
103. Martin MG, Pfrieger F, Dotti CG (2014) Cholesterol in brain disease: sometimes determinant and frequently implicated. *EMBO Rep* 15(10):1036–1052
104. Saito K, Dubreuil V, Arai Y, Wilsch-Brauninger M, Schwudke D, Saher G, Miyata T, Breier G et al (2009) Ablation of cholesterol biosynthesis in neural stem cells increases their VEGF expression and angiogenesis but causes neuron apoptosis. *Proc Natl Acad Sci U S A* 106(20):8350–8355
105. Cunningham D, DeBarber AE, Bir N, Binkley L, Merkens LS, Steiner RD, Herman GE (2015) Analysis of hedgehog signaling in cerebellar granule cell precursors in a conditional *Nsdhl* allele demonstrates an essential role for cholesterol in postnatal CNS development. *Hum Mol Genet* 24(10):2808–2825
106. Driver AM, Kratz LE, Kelley RI, Stottmann RW (2016) Altered cholesterol biosynthesis causes precocious neurogenesis in the developing mouse forebrain. *Neurobiol Dis* 91:69–82
107. Clare CE, Brassington AH, Kwong WY, Sinclair KD (2019) One-carbon metabolism: linking nutritional biochemistry to epigenetic programming of long-term development. *Annu Rev Anim Biosci* 7(1):263–287
108. Clare CE, Brassington AH, Kwong WY, Sinclair KD (2019) One-carbon metabolism: linking nutritional biochemistry to epigenetic programming of long-term development. *Annu Rev Anim Biosci* 7(1):263–287
109. Zheng Y, Cantley LC (2019) Toward a better understanding of folate metabolism in health and disease. *J Exp Med* 216(2):253–266
110. Copp AJ, Stanier P, Greene ND (2013) Neural tube defects: recent advances, unsolved questions, and controversies. *Lancet Neurol* 12(8):799–810
111. Morin I, Platt R, Weisberg I, Sabbaghian N, Wu Q, Garrow TA, Rozen R (2003) Common variant in betaine-homocysteine methyltransferase (BHMT) and risk for spina bifida. *Am J Med Genet A* 119A(2):172–176
112. Rampersaud E, Melvin EC, Siegel D, Mehltritt L, Dickerson ME, George TM, Enterline D, Nye JS, Speer MC, N.T.D.C. Group (2003) Updated investigations of the role of methylenetetrahydrofolate reductase in human neural tube defects. *Clin Genet* 63(3):210–4. <https://doi.org/10.1034/j.1399-0004.2003.00043.x>
113. Shaw GM, Lu W, Zhu H, Yang W, Briggs FB, Carmichael SL, Barcellos LF, Lammer EJ et al (2009) 118 SNPs of folate-related genes and risks of spina bifida and conotruncal heart defects. *BMC Med Genet* 10:49
114. Kruman PR II, Mouton Emokpae R, Cutler RG, Mattson MP (2005) Folate deficiency inhibits proliferation of adult hippocampal progenitors. *Neuroreport* 16(10):1055–1059
115. Zhang X, Liu H, Cong G, Tian Z, Ren D, Wilson JX, Huang G (2008) Effects of folate on notch signaling and cell proliferation in neural stem cells of neonatal rats in vitro. *J Nutr Sci Vitaminol (Tokyo)* 54(5):353–356
116. Liu H, Cao J, Zhang H, Qin S, Yu M, Zhang X, Wang X, Gao Y et al (2013) Folic acid stimulates proliferation of transplanted neural stem cells after focal cerebral ischemia in rats. *J Nutr Biochem* 24(11):1817–1822
117. Luo S, Zhang X, Yu M, Yan H, Liu H, Wilson JX, Huang G (2013) Folic acid acts through DNA methyltransferases to induce the differentiation of neural stem cells into neurons. *Cell Biochem Biophys* 66(3):559–566
118. Jerber J, Seaton DD, Cuomo ASE, Kumasaka N, Haldane J, Steer J, Patel M, Pearce D et al (2021) Population-scale single-cell RNA-seq profiling across dopaminergic neuron differentiation. *Nat Genet* 53(3):304–312
119. van de Leemput J, Boles NC, Kiehl TR, Corneo B, Lederman P, Menon V, Lee C, Martinez RA et al (2014) CORTECON: a temporal transcriptome analysis of in vitro human cerebral cortex development from human embryonic stem cells. *Neuron* 83(1):51–68
120. Ziller MJ, Edri R, Yaffe Y, Donaghey J, Pop R, Mallard W, Issner R, Gifford CA et al (2015) Dissecting neural differentiation regulatory networks through epigenetic footprinting. *Nature* 518(7539):355–359
121. Bonnefont J, Vanderhaeghen P (2021) Neuronal fate acquisition and specification: time for a change. *Curr Opin Neurobiol* 66:195–204
122. Hirabayashi Y, Gotoh Y (2010) Epigenetic control of neural precursor cell fate during development. *Nat Rev Neurosci* 11(6):377–388
123. Etchegaray JP, Mostoslavsky R (2016) Interplay between metabolism and epigenetics: a nuclear adaptation to environmental changes. *Mol Cell* 62(5):695–711
124. Dai Z, Ramesh V, Locasale JW (2020) The evolving metabolic landscape of chromatin biology and epigenetics. *Nat Rev Genet* 21(12):737–753
125. Yao B, Christian KM, He C, Jin P, Ming GL, Song H (2016) Epigenetic mechanisms in neurogenesis. *Nat Rev Neurosci* 17(9):537–549
126. Zhang M, Zhao J, Lv Y, Wang W, Feng C, Zou W, Su L, Jiao J (2020) Histone variants and histone modifications in neurogenesis. *Trends Cell Biol* 30(11):869–880
127. Niehrs C (2009) Active DNA demethylation and DNA repair. *Differentiation* 77(1):1–11
128. Cimmino L, Abdel-Wahab O, Levine RL, Aifantis I (2011) TET family proteins and their role in stem cell differentiation and transformation. *Cell Stem Cell* 9(3):193–204
129. Zou X, Ma W, Solov'yov IA, Chipot C, Schulten K (2012) Recognition of methylated DNA through methyl-CpG binding domain proteins. *Nucleic Acids Res* 40(6):2747–2758
130. Lister R, Mukamel EA, Nery JR, Urich M, Puddifoot CA, Johnson ND, Lucero J, Huang Y et al (2013) Global epigenomic reconfiguration during mammalian brain development. *Science* 341(6146):1237905

131. Ericson J, Rashbass P, Schedl A, Brenner-Morton S, Kawakami A, van Heyningen V, Jessell TM, Briscoe J (1997) Pax6 controls progenitor cell identity and neuronal fate in response to graded Shh signaling. *Cell* 90(1):169–180
132. Gao J, Wang J, Wang Y, Dai W, Lu L (2011) Regulation of Pax6 by CTCF during induction of mouse ES cell differentiation. *PLoS ONE* 6(6):e20954
133. Varela-Rey M, Iruarrizaga-Lejarreta M, Lozano JJ, Aransay AM, Fernandez AF, Lavin JL, Mosen-Ansorena D, Berdasco M et al (2014) S-adenosylmethionine levels regulate the schwann cell DNA methylome. *Neuron* 81(5):1024–1039
134. Korsmo HW, Jiang X (2021) One carbon metabolism and early development: a diet-dependent destiny. *Trends Endocrinol Metab* 32(8):579–593
135. Liu HY, Liu SM, Zhang YZ (2020) Maternal folic acid supplementation mediates offspring health via DNA methylation. *Reprod Sci* 27(4):963–976
136. Krude H, Biebermann H, Luck W, Horn R, Brabant G, Gruters A (1998) Severe early-onset obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans. *Nat Genet* 19(2):155–157
137. Kuhnen P, Handke D, Waterland RA, Hennig BJ, Silver M, Fulford AJ, Dominguez-Salas P, Moore SE et al (2016) Interindividual variation in DNA methylation at a putative POMC metastable epiallele is associated with obesity. *Cell Metab* 24(3):502–509
138. Paul BD, Sbodio JI, Snyder SH (2018) Cysteine metabolism in neuronal redox homeostasis. *Trends Pharmacol Sci* 39(5):513–524
139. Rehman T, Shabbir MA, Inam-Ur-Raheem M, Manzoor MF, Ahmad N, Liu ZW, Ahmad MH, Siddeeg A et al (2020) Cysteine and homocysteine as biomarker of various diseases. *Food Sci Nutr* 8(9):4696–4707
140. Sanosaka T, Imamura T, Hamazaki N, Chai M, Igarashi K, Ideta-Otsuka M, Miura F, Ito T et al (2017) DNA methylome analysis identifies transcription factor-based epigenomic signatures of multilineage competence in neural stem/progenitor cells. *Cell Rep* 20(12):2992–3003
141. Yu M, Li W, Luo S, Zhang Y, Liu H, Gao Y, Wang X, Wilson JX et al (2014) Folic acid stimulation of neural stem cell proliferation is associated with altered methylation profile of PI3K/Akt/CREB. *J Nutr Biochem* 25(4):496–502
142. Zhang X, Huang G, Liu H, Chang H, Wilson JX (2012) Folic acid enhances Notch signaling, hippocampal neurogenesis, and cognitive function in a rat model of cerebral ischemia. *Nutr Neurosci* 15(2):55–61
143. Wang S, Zeng Y, Pei P, He X, Liu F, Zhang T (2022) Abnormal transcriptome-wide DNA demethylation induced by folate deficiency causes neural tube defects. *Front Genet* 13:987210
144. Desai M, Han G, Li T, Ross MG (2019) Programmed epigenetic DNA methylation-mediated reduced neuroprogenitor cell proliferation and differentiation in small-for-gestational-age offspring. *Neuroscience* 412:60–71
145. Lin N, Qin S, Luo S, Cui S, Huang G, Zhang X (2014) Homocysteine induces cytotoxicity and proliferation inhibition in neural stem cells via DNA methylation in vitro. *FEBS J* 281(8):2088–2096
146. Rabaneda LG, Geribaldi-Doldan N, Murillo-Carretero M, Carrasco M, Martinez-Salas JM, Verastegui C, Castro C (2016) Altered regulation of the Spry2/Dyrk1A/PP2A triad by homocysteine impairs neural progenitor cell proliferation. *Biochim Biophys Acta* 1863(12):3015–3026
147. Shyh-Chang N, Locasale JW, Lyssiotis CA, Zheng Y, Teo RY, Ratanasirintraoort S, Zhang J, Onder T et al (2013) Influence of threonine metabolism on S-adenosylmethionine and histone methylation. *Science* 339(6116):222–226
148. Shiraki N, Shiraki Y, Tsuyama T, Obata F, Miura M, Nagae G, Aburatani H, Kume K et al (2014) Methionine metabolism regulates maintenance and differentiation of human pluripotent stem cells. *Cell Metab* 19(5):780–794
149. Sperber H, Mathieu J, Wang Y, Ferreccio A, Hesson J, Xu Z, Fischer KA, Devi A et al (2015) The metabolome regulates the epigenetic landscape during naive-to-primed human embryonic stem cell transition. *Nat Cell Biol* 17(12):1523–1535
150. Fawal MA, Jungas T, Kischel A, Audouard C, Iacovoni JS, Davy A (2018) Cross talk between one-carbon metabolism, Eph signaling, and histone methylation promotes neural stem cell differentiation. *Cell Rep* 23(10):2864–2873 e2867
151. Batie M, Frost J, Frost M, Wilson JW, Schofield P, Rocha S (2019) Hypoxia induces rapid changes to histone methylation and reprograms chromatin. *Science* 363(6432):1222–1226
152. Chakraborty AA, Laukka T, Myllykoski M, Ringel AE, Booker MA, Tolstorukov MY, Meng YJ, Meier SR et al (2019) Histone demethylase KDM6A directly senses oxygen to control chromatin and cell fate. *Science* 363(6432):1217–1222
153. Qian X, Li X, Shi Z, Bai X, Xia Y, Zheng Y, Xu D, Chen F et al (2019) KDM3A senses oxygen availability to regulate PGC-1 α -mediated mitochondrial biogenesis. *Mol Cell* 76(6):885–895 e887
154. Kong SY, Kim W, Lee HR, Kim HJ (2018) The histone demethylase KDM5A is required for the repression of astrocytogenesis and regulated by the translational machinery in neural progenitor cells. *FASEB J* 32(2):1108–1119
155. Tang QY, Zhang SF, Dai SK, Liu C, Wang YY, Du HZ, Teng ZQ, Liu CM (2020) UTX regulates human neural differentiation and dendritic morphology by resolving bivalent promoters. *Stem Cell Reports* 15(2):439–453
156. Jepsen K, Solum D, Zhou T, McEvelly RJ, Kim HJ, Glass CK, Hermanson O, Rosenfeld MG (2007) SMRT-mediated repression of an H3K27 demethylase in progression from neural stem cell to neuron. *Nature* 450(7168):415–419
157. Lin H, Zhu X, Chen G, Song L, Gao L, Khand AA, Chen Y, Lin G et al (2017) KDM3A-mediated demethylation of histone H3 lysine 9 facilitates the chromatin binding of Neurog2 during neurogenesis. *Development* 144(20):3674–3685
158. Laurent B, Ruitu L, Murn J, Hempel K, Ferrao R, Xiang Y, Liu S, Garcia BA et al (2015) A specific LSD1/KDM1A isoform regulates neuronal differentiation through H3K9 demethylation. *Mol Cell* 57(6):957–970
159. Shi Y, Lan F, Matson C, Mulligan P, Whetstine JR, Cole PA, Casero RA, Shi Y (2004) Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119(7):941–953
160. Wang J, Telese F, Tan Y, Li W, Jin C, He X, Basnet H, Ma Q et al (2015) LSD1n is an H4K20 demethylase regulating memory formation via transcriptional elongation control. *Nat Neurosci* 18(9):1256–1264
161. Sun G, Alzayady K, Stewart R, Ye P, Yang S, Li W, Shi Y (2010) Histone demethylase LSD1 regulates neural stem cell proliferation. *Mol Cell Biol* 30(8):1997–2005
162. Hirano K, Namihira M (2017) FAD influx enhances neuronal differentiation of human neural stem cells by facilitating nuclear localization of LSD1. *FEBS Open Bio* 7(12):1932–1942
163. Tapias A, Wang ZQ (2017) Lysine acetylation and deacetylation in brain development and neuropathies. *Genom Proteom Bioinform* 15(1):19–36
164. Uittenbogaard M, Brantner CA, Chiamarello A (2018) Epigenetic modifiers promote mitochondrial biogenesis and oxidative metabolism leading to enhanced differentiation of neuroprogenitor cells. *Cell Death Dis* 9(3):360
165. Farrelly LA, Zheng S, Schrode N, Topol A, Bhanu NV, Bastle RM, Ramakrishnan A, Chan JC et al (2022) Chromatin profiling

- in human neurons reveals aberrant roles for histone acetylation and BET family proteins in schizophrenia. *Nat Commun* 13(1):2195
166. Park J, Lee K, Kim K, Yi SJ (2022) The role of histone modifications: from neurodevelopment to neurodiseases. *Signal Transduct Target Ther* 7(1):217
 167. Peleg S, Sananbenesi F, Zovoilis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC, Cota P, Wittnam JL et al (2010) Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science* 328(5979):753–756
 168. Wellen KE, Hatzivassiliou G, Sachdeva UM, Bui TV, Cross JR, Thompson CB (2009) ATP-citrate lyase links cellular metabolism to histone acetylation. *Science* 324(5930):1076–1080
 169. Mews P, Donahue G, Drake AM, Luczak V, Abel T, Berger SL (2017) Acetyl-CoA synthetase regulates histone acetylation and hippocampal memory. *Nature* 546(7658):381–386
 170. Bulusu V, Tumanov S, Michalopoulou E, van den Broek NJ, MacKay G, Nixon C, Dhayade S, Schug ZT et al (2017) Acetate recapturing by nuclear acetyl-CoA synthetase 2 prevents loss of histone acetylation during oxygen and serum limitation. *Cell Rep* 18(3):647–658
 171. Li X, Yu W, Qian X, Xia Y, Zheng Y, Lee JH, Li W, Lyu J et al (2017) Nucleus-translocated ACSS2 promotes gene transcription for lysosomal biogenesis and autophagy. *Mol Cell* 66(5):684–697 e689
 172. Mews P, Egervari G, Nativio R, Sidoli S, Donahue G, Lombroso SI, Alexander DC, Riesche SL et al (2019) Alcohol metabolism contributes to brain histone acetylation. *Nature* 574(7780):717–721
 173. Bensaad K, Tsuruta A, Selak MA, Vidal MN, Nakano K, Bartrons R, Gottlieb E, Vousden KH (2006) TIGAR, a p53-inducible regulator of glycolysis and apoptosis. *Cell* 126(1):107–120
 174. Zhou W, Zhao T, Du J, Ji G, Li X, Ji S, Tian W, Wang X et al (2019) TIGAR promotes neural stem cell differentiation through acetyl-CoA-mediated histone acetylation. *Cell Death Dis* 10(3):198
 175. Hsieh J, Nakashima K, Kuwabara T, Mejia E, Gage FH (2004) Histone deacetylase inhibition-mediated neuronal differentiation of multipotent adult neural progenitor cells. *Proc Natl Acad Sci U S A* 101(47):16659–16664
 176. Yu IT, Park JY, Kim SH, Lee JS, Kim YS, Son H (2009) Valproic acid promotes neuronal differentiation by induction of proneural factors in association with H4 acetylation. *Neuropharmacology* 56(2):473–480
 177. Stein LR, Imai S (2014) Specific ablation of Nampt in adult neural stem cells recapitulates their functional defects during aging. *EMBO J* 33(12):1321–1340
 178. Zhang H, Ryu D, Wu Y, Gariani K, Wang X, Luan P, D'Amico D, Ropelle ER et al (2016) NAD(+) repletion improves mitochondrial and stem cell function and enhances life span in mice. *Science* 352(6292):1436–1443
 179. Cai Y, Xu L, Xu H, Fan X (2016) SIRT1 and neural cell fate determination. *Mol Neurobiol* 53(5):2815–2825
 180. Calvanese V, Lara E, Suarez-Alvarez B, Abu Dawud R, Vazquez-Chantada M, Martinez-Chantar ML, Embade N, Lopez-Nieva P et al (2010) Sirtuin 1 regulation of developmental genes during differentiation of stem cells. *Proc Natl Acad Sci U S A* 107(31):13736–13741
 181. Prozorovski T, Schulze-Toppfhoff U, Glumm R, Baumgart J, Schroter F, Ninnemann O, Siegert E, Bendix I, Brustle O et al (2008) Sirt1 contributes critically to the redox-dependent fate of neural progenitors. *Nat Cell Biol* 10(4):385–394
 182. Wang W, Esbensen Y, Kunke D, Suganthan R, Rachek L, Bjoras M, Eide L (2011) Mitochondrial DNA damage level determines neural stem cell differentiation fate. *J Neurosci* 31(26):9746–9751
 183. Sanchez-Ramirez E, Ung TPL, Alarcon Del Carmen A, del Toro-Rios X, Fajardo-Orduna GR, Noriega LG, Cortes-Morales VA, Tovar AR, Montesinos JJ, Orozco-Solis R et al (2022) Coordinated metabolic transitions and gene expression by NAD⁺ during adipogenesis. *J Cell Biol* 221(12). <https://doi.org/10.1083/jcb.202111137>
 184. Lanekoff I, Sharma VV, Marques C (2022) Single-cell metabolomics: where are we and where are we going? *Curr Opin Biotechnol* 75:102693
 185. Liu Q, Martínez-Jarquín S, Zenobi R (2023) Recent advances in single-cell metabolomics based on mass spectrometry. *CCS Chemistry* 5(2):310–324
 186. Qi M, Philip MC, Yang N, Sweedler JV (2018) Single cell neurometabolomics. *ACS Chem Neurosci* 9(1):40–50
 187. Saunders KDG, Lewis H-M, Beste DJV, Cexus O, Bailey MJ (2023) Spatial single cell metabolomics: current challenges and future developments. *Curr Opin Chem Biol* 75:102327
 188. Do TD, Comi TJ, Dunham SJ, Rubakhin SS, Sweedler JV (2017) Single cell profiling using ionic liquid matrix-enhanced secondary ion mass spectrometry for neuronal cell type differentiation. *Anal Chem* 89(5):3078–3086
 189. Cappuccio G, Khalil SM, Osenberg S, Li F, Maletic-Savatic M (2023) Mass spectrometry imaging as an emerging tool for studying metabolism in human brain organoids. *Front Mol Biosci* 10:1181965
 190. Zipfel WR, Williams RM, Christie R, Nikitin AY, Hyman BT, Webb WW (2003) Live tissue intrinsic emission microscopy using multiphoton-excited native fluorescence and second harmonic generation. *Proc Natl Acad Sci U S A* 100(12):7075–7080
 191. Heikal AA (2010) Intracellular coenzymes as natural biomarkers for metabolic activities and mitochondrial anomalies. *Biomark Med* 4(2):241–263
 192. Huang S, Heikal AA, Webb WW (2002) Two-photon fluorescence spectroscopy and microscopy of NAD(P)H and flavoprotein. *Biophys J* 82(5):2811–2825
 193. Chance B, Cohen P, Jobsis F, Schoener B (1962) Intracellular oxidation-reduction states in vivo. *Science* 137(3529):499–508
 194. Georgakoudi I, Quinn KP (2023) Label-free optical metabolic imaging in cells and tissues. *Annu Rev Biomed Eng* 25(1):413–443
 195. Kolenc OI, Quinn KP (2019) Evaluating cell metabolism through autofluorescence imaging of NAD(P)H and FAD. *Antioxid Redox Signal* 30(6):875–889
 196. Débarre D, Supatto W, Pena AM, Fabre A, Tordjmann T, Combettes L, Schanne-Klein MC, Beaurepaire E (2006) Imaging lipid bodies in cells and tissues using third-harmonic generation microscopy. *Nat Methods* 3(1):47–53
 197. Farrar MJ, Wise FW, Fetcho JR, Schaffer CB (2011) In vivo imaging of myelin in the vertebrate central nervous system using third harmonic generation microscopy. *Biophys J* 100(5):1362–1371
 198. Weigelin B, Bakker G-J, Friedl P (2016) Third harmonic generation microscopy of cells and tissue organization. *J Cell Sci* 129(2):245–255
 199. Bird DK, Yan L, Vrotsos KM, Eliceiri KW, Vaughan EM, Keely PJ, White JG, Ramanujam N (2005) Metabolic mapping of MCF10A human breast cells via multiphoton fluorescence lifetime imaging of the coenzyme NADH. *Cancer Res* 65(19):8766–8773
 200. Skala MC, Riching KM, Gendron-Fitzpatrick A, Eickhoff J, Eliceiri KW, White JG, Ramanujam N (2007) In vivo multiphoton microscopy of NADH and FAD redox states, fluorescence lifetimes, and cellular morphology in precancerous epithelia. *Proc Natl Acad Sci* 104(49):19494–19499
 201. Stringari C, Cinquin A, Cinquin O, Digman MA, Donovan PJ, Gratton E (2011) Phasor approach to fluorescence

- lifetime microscopy distinguishes different metabolic states of germ cells in a live tissue. *Proc Natl Acad Sci U S A* 108(33):13582–13587
202. Stringari C, Nourse JL, Flanagan LA, Gratton E (2012) Phasor fluorescence lifetime microscopy of free and protein-bound NADH reveals neural stem cell differentiation potential. *PLoS ONE* 7(11):e48014
 203. Stringari C, Sierra R, Donovan PJ, Gratton E (2012) Label-free separation of human embryonic stem cells and their differentiating progenies by phasor fluorescence lifetime microscopy. *J Biomed Opt* 17(4):046012
 204. Yaseen MA, Sakadzic S, Wu W, Becker W, Kasischke KA, Boas DA (2013) In vivo imaging of cerebral energy metabolism with two-photon fluorescence lifetime microscopy of NADH. *Biomed Opt Express* 4(2):307–321
 205. Aguilar-Arnal L, Ranjit S, Stringari C, Orozco-Solis R, Gratton E, Sassone-Corsi P (2016) Spatial dynamics of SIRT1 and the subnuclear distribution of NADH species. *Proc Natl Acad Sci U S A* 113(45):12715–12720
 206. Li D, Zheng W, Qu JY (2008) Time-resolved spectroscopic imaging reveals the fundamentals of cellular NADH fluorescence. *Opt Lett* 33(20):2365–2367
 207. Cleland NRW, Al-Juboori SI, Dobrinskikh E, Bruce KD (2021) Altered substrate metabolism in neurodegenerative disease: new insights from metabolic imaging. *J Neuroinflammation* 18(1):248
 208. König K, Uchugonova A, Gorjup E (2011) Multiphoton fluorescence lifetime imaging of 3D-stem cell spheroids during differentiation. *Microsc Res Tech* 74(1):9–17
 209. Liu Z, Pouli D, Alonzo CA, Varone A, Karaliota S, Quinn KP, Munger K, Karalis KP et al (2018) Mapping metabolic changes by noninvasive, multiparametric, high-resolution imaging using endogenous contrast. *Sci Adv* 4(3):eaap9302
 210. Meleshina AV, Dudenkova VV, Shirmanova MV, Shcheslavskiy VI, Becker W, Bystrova AS, Cherkasova EI, Zagaynova EV (2016) Probing metabolic states of differentiating stem cells using two-photon FLIM. *Sci Rep* 6:21853
 211. Smith J, Ladi E, Mayer-Proschel M, Noble M (2000) Redox state is a central modulator of the balance between self-renewal and differentiation in a dividing glial precursor cell. *Proc Natl Acad Sci U S A* 97(18):10032–10037
 212. Ito K, Hirao A, Arai F, Matsuoka S, Takubo K, Hamaguchi I, Nomiyama K, Hosokawa K et al (2004) Regulation of oxidative stress by ATM is required for self-renewal of haematopoietic stem cells. *Nature* 431(7011):997–1002
 213. Hamanaka RB, Chandel NS (2010) Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes. *Trends Biochem Sci* 35(9):505–513
 214. Pervaiz S, Taneja R, Ghaffari S (2009) Oxidative stress regulation of stem and progenitor cells. *Antioxid Redox Signal* 11(11):2777–2789
 215. Stringari C, Abdeladim L, Malkinson G, Mahou P, Solinas X, Lamarre I, Brizion S, Galey JB et al (2017) Multicolor two-photon imaging of endogenous fluorophores in living tissues by wavelength mixing. *Sci Rep* 7(1):3792
 216. Ung TPL, Lim S, Solinas X, Mahou P, Chessel A, Marionnet C, Bornschlöggl T, Beaurepaire E et al (2021) Simultaneous NAD(P)H and FAD fluorescence lifetime microscopy of long UVA-induced metabolic stress in reconstructed human skin. *Sci Rep* 11(1):22171
 217. Clow PA, Du M, Jillette N, Taghbalout A, Zhu JJ, Cheng AW (2022) CRISPR-mediated multiplexed live cell imaging of non-repetitive genomic loci with one guide RNA per locus. *Nat Commun* 13(1):1871
 218. Guo DG, Wang DB, Liu C, Lu S, Hou Y, Zhang XE (2019) CRISPR-based genomic loci labeling revealed ordered spatial organization of chromatin in living diploid human cells. *Biochim Biophys Acta Mol Cell Res* 1866(12):118518
 219. Sato Y, Kujirai T, Arai R, Asakawa H, Ohtsuki C, Horikoshi N, Yamagata K, Ueda J et al (2016) A genetically encoded probe for live-cell imaging of H4K20 monomethylation. *J Mol Biol* 428(20):3885–3902
 220. van Staalduinen J, van Staveren T, Grosveld F, Wendt KS (2023) Live-cell imaging of chromatin contacts opens a new window into chromatin dynamics. *Epigenetics Chromatin* 16(1):27

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.