



**HAL**  
open science

## Assessment of the in situ biomethanation potential of a deep aquifer used for natural gas storage

Magali Ranchou-Peyruse, Marion Guignard, Pierre Chiquet, Pierre Cézac,  
Anthony Ranchou-Peyruse

### ► To cite this version:

Magali Ranchou-Peyruse, Marion Guignard, Pierre Chiquet, Pierre Cézac, Anthony Ranchou-Peyruse. Assessment of the in situ biomethanation potential of a deep aquifer used for natural gas storage. ARPHA Conference Abstracts, 2023, 6, 10.3897/aca.6.e109175 . hal-04516919

**HAL Id: hal-04516919**

**<https://hal.science/hal-04516919>**

Submitted on 22 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Conference Abstract

# Assessment of the *in situ* biomethanation potential of a deep aquifer used for natural gas storage

Magali Ranchou-Peyruse<sup>‡</sup>, Marion Guignard<sup>§</sup>, Pierre Chiquet<sup>l</sup>, Pierre Cézac<sup>‡</sup>,  
Anthony Ranchou-Peyruse<sup>¶</sup>

<sup>‡</sup> Université de Pau et des Pays de l'Adour - LaTEP - Joint Laboratory SEnGA E2S-UPPA/Teréga, Pau, France

<sup>§</sup> Université de Pau et des Pays de l'Adour - IPREM, Pau, France

<sup>l</sup> Teréga - Geosciences department - Joint Laboratory SEnGA E2S-UPPA/Teréga, Pau, France

<sup>¶</sup> Université de Pau et des Pays de l'Adour - IPREM - Joint Laboratory SEnGA E2S-UPPA/Teréga, Pau, France

Corresponding author: Magali Ranchou-Peyruse ([magali.ranchou-peyruse@univ-pau.fr](mailto:magali.ranchou-peyruse@univ-pau.fr))

Received: 07 Jul 2023 | Published: 17 Oct 2023

Citation: Ranchou-Peyruse M, Guignard M, Chiquet P, Cézac P, Ranchou-Peyruse A (2023) Assessment of the *in situ* biomethanation potential of a deep aquifer used for natural gas storage. ARPHA Conference Abstracts 6: e109175. <https://doi.org/10.3897/aca.6.e109175>

## Abstract

In response to the challenges of sustainable development and the H<sub>2</sub> sector, it is foreseeable that H<sub>2</sub> will be stored into geological storage, such as deep aquifers. However, CO<sub>2</sub> evolves in deep aquifers because it may be naturally present there; it may also be a constituent of the stored gas mix, or could even be voluntarily stored in the context of the fight against global warming. Autochthonous microorganisms can consume them as sources of energy and carbon (methanogens, (homo)-acetogens and sulfate-reducers). This was already demonstrated in a previous experiment (Haddad 2022) and under operating conditions (Lobodice, Czech Republic; Smigan 1990).

Understanding these mechanisms and quantifying them appear necessary to assess the modifications generated by this type of microorganisms on the properties of the gas. The methanogenesis reaction ( $\text{CO}_2 \text{ gas} + 4\text{H}_2 \text{ gas} \rightarrow \text{CH}_4 \text{ gas} + 2\text{H}_2\text{O liquid}$ ) induces a lowering of pressure, since 5 gas molecules are transformed into a single gas molecule: CH<sub>4</sub> (water being condensed at subsurface conditions). *In situ* biomethanation technique could represent a potential on several scales larger than conventional catalytic or biological

methanation reactors, due to the very large reservoir volumes involved. Biomethanation in geological reservoirs would enable us to reduce our consumption of fossil fuels, so as not to emit more CO<sub>2</sub>, while meeting the growing energy needs of a region and ensuring its independence from hydrocarbon-producing countries.




Figure 1. [doi](#)

Location map of the sampled sites of the studied deep aquifer used as UGS.

A deep aquifer already used as UGS was selected for this study. Formation waters from 17 control wells in this aquifer (Fig. 1) were sampled to assess the potential activity of indigenous methanogenic populations, as well as sulfate-reducers. Despite relatively low sulfate concentrations for a deep aquifer (0.025-1.35 mM), sulfate reducers were found at all sites targeting and quantifying the *dsrB* gene, which is characteristic of this metabolic group (between  $1.8 \cdot 10^1 \pm 2.0 \cdot 10^0$  and  $1.3 \cdot 10^4 \pm 2.0 \cdot 10^3$  *dsrB* gene copy numbers.mL<sup>-1</sup>). In contrast, methanogenic archaea based on the *mcrA* gene quantification were detected at only 10 of the 17 sites (up to  $4.3 \cdot 10^2 \pm 8.3 \cdot 10^1$  *mcrA* gene copy numbers.mL<sup>-1</sup>). The choice was made to focus the rest of the study on 7 of these 10 sites. The potential for methanogenesis was assessed on cultural tests with formation water alone or supplemented with calcite (CaCO<sub>3</sub>), a mineral present in the formation. Results indicate that initial times and controls are controlled by the sulfate variable, since the latter was not consumed by sulfate-reducers. Biotic trials in the presence of calcite and H<sub>2</sub>/CO<sub>2</sub> (abiotic controls and final times) are logically characterized by higher concentrations of calcite, bicarbonate and calcium, but this is not the case for trials in the presence of H<sub>2</sub> alone. We therefore deduce that methanogenesis took place mainly *via* gaseous CO<sub>2</sub>, but that without the latter, calcite was a source of carbon for lithoautotrophs. Cultures incubated with H<sub>2</sub> as the sole gas phase have the highest methane concentrations, logically associated with the lowest sulfate concentrations (consumed by sulfate-reducers), the lowest *Eh* (probably due to the presence of sulfides) and more alkaline pH values up to 10 (which may have led to precipitation of carbonate and calcium ions). All the sites studied showed sulfate consumption and methane production. Analysis of taxonomic diversity (MiSeq; 16S *rRNA* gene V4-V5) showed the dominance of three genera of sulfate-reducers with

*Thermodesulfovibrio-Desulfovibrio-Desulfotomaculum* and methanogenic populations belonging to the *Methanobacterium* genus.

These initial results show a strong potential of *in situ* biomethanation for the deep aquifer studied. All these experiments were carried out at near-atmospheric pressure, and the results still need to be confirmed and refined in the laboratory under conditions that simulate real-life conditions as closely as possible (rock, pressure, nature of gases).

## Keywords

Deep aquifer, Underground Geological Storage, *in situ* biomethanation, H<sub>2</sub> storage

## Presenting author

Magali Ranchou-Peyruse

## Presented at

ISEB-ISSM 2023 - Poster presentation in Part 2: Impacts from Anthropogenic Activities (Gas storage)

## Acknowledgements

E2S-UPPA and Teréga are acknowledged for funding the joint Laboratory SEnGA.

## Conflicts of interest

The authors have declared that no competing interests exist.

## References

- Haddad PG, et al. (2022) Geological storage of hydrogen in deep aquifers - an experimental multidisciplinary study. *Energy Environ Sci*.
- Smigan P, et al. (1990) Methanogenic bacteria as a key factor involved in changes of town gas stored in an underground reservoir. *FEMS Microbiol Lette*.