



HAL
open science

Role of red cell mass evaluation in myeloproliferative neoplasms with splanchnic vein thrombosis and normal hemoglobin value: a study of the France Intergroupe des Syndromes myeloprolifératifs.

Jean Galtier, Louis Drevon, Yannick Le Bris, Stephane Giraudier, Mathieu Wemeau, Laurence Legros, Damien Luque Paz, François Girodon, Jean-Jacques Kiladjian, Charles Mesguich, et al.

► To cite this version:

Jean Galtier, Louis Drevon, Yannick Le Bris, Stephane Giraudier, Mathieu Wemeau, et al.. Role of red cell mass evaluation in myeloproliferative neoplasms with splanchnic vein thrombosis and normal hemoglobin value: a study of the France Intergroupe des Syndromes myeloprolifératifs.. *Haematologica*, 2024, 10.3324/haematol.2023.284488 . hal-04520499

HAL Id: hal-04520499

<https://hal.science/hal-04520499>

Submitted on 25 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Role of red cell mass evaluation in myeloproliferative neoplasms with splanchnic vein thrombosis and normal hemoglobin value: a study of the France Intergroupe des Syndromes myeloprolifératifs

by Jean Galtier, Louis Drevon, Yannick Le Bris, Stephane Giraudier, Mathieu Wemeau, Laurence Legros, Damien Luque Paz, François Girodon, Jean-Jacques Kiladjian, Charles Mesguich, Marie Parrens, Clemence Mediavilla, Lydia Roy, Alexandre Guy, Olivier Mansier, Jean-Christophe Ianotto, and Chloe James. Collaborative Groups: France Intergroupe des syndromes Myeloprolifératifs

Received: October 27, 2023.

Accepted: January 26, 2024.

Citation: Jean Galtier, Louis Drevon, Yannick Le Bris, Stephane Giraudier, Mathieu Wemeau, Laurence Legros, Damien Luque Paz, François Girodon, Jean-Jacques Kiladjian, Charles Mesguich, Marie Parrens, Clemence Mediavilla, Lydia Roy, Alexandre Guy, Olivier Mansier, Jean-Christophe Ianotto, and Chloe James. Collaborative Groups: France Intergroupe des syndromes Myeloprolifératifs. Role of red cell mass evaluation in myeloproliferative neoplasms with splanchnic vein thrombosis and normal hemoglobin value: a study of the France Intergroupe des Syndromes myeloprolifératifs Haematologica. 2024 Feb 8. doi: 10.3324/haematol.2023.284488 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.

Role of red cell mass evaluation in myeloproliferative neoplasms with splanchnic vein thrombosis and normal hemoglobin value: a study of the France Intergroupe des Syndromes myeloprolifératifs

Jean Galtier¹, Louis Drevon², Yannick Le Bris³, Stephane Giraudier², Mathieu Wemeau⁴, Laurence Legros⁵, Damien Luque Paz⁶, François Girodon⁷, Jean-Jacques Kiladjian², Charles Mesguich⁸, Marie Parrons⁹, Clémence Mediavilla¹, Lydia Roy¹⁰, Alexandre Guy¹¹, Olivier Mansier^{11,12}, Jean-Christophe Ianotto¹³, Chloe James^{11,12}

¹ Service d'hématologie et thérapie cellulaire, CHU de Bordeaux, France

² Service d'hématologie clinique, Hôpital Saint-Louis, APHP, France

³ Service d'hématologie biologique, CHU de Nantes, France

⁴ Service d'hématologie clinique, CHRU de Lille, France

⁵ Service d'hématologie clinique, APHP-Hôpital Bicêtre, France

⁶ Service d'hématologie biologique, CHU d'Angers, France

⁷ Service d'hématologie biologique, CHU de Dijon, France

⁸ Service de médecine nucléaire, CHU de Bordeaux, France

⁹ Service d'anatomie et cytologie pathologique, CHU de Bordeaux, France

¹⁰ Service d'Hématologie Clinique, Hôpital Universitaire Henri Mondor, AP-HP & Faculté de Santé, UPEC, Créteil, France

¹¹ Service d'hématologie biologique, CHU de Bordeaux, France

¹² Univ. Bordeaux, INSERM, Biologie des maladies cardiovasculaires, U1034, F-33600 Pessac, France

¹³ Service d'hématologie et d'hémostase clinique, CHU de Brest, France

Word count : 1547

Figure : 1 **Table** : 1

Correspondence :

Jean Galtier

Service d'hématologie clinique et de thérapie cellulaire,

Hôpital Haut Leveque, CHU de Bordeaux, France

Mail : jean.galtier@chu-bordeaux.fr

Disclosures : NONE

Acknowledgments : We thank the France Intergroupe des syndrome Myéloprolifératifs (FIM) for the support in the conception of the study

Data sharing statement: data can be available on direct request to the first and last author

Authors contribution: JG, LD, YLB performed the research; JG and CJ designed the research study; JG analyzed the data; JG and CJ wrote the paper; all author made substantial contributions to research design and interpretation of data

KEY POINTS :

- JAK2V617F mutation is common in splanchnic vein thrombosis despite normal hemoglobin/hematocrit value, and precise definition of underlying myeloproliferative disorder is often challenging.
- Red cell mass evaluation allows for detection of true erythrocytosis defining masked polycythemia vera, and thus may be helpful for management and follow up of these patients.

Splanchnic vein thrombosis (SVT) include portal vein thrombosis (PVT), Budd-Chiari syndrome (BCS), and splenic and/or mesenteric vein thrombosis (S/MVT). They are frequently associated with Philadelphia-negative myeloproliferative neoplasms (MPN)¹, with up to 50% of idiopathic SVT cases exhibiting the JAK2V617F mutation². Diagnosing MPN in the context of SVT is challenging because the resultant portal hypertension and subsequent splenomegaly and hypersplenism can lower blood cell counts, obscuring the elevated counts typical of MPN. The routine inclusion of JAK2V617F mutation screening in SVT patient evaluations has enabled the identification of MPNs that lack the usual hematological features, representing about 15% of BCS and PVT cases². These patients, who otherwise would remain undiagnosed, are categorized as having unclassified MPN (MPN-U), leaving the decision to initiate cytoreductive therapy unresolved. Prior to the discovery of the JAK2V617F mutation, demonstrating an increased red cell mass (RCM) through isotopic methods was a key diagnostic criterion for polycythemia vera (PV). According to the 2008 WHO criteria, hemoglobin levels above 18.5 g/dL in males and 16.5 g/dL in females were considered indicative of increased RCM, leading to the discontinuation of RCM measurement in many facilities³. We hypothesized that for patients with JAK2V617F-positive SVT and normal hemoglobin levels, RCM measurement could facilitate the diagnosis of PV, thereby guiding the initiation and adjustment of cytoreductive therapy.

A retrospective study was conducted on 71 patients diagnosed with splanchnic vein thrombosis (SVT), carrying the JAK2V617F mutation but presenting normal hemoglobin and hematocrit levels initially. Data were gathered from eight French medical centers. Patients gave written consent and the study was conducted in accordance with the Declaration of Helsinki. Key characteristics are detailed in **Table 1**. The median age at diagnosis was 44, with females comprising 48% of the cohort. The median interval between SVT diagnosis and JAK2V617F mutation detection was one month [ranges 0 - 144]. Coincident diagnosis of SVT and JAK2V617F mutation was observed in 64% of cases within a three-month margin. The mutation was detected within two years post-SVT in 14 patients, beyond two years in 9 patients, with one case's timeline unspecified. Increased RCM values over 125%, suggesting masked PV, were found in 56% of patients. Notably, average hemoglobin levels and hematocrits were comparable between the masked PV subgroup and non-PV patients. The total plasma volume value was available in 61 cases and was increased in 91% masked PV versus 46% non-PV, with a mean value of 140% vs. 114%, respectively ($p < 0.001$). Splenomegaly was palpable in a higher proportion of masked PV patients. PVT was the most common SVT, found in 78% of patients,

while BCS affected 17%, and porto-sinusoidal vascular disease was present in 4%. JAK2V617F allele burden at diagnosis and erythropoietin (EPO) levels were not significantly different between masked PV and non-PV patients. A bone marrow biopsy was performed and contributive in 32 patients (45%), i.e., 18/40 masked PV and 14/31 non-PV. Panmyelosis, the histology criterion of PV in the WHO 2016 classification, was found in 8/18 masked PV patients and, interestingly, in 2/14 other JAK2V617F patients. The final diagnoses were (masked) PV in 40 patients, ET in 10, PMF in 3, and MPN-U in 18 cases.

Upon diagnosis of SVT, therapeutic-dose anticoagulation was initiated for all but four patients. At the time the JAK2V617F mutation was identified, three patients were prescribed aspirin, and one patient with masked PV did not receive any antithrombotic treatment due to low platelet counts. After the diagnosis of MPN, 87% of patients began cytoreductive treatment after a median period of six months from their SVT diagnosis (ranges 0 – 239): Treatment started within one year of thrombosis for 37 patients, and after one year for 25 patients. The cytoreductive regimen included hydroxyurea for 43 patients, interferon for 18 patients, and ruxolitinib for one patient. Eight patients did not undergo any cytoreductive therapy; this group included two masked PV patients who were treated solely with phlebotomies. The treatment details for one patient remained undisclosed.

The average follow-up duration was 77 months, with a range from 3 to 358 months (interquartile range: 45 – 130 months). During this period, five patients initially diagnosed with conditions other than PV — two with ET and three with MPN-U — progressed to PV 1, 5, 9, and 15 years after their first diagnosis. In two cases, this progression was marked by increased red cell values, meeting the criteria for overt PV. The remaining three patients (two MPN-U and one ET) showed signs of evolving into masked PV as indicated by routine RCM assessments. Secondary myelofibrosis developed in two patients 17 and 19 years post-detection of the JAK2V617F mutation. Venous thrombosis recurred in five patients, involving the portal vein in three and the spleen in two, occurring at various times ranging from 1 to 21 years after their initial SVT diagnosis. At the time of these thrombotic events, three patients were diagnosed with masked PV with normal red cell values, one had progressed from MPN-U to overt PV, and another had MPN-U with normal red cell values. Three of them were receiving cytoreductive therapy, and 4 anticoagulation therapy at the time of the thrombosis recurrence. Eight out of 40 patients with a masked PV diagnosis underwent a second RCM assessment after at least a year of cytoreductive treatment. RCM levels remained stable in two patients, decreased but stayed

above 125% in two others (one of whom was re-evaluated due to SVT recurrence), and slightly increased in two. Only two patients achieved normal RCM levels. The total plasma volume remained elevated across the board (mean 139%, range 125 – 150%). Those with persistently high RCM were considered under-treated, prompting significant therapeutic adjustments, including the addition of phlebotomies for two patients, increased dosages of cytoreductive agents for another two, and a switch in cytoreductive agents for the remaining two. The progression of red cell values, RCM, and associated treatment modifications for these eight patients are detailed in **Figure 1A**.

Our study presents the largest known cohort of patients with JAK2-mutated SVT evaluated for RCM. A key finding is that over half of these patients were diagnosed with masked PV, despite normal hemoglobin and hematocrit levels. Among 18 patients with masked PV who underwent bone marrow biopsy, only 8 exhibited panmyelosis, which is a major WHO 2016 criterion for PV. Bone marrow biopsies are particularly challenging for JAK2V617F patients with SVT due to the need for therapeutic-dose anticoagulation and possible thrombocytopenia from portal hypertension. Therefore, we advocate for routine RCM evaluation in JAK2V617F-mutated SVT patients to accurately classify the underlying MPN and to ensure appropriate cytoreductive treatment.

Concerns have been raised in several studies about the reliability of hemoglobin and hematocrit as indicators of RCM in PV⁵⁻⁹. The notion of masked PV emerged upon observing patients with JAK2V617F mutation, increased hematocrit and hemoglobin values below WHO 2008 criteria, and heightened RCM. Hemodilution, indicated by a total plasma volume exceeding 110%, is far more prevalent in masked PV than in overt PV¹⁰. Absence of cytoreductive therapy can lead to the undertreatment of masked PV, heightening thrombosis risks compared to overt PV¹¹. These insights contributed to the revision of the WHO 2016 PV criteria, lowering the required hemoglobin levels to 16.5g/dl for men and 16g/dl for women¹². However, data indicates that 25-50% of PV patients defined by an RCM greater than 125% still present with normal red cell counts, posing a risk of PV underdiagnosis and subsequent undertreatment^{9,13}. This is especially concerning in JAK2V617F-positive SVT, where thrombosis occurrence requires cytoreductive treatment and hemodilution is frequent.

In our patient group, 91% of those with masked PV had increased plasma volume, significantly higher than that of non-PV patients, suggesting that hemodilution may play a key role in differentiating

between masked and overt PV post-SVT. This aligns with a previous cohort of JAK2-mutated SVT patients, where less than 10% had raised red cell counts per WHO 2008, yet about 60% had RCM over 125%¹⁴. In our follow-up, RCM remained high in 6 out of 8 evaluated masked PV patients despite cytoreductive therapy and normal red cell values, indicating that red cell counts alone may not suffice for adjusting cytoreductive therapy dosages. Furthermore, 3 of the 5 SVT recurrence cases occurred in patients with masked PV under anticoagulation therapy, who had normal red cell values at recurrence. Of these, one patient had subsequent RCM evaluation which exceeded the mean normal predicted value by 25%. Additionally, 3 of 4 patients with normal RCM at MPN diagnosis who had a second evaluation showed progression toward masked PV. Collectively, these findings emphasize that red cell values are not reliable indicators of RCM at diagnosis or during the follow-up of SVT-associated MPN, owing to the prevalent hemodilution. Current treatment guidelines, which do not recommend cytoreduction for patients with normal hemoglobin and advise keeping hematocrit below 45%, may inadvertently result in the undertreatment of masked PV within the SVT context¹⁵. A limitation of our study is the accessibility of isotopic RCM evaluation; however, the CO rebreathing technique, which shows promising correlation with isotopic methods, may broaden the feasibility of RCM assessment globally¹⁶.

In summary, our findings advocate for routine RCM evaluation in patients diagnosed with JAK2V617F-mutated SVT who have normal red cell values to detect masked PV. Monitoring RCM during treatment may also inform more personalized hematocrit thresholds (as depicted in Figure 1B). Further research is necessary to solidify RCM's role in managing SVT. Results from an ongoing FIM study on the effects of cytoreductive agents in MPNs with normal blood counts (NCT04539678) are forthcoming.

REFERENCES :

1. Stefano V, Qi X, Betti S, Rossi E. Splanchnic vein thrombosis and myeloproliferative neoplasms: molecular-driven diagnosis and long-term treatment. *Thromb Haemost.* 2016;115(2):240-249.
2. Smalberg JH, Arends LR, Valla DC, Kiladjian J-J, Janssen HLA, Leebeek FWG. Myeloproliferative neoplasms in Budd-Chiari syndrome and portal vein thrombosis: a meta-analysis. *Blood.* 2012;120(25):4921-4928.
3. Vardiman JW, Thiele J, Arber DA, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood.* 2009;114(5):937-951.
4. Sant'Antonio E, Guglielmelli P, Pieri L, et al. Splanchnic vein thromboses associated with myeloproliferative neoplasms: An international, retrospective study on 518 cases. *Am J Hematol.* 2020;95(2):156-166.
5. Silver RT, Chow W, Orazi A, Arles SP, Goldsmith SJ. Evaluation of WHO criteria for diagnosis of polycythemia vera: a prospective analysis. *Blood.* 2013;122(11):1881-1886.
6. Alvarez-Larrán A, Ancochea A, Angona A, et al. Red cell mass measurement in patients with clinically suspected diagnosis of polycythemia vera or essential thrombocythemia. *Haematologica.* 2012;97(11):1704-1707.
7. Johansson PL, Safai-Kutti S, Kutti J. An elevated venous haemoglobin concentration cannot be used as a surrogate marker for absolute erythrocytosis: a study of patients with polycythaemia vera and apparent polycythaemia. *Br J Haematol.* 2005;129(5):701-705.
8. Grenier M, Callegarin D, Nughe M, Gardie B, Riedinger JM, Girodon F. Can absolute polycythaemia be identified without measurement of the red cell mass? *Br J Haematol.* 2020;190(2):e107-e110.
9. Hasselbalch HC. Time for revival of the red blood cell count and red cell mass in the differential diagnosis between essential thrombocythemia and polycythemia vera? *Haematologica.* 2019;104(11):2119-2125.
10. Maslah N, Soret J, Dosquet C, et al. Masked polycythemia vera: analysis of a single center cohort of 2480 red cell masses. *Haematologica.* 2020;105(3):e95-e97.
11. Lussana F, Carobbio A, Randi ML, et al. A lower intensity of treatment may underlie the increased risk of thrombosis in young patients with masked polycythaemia vera. *Br J Haematol.* 2014;167(4):541-546.
12. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016;127(20):2391-2405.
13. Silver RT, Krichevsky S. Distinguishing essential thrombocythemia JAK2V617F from polycythemia vera: limitations of erythrocyte values. *Haematologica.* 2019;104(11):2200-2205.

14. Kiladjian J-J, Cervantes F, Leebeek FWG, et al. The impact of JAK2 and MPL mutations on diagnosis and prognosis of splanchnic vein thrombosis: a report on 241 cases. *Blood*. 2008;111(10):4922-4929.
15. Finazzi G, De Stefano V, Barbui T. Splanchnic vein thrombosis in myeloproliferative neoplasms: treatment algorithm 2018. *Blood Cancer J*. 2018;8(7):64.
16. Maaziz N, Georges M, Basille D, et al. PB2165: THE CO-REBREATHING METHOD IS A RELIABLE TEST IN DETERMINING THE RED CELL MASS: THE OLD POTS MAKE THE BEST SOUPS. *Hemasphere* 2023;7(Suppl):e993538d.

TABLE 1 : patients characteristics. RCM : red cell mass ; PV : polycythemia vera.

	Cohort (n = 71)	Normal RCM (n = 31)	Masked PV (n = 40)	p value
Median age	44	44	44	.
Female	34/71	15/31	19/40	.
Hemoglobin (g/dl), mean (ranges)	14 [9.1 – 16.3]	13.8 [12.1 – 16]	14,1 [9.1 – 16.3]	0.24
Hematocrit, mean (ranges)	42% [34 – 47.4]	41.4% [34 – 47]	42.5% [29 – 47.4]	0.15
Platelets (G/L), mean (ranges)	310 [61 – 836]	324 [199 - 639]	304 [61 – 836]	0.42
Neutrophils (G/L), mean (ranges)	5.46 [1.7 – 13]	5.86 [2.4 – 13]	5.17 [1.7 – 11.4]	0.27
RCM, mean (ranges)	135% [99 – 209]	110% [99 – 122]	149% [128 – 209]	< 0.001
Plasma volume, mean (ranges)	128% [88 – 236]	114% [88 – 180]	140% [104 – 236]	< 0.001
Palpable splenomegaly	48/67	16/29	32/38	.
BM biopsy, hyperplasia				
0-2 lineage	22/32	12/14	10/18	
3 lineages	10/32	2/14	8/18	
EPO value, mean	5.9	7.5	5	0.27
JAK2V617F allele burden, mean (ranges)	20% [1 – 68]	15% [5 – 67]	24% [1 – 68]	0.08
Type of splanchnic vein thrombosis:				
- Portal	55/70 (78%)	24/31 (77%)	31/39 (80%)	.
- Budd-Chiari	12/70 (17%)	7/31 (23%)	5/39 (12%)	.
- Porto-sinusoidal vascular disease	3/70 (4%)	0/31 (0%)	3/39 (8%)	.
				.

FIGURE 1: Role of RCM in MPN with SVT and normal red cell values **A:** Evolution of red cell mass under cytoreductive therapy for eight patients. The therapeutic adaptations that were made after second red cell mass evaluation are reported on the right **B:** Proposed strategy for diagnosis and follow-up of non-cirrhotic SVT with normal red cell value at diagnosis.


ET: essential thrombocythemia; MPN-U: myeloproliferative neoplasm – undetermined; PMF: primary myelofibrosis; PV: polycythemia vera; RCM: Red cell mass; SVT : splanchnic vein thrombosis

A.

First red cell mass

Second red cell mass

Change in treatment



Patient	Cytoreductive agent increase	Addition of phlebotomy	Cytoreductive agent switch	None
---------	------------------------------	------------------------	----------------------------	------

Patient 2	X			
-----------	---	--	--	--

Patient 4	X			
-----------	---	--	--	--

Patient 5		X		
Patient 8			X	
Patient 7		X		
Patient 3			X	

Patient 1				X
Patient 6				X

B.

