



## The stringent response is strongly activated in the antibiotic producing strain, *Streptomyces coelicolor*

Clara Lejeune, David Cornu, Laila Sago, Virginie Redeker, Marie-Joelle Virolle

### ► To cite this version:

Clara Lejeune, David Cornu, Laila Sago, Virginie Redeker, Marie-Joelle Virolle. The stringent response is strongly activated in the antibiotic producing strain, *Streptomyces coelicolor*. Research in Microbiology, In press, pp.104177. 10.1016/j.resmic.2023.104177 . hal-04522251

HAL Id: hal-04522251

<https://hal.science/hal-04522251>

Submitted on 26 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ELSEVIER

Contents lists available at ScienceDirect

## Research in Microbiology

journal homepage: [www.elsevier.com/locate/resmic](http://www.elsevier.com/locate/resmic)

## Original article

The stringent response is strongly activated in the antibiotic producing strain, *Streptomyces coelicolor*Clara Lejeune <sup>a</sup>, David Cornu <sup>a</sup>, Laila Sago <sup>a</sup>, Virginie Redeker <sup>a,b</sup>, Marie-Joelle Virolle <sup>a,\*</sup><sup>a</sup> Institute for Integrative Biology of the Cell (I2BC), Université Paris-Saclay, CEA, CNRS, Gif-sur-Yvette, France<sup>b</sup> Institut François Jacob, Molecular Imaging Center (MIRCen), Laboratory of Neurodegenerative Diseases, Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), Centre National de la Recherche Scientifique, Université Paris-Saclay, Fontenay-aux-Roses, France

## ARTICLE INFO

## Article history:

Received 19 September 2023

Accepted 19 December 2023

Available online xxx

## Keywords:

*Streptomyces*

Translational apparatus

Stringent response

ppGpp

Antibiotic biosynthesis

Regulation

## ABSTRACT

*S. lividans* and *S. coelicolor* are phylogenetically closely related strains with different abilities to produce the same specialized metabolites. Previous studies revealed that the strong antibiotic producer, *S. coelicolor*, had a lower ability to assimilate nitrogen and phosphate than the weak producer, *Streptomyces lividans*, and this resulted into a lower growth rate. A comparative proteomic dataset was used to establish the consequences of these nutritional stresses on the abundance of proteins of the translational apparatus of these strains, grown in low and high phosphate availability. Our study revealed that most proteins of the translational apparatus were less abundant in *S. coelicolor* than in *S. lividans* whereas it was the opposite for ET-Tu 3 and a TrmA-like methyltransferase. The expression of the latter being known to be under the positive control of the stringent response whereas that of the other ribosomal proteins is under its negative control, this indicated the occurrence of a strong activation of the stringent response in *S. coelicolor*. Furthermore, in *S. lividans*, ribosomal proteins were more abundant in phosphate proficiency than in phosphate limitation suggesting that a limitation in phosphate, that was also shown to trigger RelA expression, contributes to the induction of the stringent response.

© 2023 Published by Elsevier Masson SAS on behalf of Institut Pasteur.

## 1. Introduction

*S. coelicolor* (SC) and *S. lividans* (SL) are phylogenetically closely related model strains possessing the same specialized metabolite biosynthetic pathways but these are mainly highly expressed in SC [1]. Consequently, the comparative study of these two strains is useful to get a better understanding of what triggers antibiotic biosynthesis in *Streptomyces* species. Previous comparative studies revealed that SC has a lower growth rate than SL [2,3]. This might be due, at least in part, to the fact that SC has a lower ability than SL to assimilate nitrogen (N) as well as phosphate (P) and this results into nutritional stress [1,3,4]. In most microorganisms, N stress is known to trigger the stringent response that leads to the inhibition of the synthesis of ribosomal components [5]. Indeed, N limitation

leads to amino acid starvation and thus to the depletion of aminoacylated (charged) tRNA pools, resulting in the accumulation of ribosomes stalled at the ribosomal site A with non-aminoacylated (uncharged) tRNA [6,7]. The ppGpp synthetase RelA is then recruited to stalled ribosomes and activated to synthesize (p) ppGpp from GTP [8,9]. ppGpp is the mediator of the stringent response and controls at the transcriptional and translational levels a broad range of metabolic processes that allows the adaptation of the bacteria to nutritional limitation and is correlated with a reduction of growth rate.

In *Escherichia coli*, as in other microorganisms, the expression of genes involved in r-RNA and r-protein biosynthesis is down-regulated in condition of stringent control. Indeed, ribosome biogenesis and assembly involve the transcription, processing, folding, and modification of r-RNA and r-proteins and the binding of the latter to r-RNAs. The completion of functional ribosomes is facilitated by a broad range of proteins: GTPases, RNA helicases, endonucleases, modification enzymes, molecular chaperones etc ... that assist proper r-RNAs and r-proteins folding and proper r-proteins-r-RNA interactions [10]. Subsequently, the translational process itself that involves initiation [11], elongation [12], termination

Abbreviations: SC, *Streptomyces coelicolor*; SL, *Streptomyces lividans*.

\* Corresponding author.

E-mail addresses: [lejeune.clara@orange.fr](mailto:lejeune.clara@orange.fr) (C. Lejeune), [david.cornu@i2bc.paris-saclay.fr](mailto:david.cornu@i2bc.paris-saclay.fr) (D. Cornu), [laila.sago@i2bc.paris-saclay.fr](mailto:laila.sago@i2bc.paris-saclay.fr) (L. Sago), [virginie.redeker@cnrs.fr](mailto:virginie.redeker@cnrs.fr) (V. Redeker), [marie-joelle.virolle@i2bc.paris-saclay.fr](mailto:marie-joelle.virolle@i2bc.paris-saclay.fr) (M.-J. Virolle).

steps as well as the recycling of ribosomal components, requires the interaction and correct positioning of the ribosomal particles on mRNA (initiation), the synthesis of tRNAs, their charging by a specific amino acid by aminoacyl-tRNA synthetases [13] and their possible modification. The elongation step requires a correct codon–anticodon interaction and involves the peptidyl transferase center of the ribosome as well as elongation factors for the synthesis of peptidyl-tRNA. At last, when a stop codon is encountered, termination factors hydrolyze the ester bond of the peptidyl-tRNA and release the nascent protein that has to be correctly folded [14]. In *E. coli*, the synthesis of most ribosomal protein (r-protein) is coordinated with that of rRNA by mechanisms acting both at the transcriptional and translational levels [15–18]. ppGpp binds directly to the RNA polymerase [19] and in association with DksA regulates the transcriptional activity of r-proteins promoters [20]. Furthermore, the 5' region of the r-protein mRNA is often the target for translational feedback regulation [15]. ppGpp also inhibits the activity of several enzymes involved at key stages of mRNA translation including initiation, elongation and termination [21] and alters the formation of active 70S ribosomes by inhibiting small GTPases involved in ribosome maturation [22–24]. Altogether, these effects lead to a drastic reduction of the translation rate and thus of protein synthesis resulting into a reduced growth rate [25,26].

In Actinobacteria, as in other Gram+, the regulation of the stringent response slightly differs from that of *E. coli*. In Gram+ bacteria, the signal molecules ppGpp and c-di-GMP that regulate morphological and metabolic differentiation, both originate from GTP [27]. The lowering of intracellular GTP levels, linked to the biosynthesis of these molecules, might lead to a reduction of the transcription of genes requiring GTP as the initiating nucleotide, such as rRNA genes [27]. Furthermore, the biosynthesis of ppGpp was shown to be necessary for the production of antibiotics in SC as well as in other *Streptomyces* species [28–31]. Indeed, the inactivation of the two genes encoding the enzymes known to modulate the levels of ppGpp in SC, *sco1513* encoding the ppGpp synthetase RelA and *sco5794* encoding the ppGpp synthetase/hydrolase RshA/SpoT, abolishes the synthesis of ppGpp as well as that of the blue polyketide antibiotic, actinorhodin (ACT) under nitrogen-limiting conditions [32–34]. The production of these metabolites could thus be considered as part of the stringent response and the nature of the link between these two processes is discussed.

In order to determine whether the stringent response was more activated in SC than in SL, data corresponding to proteins playing a pivotal role in the triggering of the stringent response (RelA, RshA/SpoT and DksA) as well as to those of the translational apparatus were extracted from a previous proteomic dataset of SC and SL, grown in condition of low and high phosphate availability (dataset identifier PXD029263 and 10.6019/PXD029263). This analysis revealed that most proteins of the translational apparatus, whose expression is known to be under the negative control of the stringent response, were less abundant in SC than in SL with the exception of two proteins, ET-Tu 3 (SCO1321) [35] and the TrmA-like methyltransferase (SCO5901) [36], whose expression is known to be under the positive control of the stringent response. These proteins were more abundant in SC than in SL. This suggested the occurrence of a strong activation of the stringent response in SC. Interestingly our data also revealed the up-regulation of RelA expression in condition of Pi limitation and consistently the abundance of most proteins of the translational apparatus was down-regulated in Pi limitation or in other terms up-regulated in condition of Pi proficiency in *S. lividans*. However such regulation did not occur in *S. coelicolor*. These regulatory differences between these two strains are discussed and an explanatory hypothesis are proposed.

## 2. Materials and methods

### 2.1. Bacterial strains, media and culture conditions

Spores of *S. coelicolor* M145 (SC) [37] and *S. lividans* TK24 (SL) [38] used in this analysis were prepared from solid Soya Flour Mannitol (SFM) medium [39]. The 2 strains were grown, in quadruplets, on solid modified R2YE medium, with no sucrose added, on 5 cm diameter Petri dishes. The modified R2YE medium [40] contained glucose (50 mM) as major carbon source and was supplemented or not with K<sub>2</sub>HPO<sub>4</sub>. R2YE not supplemented with K<sub>2</sub>HPO<sub>4</sub> contained 1 mM free Pi originating from the constituents of the growth medium (Pi limitation) and that supplemented with K<sub>2</sub>HPO<sub>4</sub> contains 5 mM Pi (Pi proficiency), as determined with a PiBlue phosphate assay kit (Gentaur, France). Spores (10<sup>6</sup>) of the strains were plated on the surface of cellophane disks (Focus Packaging & Design Ltd, Louth, UK) laid down on the top of agar plates and incubated at 28 °C in darkness for 48 h or 60 h. The time points of 48 and 60 h were chosen as they correspond to the beginning of the production of the blue pigmented polyketide antibiotic, actinorhodin (ACT), in SC [6]. Mycelial lawns of the 4 independent biological replicates of each strain were collected with a spatula, washed twice with deionized water, lyophilized and weighted.

### 2.2. Proteins sample digestions

The 32 protein samples (2 strains x 2 media x 2 culture times x 4 biological replicates) compared in this analysis were prepared as described previously [4] for an in-depth shotgun proteomic analysis and label-free relative quantification. Briefly each protein sample was alkylated, digested using Lysyl-Endopeptidase (Wako Chemicals USA) and sequencing-grade-modified trypsin (Promega). After pre-cleaning, proteolytic peptides were concentrated under vacuum and stored.

### 2.3. Mass spectrometry-based protein identification and relative protein quantification

The mass spectrometry proteomics data reanalyzed in this paper are included in to the dataset identifier PXD029263 and 10.6019/PXD029263 deposited to the ProteomeXchange Consortium via the PRIDE partner repository [41,42]. These data were obtained from 1 µg of proteolytic peptides using nanoLC-MS/MS (liquid chromatography tandem mass spectrometry) with a nano-Elute liquid chromatography system (Bruker) coupled to a timstOF Pro mass spectrometer (Bruker) and included the conversion of raw MS and MS/MS data into mgf files as described previously [43]. Protein identifications against SC and SL protein database from UniprotKB (15012020) using the MASCOT search engine (Matrix science, London, United Kingdom) and changes in protein abundance by label-free mass spectrometry-based quantification using spectral counts or MS1 ion intensities named XIC (for extracted ion current) were performed as described previously [1,4].

### 2.4. Protein abundance changes and statistical analysis of proteins involved in translation

The data corresponding to the two strains, SC and SL, were extracted from the large dataset 10.6019/PXD029263 deposited to the ProteomeXchange. Statistical analyses of protein abundance changes in these two strains was performed in R Studio (version 1.4.17.17) using the homemade R script as described previously [1]. The present analysis thus consisted in a pairwise comparison of 28 combinations and focuses exclusively on proteins involved in the

translational process identified using the UniprotKB (15012020) protein database and the annotations of the KEGG pathway. For each of the identified proteins, a significant difference was set to 10 for spectral count and 1 Log<sub>2</sub> fold change for label-free quantitation (LFQ) values. Additionally a threshold of 4 significant pairs and an adjusted p-value of 0.05 was set to consider a protein abundance as significantly variable. The selected proteins were used for heat-map representation.

### 3. Results

#### 3.1. Proteins involved in the stringent response

Data corresponding to proteins playing a role in the stringent response were extracted from our previous a proteomic dataset of *SC* and *SL*, grown in condition of low and high phosphate availability (dataset identifier PXD029263 and 10.6019/PXD029263). DksA (SCO2075) [44] was not detected in our study but SpoT/RshA (SCO5794) [45] and RelA (SCO1513) were identified. The abundance of SpoT/RshA (SCO5794) was similar in all the conditions tested whereas an up-regulation (1.7 fold) of RelA abundance was noticed at 48 h and 60 h in *SL* and *SC*, respectively, in condition of Pi limitation. This triggering was correlated, as expected, with the down-regulation of most r-proteins in *SL* in Pi limitation whereas the abundance of r-proteins remained low in both Pi conditions in *SC*. The up-regulation of RelA abundance in Pi limitation suggested that its expression might be under the positive control of the two components system PhoR/PhoP (SCO4229/SCO4230) and thus belongs to the Pho regulon [46]. A close inspection of the *relA* promoter region indeed revealed the presence of a rather well conserved Pho box (Fig. S1). This suggests that a Pi limitation contributes to the triggering of the stringent response via the up-regulation of RelA abundance in Pi limitation. A contribution of phosphate limitation in the triggering of the stringent response was previously reported in another Actinobacteria, *Mycobacterium tuberculosis* [47].

#### 3.2. Proteins involved in rRNA modification and ribosome assembly

The 26 proteins, detected in our study that are involved in rRNA modifications, fall into 2 groups (A and B) as shown in Fig. 1:


- Group A includes 7 proteins that had a similar abundance in the 2 strains and whose abundance did not vary significantly with Pi availability nor with time point. This group includes 2 GTPases RsgA-like (SCO5211 and SCO6149) involved in the late steps of the maturation of the functional core of the 30S ribosomal subunit [48] as well as the GTPase HflX (SCO5796) that acts as a ribosome splitting factor that disassembles the 70S ribosomes into its subunits [49]; a methyltransferase type G (SCO3885) of the 16S rRNA that in *Bacillus subtilis* specifically methylates the N7 position of guanine in position 527 of 16S rRNA [50]; a methylase of polypeptide chain release factors (SCO5361) that is required for normal translation termination *in vivo* [51], the ribosome maturation factor RimM (SCO5593) involved in the late assembly stage of the 30S ribosomal subunit [52] and the ribosome silencing factor RsfS (SCO2577) that slows down cell growth by inhibiting protein synthesis during periods of reduced nutrient availability [53].
- Group B includes 19 proteins. Most of these proteins were more abundant in *SL* than in *SC* in both Pi conditions and were up-regulated in Pi proficiency in *SL* but not in *SC*. One noticeable exception was the TrmA-like methyltransferase (SCO5901) that catalyzes the S-adenosylmethionine-dependent methylation of the uracil in position 54 in all *E. coli* tRNAs and is essential for viability [54], whose abundance was higher in *SC* than in *SL*.

Otherwise this group includes 9 methyltransferases [55], 3 GTPases [56], the methylthiotransferase RimO (SCO5752) involved in the methylthiolation of the ribosomal protein S12 [57], 2 proteins involved in the maturation of the 30S sub-unit RimP (SCO5703) [52] and RbfA (SCO5708) [58], the ribosome associated translation inhibitor RaiA (SCO3009) that may play a role in translation fidelity [59] and the ribosome binding ATPase YchF (SCO5627), involved in ribosome recycling [60]. Interestingly the absence of YchF, was shown to enhance the translation of leaderless mRNA in stressful conditions in *E. coli* [61]. The 9 methyltransferases that might be involved in the fine tuning of ribosomal decoding center [62] include 4 putative tRNA/rRNA methyltransferases of the SpoU family (SCO3836, SCO1552, SCO1597 and SCO4236) [63], 4 methyl transferases of the small subunit of the rRNA: RsmC-like (SCO1041) [64], RsmE-like (SCO2552) [65], RsmH-like (SCO2092) [66] and RsmI-like (SCO3253) [67] as well as a rRNA adenine dimethylase of the KsgA family (SCO3149) [68]. The 3 GTPases involved in ribosome assembly and homeostasis [69] belong to the Era (SCO2539) [70], Der/EngA (SCO1758) [71] or Obg (SCO2595) [72] families. Interestingly, in condition of over-expression Obg has a negative impact on aerial mycelium formation in *SC* [73].

#### 3.3. Ribosomal proteins

Ribosomes are made of several rRNA molecules and of 50–80 proteins that play a role in assembly, structure and function of the ribosome [74,75]. The 58 ribosomal proteins detected in our study fall into 4 groups (A to D) as shown in Fig. 2:

- Group A includes 15 proteins (25,8% of total ribosomal proteins) that were more abundant in *SL* than in *SC* especially in Pi proficiency at both time points as well as in Pi limitation at 48 h. This group includes 7 and 8 proteins belonging to the 50S (L4, L5, L10, L19, L23, L27, L28) and 30S (S1, S2, S4, S5, S7, S13, S16, S17) subunits, respectively. Interestingly, the S1 protein was far more abundant in Pi limitation than in Pi proficiency at 48 h in *SL*. In Pi limitation, this protein might play a role in mRNA decay besides its role in translation [76,77].
- Group B includes 22 proteins (37,9%) that were up-regulated in Pi proficiency in *SL* but not in *SC*. These include 16 and 7 proteins belonging to the 50S (L1, L2, L3, L6, L9, L13, L14, L15, L16, L18, L20, L21, L22, L24, L35) and 30S (S3, S8, S9, S11, S12, S17, S19) subunits, respectively.
- Group C includes 8 proteins (13,7%) that were up-regulated in Pi proficiency in a similar way in both strains. These includes 3 and 5 proteins belonging to the 30S (S6, S14 and S18) and 50S (L7/12, L28, L31, L32.2 and L33.1) subunits, respectively.
- Group D includes 11 proteins (18,96%) that had a similar abundance in the two strains and their abundance did not significantly vary with Pi availability. These includes 5 and 6 proteins belonging to the 30S (S8, S10, S15, S18 and S20) and 50S (L11, L25, L28, L29, L30, L33.2) subunits, respectively. The expression pattern of L31 (SCO5359) did not fit into any of the 4 groups. L31 was far more abundant in *SC* than in *SL* in Pi limitation, at both time points. L31 is thought to contribute to sub-units association [78] and to enhance initiation of translation efficiency as well as to reduce frameshifting and miscoding [79]. In *SC* the expression of L31 was shown to be under the positive control of Sigma R [80] that controls the expression of genes involved in the resistance to oxidative/disulfide stress [81]. So the high expression of L31 in *SC* is likely to be due to high oxidative stress linked to the previously reported strong activation of the oxidative metabolism of this strain [1,2,82].



**Fig. 1.** Heatmap representation of the abundance of proteins involved in rRNA modification of *SL* and *SC* grown for 48 h or 60 h in R2YE medium either limited (-Pi, 1 mM) or proficient (+Pi, 5 mM) in phosphate. This heatmap is divided in 2 sub-clusters A and B. Cluster A includes 7 proteins that have a similar abundance in the two strains, conditions and time points. Cluster B includes 19 proteins more abundant in *SL* than in *SC* and up-regulated in Pi proficiency in *SL* but not in *SC*.

The abundance of proteins of group A and B (65.5% of total ribosomal proteins) was up-regulated in condition of Pi proficiency in *SL* but not in *SC*. A similar absence of up-regulation in Pi proficiency was previously reported for some proteins belonging to C and N metabolism in *SC* [1,3,4]. However, obviously proteins of groups C and D escape this regulation and these might constitute a minimal ribosome machinery abundant in *SC* that would be able to translate a subset of proteins involved in the alleviation of the multiple stresses this bacteria is suffering from.


#### 3.4. Enzymes involved in tRNA synthesis or modification

The 48 proteins, detected in our study that are involved in tRNA synthesis or modification, fall into 6 groups (A to F) as shown in Fig. 3:

- Group A includes 6 proteins that were mainly expressed in *SC* in both Pi conditions and at both time points. This group includes Cys, Arg, Phe, Lys and Val tRNA synthetases, enzymes that catalyze the esterification of the corresponding amino acid on the 3' end of tRNA [83], as well as the Glu-tRNAGln amidotransferase subunit B (SCO5501) that allows the formation of correctly charged Gln-tRNA(Gln) through the transamidation of misacylated Glu-tRNA(Gln) when glutaminyl-tRNA synthetase is missing [84].
- Group B includes 4 proteins mainly expressed in *SC* in Pi proficiency at both time points as well as in *SL* in Pi limitation but mainly at 60 h. Its includes the Glu-tRNAGln amidotransferase subunit C (SCO5498), the deacylase Ykak-like (SCO1311) involved in the tRNA editing of mis-charged tRNAs [85] as well as 2 tRNA-specific adenosine deaminases (SCO4038 and

SCO4039). These enzymes catalyze the editing of adenosine into inosine at the wobble position of tRNA<sup>Arg</sup> in prokaryotes [86–88]. This modification is necessary to allow a single tRNA specie to recognize synonymous codons when tRNA specie recognizing the precise codon is poorly available [89].

- Group C includes 7 proteins with a similar abundance in the two strains, Pi conditions and time points. This group includes Thr, Arg, Met and Ileu tRNA synthetases as well as the following tRNA modifying enzymes: tRNA N6-adenosine threonylcarbamoyl-transferase (SCO4752) [90], tRNA (guanine-N(7))-methyltransferase (SCO4111) [91], Met-tRNA formyl transferase (SCO1473) [92] and methionyl-tRNA formyltransferase (SCO5791) [93].
- Group D includes 9 proteins strongly up-regulated in Pi proficiency in *SL* but not in *SC*. It includes Tyr and Lys tRNA synthetases [94], putative archaeal-like Glu-tRNAGln amidotransferase subunit E (SCO3581) and the following tRNA modifying enzymes: the pseudouridine tRNA synthase (SCO1768) that catalyzes a post-translational modification of uridine in some tRNA that might stabilize the correct anticodon-codon pairing during translation [95], tRNA threonylcarbamoyl adenosine modification protein YjeE-like (SCO4747) [90], tRNA (guanine-N1)-methyltransferase (SCO5594) which may which prevent frameshifting [96], ribosomal RNA large subunit methyltransferase N (SCO5645) [97] and tRNA (5-methylaminomethyl-2-thiouridylate)-methyltransferase (SCO5488) [98].
- Group E includes 13 proteins far more abundant in *SL* than in *SC* in both Pi conditions and at both time points. This group includes Pro, Thr, 2 Trp, Lys, Gly, Leu, Ala and Met tRNA-synthetases as well as the following tRNA modifying enzymes: pseudouridine tRNA-synthetase (SCO5709) [95], d-tyrosyl-tRNA(Tyr) deacylase (SCO4182) that removes some d-amino



**Fig. 2.** Heatmap representation of the abundance of ribosomal proteins of *SL* and *SC* grown for 48 h or 60 h in R2YE medium either limited (-Pi, 1 mM) or proficient (+Pi, 5 mM) in phosphate. This heatmap is divided in 4 sub-clusters. Cluster A includes 15 proteins more abundant in *SL* than in *SC* and especially abundant in *SL* at 48 h and 60 h in Pi limitation and proficiency, respectively. Cluster B includes 21 proteins more abundant in *SL* than in *SC* and up-regulated in Pi proficiency in both strains. Cluster C includes 8 proteins similarly abundant in *SL* than in *SC* and up-regulated in Pi proficiency in both strains. Cluster D includes 11 proteins that have a similar abundance in the two strains, conditions and time points. Two proteins S14 type Z (SCO4715) and L31 (SCO5359) do not fit in any of these clusters.

acids from mischarged tRNAs allowing their replacement by the L form used in translation [99,100] and a dimethylallyl adenosine tRNA methylthiotransferase *MiaB* (SCO5787) thought to be involved in improvement of reading frame maintenance [79,80]. Mutation in *miaB* were shown to affect morphological differentiation and secondary metabolism in Streptomyces [81–84]. - Group F includes 6 proteins more abundant in Pi limitation than in Pi proficiency in both strains and more abundant at 48 h and 60 h in *SL* and *SC*, respectively. These proteins were also more abundant in Pi proficiency in *SL* than in *SC*. This group includes Ser, Phe, Glu, Asp, His tRNA synthetases and the pseudouridine synthetase (SCO4731) [95].


The regulation of the expression of aminoacyl-tRNA synthetases in prokaryotes proved to be complex and diverse [101]. Our results indicate that tRNA synthetases of Fig. 3/groups D and E were more abundant in *SL* than in *SC* but in contrast tRNA synthetases of Fig. 3/group A (cys-tRNA ligase (SCO4235), the-tRNA ligase beta subunit (SCO1594), val-tRNA ligase (SCO2615), arg-tRNA ligase (SCO3304) and the Glu-tRNAGln amidotransferase (SCO5501) were more abundant in *SC* than in *SL*.

Even if the expression of most aminoacyl-tRNA ligases/synthetases is coordinated with the synthesis/availability of their cognate amino acids [102] and increases with growth rate [101], up-regulation of tRNA synthetases abundance has been reported in *B. subtilis* when the availability of their substrates is low [101]. In *SC*, a similar regulation might occur indicating the low availability of some amino acids due to the reduced ability of this strain to up-take amino acids present in the R2YE medium.

### 3.5. Proteins involved in translational initiation, elongation and termination processes

The 13 proteins, detected in our study that are involved in translational initiation, elongation and termination, fall into 2 groups (A and B) as shown in Fig. 4:

- The 2 proteins of group A had a similar abundance in the two strains and their abundance did not vary with Pi availability nor with time point. This group includes the translational initiation factors IF1 [103] and IF3 [104,105].



**Fig. 3.** Heatmap representation of the abundance of enzymes involved in tRNA synthesis or modification of *SL* and *SC* grown for 48 h or 60 h in R2YE medium either limited (-Pi, 1 mM) or proficient (+Pi, 5 mM) in phosphate. This heatmap is divided in 6 sub-clusters A to F. Cluster A includes 6 proteins more abundant in *SC* than in *SL* in both Pi conditions and at both time points. Cluster B includes 4 proteins more abundant in *SC* than in *SL* and up-regulated in Pi proficiency in *SC* but not in *SL*. Cluster C includes 7 proteins that have a similar abundance in the two strains, conditions and time points. Cluster D includes 9 proteins more abundant in *SL* than in *SC* and up-regulated in Pi proficiency in *SL* but not in *SC*. Cluster E includes 13 proteins more abundant in *SL* than in *SC* in both Pi conditions and at both time points. Cluster F includes 6 proteins more abundant in *SL* than in *SC* and up-regulated in Pi limitation in both strains.

- Most of the 11 proteins of group B were more abundant in *SL* than in *SC* in both Pi conditions. This group includes the translation initiation factor IF2 [106], 7 elongation factors [12], 2 peptide chain release factors [107] and one peptidyl tRNA hydrolase [108]. However, the elongation factor EF-Tu3 (SCO1321) escapes this rule since that it was far more abundant in *SC* than in *SL* in Pi limitation at both time points.


### 3.6. Proteins involved in protein folding and modification

The 20 proteins, detected in our study and involved in protein folding and modification, fall into 5 groups (A to E) as shown in Fig. 5:


- Group A includes two chaperones DnaJ2 (SCO2554) and HtpG (SCO7516) [109,110] and a peptidyl-prolyl cis-trans isomerase (PPIase, SCO1639) [111] that were up-regulated in Pi limitation in both strains. PPIases assist protein folding via the catalysis of cis/trans isomerization of peptide bonds preceding prolyl residues [112].
- Group B includes 4 proteins, the chaperone (SCO4761), two Heat Shock Proteins (HSP) HspR (SCO3668) [113] and SCO1405 (Hsp90 family) [114] and a peptide deformylase (SCO0883)

[115]. These proteins were up-regulated in Pi limitation in both strains but were far more abundant in *SC* than in *SL*. Peptide deformylases remove the formyl group from the formylmethionine present at the N-terminus end of all nascent peptides [116]. These proteins are thought to contribute to the correct folding of membrane proteins and thus to maintain proper structure and function of the membrane [116]. The over-expression of SCO0883 in *SC* might contribute to fight membrane stress that occurs in *SC* [1,4].

- Group C includes 6 proteins, two chaperones, DnaK (SCO3671) [117] and the ATP-dependent chaperon ClpB (SCO3661) [118], three PPIases (SCO3856, SCO1638 and SCO7510) and a peptide deformylase (SCO5221). These proteins were up-regulated in *SL* in Pi limitation but not in *SC*.
- Group D includes 4 proteins, two chaperones, DnaJ (SCO3669) and SCO2899, the heat shock protein GrpE-like (SCO3670) [119] and a second PPIase (SCO1510). These proteins had a similar abundance in the two strains and their abundance did not vary with Pi availability nor time points.
- Group E includes 3 proteins, the heat shock protein GrpE-like (SCO1771) and two 60 KDa chaperones, GroEL2 (SCO4296) and GroEL1 (SCO4762) [120]. These proteins were up-regulated in *SC* in Pi proficiency at both time points but not in *SL*.



**Fig. 4.** Heatmap representation of the abundance of proteins involved in translation initiation, elongation and termination processes of SL and SC grown for 48 h or 60 h in R2YE medium either limited (-Pi, 1 mM) or proficient (+Pi, 5 mM) in phosphate. This heatmap is divided in 2 sub-clusters A and B. Cluster A includes 2 proteins that have a similar abundance in the two strains, conditions and time points. Cluster B includes 11 proteins more abundant in SL than in SC in both Pi conditions and at the two time points.



**Fig. 5.** Heatmap representation of the abundance of proteins involved in proteins folding and modification enzymes of SL and SC grown for 48 h or 60 h in R2YE medium either limited (-Pi, 1 mM) or proficient (+Pi, 5 mM) in phosphate. This heatmap is divided in 5 sub-clusters A to E. Cluster A includes 3 proteins more abundant in SL than in SC and up-regulated in Pi limitation. Cluster B includes 4 proteins more abundant in SC than in SL and up-regulated in Pi limitation in SC. Cluster C includes 6 proteins more abundant in SL than in SC and up-regulated in Pi limitation in SL but not in SC. Cluster D includes 4 proteins that have a similar abundance in the two strains, conditions and time points. Cluster E includes 3 proteins up-regulated in Pi proficiency in SC but not in SL.

The higher abundance of proteins of groups A and B in both strains or of proteins of group C in SL, in Pi limitation, suggest that these proteins may belong to the Pho regulon.

### 3.7. Relative abundance of proteins translated from leaderless RNA in *S. coelicolor* and *S. lividans*

Our data revealed that most elements of the translational apparatus were less abundant in SC than in SL whereas a few were similarly abundant in the two strains. These few proteins

might thus constitute the core of minimal ribosomal particles that would be more abundant in SC than in SL. Since some reports in the literature mention that minimal ribosomal particles might preferentially translates leaderless mRNAs (lRNAs) [121–123], we uploaded the list of SC's lRNAs from Jeong et al. [124] in order to determine whether they would be more efficiently translated in SC than in SL. Among the 713 lRNAs mentioned in Jeong et al. [124], 368 were detected in our study. These lRNAs could be classified into 6 groups (Figure S2A–F).

Proteins translated from mRNAs of groups A1 (9.5%), A3 (5.1%) and A4 (4.6%) were more abundant in Pi limitation than in Pi proficiency, especially in SC and proteins of group A5 (3.8%) were more abundant in SC than in SL in Pi proficiency. Proteins of group A2 (2.17%) were similarly up-regulated in Pi limitation in both strains. Group A2 includes proteins belonging to the Pho regulon [46,125]: the sensor kinase PhoR (SCO4229), the polyphosphate kinase Ppk (SCO4145) [126] as well as 6 genes encoding proteins involved in the biosynthesis of teichulosonic acid cell wall glycopolymers (SCO4877 to SCO4882) [127,128]. Phosphate free teichulosonic and teichuronic acids cell wall glycopolymers are known to replace phosphate rich teichoic acids in Gram + bacteria in condition of Pi limitation [129].

Proteins translated from mRNAs of groups B and C (35.6% of total mRNAs) were clearly more abundant in SC than in SL. Proteins of group B were more abundant in SC than in SL in both Pi conditions whereas proteins of group C were mainly up-regulated in Pi limitation in SC.

Proteins translated from mRNAs of group D1 (5.1%) were more abundant in Pi proficiency than in Pi limitation in both strains but rather more abundant in SL than in SC. Proteins of group D2 (2.4%) were similarly up-regulated in Pi proficiency in both strains. Proteins of group D3 (9.78%) were also up-regulated in Pi proficiency but mainly in SC.

In contrast proteins belonging to groups E and F (21.46% of total mRNAs) were clearly more abundant in SL than in SC. Proteins of group E (16%) were more abundant in SL than in SC in both Pi conditions. The 8 proteins of group F1 (2.17%) were clearly up-regulated in Pi proficiency in SL but not in SC and the 12 proteins of group F2 (3.26%) were up-regulated in Pi limitation in SL but not in SC. Group F1 includes 3 genes encoding large (SCO2486 and SCO2487) and small (SCO2488) subunits of the nitrite reductase, the heme peroxidase (SCO6042), two transcriptional regulators (SCO3810 and SCO4766) and the phosphoribosyltransferase (SCO3677) that belongs to the purine salvage pathway that is usually up-regulated during the stringent response [130]. Group F2 includes the response regulators SCO3013, located downstream the translation initiation factor SCO3014 and PhoP (SCO4230) that governs adaptation to Pi limitation, the inositol monophosphatase (SCO6445) that might be involved in a phosphophatidylinositol signaling pathway controlling cell growth in *Streptomyces* [131] and the ribokinase (SCO2158) that is involved in the biosynthesis of ribose 5 phosphate, a metabolite proposed to link nucleotide and amino acids metabolism in condition of stringent control [132].

Our data indicate that only 7% of total proteins belonging to groups A2 (up-regulated in Pi limitation), D1 and D2 (up-regulated in Pi proficiency) were similarly regulated in both strains whereas 54.6% of the proteins translated from mRNAs were more abundant in SC than in SL (72 and 59 proteins from groups B and C, 33 proteins from groups A3/A5 and 37 proteins of group D3) when only 27.4 % of the proteins translated from mRNAs were more abundant in SL than in SC (59 and 20 proteins of groups E and F and 14 and 8 proteins of groups D1 and D3). Proteins translated from mRNAs are thus two fold more abundant in SC than in SL but it cannot be concluded with certainty whether this is due to higher transcriptional level of mRNAs or to higher translational rate of the latter's, that might be linked to the higher abundance of minimal ribosomal particles, in SC than in SL.

#### 4. Discussion

The regulatory process called "stringent response" that is aimed at the adaptation of bacterial growth to nutrient limitation is known to be highly conserved throughout the bacterial domain of life [17]. Stringent control is known to regulate negatively the

expression of most ribosomal proteins [25,26]. Our data indicated that the majority of ribosomal proteins (Fig. 2, groups A and B), of proteins involved in rRNA modifications (Fig. 1, group B), of enzymes involved in tRNA synthesis and modification (Fig. 3, groups D and E) and of proteins involved in translation initiation, elongation and termination processes (Fig. 4, group B) were less abundant in SC than in SL. In consequence, the reduced abundance of these proteins in SC is proposed to be due to the strong activation of the stringent response in this strain that is most likely linked to the reduced ability of this strain to assimilate nitrogen as well as phosphate, to a lesser extent [1,3,4]. This hypothesis is supported by the observation that two proteins whose expression is known to be under the positive control of stringent response in other bacteria were over-expressed in SC compared to SL. These are the tRNAM5(U34) methyltransferase TrmA-like (SCO5901) (Fig. 1) [36] and the elongation factor EF-Tu3 (SCO1321) (Fig. 4) [35]. The protein L31 (SCO5359) was also found more abundant in SC than in SL at 48 h and 60 h in Pi limitation. The expression of this protein was shown to be under the positive control of Sigma R that controls the expression of genes involved in the resistance to oxidative/disulfide stress in *Streptomyces* [Paget, 1998 #1191] [80]. So the high expression of L31 in SC ought to be related to high oxidative stress linked to the previously reported strong activation of the oxidative metabolism of this strain [1,2,82]. Interestingly, it was recently demonstrated that ppGpp is also involved in the response to oxidative [133] and heat stresses [134]. ppGpp might thus be involved in the response to any stresses that would lead to a growth slow down mediated by a downregulation of the translational process. Such regulation would also reduce the load on the protein quality control system (including numerous chaperones and proteases) that is induced by these stresses.

Interestingly, the abundance of 31 proteins, 8.4% of total proteins detected (Fig. 1, 7 of group A; Fig. 2, 11 of group D; Fig. 3, 7 of group C; Fig. 4, 2 of group A and Fig. 5, 4 of group D), was similar in the two strains in both Pi conditions. These proteins might thus constitute the core of a minimal translational apparatus that would be relatively more abundant in SC than in SL and might be responsible for the more efficient translation of mRNAs in SC than in SL.

Furthermore, the abundance of most ribosomal proteins (Fig. 2, group B), of proteins involved in rRNA modifications (Fig. 1, group B), of enzymes involved in tRNA synthesis and modification (Fig. 3, groups D and E) and of proteins involved in translation initiation, elongation and termination processes (Fig. 4, group B) was up-regulated in Pi proficiency in SL but such up-regulation did not occur in SC except for group C of ribosomal proteins (Fig. 2). These observations indicate that high Pi availability reduces the intensity of the stringent response in SL or in other terms that Pi limitation contributes to the stringent response. However this regulatory process does not occur in SC since the abundance of most proteins of the translational apparatus remains similarly low in both Pi conditions. We propose that the absence in SC of this phosphate-mediated up-regulation of ribosomal proteins in Pi proficiency might be due a default in external phosphate sensing and/or uptake. This default itself might be due to the strong activation of the stringent response occurring in this strain that would control negatively phosphate assimilation and phosphate-mediated regulation. Indeed, the assimilation of nitrogen (N), phosphate (P) and carbon (C), ought to be coordinated in *Streptomyces* [135], as in all living organisms. The poor ability of SC to assimilate N is correlated with the reduced abundance of proteins playing a role in C up-take and assimilation such as the glucose permease (SCO5578) (Fig. 16B of [1]) and all glycolytic enzymes in SC compared to SL (Fig. 2 of [1]) as well as of proteins of the Pho regulon [3]. The low affinity phosphate transporter PitH1 (SCO4138) is also poorly expressed in

Pi proficiency in SC compared to SL (Group D of Fig. 6 of [1]) and consequently some proteins involved in Pi re-cycling such as the nucleases SCO4152 and SCO1908, the phospholipases SCO6691 and SCO1048, the phosphodiesterases SCO1968 and SCO1565 and the phosphatase SCO2068 were up-regulated in SC compared to SL, in condition of Pi proficiency (Groups A and C of Fig. 6 of [1]). This suggests that despite the abundance of Pi in the growth medium, SC does not up-take it efficiently. SC is thus limited in phosphate, even when Pi is abundant in the growth medium, and is forced to degrade its phosphate-rich constituents in order to recycle phosphate. Altogether our data suggest that nitrogen and phosphate limitation both contribute to the triggering of the stringent response that in turn controls negatively nitrogen, phosphate as well as carbon metabolism to achieve growth reduction in SC. In contrast the stringent response obviously controls positively the expression of some specialized metabolites biosynthetic pathways [28]. Similar regulations were shown to occur in *Pseudomonas protegens* [136].

The production of these specialized metabolites could thus be considered as part of the stringent response. Since the main aim of the stringent response is to reduce growth when essential nutrients, such as N or P, are poorly available, we can consider that some specialized metabolites also contribute to growth slow down. Interestingly ACT, as well as other specialized metabolites of the polyketide family, were proposed to be able to capture electrons of ROS/NOS as well as of the respiratory chain, thanks to their quinone groups [137]. This ability would confer them anti-oxidant and anti-respiratory properties leading to a reduction of ATP generation [137]. Consistently, the onset of ACT production was shown to coincide with an abrupt drop in the intracellular ATP concentration [2] that might contribute to growth slow down.

Further studies are needed to elucidate the complex regulatory network governed by ppGpp in *Streptomyces* but it should be stressed that these studies have great fundamental as well as practical interest since the stringent response has a strong positive impact on the production of bio-active specialized metabolites that have considerable utility in the medical and agricultural fields.

## Declaration of competing interest

The authors declare no conflict of interest.

## Acknowledgments

This research was funded by public Institutions (CNRS and University Paris-Saclay) as well as by the program "Investment for the Future Biotechnologies et Bioresources" n°11-BTBR-0003 (PROBIO3), the ANR BioSound-IR (ANR-15-CE09-0002) and the ANR-17-ASTR-0018 INNOVANTIBIO from the "Direction Générale des Armées" (DGA) et "l'Agence de l'Innovation de Défense" (AID). Proteomic analysis were performed on the proteomic platform of I2BC (Proteomic-Gif, SICAPS) supported by IBISA, Ile de France Région, Plan Cancer, CNRS and Paris-Saclay University.

The authors wish to thank Céline Fabret and Olivier Namy (I2BC) for stimulating discussion and critical reading of the manuscript.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.resmic.2023.104177>.

## References

- [1] Lejeune C, Sago L, Cornu D, Redeker V, Virolle MJ. A proteomic analysis indicates that oxidative stress is the common feature triggering antibiotic production in *Streptomyces coelicolor* and in the *ppmA* mutant of *Streptomyces lividans*. *Front Microbiol* 2022;12:813993.
- [2] Esnault C, Dulermo T, Smirnov A, Askora A, David M, Deniset-Besseau A, et al. Strong antibiotic production is correlated with highly active oxidative metabolism in *Streptomyces coelicolor* M145. *Sci Rep* 2017;7:200.
- [3] Millan-Oropeza A, Henry C, Lejeune C, David M, Virolle MJ. Expression of genes of the *Pho* regulon is altered in *Streptomyces coelicolor*. *Sci Rep* 2020;10:8492.
- [4] Lejeune C, Cornu D, Sago L, Redeker V, Virolle M-J. Comparative proteomic analysis of transcriptional and regulatory proteins abundances in *S. Lividans* and *S. Coelicolor* suggests a link between various stresses and antibiotic production. *Int J Mol Sci* 2022;23(23):14792.
- [5] Brown DR, Barton G, Pan Z, Buck M, Wigneshweraraj S. Nitrogen stress response and stringent response are coupled in *Escherichia coli*. *Nat Commun* 2014;5:4115.
- [6] Goldman E, Jakubowski H. Uncharged tRNA, protein synthesis, and the bacterial stringent response. *Mol Microbiol* 1990;4:2035–40.
- [7] Winther KS, Roghman M, Gerdes K. Activation of the stringent response by loading of RelA-tRNA complexes at the ribosomal A-site. *Mol Cell* 2018;70:95–105 e4.
- [8] Dworkin J. Understanding the stringent response: experimental context matters. *mBio* 2023;14:e0340422.
- [9] Steinchen W, Zegarra V, Bange G. (p)ppGpp: magic modulators of bacterial physiology and metabolism. *Front Microbiol* 2020;11:2072.
- [10] Maksimova E, Kravchenko O, Korepanov A, Stolboushkina E. Protein assistants of small ribosomal subunit biogenesis in bacteria. *Microorganisms* 2022;10.
- [11] Gualerzi CO, Pon CL. Initiation of mRNA translation in bacteria: structural and dynamic aspects. *Cell Mol Life Sci* 2015;72:4341–67.
- [12] Kraal B, Bosch L, Mesters JR, de Graaf JM, Woudt LP, Vijgenboom E, et al. Elongation factors in protein synthesis. *Ciba Found Symp* 1993;176:28–43. discussion 44–52.
- [13] Rubio Gomez MA, Ibba M. Aminoacyl-tRNA synthetases. *RNA* 2020;26:910–36.
- [14] Rodnina MV. Translation in prokaryotes. *Cold Spring Harb Perspect Biol* 2018;10.
- [15] Burgos HL, O'Connor K, Sanchez-Vazquez P, Gourse RL. Roles of transcriptional and translational control mechanisms in regulation of ribosomal protein synthesis in *Escherichia coli*. *J Bacteriol* 2017;199.
- [16] Petibon C, Malik Ghulam M, Catala M, Abou Elela S. Regulation of ribosomal protein genes: an ordered anarchy. *Wiley Interdiscip Rev RNA* 12; 2021. p. e1632.
- [17] Zhu M, Dai X. Stringent response ensures the timely adaptation of bacterial growth to nutrient downshift. *Nat Commun* 2023;14:467.
- [18] Wagner R. Translational components in prokaryotes: genetics and regulation of ribosomes. *Encyclopedia of life sciences*. John Wiley & Sons, Ltd.; 2001. <https://doi.org/10.1002/047001590X>.
- [19] Liu K, Bittner AN, Wang JD. Diversity in (p)ppGpp metabolism and effectors. *Curr Opin Microbiol* 2015;24:72–9.
- [20] Lemke JJ, Sanchez-Vazquez P, Burgos HL, Hedberg G, Ross W, Gourse RL. Direct regulation of *Escherichia coli* ribosomal protein promoters by the transcription factors ppGpp and DksA. *Proc Natl Acad Sci U S A* 2011;108:5712–7.
- [21] Irving SE, Choudhury NR, Corrigan RM. The stringent response and physiological roles of (pp)pGpp in bacteria. *Nat Rev Microbiol* 2021;19:256–71.
- [22] Bennison DJ, Irving SE, Corrigan RM. The impact of the stringent response on TRAFAC GTPases and prokaryotic ribosome assembly. *Cells* 2019;8.
- [23] Zhang X, Yan K, Zhang Y, Li N, Ma C, Li Z, et al. Structural insights into the function of a unique tandem GTPase EngA in bacterial ribosome assembly. *Nucleic Acids Res* 2014;42:13430–9.
- [24] Corrigan RM, Bellows LE, Wood A, Grundling A. ppGpp negatively impacts ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive bacteria. *Proc Natl Acad Sci U S A* 2016;113:E1710–9.
- [25] Burkovski A. Ammonium assimilation and nitrogen control in *Corynebacterium glutamicum* and its relatives: an example for new regulatory mechanisms in actinomycetes. *FEMS Microbiol Rev* 2003;27:617–28.
- [26] Boutte CC, Crosson S. Bacterial lifestyle shapes stringent response activation.  *Trends Microbiol* 2013;21:174–80.
- [27] Sivapragasam S, Grove A. The link between purine metabolism and production of antibiotics in *Streptomyces*. *Antibiotics* 2019;8.
- [28] Hesketh A, Sun J, Bibb M. Induction of ppGpp synthesis in *Streptomyces coelicolor* A3(2) grown under conditions of nutritional sufficiency elicits actII-ORF4 transcription and actinorhodin biosynthesis. *Mol Microbiol* 2001;39:136–44.
- [29] Kang SG, Jin W, Bibb M, Lee KJ. Actinorhodin and undecylprodigiosin production in wild-type and *relA* mutant strains of *Streptomyces coelicolor* A3(2) grown in continuous culture. *FEMS Microbiol Lett* 1998;168:221–6.
- [30] Song Y, Zhang X, Zhang Z, Shentu X, Yu X. Physiology and transcriptional analysis of ppGpp-related regulatory effects in *Streptomyces dia-*statochromogenes 1628. *Microbiol Spectr* 2023;11:e0120022.
- [31] Xu Z, You D, Tang LY, Zhou Y, Ye BC. Metabolic engineering strategies based on secondary messengers (p)ppGpp and C-di-GMP to increase erythromycin yield in *saccharopolyspora erythraea*. *ACS Synth Biol* 2019;8:332–45.
- [32] Chakraburty R, Bibb M. The ppGpp synthetase gene (*relA*) of *Streptomyces coelicolor* A3(2) plays a conditional role in antibiotic production and morphological differentiation. *J Bacteriol* 1997;179:5854–61.

- [33] Ryu YG, Kim ES, Kim DW, Kim SK, Lee KJ. Differential stringent responses of *Streptomyces coelicolor* M600 to starvation of specific nutrients. *J Microbiol Biotechnol* 2007;17:305–12.
- [34] Strauch E, Takano E, Baylis HA, Bibb MJ. The stringent response in *Streptomyces coelicolor* A3(2). *Mol Microbiol* 1991;5:289–98.
- [35] van Wezel GP, Takano E, Vijgenboom E, Bosch L, Bibb MJ. The *tuf3* gene of *Streptomyces coelicolor* A3(2) encodes an inessential elongation factor Tu that is apparently subject to positive stringent control. *Microbiology (Read 1995)*;141(Pt 10):2519–28.
- [36] Ny T, Bjork GR. Stringent regulation of the synthesis of a transfer ribonucleic acid biosynthetic enzyme: transfer ribonucleic acid(m5U)methyltransferase from *Escherichia coli*. *J Bacteriol* 1977;130:635–41.
- [37] Bentley SD, Chater KF, Cerdeno-Tarraga AM, Challis GL, Thomson NR, James KD, et al. Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* 2002;417:141–7.
- [38] Ruckert C, Albersmeier A, Busche T, Jaenicke S, Winkler A, Friethjonsen OH, et al. Complete genome sequence of *Streptomyces lividans* TK24. *J Biotechnol* 2015;199:21–2.
- [39] Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA, editors. Practical *Streptomyces* genetics. John Innes Foundation; 2000.
- [40] Hopwood DA BM, Chater KF, Kieser T, Bruton CJ, Kieser HM, Lydiate DJ, et al. Genetic manipulation of *Streptomyces*: a Laboratory manual. Norwich: John Innes Foundation.; 1985.
- [41] Deutsch EW, Bandeira N, Sharma V, Perez-Riverol Y, Carver JJ, Kundu DJ, et al. The ProteomeXchange consortium in 2020: enabling ‘big data’ approaches in proteomics. *Nucleic Acids Res* 2020;48:D1145–52.
- [42] Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al. The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* 2019;47:D442–50.
- [43] Shikura N, Darbon E, Esnault C, Deniset-Besseau A, Xu D, Lejeune C, et al. The phosin PptA plays a negative role in the regulation of antibiotic production in *Streptomyces lividans*. *Antibiotics* 2021;10.
- [44] Aldridge M, Facey P, Francis L, Bayliss S, Del Sol R, Dyson P. A novel bifunctional histone protein in *Streptomyces*: a candidate for structural coupling between DNA conformation and transcription during development and stress? *Nucleic Acids Res* 2013;41:4813–24.
- [45] Sun J, Hesketh A, Bibb M. Functional analysis of relA and rshA, two relA/spoT homologues of *Streptomyces coelicolor* A3(2). *J Bacteriol* 2001;183:3488–98.
- [46] Allenby NE, Laing E, Bucca G, Kierzek AM, Smith CP. Diverse control of metabolism and other cellular processes in *Streptomyces coelicolor* by the PhoP transcription factor: genome-wide identification of in vivo targets. *Nucleic Acids Res* 2012;40:9543–56.
- [47] Rifat D, Bishai WR, Karakousis PC. Phosphate depletion: a novel trigger for *Mycobacterium tuberculosis* persistence. *J Infect Dis* 2009;200:1126–35.
- [48] Lopez-Alonso JP, Kaminiishi T, Kikuchi T, Hirata Y, Iturrioz I, Dhimole N, et al. RsgA couples the maturation state of the 30S ribosomal decoding center to activation of its GTPase pocket. *Nucleic Acids Res* 2017;45:6945–59.
- [49] Coatham ML, Brandon HE, Fischer JJ, Schummer T, Wieden HJ. The conserved GTPase HflX is a ribosome splitting factor that binds to the E-site of the bacterial ribosome. *Nucleic Acids Res* 2016;44:1952–61.
- [50] Nishimura K, Johansen SK, Inaoka T, Hosaka T, Tokuyama S, Tahara Y, et al. Identification of the RsmG methyltransferase target as 16S rRNA nucleotide G527 and characterization of *Bacillus subtilis* rsmG mutants. *J Bacteriol* 2007;189:6068–73.
- [51] Mora L, Heurgue-Hamard V, de Zamaroczy M, Kervestin S, Buckingham RH. Methylation of bacterial release factors RF1 and RF2 is required for normal translation termination in vivo. *J Biol Chem* 2007;282:35638–45.
- [52] Nord S, Bylund GO, Lovgren JM, Wikstrom PM. The RimP protein is important for maturation of the 30S ribosomal subunit. *J Mol Biol* 2009;386:742–53.
- [53] Li X, Sun Q, Jiang C, Yang K, Hung LW, Zhang J, et al. Structure of ribosomal silencing factor bound to *Mycobacterium tuberculosis* ribosome. *Structure* 2015;23:2387.
- [54] Persson BC, Gustafsson C, Berg DE, Bjork GR. The gene for a tRNA modifying enzyme, m5U54-methyltransferase, is essential for viability in *Escherichia coli*. *Proc Natl Acad Sci U S A* 1992;89:3995–8.
- [55] Sergiev PV, Alekashin NA, Chugunova AA, Polikanov YS, Dontsova OA. Structural and evolutionary insights into ribosomal RNA methylation. *Nat Chem Biol* 2018;14:226–35.
- [56] Britton RA. Role of GTPases in bacterial ribosome assembly. *Annu Rev Microbiol* 2009;63:155–76.
- [57] Anton BP, Saleh L, Benner JS, Raleigh EA, Kasif S, Roberts RJ. RimO, a MiaB-like enzyme, methylthiolates the universally conserved Asp88 residue of ribosomal protein S12 in *Escherichia coli*. *Proc Natl Acad Sci U S A* 2008;105:1826–31.
- [58] Datta PP, Wilson DN, Kawazoe M, Swami NK, Kaminiishi T, Sharma MR, et al. Structural aspects of RbfA action during small ribosomal subunit assembly. *Mol Cell* 2007;28:434–45.
- [59] Agafonov DE, Spirin AS. The ribosome-associated inhibitor A reduces translation errors. *Biochem Biophys Res Commun* 2004;320:354–8.
- [60] Kiel MC, Kaji H, Kaji A. Ribosome recycling: an essential process of protein synthesis. *Biochem Mol Biol Educ* 2007;35:40–4.
- [61] Landwehr V, Milanov M, Angebauer L, Hong J, Jungert G, Hiersemelz A, et al. The universally conserved ATPase YchF regulates translation of leaderless mRNA in response to stress conditions. *Front Mol Biosci* 2021;8:643696.
- [62] Kimura S, Suzuki T. Fine-tuning of the ribosomal decoding center by conserved methyl-modifications in the *Escherichia coli* 16S rRNA. *Nucleic Acids Res* 2010;38:1341–52.
- [63] Trassler SE, Bowles IE, Dey D, Jackman JE, Conn GL. Tied up in knots: untangling substrate recognition by the SPOUT methyltransferases. *J Biol Chem* 2022;298:102393.
- [64] Gc K, Gyawali P, Balci H, Abeysirigunawardena S. Ribosomal RNA methyltransferase RsmC moonlights as an RNA chaperone. *Chembiochem* 2020;21:1885–92.
- [65] Basturea GN, Deutscher MP. Substrate specificity and properties of the *Escherichia coli* 16S rRNA methyltransferase, RsmE. *RNA* 2007;13:1969–76.
- [66] Wei Y, Zhang H, Gao ZQ, Wang WJ, Shtykova EV, Xu JH, et al. Crystal and solution structures of methyltransferase RsmH provide basis for methylation of C1402 in 16S rRNA. *J Struct Biol* 2012;179:29–40.
- [67] Zhao M, Zhang H, Liu G, Wang L, Wang J, Gao Z, et al. Structural insights into the methylation of C1402 in 16S rRNA by methyltransferase RsmI. *PLoS One* 2016;11:e0163816.
- [68] Stephan NC, Ries AB, Boehringer D, Ban N. Structural basis of successive adenosine modifications by the conserved ribosomal methyltransferase KsgA. *Nucleic Acids Res* 2021;49:6389–98.
- [69] Karbstein K. Role of GTPases in ribosome assembly. *Biopolymers* 2007;87:1–11.
- [70] Gruffaz C, Smirnov A. GTPase Era at the heart of ribosome assembly. *Front Mol Biosci* 2023;10:1263433.
- [71] Hwang J, Inouye M. The tandem GTPase, Der, is essential for the biogenesis of 50S ribosomal subunits in *Escherichia coli*. *Mol Microbiol* 2006;61:1660–72.
- [72] Chakraborty A, Halder S, Kishore P, Saha D, Saha S, Sikder K, et al. The structure-function analysis of Obg-like GTPase proteins along the evolutionary tree from bacteria to humans. *Genes Cells* 2022;27:469–81.
- [73] Okamoto S, Ochi K. An essential GTP-binding protein functions as a regulator for differentiation in *Streptomyces coelicolor*. *Mol Microbiol* 1998;30:107–19.
- [74] Klein DJ, Moore PB, Steitz TA. The roles of ribosomal proteins in the structure assembly, and evolution of the large ribosomal subunit. *J Mol Biol* 2004;340:141–77.
- [75] Shahani Z, Sykes MT, Williamson JR. Assembly of bacterial ribosomes. *Annu Rev Biochem* 2011;80:501–26.
- [76] Delvillani F, Papiani G, Deho G, Briani F. S1 ribosomal protein and the interplay between translation and mRNA decay. *Nucleic Acids Res* 2011;39:7702–15.
- [77] Duval M, Korepanov A, Fuchsbaumer O, Fechter P, Haller A, Fabbretti A, et al. *Escherichia coli* ribosomal protein S1 unfolds structured mRNAs onto the ribosome for active translation initiation. *PLoS Biol* 2013;11:e1001731.
- [78] Ueta M, Wada C, Bessho Y, Maeda M, Wada A. Ribosomal protein L31 in *Escherichia coli* contributes to ribosome subunit association and translation, whereas short L31 cleaved by protease 7 reduces both activities. *Genes Cells* 2017;22:452–71.
- [79] Lilleorg S, Reier K, Remme J, Liiv A. The intersubunit bridge B1b of the bacterial ribosome facilitates initiation of protein synthesis and maintenance of translational fidelity. *J Mol Biol* 2017;429:1067–80.
- [80] Shin JH, Oh SY, Kim SJ, Roe JH. The zinc-responsive regulator Zur controls a zinc uptake system and some ribosomal proteins in *Streptomyces coelicolor* A3(2). *J Bacteriol* 2007;189:4070–7.
- [81] Paget MS, Kang JG, Roe JH, Buttner MJ. sigmaR, an RNA polymerase sigma factor that modulates expression of the thioredoxin system in response to oxidative stress in *Streptomyces coelicolor* A3(2). *EMBO J* 1998;17:5776–82.
- [82] Millan-Oropeza A, Henry C, Blein-Nicolas M, Aubert-Frambourg A, Moussa F, Bleton J, et al. Quantitative proteomics analysis confirmed oxidative metabolism predominates in *Streptomyces coelicolor* versus glycolytic metabolism in *Streptomyces lividans*. *J Proteome Res* 2017;16:2597–613.
- [83] Giege R, Springer M. Aminoacyl-tRNA synthetases in the bacterial world. *EcoSal Plus* 2016;7.
- [84] Curnow AW, Hong K, Yuan R, Kim S, Martins O, Winkler W, et al. Glut-tRNAGln amidotransferase: a novel heterotrimeric enzyme required for correct decoding of glutamine codons during translation. *Proc Natl Acad Sci U S A* 1997;94:11819–26.
- [85] Abel I, Korencic D, Ibbi M, Soll D. Trans-editing of mischarged tRNAs. *Proc Natl Acad Sci U S A* 2003;100:15422–7.
- [86] Gerber AP, Keller W. An adenosine deaminase that generates inosine at the wobble position of tRNAs. *Science* 1999;286:1146–9.
- [87] Rubio MA, Pastor I, Gaston KW, Ragone FL, Janzen CJ, Cross GA, et al. An adenosine-to-inosine tRNA-editing enzyme that can perform C-to-U deamination of DNA. *Proc Natl Acad Sci U S A* 2007;104:7821–6.
- [88] Torres AG, Pineyro D, Filonava L, Stracker TH, Batlle E, Ribas de Pouplana L. A-to-I editing on tRNAs: biochemical, biological and evolutionary implications. *FEBS Lett* 2014;588:4279–86.
- [89] Agris PF, Narendra A, Sarachan K, Vare VVP, Erusalys E. The importance of being modified: the role of RNA modifications in translational fidelity. *Enzymes* 2017;41:1–50.
- [90] Su C, Jin M, Zhang W. Conservation and diversification of tRNA t(6)-a-modifying enzymes across the three domains of life. *Int J Mol Sci* 2022;23.
- [91] Hori H. Methylated nucleosides in tRNA and tRNA methyltransferases. *Front Genet* 2014;5:144.
- [92] Guillou JM, Mechulam Y, Schmitter JM, Blanquet S, Fayat G. Disruption of the gene for Met-tRNA(fMet) formyltransferase severely impairs growth of *Escherichia coli*. *J Bacteriol* 1992;174:4294–301.

- [93] Newton DT, Creuzenet C, Mangroo D. Formylation is not essential for initiation of protein synthesis in all eubacteria. *J Biol Chem* 1999;274:22143–6.
- [94] Ofengand J, Malhotra A, Remme J, Gutgsell NS, Del Campo M, Jean-Charles S, et al. Pseudouridines and pseudouridine synthases of the ribosome. *Cold Spring Harb Symp Quant Biol* 2001;66:147–59.
- [95] Lin TY, Mehta R, Glatt S. Pseudouridines in RNAs: switching atoms means shifting paradigms. *FEBS Lett* 2021;595:2310–22.
- [96] Bjork GR, Jacobsson K, Nilsson K, Johansson MJ, Bystrom AS, Persson OP. A primordial tRNA modification required for the evolution of life? *EMBO J* 2001;20:231–9.
- [97] Stojkovic V, Fujimori DG. Radical SAM-mediated methylation of ribosomal RNA. *Methods Enzymol* 2015;560:355–76.
- [98] Hagervall TG, Edmonds CG, McCloskey JA, Bjork GR. Transfer RNA(5-methylaminomethyl-2-thiouridine)-methyltransferase from Escherichia coli K-12 has two enzymatic activities. *J Biol Chem* 1987;262:8488–95.
- [99] Soutourina O, Soutourina J, Blanquet S, Plateau P. Formation of D-tyrosyl-tRNATyr accounts for the toxicity of D-tyrosine toward Escherichia coli. *J Biol Chem* 2004;279:42560–5.
- [100] Calendar R. Deacylase d-Tyrosyl-tRNA. A new function. *Trends Biochem Sci* 2017;42:684–6.
- [101] H.a.L. Putzer S. Regulation of the expression of aminoacyl-tRNA synthetases and translation factors. *Madame curie bioscience database*. Austin (TX): Landes Bioscience; 2000–2013. p. 2000–13.
- [102] Yu YC, Han JM, Kim S. Aminoacyl-tRNA synthetases and amino acid signaling. *Biochim Biophys Acta Mol Cell Res* 2021;1868:118889.
- [103] Dahlquist KD, Puglisi JD. Interaction of translation initiation factor IF1 with the E. coli ribosomal A site. *J Mol Biol* 2000;299:1–15.
- [104] Nakamoto JA, Evangelista W, Vinogradova DS, Konevega AL, Spurio R, Fabbretti A, et al. The dynamic cycle of bacterial translation initiation factor IF3. *Nucleic Acids Res* 2021;49:6958–70.
- [105] Day JM, Janssen GR. Isolation and characterization of ribosomes and translation initiation factors from the gram-positive soil bacterium Streptomyces lividans. *J Bacteriol* 2004;186:6864–75.
- [106] Milon P, Carotti M, Konevega AL, Wintermeyer W, Rodnina MV, Gualerzi CO. The ribosome-bound initiation factor 2 recruits initiator tRNA to the 30S initiation complex. *EMBO Rep* 2010;11:312–6.
- [107] Adio S, Sharma H, Senyushkina T, Karki P, Maracci C, Wohlgemuth I, et al. Dynamics of ribosomes and release factors during translation termination in E. coli. *Elife* 2018;7.
- [108] Das G, Varshney U. Peptidyl-tRNA hydrolase and its critical role in protein biosynthesis. *Microbiol (Read)* 2006;152:2191–5.
- [109] Nakamoto H, Fujita K, Ohtaki A, Watanabe S, Narumi S, Maruyama T, et al. Physical interaction between bacterial heat shock protein (Hsp) 90 and Hsp70 chaperones mediates their cooperative action to refold denatured proteins. *J Biol Chem* 2014;289:6110–9.
- [110] Grandvalet C, Rapoport G, Mazodier P. hrcA, encoding the repressor of the groEL genes in Streptomyces albus G, is associated with a second dnaJ gene. *J Bacteriol* 1998;180:5129–34.
- [111] Stoller C, Rucknagel KP, Nierhaus KH, Schmid FX, Fischer G, Rahfeld JU. A ribosome-associated peptidyl-prolyl cis/trans isomerase identified as the trigger factor. *EMBO J* 1995;14:4939–48.
- [112] Gothen SF, Marahiel MA. Peptidyl-prolyl cis-trans isomerases, a superfamily of ubiquitous folding catalysts. *Cell Mol Life Sci* 1999;55:423–36.
- [113] Bucca G, Hindle Z, Smith CP. Regulation of the dnaK operon of Streptomyces coelicolor A3(2) is governed by HspR, an autoregulatory repressor protein. *J Bacteriol* 1997;179:5999–6004.
- [114] Stechmann A, Cavalier-Smith T. Evolutionary origins of Hsp90 chaperones and a deep paralogy in their bacterial ancestors. *J Eukaryot Microbiol* 2004;51:364–73.
- [115] Boge holz LAK, Mercier E, Wintermeyer W, Rodnina MV. Deformylation of nascent peptide chains on the ribosome. *Methods Enzymol* 2023;684:39–70.
- [116] Yang CI, Zhu Z, Jones JJ, Lomenick B, Chou TF, Shan SO. System-wide analyses reveal essential roles of N-terminal protein modification in bacterial membrane integrity. *iScience* 2022;25:104756.
- [117] Bucca G, Brassington AM, Hotchkiss G, Merninas V, Smith CP. Negative feedback regulation of dnaK, clpB and lon expression by the DnaK chaperone machine in Streptomyces coelicolor, identified by transcriptome and in vivo DnaK-depletion analysis. *Mol Microbiol* 2003;50:153–66.
- [118] Grandvalet C, de Crecy-Lagard V, Mazodier P. The ClpB ATPase of Streptomyces albus G belongs to the HspR heat shock regulon. *Mol Microbiol* 1999;31:521–32.
- [119] Ahmad S, Selvapandian A, Bhatnagar RK. Phylogenetic analysis of gram-positive bacteria based on grpE, encoded by the dnaK operon. *Int J Syst Evol Microbiol* 2000;50 Pt 5:1761–6.
- [120] Duchene AM, Kieser HM, Hopwood DA, Thompson CJ, Mazodier P. Characterization of two groEL genes in Streptomyces coelicolor A3(2). *Gene* 1994;144:97–101.
- [121] Wilson DN. Less is more for leaderless mRNA translation. *Mol Cell* 2009;33:141–2.
- [122] Vesper O, Amitai S, Belitsky M, Byrgazov K, Kaberdina AC, Engelberg-Kulka H, et al. Selective translation of leaderless mRNAs by specialized ribosomes generated by MazF in Escherichia coli. *Cell* 2011;147:147–57.
- [123] Leiva LE, Katz A. Regulation of leaderless mRNA translation in bacteria. *Microrganisms* 2022;10.
- [124] Jeong Y, Kim JN, Kim MW, Bucca G, Cho S, Yoon YJ, et al. The dynamic transcriptional and translational landscape of the model antibiotic producer Streptomyces coelicolor A3(2). *Nat Commun* 2016;7:11605.
- [125] Santos-Beneit F. The Pho regulon: a huge regulatory network in bacteria. *Front Microbiol* 2015;6:402.
- [126] Chouayekh H, Virolle MJ. The polyphosphate kinase plays a negative role in the control of antibiotic production in Streptomyces lividans. *Mol Microbiol* 2002;43:919–30.
- [127] Tul'skaya EM, Shashkov AS, Streshinskaya GM, Senchenkova SN, Potekhina NV, Kozlova YI, et al. Teichuronic and teichulosonic acids of actinomycetes. *Biochemistry (Mosc)* 2011;76:736–44.
- [128] Ostash B, Shashkov A, Streshinskaya G, Tul'skaya E, Baryshnikova L, Dmitrenok A, et al. Identification of Streptomyces coelicolor M145 genomic region involved in biosynthesis of teichulosonic acid-cell wall glycopolymer. *Folia Microbiol (Praha)* 2014;59:355–60.
- [129] Lang WK, Glassey K, Archibald AR. Influence of phosphate supply on teichoic acid and teichuronic acid content of *Bacillus subtilis* cell walls. *J Bacteriol* 1982;151:367–75.
- [130] Sivapragasam S, Grove A. Streptomyces coelicolor XdhR is a direct target of (p)ppGpp that controls expression of genes encoding xanthine dehydrogenase to promote purine salvage. *Mol Microbiol* 2016;100:701–18.
- [131] Chouayekh H, Nothaft H, Delaunay S, Linder M, Payrastre B, Seghezzi N, et al. Phosphoinositides are involved in control of the glucose-dependent growth resumption that follows the transition phase in Streptomyces lividans. *J Bacteriol* 2007;189:741–9.
- [132] Grucela PK, Fuhrer T, Sauer U, Chao Y, Zhang YE. Ribose 5-phosphate: the key metabolite bridging the metabolisms of nucleotides and amino acids during stringent response in *Escherichia coli*? *Microb Cell* 2023;10:141–4.
- [133] Fritsch VN, Loi VV, Busche T, Tung QN, Lill R, Horvatek P, et al. The alarmone (p)ppGpp confers tolerance to oxidative stress during the stationary phase by maintenance of redox and iron homeostasis in *Staphylococcus aureus*. *Free Radic Biol Med* 2020;161:351–64.
- [134] Driller K, Cornejo FA, Turgay K. (p)ppGpp – an important player during heat shock response. *Microlife* 2023;4:uqad017.
- [135] Romero-Rodriguez A, Maldonado-Carmona N, Ruiz-Villafan B, Koiala N, Rocha D, Sanchez S. Interplay between carbon, nitrogen and phosphate utilization in the control of secondary metabolite production in Streptomyces. *Antonie Van Leeuwenhoek* 2018;111:761–81.
- [136] Wu L, Wang Z, Guan Y, Huang X, Shi H, Liu Y, et al. The (p)ppGpp-mediated stringent response regulatory system globally inhibits primary metabolism and activates secondary metabolism in *Pseudomonas protegens* H78. *Appl Microbiol Biotechnol* 2020;104:3061–79.
- [137] Virolle MJ. A Challenging View. Antibiotics play a role in the regulation of the energetic metabolism of the producing bacteria antibiotics. *Basel* 2020;9.