



HAL
open science

Self-avoiding space-filling folding curves in dimension 3

Francis Oger

► **To cite this version:**

| Francis Oger. Self-avoiding space-filling folding curves in dimension 3. 2024. hal-04524150

HAL Id: hal-04524150

<https://hal.science/hal-04524150>

Preprint submitted on 27 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Self-avoiding space-filling folding curves in dimension 3

F. Oger

Various examples of folding curves have been considered in the plane. For each integer $n \geq 1$, a square n -curve such as the dragon curve is obtained by folding n times a strip of paper in 2, then unfolding it, leaving a $\pi/2$ angle at each fold. A triangular n -curve such as the terdragon curve is obtained by folding n times a strip of paper in 3, then unfolding it, leaving a $\pi/3$ angle at each fold. Peano-Gosper curves, which are based on regular hexagons, have similar properties.

Folding curves were used to construct fractals. They have two characteristic properties: First, they are self-avoiding. Second, n -curves of the same type completely cover larger and larger regions of the plane when n becomes larger.

Square (resp. triangular) folding curves are finite or infinite sequences of consecutive segments taken among the edges of squares (resp. equilateral triangles) which form a regular tiling of the plane. They are obtained from 1 segment by applying repeatedly an inflation process: multiply the length of each segment by k , then replace all the segments with the same curve which consists of k^2 segments. The same process can be applied uniformly, starting with a covering of the plane by segments which are the edges of squares (resp. equilateral triangles) in a regular tiling.

According to [3], [4] and [5], the inductive limit of such a process gives a covering of the plane by curves without endpoint which satisfies the local isomorphism property. Any infinite folding curve can be extended into such a covering in an essentially unique way. Two coverings obtained by the same process are locally isomorphic.

Such coverings of the plane consist of a small number of curves, at most 6 in the examples that we considered until present. Each local isomorphism class contains some coverings which consist of only 1 curve.

We also proved in [5] that the same results are true for Peano-Gosper curves.

In order to obtain similar examples in dimension 3, one natural idea was to fold a wire in 2 several times according to the 3 directions, then unfold it, leaving a $\pi/2$ angle at each fold. However, it was proved in [2] that the

curves obtained in that way are generally not self-avoiding. Another possible direction of research was mentioned in [1].

Presently, we have an example with the properties above in dimension 3. Here, the curves consist of consecutive segments of length 1 taken among the edges of cubes which form a regular tiling of the space. The inflation process is as follows: multiply the length of the segments by 2, then replace each segment with the image through a positive isometry of the curve with 8 segments passing through the points $(0, 0, 0)$, $(0, \bar{1}, 0)$, $(0, \bar{1}, \bar{1})$, $(0, 0, \bar{1})$, $(1, 0, \bar{1})$, $(1, \bar{1}, \bar{1})$, $(1, \bar{1}, 0)$, $(1, 0, 0)$, $(2, 0, 0)$, where $\bar{1} = -1$.

References

- [1] S.I. Ben-Abraham, A. Quandt and D. Shapira, Multidimensional paperfolding systems, *Acta Cryst. A* 69 (2013), 123-130.
- [2] M. Mendès France and O. Shallit, Wire Bending, *J. Combinatorial Theory* 50 (1989), 1-23.
- [3] F. Oger, Paperfolding sequences, paperfolding curves and local isomorphism, *Hiroshima Math. Journal* 42 (2012), 37-75.
- [4] F. Oger, The number of paperfolding curves in a covering of the plane, *Hiroshima Math. Journal* 47 (2017), 1-14.
- [5] F. Oger, Coverings of the plane by self-avoiding curves which satisfy the local isomorphism property, arXiv:2310.19364.