



HAL
open science

Invariant kernels on the space of complex covariance matrices

Cyrus Mostajeran, Salem Said

► **To cite this version:**

Cyrus Mostajeran, Salem Said. Invariant kernels on the space of complex covariance matrices. 2024. hal-04525081v2

HAL Id: hal-04525081

<https://hal.science/hal-04525081v2>

Preprint submitted on 25 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Invariant kernels on the space of complex covariance matrices

Cyrus Mostajeran and Salem Said

Abstract

The present work develops certain analytical tools required to construct and compute invariant kernels on the space of complex covariance matrices. The main result is the L^1 -Godement theorem, which states that any invariant kernel, which is (in a certain natural sense) also integrable, can be computed by taking the inverse spherical transform of a positive function. General expressions for inverse spherical transforms are then provided, which can be used to explore new families of invariant kernels, at a rather moderate computational cost. A further, alternative approach for constructing new invariant kernels is also introduced, based on Ramanujan's master theorem for symmetric cones.

Index Terms

positive definite kernel, covariance matrix, Bochner's theorem, symmetric space, spherical transform, Ramanujan's master theorem

I. INTRODUCTION

Positive definite kernels, and especially invariant positive definite kernels, play a prominent role across probability, statistics, and machine learning. Bochner's theorem characterises the class of invariant positive definite kernels or (equivalently) the class of positive definite functions: a function $f : \mathbb{R}^N \rightarrow \mathbb{C}$ is positive definite if and only if it is the inverse Fourier transform of some finite positive measure.

Recent research in machine learning and data science has focused on invariant positive definite kernels which are defined on non-Euclidean spaces (rather than on \mathbb{R}^N as in Bochner's theorem) [1] [2] [3] [4]. In turn, this revived interest in the non-Euclidean generalisations of Bochner's theorem, due to Godement, Gelfand, and others [5] [6]. In [3], the L^p -Godement theorems ($p = 1, 2$) were introduced, in order to deal with integrable invariant kernels defined on Riemannian symmetric spaces of non-compact type. These are somewhat similar to Bochner's theorem, with the role of the Fourier transform played by the so-called spherical transform.

The aim of the present work is to apply the L^1 -Godement theorem, in order to construct and compute positive definite functions on the space of complex covariance matrices. This requires developing certain analytical tools which make it possible to work with spherical transforms on this space. These are the main focus in the following.

Necessary background is given in Section II. This begins with two definitions, invariant positive definite kernels and U -invariant positive definite functions. Proposition 1 shows that these two concepts are in fact equivalent (below, for brevity, the term "kernel" will mean positive definite kernel). Section II also discusses the Riemannian geometry of the space of complex covariance matrices. This is indeed a Riemannian symmetric space, when equipped with its well-known affine invariant metric.

Section III is concerned with the spherical transform on this space. Roughly, this is an expansion of integrable U -invariant functions in terms of eigenfunctions of the Laplace-Beltrami operator, called spherical functions [7] [8]. Proposition 2 provides a closed-form determinantal formula for these spherical functions, a slight generalisation of the classic Gelfand-Naimark formula [9], while Proposition 3 expresses the spherical transform itself, along with its inverse, in the form of a multiple integral. The following Proposition 4 is the first application of Proposition 3. It computes explicitly the spherical transform of a Gaussian function.

Section IV comes to the main issue of applying the L^1 -Godement theorem. In [3], this theorem was stated under the assumption that the underlying symmetric space should be of non-compact type. This assumption is not satisfied for the space of complex covariance matrices. Theorem 1 (proved in Appendix C) makes up for this problem. Proposition 5 shows how Theorem 1 can be used to construct and compute U -invariant positive definite functions. Specifically, the integrable U -invariant positive definite functions are exactly the inverse spherical transforms of a certain class of positive functions (described in Theorem 1). Formulas (23) and (24) provide general expressions for these inverse spherical transforms. To generate a new positive definite function, it is enough to plug a suitable positive function into one of these two formulas.

Section IV closes with three examples. The first one showcases an application of Formula (23). The second one uses (24) in order to compute the heat kernel (on the space of complex covariance matrices) in closed form. The last one combines Proposition 4 and Theorem 1 to prove that a Gaussian function is never positive definite (this means there exists no value of its “variance parameter” for which it is positive definite). This provides a new, purely analytical, way of proving this statement, alternative to the geometric proof laid out in [3].

Section V features Proposition 6, which shows how U -invariant positive definite functions can be obtained in the form of spherical power series, through an application of Ramanujan’s master theorem for symmetric cones [10]. An example of a positive definite function obtained in this way is the “Beta-prime” function (see Expression (33)).

Most of the results given in the present work provide analytical expressions which can be evaluated rather directly (*e.g.* they require computing $N \times N$ -size determinants, when working on the space of $N \times N$ covariance matrices). This nice situation is due to the closed-form expression for spherical functions, in Proposition 2. Generally speaking, such an expression is always available on any symmetric space whose group of isometries is a complex Lie group. Symmetric spaces with this property are called symmetric spaces of type IV [7] [11]. These are exactly the following (a) spaces of complex covariance matrices, (b) spaces of complex covariance matrices which are also orthogonal, (c) spaces of complex covariance matrices which are also symplectic, (d) certain other so-called exceptional spaces. These are the non-compact duals of the compact Lie groups (unitary, orthogonal, symplectic, and exceptional). The results given in the present work (with no exception) readily extend to any symmetric space of type IV.

On the other hand, several spaces of covariance matrices (real, quaternion, block-Toeplitz, *etc.*) can be embedded into spaces of complex covariance matrices (eventually of larger matrix size), in a way which preserves all the fundamental symmetry and invariance properties. Therefore, any invariant kernel constructed using the methods proposed in the present work immediately yields an invariant kernel on any of the above-mentioned spaces. In this way, the contribution of the present work is to introduce a general means of generating invariant kernels which admit analytical expressions and can be used on most of the usual spaces of covariance matrices.

II. GENERAL BACKGROUND

A. Positive definite functions

Denote by M the space of $N \times N$ complex covariance matrices. Specifically, these are $N \times N$ Hermitian positive definite matrices. Moreover, denote by G the group of $N \times N$ invertible complex matrices, and by U the group of $N \times N$ unitary matrices, a subgroup of G .

Recall that G acts transitively on M in the following way: $g \cdot x = gxg^\dagger$ for $g \in G$ and $x \in M$ (where \dagger denotes the conjugate-transpose) [12] [13]. For this action, U is the stabiliser of the identity matrix $\text{id} \in M$. In other words, $g \cdot \text{id} = \text{id}$ if and only if $g \in U$.

A kernel \mathcal{K} is a continuous function $\mathcal{K} : M \times M \rightarrow \mathbb{C}$, such that for any $x_1, \dots, x_n \in M$ (here, $n = 2, 3, \dots$), the $n \times n$ matrix with elements $\mathcal{K}(x_i, x_j)$ is Hermitian positive semidefinite. The focus of the present work is on invariant kernels. These are kernels which satisfy $\mathcal{K}(g \cdot x, g \cdot y) = \mathcal{K}(x, y)$ for all $g \in G$ and all $x, y \in M$ [1] [3].

It is convenient to study invariant kernels indirectly, by studying U -invariant positive definite functions [3]. A function $f : M \rightarrow \mathbb{C}$ is called U -invariant if $f(u \cdot x) = f(x)$ for all $u \in U$ and $x \in M$. This means that $f(x) = f_o(\rho)$ where f_o is a symmetric function and $\rho = (\rho_1, \dots, \rho_N)$ are the eigenvalues of x : $f_o(\rho)$ remains unchanged after any permutation of (ρ_1, \dots, ρ_N) .

If f is continuous, then it is called positive definite if, for any $x_1, \dots, x_n \in M$ (where $n = 2, 3, \dots$), the $n \times n$ matrix with elements $f(x_i^{-1/2} x_j x_i^{-1/2})$ is Hermitian positive semidefinite.

The two concepts (invariant kernel and U -invariant positive definite function) are equivalent.

Proposition 1: If \mathcal{K} is an invariant kernel, then the function $f(x) = \mathcal{K}(x, \text{id})$ is U -invariant and positive definite. If f is a U -invariant and positive definite function, $\mathcal{K}(x, y) = f(y^{-1/2} x y^{-1/2})$ defines an invariant kernel.

Proof: let $f(x) = \mathcal{K}(x, \text{id})$. If $u \in U$, then $f(u \cdot x) = \mathcal{K}(u \cdot x, \text{id}) = \mathcal{K}(u \cdot x, u \cdot \text{id})$ because $u \cdot \text{id} = \text{id}$. However, if \mathcal{K} is invariant, then $\mathcal{K}(u \cdot x, u \cdot \text{id}) = \mathcal{K}(x, \text{id})$. Therefore, $f(u \cdot x) = f(x)$ and f is U -invariant. To see that f is positive definite, it is enough to note that

$$f(y^{-1/2} x y^{-1/2}) = \mathcal{K}(y^{-1/2} \cdot x, \text{id}) = \mathcal{K}(x, y^{1/2} \cdot \text{id}) = \mathcal{K}(x, y)$$

where the first equality follows from the definition of $g \cdot x$, by taking $g = y^{-1/2}$, and the second equality because \mathcal{K} is invariant. Thus, for any $x_1, \dots, x_n \in M$, the matrix with elements $f(x_i^{-1/2} x_j x_i^{-1/2})$ is the same as the matrix with elements $\mathcal{K}(x_j, x_i)$, which is positive semidefinite because \mathcal{K} is a kernel. In addition, continuity of f follows from continuity of \mathcal{K} , and this ensures f is positive definite.

Conversely, let f be U -invariant and positive definite. This clearly implies that $\mathcal{K}(x, y) = f(y^{-1/2} x y^{-1/2})$ is a kernel. To see that this \mathcal{K} is invariant, note that $\mathcal{K}(x, y) = f_o(\rho)$ where $\rho = (\rho_1, \dots, \rho_N)$ are the eigenvalues of $y^{-1/2} x y^{-1/2}$. These are the same as the eigenvalues of $y^{-1} x$ because the two matrices are similar: $y^{-1} x = y^{-1/2} (y^{-1/2} x y^{-1/2}) y^{1/2}$. By the same argument, $\mathcal{K}(g \cdot x, g \cdot y) = f_o(\rho')$ where $\rho' = (\rho'_1, \dots, \rho'_N)$ are the eigenvalues of $(g \cdot y)^{-1} (g \cdot x)$. However, this last matrix is similar to $y^{-1} x$,

$$(g \cdot y)^{-1} (g \cdot x) = (g^\dagger)^{-1} (y^{-1} x) (g^\dagger)$$

Therefore, $\rho' = \rho$ and $\mathcal{K}(g \cdot x, g \cdot y) = \mathcal{K}(x, y)$, as required. ■

B. Riemannian geometry

An explicit description of U -invariant positive definite functions relies on the Riemannian geometry of M . Precisely, it relies on the fact that M is a Riemannian symmetric space [8] [11].

Note that M is an open subset of H , the real vector space of $N \times N$ Hermitian matrices. Therefore, M is a differentiable manifold with its tangent space at any $x \in M$ naturally isomorphic to H . Now, with this in mind, consider the Riemannian metric on M ,

$$\langle v, w \rangle_x = \operatorname{Re} [\operatorname{tr}(x^{-1} v x^{-1} w)] \quad v, w \in H \quad (1)$$

where Re denotes the real part and tr the trace. This is the affine-invariant metric, which was made popular by [14].

For any $g \in G$, the map $x \mapsto g \cdot x$ is an isometry of the metric (1). The same is true for inversion $x \mapsto x^{-1}$. These two facts together show that M satisfies the definition of a Riemannian symmetric space (for details, see [13]).

The class of U -invariant functions behaves in a special way with respect to the metric (1). For example, let vol denote the Riemannian volume element of this metric. If $f : M \rightarrow \mathbb{C}$ is an integrable U -invariant function [13],

$$\int_M f(x) \operatorname{vol}(dx) = \frac{C_N}{N!} \int_{\mathbb{R}_+^N} f_o(\rho) (V(\rho))^2 \prod_{k=1}^N \rho_k^{-N} d\rho_k \quad (2)$$

where $f(x) = f_o(\rho)$ is a symmetric function of the eigenvalues (ρ_1, \dots, ρ_N) of x , and where V stands for the Vandermonde polynomial. Here, and throughout the following, C_N denotes a positive constant that only depends on N and whose value is allowed to differ from one formula to another.

Moreover, if L is the Laplace-Beltrami operator of the metric (1), and f is smooth and U -invariant, then [7] [8],

$$Lf = \sum_{k=1}^N \rho_k^2 \frac{\partial^2 f_o}{\partial \rho_k^2} + 2 \sum_{k < \ell} \frac{\rho_k \rho_\ell}{\rho_k - \rho_\ell} \left(\frac{\partial f_o}{\partial \rho_k} - \frac{\partial f_o}{\partial \rho_\ell} \right) + N \sum_{k=1}^N \rho_k \frac{\partial f_o}{\partial \rho_k} \quad (3)$$

Formulas (2) and (3) arise systematically from the Riemannian geometry of M , but they are also familiar in certain problems of multivariate statistics and random matrix theory [15] [16] [17].

To close the present section, consider a special case of the integral formula (2). Assume that $f_o(\rho)$ factors into $f_o(\rho) = w(\rho_1) \dots w(\rho_N)$ where w is an integrable function such that

$$\int_0^\infty |w(\rho)| \rho^{-N} d\rho < \infty \quad \text{and} \quad \int_0^\infty |w(\rho)| \rho^{N-2} d\rho < \infty$$

Then, the volume integral in (2) is convergent and admits a determinantal expression

$$\int_M f(x) \operatorname{vol}(dx) = C_N \det \left[\int_0^\infty w(\rho) \rho^{k+\ell-N} d\rho \right]_{k,\ell=0}^{N-1} \quad (4)$$

This is an application of the Andréief identity, widely used in random matrix theory, and was pointed out in [18].

Example : the expression (4) can be used to compute the Gaussian integral of [19] [20]

$$Z(\sigma) = \int_M \exp \left[-\frac{d^2(x, \operatorname{id})}{2\sigma^2} \right] \operatorname{vol}(dx) \quad (5)$$

where $d(\cdot, \cdot)$ denotes the Riemannian distance induced on M by the metric (1). In (4), this integral corresponds to $w(\rho) = \exp[-\log^2(\rho)/2\sigma^2]$, which yields

$$Z(\sigma) = C_N \det \left[\sigma e^{(\sigma^2/2)(k+\ell-N-1)^2} \right]_{k,\ell=1}^N \quad (6)$$

a formula due to [18], which will be both simplified and generalised in the following section (see Proposition 4).

III. THE SPHERICAL TRANSFORM

The key ingredient which will be employed in constructing and computing U -invariant positive definite functions is the spherical transform. Roughly, this provides an expansion of any well-behaved U -invariant function, in terms of eigenfunctions of the Laplace-Beltrami operator (3), which are known as spherical functions [7] [8].

The set of all spherical functions is described as follows [8]. Consider first the power function, $\Delta_s : M \rightarrow \mathbb{C}$ where $s = (s_1, \dots, s_N)$ belongs to \mathbb{C}^N . This is

$$\Delta_s(x) = (\Delta_1(x))^{s_1 - s_2} (\Delta_2(x))^{s_2 - s_3} \dots (\Delta_N(x))^{s_N} \quad (7)$$

where $\Delta_k(x)$ is the k -th leading principal minor of $x \in M$. A spherical function is a function of the form

$$\Phi_\lambda(x) = \int_U \Delta_{\lambda + \delta}(u \cdot x) du \quad (8)$$

where $\lambda \in \mathbb{C}^N$ and $\delta_k = \frac{1}{2}(2k - N - 1)$, while du denotes the normalised Haar measure on the unitary group U . Two functions Φ_λ and $\Phi_{\lambda'}$ are identical if and only if $(\lambda_1, \dots, \lambda_N)$ is a permutation of $(\lambda'_1, \dots, \lambda'_N)$.

Each Φ_λ is U -invariant and an eigenfunction of the Laplace-Beltrami operator (3), with eigenvalue $(\lambda, \lambda) - (\delta, \delta)$ [8] (Theorem XIV.3.1). Here, and throughout the following, $(\mu, \nu) = \sum_{k=1}^N \mu_k \nu_k$ for $\mu, \nu \in \mathbb{C}^N$.

The first claim of the present section is that the spherical functions Φ_λ , while initially given by the integral formula (8), admit the following determinantal expression.

Proposition 2: If $\lambda \in \mathbb{C}^N$ and Φ_λ is given by (8), then

$$\Phi_\lambda(x) = \prod_{k=1}^{N-1} k! \times \frac{\det \left[\rho_k^{\lambda_\ell + (N-1)/2} \right]_{k,\ell=1}^N}{V(\lambda)V(\rho)} \quad (9)$$

where (ρ_1, \dots, ρ_N) are the eigenvalues of x and V stands for the Vandermonde polynomial.

The proof of Proposition 2 will be given in Appendix A. Formula (9) will be called the Gelfand-Naimark formula, as it is a slight generalisation of the formula introduced by Gelfand and Naimark [9] in 1950. The reason why the spherical functions (8) admit the determinantal expression (9) is that the group G is here a complex Lie group. This fact is the foundation of the proof in Appendix A.

Now, let $f : M \rightarrow \mathbb{C}$ be an integrable U -invariant function (integrable means with respect to vol , as in (2)). Its spherical transform is the function $\hat{f} : \mathbb{R}^N \rightarrow \mathbb{C}$,

$$\hat{f}(t) = \int_M f(x) \Phi_{-it}(x) \text{vol}(dx) \quad (10)$$

where $i = \sqrt{-1}$ [8]. An inversion theorem for the spherical transform (10) is given in [8] (Theorem XIV.5.3). Specifically, if \hat{f} satisfies the integrability condition

$$\int_{\mathbb{R}^N} |\hat{f}(t)| (V(t))^2 dt < \infty \quad (11)$$

then the following inversion formula holds,

$$f(x) = C_N \int_{\mathbb{R}^N} \hat{f}(t) \Phi_{it}(x) (V(t))^2 dt \quad (12)$$

After substituting (2) and (9) into (10) and (12), the following is obtained.

Proposition 3: For the spherical transform pair (10)–(12),

$$\hat{f}(t) = \frac{C_N}{V(-it)} \times \frac{1}{N!} \int_{\mathbb{R}_+^N} f_o(\rho) V(\rho) \det \left[\rho_k^{-it_\ell - (N+1)/2} \right] d\rho \quad (13)$$

$$f(x) = \frac{C_N}{V(it)} \times \frac{1}{N!} \int_{\mathbb{R}^N} \hat{f}(t) V(t) \det \left[\rho_k^{it_\ell + (N-1)/2} \right] dt \quad (14)$$

where $f(x) = f_o(\rho)$ is a symmetric function of the eigenvalues (ρ_1, \dots, ρ_N) of x .

The proof of this proposition will not be given in detail, as it merely consists of performing straightforward algebraic simplifications.

It is remarkable that the spherical transform does not involve all the spherical functions Φ_λ but only the functions Φ_{it} where $t \in \mathbb{R}^N$. These functions have in common the property that they correspond to real, negative eigenvalues of the Laplace-Beltrami operator (3): each Φ_{it} corresponds to the eigenvalue $-(t, t) - (\delta, \delta)$. The spherical functions that do not appear in the spherical transform are interesting in their own right. For instance, if $\lambda + \delta = m$ where (m_1, \dots, m_N) are positive integers arranged in decreasing order, then one has

$$\Phi_\lambda(x) = \prod_{k=1}^{N-1} k! \times S_m(\rho)/V(\lambda) \quad (15)$$

where S_m denotes the Schur polynomial corresponding to (m_1, \dots, m_N) . Schur polynomials are very important in the study of circular and unitary-invariant random matrix ensembles [16] [17] [21] (because they provide the irreducible characters of the unitary group). In Section V, below, they will appear within the framework of Ramanujan's master theorem.

The following proposition is motivated by the study of the Gaussian integral (5). Specifically, consider the integrals

$$Z(\sigma, \lambda) = \int_M \exp \left[-\frac{d^2(x, \text{id})}{2\sigma^2} \right] \Phi_\lambda(x) \text{vol}(dx) \quad (16)$$

where Φ_λ was defined in (8). If $\lambda = -\delta$ then $Z(\sigma, \lambda)$ is just $Z(\sigma)$ from (5). On the other hand, note that $Z(\sigma, -it) = \hat{f}(t)$ where $f(x) = \exp[-d^2(x, \text{id})/2\sigma^2]$.

Proposition 4: The integrals (16) admit the following expression

$$\begin{aligned} Z(\sigma, \lambda) &= \frac{C_N}{V(\lambda)} \times \det \left[\sigma \exp \left((\sigma^2/2)(\delta_k + \lambda_\ell)^2 \right) \right]_{k, \ell=1}^N \\ &= C_N \sigma^{N^2} e^{\frac{\sigma^2}{2}((\lambda, \lambda) + (\delta, \delta))} \prod_{k < \ell} \text{sch}((\sigma^2/2)(\lambda_\ell - \lambda_k)) \end{aligned} \quad (17)$$

where $\text{sch}(a) = \sinh(a)/a$.

The proof of Proposition 4 will be given in Appendix B. This proposition provides a simplified form of (6),

$$Z(\sigma) = C_N \sigma^N e^{\sigma^2(\delta, \delta)} \prod_{k < \ell} \sinh((\sigma^2/2)(\ell - k)) \quad (18)$$

as follows by replacing $\lambda = -\delta$ into (17). In addition, for the Gaussian function f defined before the proposition, putting $\lambda = -it$ gives the spherical transform

$$\hat{f}(t) = C_N \sigma^{N^2} e^{\frac{\sigma^2}{2}((\delta, \delta) - (t, t))} \prod_{k < \ell} \text{sc}((\sigma^2/2)(t_\ell - t_k)) \quad (19)$$

where $\text{sc}(a) = \sin(a)/a$. This formula will be used in the example at the end of the following section, in order to show that the Gaussian function is not positive definite (there exists no value of σ for which it is positive definite).

IV. CONSTRUCTING INVARIANT KERNELS

The L^1 –Godement theorem was introduced in [3]. Roughly, this theorem shows that U -invariant positive definite functions can be obtained by taking inverse spherical transforms of positive symmetric functions.

Theorem 1: Let $f : M \rightarrow \mathbb{C}$ be an integrable U -invariant function (integrable means with respect to vol). Then, f is positive definite if and only if

$$f(x) = \int_{\mathbb{R}^N} g(t) \Phi_{it}(x) (V(t))^2 dt \quad (20)$$

where the function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ is positive ($g(t) \geq 0$ for all t), symmetric, and satisfies the integrability condition

$$\int_{\mathbb{R}^N} g(t) (V(t))^2 dt < \infty \quad (21)$$

Moreover, this function g is then unique — recall that g is said to be symmetric if $g(t)$ remains unchanged after any permutation of (t_1, \dots, t_N) .

The proof of Theorem 1 will be given in Appendix C.

As explained in [3], the L^1 –Godement theorem is based on the celebrated Godement theorem, which generalises Bochner’s theorem to the context of symmetric spaces [5]. The only-if part of this theorem can be used to check whether a given function f is positive definite or not. On the other hand, the if part can be used to construct and compute positive definite functions.

Indeed, note that (20) is essentially an inverse spherical transform as in (12), with $g(t)$ instead of $\hat{f}(t)$. Therefore, just as in (14), it is possible to rewrite (20),

$$f(x) = \frac{1}{V(i\rho)} \times \frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell + (N-1)/2} \right] dt \quad (22)$$

where (ρ_1, \dots, ρ_N) are the eigenvalues of x . To obtain a positive definite function f , it is then enough to choose a suitable positive function g and then evaluate the integral (22). This is considered in the following proposition.

Proposition 5: Let $g : \mathbb{R}^N \rightarrow \mathbb{R}$ satisfy the conditions of Theorem 1.

(a) Assume that $g(t)$ factors into $g(t) = \gamma(t_1) \dots \gamma(t_N)$, where γ is a positive function. It follows from (22) that

$$f(x) = \frac{(\det(x))^{(N-1)/2}}{V(i\rho)} \times \det \left[\int_{\mathbb{R}} \gamma(t) t^{k-1} e^{its_\ell} dt \right] \quad (23)$$

whenever the integrals under the determinant exist. Here, $s_\ell = \log(\rho_\ell)$ for $\ell = 1, \dots, N$.

(b) Assume that the inverse Fourier transform

$$\tilde{g}(s) = \int_{\mathbb{R}^N} g(t) e^{i(s,t)} dt \quad (s_1, \dots, s_N) \in \mathbb{R}^N$$

is smooth. It follows from (22) that

$$f(x) = \frac{(\det(x))^{(N-1)/2}}{V(\rho)} \times V \left(-\frac{\partial}{\partial s} \right) \tilde{g}(s) \Big|_{s_\ell = \log(\rho_\ell)} \quad (24)$$

where $V(\partial/\partial s)$ is the Vandermonde operator $V(\partial/\partial s) = \prod_{i < j} (\partial/\partial s_j - \partial/\partial s_i)$.

The proof of Proposition 5 is given in Appendix D.

Example : as an application of Proposition 5-(a), choose $g(t) = \gamma(t_1) \dots \gamma(t_N)$ where $\gamma(t) = (\kappa/2) \exp(-\kappa|t|)$ for $\kappa > 0$. Replacing into (23) and using elementary properties of the Fourier transform,

$$f(x) = \frac{(\det(x))^{(N-1)/2}}{V(\rho)} \times \det \left[-\tilde{\gamma}^{(k-1)}(\log(\rho_\ell)) \right] \quad (25)$$

where $\tilde{\gamma}^{(k-1)}$ is the $(k-1)$ -th derivative of $\tilde{\gamma}(s) = (\kappa^2 + s^2)^{-1}$. Theorem 1 now says that this f is a positive definite function. In fact, (25) provides a whole family of positive definite functions, parameterised by κ .

Example : while it looks a bit too complicated, Proposition 5-(b) can be used to compute the heat kernel of M in closed form [7] [22]. This corresponds to $g(t) = \exp[-\kappa((t, t) + (\delta, \delta))]$ with $\kappa > 0$. Now, for this choice of $g(t)$, one has the following inverse Fourier transform

$$\tilde{g}(s) = \left((\pi/\kappa)^{N/2} e^{-\kappa(\delta, \delta)} \right) \exp[-(s, s)/4\kappa]$$

and one may use a beautiful identity from [22] (Chapter XII, Page 405)

$$V \left(-\frac{\partial}{\partial s} \right) \exp[-(s, s)/4\kappa] = (1/2\kappa)^{N(N-1)/2} V(s) \exp[-(s, s)/4\kappa] \quad (26)$$

Replacing this into (24) yields the heat kernel (rather, $f(x) = \mathcal{K}(x, \text{id})$ where \mathcal{K} is the heat kernel)

$$f(x) = C_\kappa (\det(x))^{(N-1)/2} \times (V(\log(\rho))/V(\rho)) \exp[-|\log(\rho)|^2/4\kappa] \quad (27)$$

where $C_\kappa > 0$ is a constant and $|\log(\rho)|^2 = (\log(\rho), \log(\rho))$. Of course, this f is a positive definite function for each $\kappa > 0$, thanks to Theorem 1.

Example : recall the Gaussian function f defined before Proposition 4. Unlike the functions in the two previous examples, this one is not positive definite. Precisely, there exists no value of σ for which it is positive definite. This can be seen using the only-if part of Theorem 1. According to this theorem, since f is integrable and U -invariant, if f were positive definite, its spherical transform would be identical to the positive function g in (20), up to a constant factor. Indeed, because the spherical transform \hat{f} in (19) satisfies the integrability condition (11), the inversion formula (12) holds true. The fact that $g = C_N \hat{f}$ then follows by injectivity of the inverse spherical transform (after comparing (12) and (20)). Now, to show that f is not positive definite, it is enough to show that $\hat{f}(t) < 0$ for some $t \in \mathbb{R}^N$. Choosing t such that $t_1 < \dots < t_N$, note that $t_\ell - t_k \leq t_N - t_1$ for any $k < \ell$, with equality only if $(k, \ell) = (1, N)$. If $t_N - t_1 \leq \pi$ then $\hat{f}(t) \geq 0$, but as soon as $t_N - t_1 > \pi$, then $\text{sc}((\sigma^2/2)(t_N - t_1))$ becomes the only negative term in the product on the right-hand side of (19), and it then follows that $\hat{f}(t) < 0$.

V. APPLYING RAMANUJAN'S THEOREM

Ramanujan's master theorem for symmetric cones was stated and proved in [10]. Roughly, this theorem converts so-called spherical power series into spherical transforms (or inverse spherical transforms). It is a generalisation of the theorem obtained by Ramanujan, which deals with classical (one-variable) power series and Mellin transforms [23].

Here, the aim is to examine how this theorem can be used in order to construct U -invariant positive definite functions (as in the previous paragraph). In the first place, this is possible because the space M of $N \times N$ complex covariance matrices is indeed a symmetric cone [8].

Spherical power series will be expressed in terms of complex zonal polynomials [24]. The Schur polynomials S_m from (15) in Section III can be normalised (multiplied by certain positive constants) to obtain new polynomials Z_m with the property that

$$(\text{tr}(x))^n = \sum_{[m]=n} Z_m(\rho) \quad \text{where } [m] = m_1 + \dots + m_N$$

for any positive integer n . These are the complex zonal polynomials, and a series of the following form

$$F(x) = \sum_m \frac{(-1)^{[m]}}{[m]!} a(m) Z_m(x) \quad (28)$$

will be called a spherical power series (the sum is over positive integers $m_1 \geq \dots \geq m_N \geq 0$).

Ramanujan's master theorem can be used to prove the following Proposition 6. In this proposition, Γ_M denotes the Gamma function of the symmetric cone M ,

$$\Gamma_M(\lambda) = (2\pi)^{N(N-1)/2} \prod_{k=1}^N \Gamma(\lambda_k - k + 1) \quad \lambda \in \mathbb{C}^N \quad (29)$$

where Γ is Euler's Gamma function of a complex variable [8] [10]. Moreover, the variable x in (28) ranges over the real vector space H of $N \times N$ Hermitian matrices (M is an open cone within H).

Proposition 6: Let $\alpha > N - 1$ and assume that the coefficients $a(m)$ in (28) are given by a function $a : \mathbb{C}^N \rightarrow \mathbb{C}$, of the form

$$a(\lambda) = \Gamma_M(2\alpha + \lambda) \psi(\lambda - \delta) \quad (30)$$

where $\psi : \mathbb{C}^N \rightarrow \mathbb{C}$ is symmetric, holomorphic for $\text{Re}(\lambda_k) > (N - 1)/2 - 2\alpha$, and satisfies the growth condition

$$|\psi(\lambda)| \leq C_N \prod_{k=1}^N e^{P \text{Re}(\lambda_k)} \times e^{A |\text{Im}(\lambda_k)|} \quad (31)$$

where $P, A > 0$ and $A < \pi$. Then, the series (28) converges in a neighborhood of $x = 0$, where it defines a real-analytic function F . Moreover, this function extends analytically to all of M , in such a way that $\Delta^\alpha(x)F(x)$ defines an integrable U -invariant function of $x \in M$ ($\Delta(x) = \det(x)$), which is positive definite if and only if $\psi(it - \alpha) \geq 0$ for all $t \in \mathbb{R}^N$.

The proof of Proposition 6 is given in Appendix E. The idea is to show that $f(x) = \Delta^\alpha(x)F(x)$ defines an integrable U -invariant function with spherical transform

$$\hat{f}(t) = |\Gamma_M(\alpha + \delta + it)|^2 \psi(it - \alpha) \quad (32)$$

and that this $\hat{f}(t)$ satisfies the integrability condition (11), so that the inversion formula (12) is also satisfied.

The statement about positive-definiteness of f then follows by an application of Theorem 1.

Example: in (30), if ψ is a constant function equal to 1, the series (28) becomes a generalised binomial series [8] (Proposition XII.1.3), which converges to $F(x) = \Gamma_M(2\alpha) \Delta^{-2\alpha}(\text{id} + x)$ for any x whose eigenvalues are all < 1 . Proposition 6 then says that, for each $\alpha > N - 1$,

$$f(x) = \Gamma_M(2\alpha) \Delta^\alpha (x(\text{id} + x)^{-2}) \quad (33)$$

is an integrable U -invariant positive definite function, whose spherical transform can be read from (32), by setting $\psi \equiv 1$. In the one-dimensional case ($N = 1$), this function reduces to the density of the Beta-prime distribution [25].

REFERENCES

- [1] A. Azangulov, A. Smolensky, A. Terenin, and S. Borovitsky, “Stationary kernels and Gaussian processes on Lie groups and their homogeneous spaces II: non-compact symmetric spaces,” Apr. 2023. [Online]. Available: arxiv:2301.13088
- [2] —, “Stationary kernels and Gaussian processes on Lie groups and their homogeneous spaces I: the compact case,” Aug. 2022. [Online]. Available: arxiv:2208.14960
- [3] N. Da Costa, C. Mostajeran, J. P. Ortega, and S. Said, “Invariant kernels on Riemannian symmetric spaces: a harmonic-analytic approach,” Oct. 2023. [Online]. Available: arxiv:2310.19270
- [4] —, “Geometric learning with positively decomposable kernels,” Oct. 2023. [Online]. Available: arxiv:2310.13821
- [5] R. Godement, “Introduction aux travaux de Selberg,” *Séminaire Bourbaki*, (no. 144), pp. 95–110, 1958.
- [6] A. M. Yaglom, “Second-order homogeneous random fields,” in *Berkeley Symposium on Mathematical Statistics and Probability*, J. Neyman, Ed., 1961, pp. 593–622.
- [7] S. Helgason, *Groups and geometric analysis: integral geometry, invariant differential operators, and spherical functions*. Academic Press, 1984.
- [8] J. Faraut and A. Korányi, *Analysis on symmetric cones*. Claredon Press, 1994.
- [9] I. M. Gelfand and M. A. Naimark, “Unitary representations of the classical groups,” *Trudy Mat. Inst. Steklov*, vol. 36, pp. 3–288, 1950.
- [10] H. Ding, K. I. Gross, and D. Richards, “Ramanujan’s master theorem for symmetric cones,” *Pacific Journal of Mathematics*, vol. 175, no. 2, pp. 447–490, 1996.
- [11] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*. American Mathematical Society, 2001.
- [12] N. Guigui, N. Miolane, and X. Pennec, *Introduction to Riemannian Geometry and Geometric Statistics*. Now Publishers Inc., 2023.
- [13] S. Said, “Statistical models and probabilistic methods on Riemannian manifolds,” 2021, Université de Bordeaux, Habilitation thesis (HDR).
- [14] X. Pennec, “Intrinsic statistics on Riemannian manifolds: Basic tools for geometric measurements,” *Journal of Mathematical Imaging and Vision*, vol. 25, no. 1, pp. 127–154, 2006.
- [15] R. I. Muirhead, *Aspects of multivariate statistical theory*. Wiley Interscience, 1982.
- [16] P. Mergny, “Spherical integrals and their applications to random matrix theory,” Ph.D. dissertation, Université Paris-Saclay, 2022.
- [17] B. Jonnadula, J. P. Keating, and F. Mezzadri, “Symmetric function theory and unitary invariant ensembles,” *Journal of Mathematical Physics*, vol. 62, 2021.
- [18] L. Santilli and M. Tierz, “Riemannian Gaussian distributions, random matrix ensembles and diffusion kernels,” *Nucl. Phys. B*, vol. 973, p. 115582, 2021.
- [19] S. Said, H. Hajri, L. Bombrun, and B. C. Vemuri, “Gaussian distributions on Riemannian symmetric spaces: statistical learning with structured covariance matrices,” *IEEE Transactions on Information Theory*, vol. 64, no. 2, pp. 752–772, 2018.
- [20] S. Said, S. Heuveline, and C. Mostajeran, “Riemannian statistics meets random matrix theory: toward learning from high-dimensional covariance matrices,” *IEEE Transactions on Information Theory*, vol. 69, no. 1, pp. 472–481, 2023.
- [21] E. S. Meckes, *The random matrix theory of the classical compact groups*. Cambridge University Press, 2019.
- [22] S. Lang and J. Jorgenson, *Spherical inversion on $SL_n(\mathbb{R})$* . Springer-Verlag, 2001.
- [23] G. H. Hardy, *Ramanujan: twelve lectures on subjects suggested by his life and work*. Chelsea Publishing Company, 1978.
- [24] A. Takemura, *Zonal Polynomials*, ser. Institute of Mathematical Statistics Lecture Notes—Monograph Series, 4, 1984.
- [25] N. Balakrishnan, S. Kotz, and N. L. Johnson, *Continuous univariate distributions (Volume 2)*. Wiley interscience, 1995.
- [26] A. W. Knap, *Lie groups beyond an introduction (2nd edition)*. Birkhauser, 2002.
- [27] G. Livan, M. Novaes, and P. Vivo, *Introduction to random matrices: theory and practice*. Springer Cham, 2018.
- [28] F. W. J. Olver, *Asymptotics and special functions*. A. K. Peters Ltd, 1997.
- [29] R. Beals and R. Wong, *Special functions, a graduate text*. Cambridge University Press, 2010.
- [30] S. Bochner and W. T. Martin, *Several Complex Variables*. Princeton University Press, 1948.

APPENDIX A
PROOF OF PROPOSITION 2

The proof starts from (8) and shows that it is equivalent to (9). The first step is to reduce to the case where x has unit determinant. Let $x = r\bar{x}$ where $\det(\bar{x}) = 1$. By a direct calculation, it follows from (7) that $\Delta_s(x) = r^{(1,s)}\Delta_s(\bar{x})$ where $(1, s) = \sum_{k=1}^N s_k$. In particular, $s = \lambda + \delta$ implies that $\Delta_s(x) = r^{(1,\lambda)}\Delta_s(\bar{x})$, because $(1, \delta) = 0$. Replacing into (8) yields the following identity

$$\Phi_\lambda(x) = r^{(1,\lambda)}\Phi_\lambda(\bar{x}) \quad (34)$$

Now, returning to (8), note that

$$\Phi_\lambda(\bar{x}) = \int_U \Delta_{\lambda+\delta}(u \cdot \bar{x}) du \quad (35)$$

This integral can be restricted to SU , the special unitary subgroup of U (SU is the set of $u \in U$ with $\det(u) = 1$). Indeed, replacing u by $e^{i\theta}u$ (with θ real) does not change $u \cdot \bar{x}$. Moreover, the normalised Haar measure on U descends to the normalised Haar measure on SU [26] (see Theorem 8.32). Therefore, (35) is equivalent to

$$\Phi_\lambda(\bar{x}) = \int_{SU} \Delta_{\lambda+\delta}(u \cdot \bar{x}) du \quad (36)$$

The next step of the proof is to show that (36) is the same as the following Harish-Chandra integral [7] (Page 418)

$$\Phi_\lambda(\bar{x}) = \int_{SU} \exp[(2\lambda - 2\delta, \log a(u\bar{\rho}^{1/2}))] du \quad (37)$$

Here, $\bar{\rho} = (\bar{\rho}_1, \dots, \bar{\rho}_N)$ are the eigenvalues of \bar{x} , and $a(u\bar{\rho}^{1/2})$ is the diagonal matrix with positive entries, such that $u\bar{\rho}^{1/2} = na(u\bar{\rho}^{1/2})h$ where the matrix n is upper triangular with ones on its diagonal, and h belongs to SU (vectors such as $\bar{\rho}$ will be identified with diagonal matrices, in a self-evident way, whenever that is convenient).

Because Φ_λ is U -invariant, it is enough to prove (37) when $\bar{x} = \bar{\rho}$. It will be helpful to apply the identity $\Phi_\lambda(x) = \Phi_{-\lambda}(x^{-1})$ [8] (Theorem XIV.3.1). For short, let $a = a(u\bar{\rho}^{1/2})$, so $u \cdot \bar{\rho} = (u\bar{\rho}^{1/2}) \cdot \text{id}$ is equal to na^2n^\dagger . Using the identity just mentioned,

$$\Phi_\lambda(\bar{\rho}) = \Phi_{-\lambda}(\bar{\rho}^{-1}) = \int_{SU} \Delta_{-\lambda+\delta}(u \cdot \bar{\rho}^{-1}) du = \int_{SU} \Delta_{-\lambda+\delta}((n^\dagger)^{-1} \cdot a^{-2}) du \quad (38)$$

where the second equality follows from (36) and the third equality holds because $u \cdot \bar{\rho}^{-1} = (u \cdot \bar{\rho})^{-1}$ and $u \cdot \bar{\rho} = na^2n^\dagger$. However, since $(n^\dagger)^{-1}$ is lower triangular with ones on its diagonal, it is easy to see that $\Delta_{-\lambda+\delta}((n^\dagger)^{-1} \cdot a^{-2}) = \Delta_{-\lambda+\delta}(a^{-2})$. Then, from (7) and the fact that a is diagonal,

$$\Delta_{-\lambda+\delta}((n^\dagger)^{-1} \cdot a^{-2}) = \prod_{k=1}^N a_k^{2(\lambda_k - \delta_k)} = \exp[(2\lambda - 2\delta, \log a)]$$

and (37) follows immediately by replacing this into (38). The final step of the proof exploits the fact that G is a complex Lie group. In this case, the Harish-Chandra integral (37) admits a closed-form expression [7] (Page 432),

$$\Phi_\lambda(\bar{x}) = \frac{\Pi(-\delta)}{\Pi(\lambda)} \times \frac{\sum_{w \in S_N} \varepsilon(w) e^{(\lambda, w\bar{\rho})}}{\sum_{w \in S_N} \varepsilon(w) e^{(\delta, w\bar{\rho})}} \quad (39)$$

Here, $\Pi(\mu) = \prod_{k < \ell} (\mu_k - \mu_\ell)$ for $\mu \in \mathbb{C}^N$, S_N is the symmetric group (group of permutations of N objects), and $\varepsilon(w)$ is the signature of the permutation w , while $w\bar{\rho}$ denotes the action of that permutation on $(\bar{\rho}_1, \dots, \bar{\rho}_N)$.

Clearly, the polynomial Π is the Vandermonde polynomial, up to sign, while the sums in the second fraction are Leibniz expansions of certain determinants. Using these observations, and performing some basic simplifications,

$$\Phi_\lambda(\bar{x}) = \frac{\prod_{k=1}^{N-1} k!}{V(-\lambda)} \times \frac{\det[\bar{\rho}_k^{\lambda_\ell}]}{\det[\bar{\rho}_k^{N-\ell}]} = \frac{\prod_{k=1}^{N-1} k!}{V(-\lambda)} \times \frac{\det[\bar{\rho}_k^{\lambda_\ell}]}{V(-\bar{\rho})} = \prod_{k=1}^{N-1} k! \times \frac{\det[\bar{\rho}_k^{\lambda_\ell}]}{V(\lambda)V(\bar{\rho})} \quad (40)$$

Now, (9) can be retrieved from (34) and (40). To do so, it is enough to note that $\rho = r\bar{\rho}$ and that this implies (recall $\rho = (\rho_1, \dots, \rho_N)$ are the eigenvalues of x)

$$r^{(1,\lambda)} \det[\bar{\rho}_k^{\lambda_\ell}] = \det[\rho_k^{\lambda_\ell}] \quad \text{and} \quad V(\bar{\rho}) = r^{-N(N-1)/2} V(\rho)$$

Then, since $r^N = \det(x)$ (which is the product of the eigenvalues ρ_k),

$$r^{(1,\lambda)} \frac{\det[\bar{\rho}_k^{\lambda_\ell}]}{V(\bar{\rho})} = r^{N(N-1)/2} \frac{\det[\rho_k^{\lambda_\ell}]}{V(\rho)} = \frac{\det[\rho_k^{\lambda_\ell + (N-1)/2}]}{V(\rho)} \quad (41)$$

Therefore, performing the multiplication in (34), with the help of (40) and (41), yields the required (9).

APPENDIX B

PROOF OF PROPOSITION 4

Let $f(x) = \exp[-d^2(x, \text{id})/2\sigma^2]$. Then, note that (16) reads

$$Z(\sigma, \lambda) = \int_M f(x) \Phi_\lambda(x) \text{vol}(dx) \quad (42)$$

Since both f and Φ_λ are U -invariant, this can be evaluated using (2). In terms of the eigenvalues (ρ_1, \dots, ρ_N) , $f(x) = w(\rho_1) \dots w(\rho_N)$ where $w(\rho) = \exp[-\log^2(\rho)/2\sigma^2]$ [18] [20]. On the other hand, $\Phi_\lambda(x)$ is given by (9). Replacing this into (2), it follows that

$$Z(\sigma, \lambda) = \frac{C_N}{V(\lambda)} \times \frac{1}{N!} \int_{\mathbb{R}_+^N} V(\rho) \det[\rho_k^{\lambda_\ell - (N+1)/2}] \prod_k w(\rho_k) d\rho_k$$

Recalling that $V(\rho) = \det[\rho_k^{\ell-1}]$, and applying the Andréief identity (as stated in [27], Chapter 11), this becomes

$$Z(\sigma, \lambda) = \frac{C_N}{V(\lambda)} \times \det \left[\int_0^\infty w(\rho) \rho^{k-1 + \lambda_\ell - (N+1)/2} d\rho \right]_{k,\ell=1}^N$$

By the definition of δ_k (right after Formula (8) in Section III), this is the same as

$$Z(\sigma, \lambda) = \frac{C_N}{V(\lambda)} \times \det \left[\int_0^\infty w(\rho) \rho^{\delta_k + \lambda_\ell - 1} d\rho \right]_{k,\ell=1}^N$$

The integrals inside the determinant can be expressed in terms of the moments of a log-normal probability density.

This gives the first line in (17),

$$Z(\sigma, \lambda) = \frac{C_N}{V(\lambda)} \times \det \left[\sigma \exp \left((\sigma^2/2)(\delta_k + \lambda_\ell)^2 \right) \right]_{k,\ell=1}^N \quad (43)$$

To complete the proof of (17), it is enough to use elementary properties of the determinant,

$$\begin{aligned} \det \left[\exp \left((\sigma^2/2)(\delta_k + \lambda_\ell)^2 \right) \right] &= \det \left[\exp \left((\sigma^2/2)(\delta_k^2 + \lambda_\ell^2 + 2\delta_k \lambda_\ell) \right) \right] \\ &= \prod_{k=1}^N e^{\frac{\sigma^2}{2}(\delta_k^2 + \lambda_k^2)} \times \det \left[\exp \left(\sigma^2 \delta_k \lambda_\ell \right) \right] \end{aligned} \quad (44)$$

Indeed, it is clear that

$$\prod_{k=1}^N e^{\frac{\sigma^2}{2}(\delta_k^2 + \lambda_k^2)} = e^{\frac{\sigma^2}{2}((\lambda, \lambda) + (\delta, \delta))} \quad (45)$$

Moreover, using the definition of δ_k and performing some straightforward simplifications

$$\begin{aligned} \det[\exp(\sigma^2 \delta_k \lambda_\ell)] &= \prod_{k=1}^N e^{\frac{-\sigma^2}{2}(N-1)\lambda_k} \times V(e^{\sigma^2 \lambda}) \\ &= \prod_{k < \ell} 2 \sinh((\sigma^2/2)(\lambda_\ell - \lambda_k)) \end{aligned} \quad (46)$$

Therefore, replacing (45) and (46) into (44),

$$\det\left[\exp\left((\sigma^2/2)(\delta_k + \lambda_\ell)^2\right)\right] = e^{\frac{\sigma^2}{2}((\lambda, \lambda) + (\delta, \delta))} \prod_{k < \ell} 2 \sinh((\sigma^2/2)(\lambda_\ell - \lambda_k))$$

In turn, replacing this into (43) directly yields the second line in (17).

APPENDIX C

PROOF OF THEOREM 1

if part: roughly, the idea of the proof is to show that the spherical functions Φ_{it} are positive definite functions. Then, (20) says that the function f is a positive linear combination of these Φ_{it} and is therefore positive definite. For $x \in M$, let $x = \exp(s)\bar{x}$ where $s = \log \det(x)$ so that $\det(\bar{x}) = 1$. From (34) (putting $\lambda = it$ and $\tau = (1, t)$),

$$\Phi_{it}(x) = e^{i\tau s} \Phi_{it}(\bar{x}) \quad (47)$$

Thinking of s and \bar{x} as functions of x , let $\varphi^\tau(x) = e^{i\tau s}$ and $\bar{\varphi}(x) = \Phi_{it}(\bar{x})$, so that $\Phi_{it}(x) = \varphi^\tau(x)\bar{\varphi}(x)$. Now, recalling the well-known fact that a product of positive definite functions is positive definite, it is enough to show that φ^τ and $\bar{\varphi}$ are both positive definite. For any $x_1, \dots, x_n \in M$, note that

$$\varphi^\tau(x_i^{-1/2} x_j x_i^{-1/2}) = e^{i\tau(s_j - s_i)} \quad (48)$$

where $s_i = \log \det(x_i)$. Therefore, the matrix with elements $\varphi^\tau(x_i^{-1/2} x_j x_i^{-1/2})$ is the same as the matrix with elements $e^{i\tau(s_j - s_i)}$, which is Hermitian non-negative definite (of rank 1). This shows that φ^τ is positive definite.

To see that $\bar{\varphi}$ is also positive definite, note that

$$\bar{\varphi}(x_i^{-1/2} x_j x_i^{-1/2}) = \Phi_{it}(\bar{x}_i^{-1/2} \bar{x}_j \bar{x}_i^{-1/2}) \quad (49)$$

However, according to [7] (Page 484), the restriction of Φ_{it} to the unit-determinant hypersurface (that is to the set of $x \in M$ with $\det(x) = 1$), which is given by the Harish-Chandra integral (37), is a positive definite function.

In particular, the matrix whose elements appear in (49) is Hermitian non-negative definite. This shows that $\bar{\varphi}$ is positive definite. Thus, being a product of positive definite functions, Φ_{it} is positive definite (for any t) as required.

To conclude, recall Godement's theorem [3] [5] (in particular, Formula (2.4) in [3]). This gives a rigorous justification of the claim that f is positive definite because it is a positive linear combination of positive definite functions.

The last step of the proof is thus a direct application of Godement's theorem.

only-if part: as stated in [3], the L^1 -Godement theorem says that an integrable function f is positive definite if and only if its spherical transform is positive and integrable (details can be found in [3], Section 2 and Appendix A).

In [3], it is required from the outset that the underlying symmetric space M should be of non-compact type (in particular, the group G should be semisimple). This requirement is clearly not satisfied, in the present context. The aim here is to explain that the L^1 -Godement theorem can still be applied.

In [3], the requirement that M should be of non-compact type was introduced only in order to ensure that spherical functions on M are given by Harish-Chandra integrals (integrals of the form (37) in the proof of Proposition 2). The proof of the L^1 -Godement theorem (see [3], Appendix A) relies on Godement's (much older) theorem [5], which applies to any symmetric space and in particular to the space M of complex covariance matrices. Specifically, the convolution product of two compactly-supported continuous U -invariant functions on M is commutative [8] (Proposition XIV.4.1), and this is the only hypothesis needed for Godement's theorem.

With this in mind, the proof is just an application of the L^1 -Godement theorem as stated in [3]. Precisely, f is positive definite only if \hat{f} is positive and integrable (which here means it satisfies the integrability condition (11)). Then, f is given by the inversion formula (12), which is the same as (20) with $g = C_N \hat{f}$.

To finish the proof, note that uniqueness of g follows by injectivity of the inverse spherical transform (the linear map that takes \hat{f} to f according to (12)).

APPENDIX D

PROOF OF PROPOSITION 5

Part (a): in order to prove (23), it is enough to show that, on the right-hand side of (22),

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell + (N-1)/2} \right] dt = (\det(x))^{(N-1)/2} \det \left[\int_{\mathbb{R}} \gamma(t) t^{k-1} e^{its_\ell} dt \right] \quad (50)$$

where $g(t) = \gamma(t_1) \dots \gamma(t_N)$ and $s_\ell = \log(\rho_\ell)$. Note first that

$$\det \left[\rho_k^{it_\ell + (N-1)/2} \right] = \prod_{k=1}^N \rho_k^{(N-1)/2} \times \det \left[\rho_k^{it_\ell} \right] = (\det(x))^{(N-1)/2} \times \det \left[\rho_k^{it_\ell} \right]$$

This implies that (50) is equivalent to

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell} \right] dt = \det \left[\int_{\mathbb{R}} \gamma(t) t^{k-1} e^{its_\ell} dt \right]$$

or, what is the same if $V(t)$ is expressed as a determinant,

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) \det [t_\ell^{k-1}] \det \left[\rho_k^{it_\ell} \right] dt = \det \left[\int_{\mathbb{R}} \gamma(t) t^{k-1} e^{its_\ell} dt \right] \quad (51)$$

Here, using the Andréief identity [27] (Chapter 11, Page 75), the left-hand side is equal to

$$\frac{1}{N!} \int_{\mathbb{R}^N} \det [t_\ell^{k-1}] \det \left[\rho_k^{it_\ell} \right] \prod_{\ell=1}^N \gamma(t_\ell) dt_\ell = \det \left[\int_{\mathbb{R}} \gamma(t) t^{k-1} \rho_\ell^{it} dt \right]$$

which is the same as the right-hand side (by definition of s_ℓ). Thus, (51) (equivalent to (50)) has been proven true.

Part (b): comparing (22) and (24), it becomes clear that one must show

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell + (N-1)/2} \right] dt = (\det(x))^{(N-1)/2} i^{N(N-1)/2} V(-\partial/\partial s) \tilde{g}(s) \quad (52)$$

However, as in the proof of Part (a), this is equivalent to

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell} \right] dt = i^{N(N-1)/2} V(-\partial/\partial s) \tilde{g}(s) \quad (53)$$

After writing the Leibniz expansion of the determinant, the left-hand side is equal to

$$\frac{1}{N!} \sum_{w \in S_N} \varepsilon(w) \int_{\mathbb{R}^N} g(t) V(t) \prod_{k=1}^N \rho_k^{it_{w(k)}} dt = \frac{1}{N!} \sum_{w \in S_N} \varepsilon(w) \int_{\mathbb{R}^N} g(t) V(t) e^{i(s, wt)} dt$$

Here, S_N is the group of permutations of $\{1, \dots, N\}$ and $\varepsilon(w)$ is the signature of the permutation w . Moreover, on the right-hand side $s_\ell = \log(\rho_\ell)$ and wt denotes the action of the permutation w on (t_1, \dots, t_N) . By introducing a new variable of integration $u = wt$ in each one of the integrals under the sum,

$$\frac{1}{N!} \sum_{w \in S_N} \varepsilon(w) \int_{\mathbb{R}^N} g(t) V(t) e^{i(s, wt)} dt = \frac{1}{N!} \sum_{w \in S_N} \varepsilon(w) \int_{\mathbb{R}^N} g(w^{-1}u) V(w^{-1}u) e^{i(s, u)} du$$

But the function g is symmetric, while the Vandermonde polynomial V is alternating — for any permutation w , $g(wu) = g(u)$ and $V(wu) = \varepsilon(w)V(u)$. Therefore, the above sum is equal to

$$\frac{1}{N!} \sum_{w \in S_N} \int_{\mathbb{R}^N} g(u) V(u) e^{i(s, u)} du = \int_{\mathbb{R}^N} g(u) V(u) e^{i(s, u)} du$$

and it now follows that the left-hand side of (53) is

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell} \right] dt = \int_{\mathbb{R}^N} g(u) V(u) e^{i(s, u)} du \quad (54)$$

Finally, recalling the definition of the inverse Fourier transform $\tilde{g}(s)$, and differentiating under the integral, one has

$$V(\partial/\partial s) \tilde{g}(s) = i^{N(N-1)/2} \int_{\mathbb{R}^N} g(u) V(u) e^{i(s, u)} du$$

which can be replaced back into (54) to obtain

$$\frac{1}{N!} \int_{\mathbb{R}^N} g(t) V(t) \det \left[\rho_k^{it_\ell} \right] dt = (-i)^{N(N-1)/2} V(\partial/\partial s) \tilde{g}(s) = i^{N(N-1)/2} V(-\partial/\partial s) \tilde{g}(s)$$

which is identical to (53), as required.

APPENDIX E

PROOF OF PROPOSITION 6

The proof relies heavily on Ramanujan's theorem for symmetric cones, as given in [10] (Page 450). This states that if the coefficients $a(m)$ in (28) are of the form $a(m) = q(m - \delta)$ for a function $q : \mathbb{C}^N \rightarrow \mathbb{C}$, which is symmetric, holomorphic for $\text{Re}(\lambda_k) > -L$, where $L > 3(N-1)/2$, and satisfies the growth condition

$$|q(\lambda)| \leq C_N |\Gamma_M(N + \lambda + \delta)| \prod_{k=1}^N e^{P \text{Re}(\lambda_k)} \times e^{A |\text{Im}(\lambda_k)|} \quad (55)$$

where $P, A > 0$ and $A < \pi$, then the following hold.

- (a) the series (28) converges in a neighborhood of $x = 0$, where it defines a real-analytic function F .
- (b) this function extends continuously to all of M , by the following absolutely convergent integral

$$F(x) = C_N \int_{\mathbb{R}^N} \tilde{F}(\sigma + it) \Phi_{\sigma+it}(x) (V(t))^2 dt \quad (56)$$

where $-L + (N-1)/2 < \sigma < -(N-1)$ and $\tilde{F}(\lambda) = \Gamma_M(\delta - \lambda) q(\lambda)$.

- (c) for any σ as above, ω in the convex hull of $w\delta$, where w ranges over the symmetric group S_N ,

$$\tilde{F}(\lambda) = \int_M F(x) \Phi_{-\lambda}(x) \text{vol}(dx) \quad (57)$$

is an absolutely convergent integral, whenever $\lambda = \sigma + \omega + it$ with $t \in \mathbb{R}^N$.

Here, the notation $\sigma + \lambda$, where σ is a number and $\lambda \in \mathbb{C}^N$, means σ is added to each component of λ .

For the proof of Proposition 6, in view of (30), take

$$q(\lambda) = \Gamma_M(2\alpha + \lambda + \delta)\psi(\lambda) \quad (58)$$

which is then symmetric, because ψ is symmetric by assumption and $\Gamma_M(\lambda + \delta)$ is a symmetric function of λ , as can be seen after replacing $\delta_k = k - (N + 1)/2$ into (29), and holomorphic for $\text{Re}(\lambda_k) > -L$, where $L = 2\alpha - (N - 1)/2 > 3(N - 1)/2$ (recall that $\alpha > N - 1$), again by the assumptions made regarding ψ and the fact that the Gamma function $\Gamma_M(\lambda)$ is holomorphic for $\text{Re}(\lambda_k) > N - 1$, as one may see from (29). Moreover, this q satisfies the growth condition (55), since (31) implies

$$|q(\lambda)| \leq C_N |\Gamma_M(2\alpha + \lambda + \delta)| \prod_{k=1}^N e^{P\text{Re}(\lambda_k)} \times e^{A|\text{Im}(\lambda_k)|} \quad (59)$$

and since, from (29), after putting $z_k = \lambda_k + 2\alpha - (N - 1)/2$,

$$\frac{\Gamma_M(2\alpha + \lambda + \delta)}{\Gamma_M(N + \lambda + \delta)} = \prod_{k=1}^N \frac{\Gamma(z_k)}{\Gamma(z_k + N - 2\alpha)} = \prod_{k=1}^N z_k^{2\alpha - N} (1 + O(|z_k|^{-1})) \quad (60)$$

where the second equality follows from the asymptotic form for the ratio of two Gamma functions [28] (Page 119).

Indeed, by (59) and (60),

$$|q(\lambda)| \leq C_N \times \prod_{k=1}^N z_k^{2\alpha - N} (1 + O(|z_k|^{-1})) \times |\Gamma_M(N + \lambda + \delta)| \prod_{k=1}^N e^{P\text{Re}(\lambda_k)} \times e^{A|\text{Im}(\lambda_k)|}$$

and this implies (55) because $|z|^{2\alpha - N} = o(e^{\varepsilon(\text{Re}(z) + \text{Im}(z))})$ for any $\varepsilon > 0$, in the limit where $|z| \rightarrow \infty$. Thus, the conditions of Ramanujan's master theorem are all verified for q given by (58), so items (a) to (c) above can be applied in the context of Proposition 6. First, (a) implies the spherical series (28) defines a real-analytic function F in the neighborhood of $x = 0$. Moreover, (b) and (c) imply that this F extends continuously to all of M , in such a way that (56) and (57) are satisfied. Let $\sigma = -\alpha$ in (56) and note that (8) implies $\Phi_{it - \alpha}(x) = \Delta^{-\alpha}(x)\Phi_{it}(x)$. Then, (56) and the subsequent definition of $\tilde{F}(\lambda)$, applied with q as in (58), show that

$$\Delta^\alpha(x)F(x) = C_N \int_{\mathbb{R}^N} |\Gamma_M(\alpha + \delta + it)|^2 \psi(it - \alpha)\Phi_{it}(x)(V(t))^2 dt \quad (61)$$

Now, letting $f(x) = \Delta^\alpha(x)F(x)$, it becomes clear f is U -invariant, since each spherical function Φ_{it} is U -invariant. On the other hand, choosing $\sigma = -\alpha$, $\omega = \delta$, and $t = 0$ in (57), and noting that $\Phi_{-\delta}(x) = 1$ is a constant function, it follows that f is integrable. The spherical transform of f is found from (57), with $\lambda = -\alpha + it$, which shows that this spherical transform is equal to \hat{f} in (32).

Before pursuing the final stage of the proof, note that \hat{f} (given by (32)) satisfies the integrability condition (11). Indeed, (31) and (32) imply that

$$|\hat{f}(t)| = |\Gamma_M(\alpha + \delta + it)|^2 |\psi(it - \alpha)| \leq |\Gamma_M(\alpha + \delta + it)|^2 \times C_N e^{-NP\alpha} \prod_{k=1}^N e^{A|t_k|}$$

However, by (29) and [29] (Corollary 2.5.3)

$$\begin{aligned} |\Gamma_M(\alpha + \delta + it)|^2 &= (2\pi)^{N(N-1)} \prod_{k=1}^N |\Gamma(it_k + \alpha - (N-1)/2)|^2 \\ &= (2\pi)^{N^2} \prod_{k=1}^N |t_k|^{\alpha-N/2} e^{-\pi|t_k|} (1 + O(|t_k|^{-1})) \end{aligned}$$

and it therefore follows

$$|\hat{f}(t)| \leq C_N e^{-NP\alpha} \prod_{k=1}^N |t_k|^{\alpha-N/2} e^{(A-\pi)|t_k|} \quad (62)$$

But since $A < \pi$, this is exponentially small as the $|t_k|$ become large. In turn, this shows that (11) is satisfied.

To complete the proof of Proposition 6, it remains to show that F in (61) is real-analytic (so that it gives an analytic extension, rather than just the continuous extension stated in item (b)), and also that f is positive definite if and only if $\psi(it - \alpha) \geq 0$ for all $t \in \mathbb{R}^N$. To do so, rewrite (61) as follows

$$\Delta^\alpha(x)F(x) = C_N \int_{\mathbb{R}^N} \hat{f}(t) \Phi_{it}(x) (V(t))^2 dt = \frac{C_N}{V(i\rho)} \int_{\mathbb{R}^N} \hat{f}(t) V(t) \det[\rho_k^{it_\ell + (N-1)/2}] dt$$

where the second equality follows from (12) and (14) (here, the factor $1/N!$ from (14) has been absorbed into C_N).

This easily simplifies to

$$\Delta^{\alpha-(N-1)/2}(x)F(x) = \frac{C_N}{V(i\rho)} \int_{\mathbb{R}^N} \hat{f}(t) V(t) \det[\rho_k^{it_\ell}] dt \quad (63)$$

Now, replace ρ_k with the complex variable $\chi_k = \rho_k e^{i\varphi_k}$ where φ_k is real and $|\varphi_k| < \varepsilon$ with $\varepsilon < \pi/2$ and $(\pi - A)/2$ (here, $A < \pi$ is the constant in (62)). With this replacement, the right-hand side of (63) becomes

$$\Psi(\chi) = \frac{C_N}{V(i\chi)} \int_{\mathbb{R}^N} \hat{f}(t) V(t) \det[\chi_k^{it_\ell}] dt \quad (64)$$

The aim is to show that Ψ is a holomorphic function of χ (each χ_k having its argument φ_k subject to $|\varphi_k| < \varepsilon$). It will then follow that (63) defines a real-analytic function of ρ (the restriction of a holomorphic function of χ). Moreover, since $\Delta^{\alpha-(N-1)/2}(x)$ is analytic and non-zero, as a function of ρ , this will show that F is real-analytic. To see that Ψ is holomorphic, note that the integral in (64) defines a holomorphic function of χ – call this $I(\chi)$. Indeed, the function under that integral is holomorphic for each fixed t , and the integral converges uniformly in χ , due to the following upper bound (the fact that the integral in (64) is holomorphic then follows from [30] (Page 41))

$$\left| \hat{f}(t) V(t) \det[\chi_k^{it_\ell}] \right| \leq |P(t)| \prod_{k=1}^N e^{(A-\pi)|t_k|} \times \left| \det[\chi_k^{it_\ell}] \right| \leq |P(t)| \prod_{k=1}^N e^{(A-\pi+\varepsilon)|t_k|}$$

where $P(t)$ is some polynomial and $A - \pi + \varepsilon < 0$. Here, the first inequality follows from (62), and the second one since $|\chi_k^{it_\ell}| = e^{-t_\ell \varphi_k}$. Therefore, $\Psi(\chi) = C_N I(\chi)/V(i\chi)$ is a ratio of holomorphic functions, and is holomorphic because any zero of $V(i\chi)$ is also a zero of $I(\chi)$ (note that zeros of $V(i\chi)$ occur when $\chi_k = \chi_\ell$ for some $k < \ell$).

The final part of the proof requires showing that f (defined after (61)) is positive definite if and only if $\psi(it - \alpha) \geq 0$ for all $t \in \mathbb{R}^N$. Recall that f is integrable and its spherical transform is \hat{f} given in (32). Recall also that \hat{f} satisfies the integrability condition (11), as shown above, using (62). Therefore, f is given by the inversion formula (12). Accordingly, from (12) and (20) of Theorem 1, by injectivity of the inverse spherical transform, f is positive definite if and only if $\hat{f}(t) \geq 0$ for all $t \in \mathbb{R}^N$. This happens precisely when $\psi(it - \alpha) \geq 0$ for all $t \in \mathbb{R}^N$ (this is by (32)).