



New objects of study for mass spectrometry by secondary-ion and laser desorption-ionization: Acritarchs, a pilot study of the species " *Gloeocapsomorpha prisca*"

Mathilde Bon, Kevin Lepot, Yvain Carpentier, Fabrice Bray, Armelle Riboulleau, François Baudin, Nicolas Nuns, Maxime Bridoux, Christian Rolando, Philippe Steemans, et al.

► To cite this version:

Mathilde Bon, Kevin Lepot, Yvain Carpentier, Fabrice Bray, Armelle Riboulleau, et al.. New objects of study for mass spectrometry by secondary-ion and laser desorption-ionization: Acritarchs, a pilot study of the species " *Gloeocapsomorpha prisca* ". Goldschmidt2023, Jul 2023, Lyon, France. 10.7185/gold2023.19548 . hal-04525545

HAL Id: hal-04525545

<https://hal.science/hal-04525545>

Submitted on 28 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New objects of study for mass spectrometry by secondary-ion and laser desorption-ionization: Acritarchs, a pilot study of the species "*Gloeocapsomorpha prisca*"

MATHILDE BON^{1,2,3}, KEVIN LEPOT^{1,4}, YVAIN CARPENTIER², FABRICE BRAY⁵, ARNELLE RIBOULLEAU¹, FRANÇOIS BAUDIN⁶, NICOLAS NUNS⁷, MAXIME CYRIL BRIDOUX⁸, CHRISTIAN ROLANDO⁵, PHILIPPE STEEMANS⁹ AND THIJS R. A. VANDENBROUCKE³

¹Univ. Lille, CNRS, Univ. Littoral Côte d'Opale, UMR 8187 – LOG – Laboratoire d'Océanologie et de Géosciences

²Univ. Lille, CNRS, UMR 8523 – PhLAM – Physique des Lasers Atomes et Molécules

³Department of Geology (WE13), Ghent University

⁴Institut Universitaire de France (IUF)

⁵Univ. Lille, CNRS, USR 3290 – MSAP – Miniaturisation pour la Synthèse, l'Analyse et la Protéomique

⁶SU CNRS, ISTeP UMR 7193

⁷Univ. Lille, CNRS, Centrale Lille, Univ. Artois, UMR 8181 - UCCS - Unité de Catalyse et Chimie du Solide

⁸CEA

⁹EDDY Lab/Palaeobotany and Palaeopalynology, Univ. Liège, 4000 Liège, Belgium

Presenting Author: mathilde.bon2@univ-lille.fr

Acritarchs are organic-walled microfossils of uncertain biological affinity. The species *G. prisca* (~10 µm), a probable cyanobacterium, comprises the bulk of the insoluble organic matter (kerogen) from a rock type named “kukersite”^{1–5}. *G. prisca* has been studied by gas chromatography-mass spectrometry (GC-MS) using extraction, chemolysis, and pyrolysis^{1–4}.

Here we test the potential of laser-assisted mass spectrometry techniques to analyze the molecular composition of single microfossils. Bitumen and kerogen (dominated by the *G. prisca* microfossils) were analyzed after organic solvent extractions and demineralization of a sample from a 460 million year-old deposit of north-western Russia⁵. Using secondary ion time-of-flight mass spectrometry (ToF-SIMS), we carry out mass spectrometric imaging with ~1 µm spatial resolution and high surface sensitivity. We perform high resolution two-step laser desorption-ionization mass spectrometry (HR-µL2-MS) with a 140 µm probe and laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry (LDI-FT-ICR-MS) with laser probes down to less than 20 µm.

The ToF-SIMS and HR-µL2-MS analysis of microfossils are dominated by abundant (poly)aromatic hydrocarbons. Unique formula assignments were possible for the ions that dominate the signal between ca. 150–450 m/z in LDI-FT-ICR-MS. Thus, we identify a few thousands of molecular formulae and show

aromatic hydrocarbons, oxygenated and nitrogenous compounds, with a major contribution of the O₂–₄ compounds. This predominance is consistent with the organic composition of the *G. prisca* wall deduced from pyrolysis- and chemolysis-assisted GC-MS analyses^{1–4}. Importantly, the bitumen and the microfossils showed distinct molecular signatures with all techniques.

The potential effects of desorption lasers in LDI-FT-ICR-MS, such as fragmentation and pyrolysis, were examined using principal component analysis. Our analytical method and data analysis workflow is expected to help future molecular discrimination of microfossils from heterogenous assemblages, and can help the search for molecular signatures for exobiology⁶.

¹ Blokker P *et al.* (2001) *Geochimica et Cosmochimica Acta* 65.

² da Silva TF *et al.* (2016) *International Journal of Coal Geology* 168.

³ Derenne S *et al.* (1990) *Organic Geochemistry* 16.

⁴ Derenne S *et al.* (1992) *Organic Geochemistry* 19.

⁵ Raevkaya E *et al.* (2004) *IGCP* 503.

⁶ Goesmann F *et al.* (2017) *Astrobiology* 17.

