



HAL
open science

Biomolecular modifications during keratinocyte differentiation: Raman spectroscopy and chromatographic techniques

Joudi Bakar, Rime Michael-Jubeli, Sana Tfaili, Ali Assi, Arlette Baillet-Guffroy, Ali Tfayli

► To cite this version:

Joudi Bakar, Rime Michael-Jubeli, Sana Tfaili, Ali Assi, Arlette Baillet-Guffroy, et al.. Biomolecular modifications during keratinocyte differentiation: Raman spectroscopy and chromatographic techniques. *Analyst*, 2021, 146 (9), pp.2965-2973. <10.1039/d1an00231g>. <hal-04528250>

HAL Id: hal-04528250

<https://hal.science/hal-04528250v1>

Submitted on 1 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Analyst

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: J. bakar, R. Michael-Jubeli, S. Tfaili, A. ASSI, A. Baillet-Guffroy and A. Tfayli, *Analyst*, 2021, DOI: 10.1039/D1AN00231G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Biomolecular modifications during the keratinocytes differentiation: Raman spectroscopy and chromatographic techniques.

Joudi Bakar, Rime Michael-Jubeli*, Sana Tfaili*, Ali Assi, Arlette Baillet-Guffroy, Ali Tfayli

Lipides : systèmes analytiques et biologiques, Université Paris-Saclay, 92296 Châtenay-Malabry, France.

Author(s):

- 1- Joudi Bakar : joudi.bakar@u-psud.fr, Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France.
- 2- Rime Michael-Jubeli : **This author accepts correspondance and proofs**, rime.michael-jubeli@u-psud.fr, Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France. Orcid ID : 0000-0002-2944-5295.
- 3- Sana Tfaili : **This author accepts correspondance and proofs**, sana.tfaili@u-psud.fr, Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France. Orcid ID : 0000-0002-6256-6777
- 4- Ali Assi: ali.assi@u-psud.fr Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France.
- 5- Arlette Baillet-Guffroy: arlette.baillet-guffroy@u-psud.fr, Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France.
- 6- Ali Tfayli : ali.tfayli@u-psud.fr, Université Paris-Saclay, Lipides : systèmes analytiques et biologiques, 92296, Châtenay-Malabry, France.

***Corresponding authors:** Sana Tfaili, 5 rue Jean-Baptiste Clément - 92290 Châtenay-Malabry – France, Tel: +33146835463, fax: +33 Fax: +33149835458, sana.tfaili@u-psud.fr

Rime Michael-Jubeli, 5 rue Jean-Baptiste Clément - 92290 Châtenay-Malabry – France, , Tel: +33146835904, fax: +33 Fax: +33149835458, rime.michael-jubeli@u-psud.fr

Abstract

From basal layer until *stratum corneum*, lipid and protein biomarkers associated to morphological changes denote keratinocytes differentiation and characterize each epidermis layer. Herein, we followed keratinocytes differentiation in early stages on HaCaT cells over a period of two weeks by two complementary analytical techniques: Raman microspectroscopy and High-Performance Liquid Chromatography coupled to High resolution Mass Spectrometry. High concentration of calcium in the medium induced HaCaT differentiation *in vitro*. Results from both techniques underlined keratinocytes passage from granular layer (Day 9) to *stratum corneum* layer (Day 13). After 13 days of differentiation, we observed a strong increase in lipids content, decrease in proteins, decrease in DNA, and a decrease in glucosylceramides/ceramide and sphingomyelins /ceramides ratios.

Keywords: Keratinocytes differentiation, Raman microspectroscopy, HPLC/HR-MS, lipid biomarkers.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abbreviation

APPI: Atmospheric Pressure PhotoIonization

CER: Ceramide

D: Differentiation Day

FFA: Free Fatty Acids

HPLC: High-Performance Liquid Chromatography

HR-MS: High-Resolution Mass Spectrometry

NPLC: Normal-Phase Liquid Chromatography

SC: *Stratum Corneum*

SG: *Stratum Granulosum*

Analyst Accepted Manuscript

Introduction:

Epidermis – the outermost layer of the skin – consists of several stratified layers of keratinocytes: basal, spinous, granular and cornified layers¹. These layers differ by their thickness, their cells morphology, the proliferative activity and the expression of different differentiation markers². The basal layer (*stratum basale*) consists of large and columnar cells – epidermal stem cells – attached to the basement membrane by hemidesmosomes³. Stem cells have a proliferative activity and give rise to cells that can migrate and differentiate through the three other layers⁴. The spinous layer (*stratum spinosum*) is a suprabasal layer. Cells of this layer are attached together by desmosomes⁵. In the granular layer (*stratum granulosum*), keratohyalin granules and lamellar bodies appear, the latter are rich in lipids and catabolic enzymes. Lamellar bodies fuse with plasma membrane and release their lipid content at the interface between the granular and cornified layers. The secreted lipids are rearranged and assembled into lamellar structure. These structures fill the intercellular region of *stratum corneum* (SC) and surround the corneocytes to form a compact intercorneocytes matrix which plays a key role in skin barrier function^{3,6}. When the keratinocytes reach the uppermost layer of the skin (SC), they lose their capacity for cell division, lose their nucleus and acquire a flatter shape to become corneocytes. Corneocytes are packed with keratin filaments and embedded in a lipid matrix organized in lamellae consisting mainly of long-chain ceramides (CER), free fatty acids (FFA), and cholesterol^{6,7}.

Keratinocytes differentiation stages go together with morphological changes. These changes are generally confirmed biochemically by immunoblotting and immunofluorescence of proteins and genes known as differentiation markers in the epidermis such as keratin 1 involucrin, and filaggrin⁸. In parallel with protein modifications and morphological changes, lipids composition and concentrations evolve along differentiation. In fact, epidermal lipid biosynthesis is initiated at early stage of keratinocytes differentiation. Ceramides are generated in the endoplasmic reticulum of *Stratum Granulosum* (SG) cells, they consist of fatty acid chains linked to sphingoid bases via amide bonds⁹⁻¹¹.

1
2
3 Ceramide biosynthesis increases along keratinocytes differentiation, ceramides are subjected to changes –
4 probably to avoid their intracellular cytotoxicity level – in the Golgi apparatus; *i.e.*/ glycosylation to give
5 the glucosylceramides and phosphorylation to generate Sphingomyelins¹¹. The glucosylceramides and
6 Sphingomyelins are then packaged into epidermal lamellar bodies (LB).

7
8
9
10 At the interface between the SG and SC, lamellar bodies release their lipid content of glucosylceramides
11 and sphingomyelins. b-glucocerebrosidase and sphingomyelinase convert back respectively
12 glucosylceramides and sphingomyelins to their ceramides species^{3,6}. Therefore, ceramides are identified as
13 SC markers, and glucosylceramides and sphingomyelins are identified as SG markers in keratinocytes
14 differentiation stages.

15
16
17
18
19
20
21
22
23
24
25 Spontaneously immortalized Human cell line HaCaT – often used as a keratinocytes model – are near to
26 normal phenotype, possess the capacity of proliferating and differentiating, and exhibit all major
27 differentiation surface markers^{8, 12}. Primary human keratinocytes are perfect to study differentiation; but
28 some drawbacks such as slow growth rate complicate cell culture and prevent long-term investigation of
29 differentiation signals¹²⁻¹⁴. HaCaT cells have the advantage to permit following differentiation process
30 during a long period (two weeks).

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Several modifications of culture medium conditions are described to stimulate keratinocytes differentiation
*in vitro*¹⁵. Ionic calcium has been shown to be a major agent in controlling keratinocytes differentiation *in*
vitro and *in vivo*, it stimulates lamellar bodies migration which secret their lipid content at interface between
the SG and SC^{6, 16}.

59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

In this work, HaCaT cells were first cultured in a low calcium medium to induce dedifferentiation then they
were subjected to a high calcium concentration (2.8 mM) medium to induce their differentiation during a
period of 13 days *in vitro*. The differentiation process was followed using Raman spectroscopy which
enables a direct molecular characterization with no labeling¹⁷. Raman spectroscopy enables following cells
differentiation at single cell level. In addition to the advantage of Raman as a free label diagnostic technique,

1
2
3 it provides a phenotype-specific biomolecular fingerprint for the different molecular components in the
4 sample. Complementary information was obtained using High-performance Liquid Chromatography
5 coupled to High Resolution Mass Spectrometry (HPLC/HR-MS) and (HPLC/Corona) permitted to follow
6 lipid biomarkers evolution classes (ceramide, glucosylceramides and sphingomyelins). These two analytical
7 methods are complementary and constitute an efficient methodology to study differentiation biomolecular
8 changes and to determine keratinocytes differentiation stages.
9

Materials and methods

1- Cell culture

10
11 HaCaT cells were cultured in DMEM High Glucose, calcium and serum free media (Life Technologies),
12 and supplemented with 10% fetal bovine serum (sigma) treated with Chelex 100 to remove endogenous
13 calcium, 100 U/mL of penicillin, 100 µg/mL of streptomycin, 2% glutamine solution 200mM (Sigma) and
14 1% of calcium chloride solution 3mM (Sigma) to obtain a final concentration of 0.03mM. HaCaT cells were
15 maintained for at least 3 weeks in low calcium media to dedifferentiate to the basal phenotype¹⁸. High
16 calcium concentration induces a rapid morphological change of keratinocytes *in vitro*^{1, 12}. To induce the
17 differentiation, the low-calcium medium (0.03mM) was replaced by a high-calcium medium (2.8 mM as
18 final concentration). Cultures were maintained at 37C° in a humidified atmosphere with 5% CO₂. Since the
19 HaCaT cells are very sensitive to cell density, cells in low calcium medium should be detached using
20 trypsin-EDTA when they reach 70% confluence, otherwise they may start to differentiate at higher density⁸.
21 The HaCaT cells were graciously obtained from Nanomédicaments et nanosondes laboratory, Tours
22 University; France (Source: Thermo Fisher scientific, Les Ulis, France).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1.1- Cell culture preparation for Raman microspectroscopy analyses

The differentiation process of HaCaT keratinocytes was followed over 13 days. For Raman analysis, CaF₂
windows were introduced in well plates during cell culture. To reach the required cell density at each
differentiation day, basal HaCaT cells were seeded in a 24 well plate on sterile CaF₂ windows. Cells were

deposited in a descending order: reducing number of cells compensates continuing cell proliferation occurring during differentiation process and permits to harvest cells subsequently.

HaCaT were first cultured for 24 h under a low calcium medium to allow cells adhesion. After 24 h, the low-calcium medium was replaced with high-calcium medium to initiate the differentiation process. The medium was changed every three days. At each collection step 2 hours, 3, 6, 9 and 13 days, HaCaT cells were washed twice with phosphate buffered saline (PBS) and then fixed with formalin (4%, 15 min). Cells fixed on CaF₂ window were then washed again with PBS and water.

1.2- Cell culture preparation for HPLLC/HR-MS, HPLC/corona analyses

A decreased density of cells was seeded directly – without CaF₂ windows – into 75-cm² flasks in presence of low calcium medium and incubated for 24h in 5% CO₂, 37C°. Decreased cells density permits to not exceed 60-70% of confluence at each collection step. To follow the changes in lipid composition during keratinocytes differentiation process, the low calcium medium was replaced after 24h by the high one and maintained for 13 days.

2- Raman microspectroscopy

2.1-Instrumentation Raman spectrometer

Raman spectral acquisition and spectral imaging were collected using a Labram HR microspectrometer (Horiba scientific, Palaiseau, France). A 633 nm He:Ne laser (Toptica Photonics, Munich, Germany) giving a 10 mW power on the sample was used. The laser beam was focused using a 100X short focal microscope objective (Olympus, Tokyo, Japan). The collected light was filtered with a Edge filter and dispersed with a 4 cm⁻¹ spectral resolution using a 300 μm slit and 300 grooves/mm holographic grating. The Raman Stokes signal was recorded with a Synapse Charge-Coupled Device detector: CCD camera (Andor Technology, Belfast, UK) containing 1024x256 pixels. Spectral acquisition was performed using Labspec 6 software (Horiba Scientific, Lille, France). Raman measurements were performed on the SC surface in the 400–3800 cm⁻¹ spectral range. For each spectrum the acquisition time was 2 x 5 seconds. The used experimental set-up enables to reach spatial resolution down to 0.5x0.5 μm². The resolution limit depends on the Rayleigh

1
2
3 diffraction criterion and can be defined by the laser wavelength and the numerical aperture of the microscope
4 objective. The other determining factor for the spatial resolution is the chosen step size for mapping. In this
5 study, we have used a $1 \times 1 \mu\text{m}^2$ (for X and Y lateral steps).
6
7
8
9

10 11 **2.2- Raman spectral processing**

12 Acquisition and pre-processing of all spectra were performed using a Labspec 6 software (Horiba Scientific,
13 Palaiseau, France). All spectra were subjected to the same pre-processing protocol. All spectra were
14 smoothed using Savitzky-Golay algorithm (4th order, 9 points), normalized by applying unit vector method
15 and baseline corrected using an automatic polynomial function. To follow the significant variations in cells
16 spectra, Kruskal-wallis (a non-parametric test) was applied using in house developed software with Matlab
17 (Mathworks, Natick, MA, USA).
18
19

20 For spectral imaging, single cells were scanned with $1 \mu\text{m}$ step (X and Y) with an acquisition time of 2x5
21 seconds for each point.
22
23

24 **3- Lipid extraction**

25 Cells were dispersed using trypsin-EDTA, then cell pellets were dissolved in 600 μL of distilled water,
26 mixed with a magnetic stirrer for 10 minutes at room temperature. Total lipids were extracted by the
27 modified method of Bligh and Dyer¹⁹. Briefly, 1.5 mL of methanol was added and vortexed for 2 minutes.
28 Then, 3 mL of chloroform containing 1-Eicosanol as an internal standard (0.3 mg/mL on chloroform) was
29 also added, the mixture was incubated for 1 hour at room temperature in the dark. The organic phase was
30 collected and evaporated under a stream of nitrogen, then lipid residues were resuspended with 100 μL of
31 chloroform^{20, 21}.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4- Instrumentation and analytical method (HPLC/corona and HPLC/HR-MS) for biomarkers analysis

The analysis of the extracted lipid was conducted by HPLC (Dionex Ultimate 3000, ThermoFisher Scientific, San Jose, CA). Separation of lipid occurred in NPLC using (PVA)-Sil column (PVA-bonded column; 5 μm particle size, 150 \times 4.6 mm) purchased from YMC (Kyoto, Japan). A solvent gradient of Heptane/ethyl acetate (99.8: 0.2 v/v) (Solvent A), acetone and ethyl acetate (2: 1 v/v) with acetic acid (0.02% v/v) (Solvent B) and 2-propanol/ water (85:15v/v) (with acetic acid and ethanolamine each at 0.025% v/v) (solvent C) were used. Flow rate was set to 1 mL/min and injected volume was 10 μL . Detection was performed using two detectors simultaneously: a Corona CAD[®] and a hybrid mass spectrometer LTQ-Orbitrap Velos Pro (Thermo Fisher Scientific, San Jose, CA) equipped with an Atmospheric Pressure Photoionization (APPI) source. APPI positive ion mode was used to analyze the extracted lipid. The mass scan range was 220-1800 atomic mass units (amu). A full scan detection mode and fragmentation was performed by the orbital trap at a resolution of 100 000. The analytical conditions are detailed in a previous work²¹.

Results:

1- Morphological evolution

The first step of the study consisted of the dedifferentiation of the HaCaT cells by incubating them in a low calcium concentration medium (0.03) mM for 3 weeks. Non differentiated keratinocytes (ND) had a morphology closer to the spindle shape. They are less compacted and with no contact between cells as shown in figure 1(A). In order to induce HaCaT cells differentiation process, medium was replaced by a high calcium concentration medium (2.8 mM). As shown in figure 1 (B), morphological changes were observed following the keratinocytes differentiation. After 2H of exposure, cells began to exhibit a more cuboidal shape with close packing. Fusiform cells were detected at D9 and D13 days (orange arrows in figure 1 (B)). Stratified layers appeared from D9 and became thicker at D13. Cells overlapped at D13 and made impossible to determine the limit at individual cell level (Day 13 in figure 1 (B) and figure 3).




Figure 1: Morphological evolution of HaCaT cells during the differentiation process. A) Microscopic image after 3 weeks of incubation in low calcium concentration medium (0.03 mM). B) White light images (x10, and x100) during different period (2H, 3, 6, 9, and 13 days) of incubation in high calcium concentration medium (2.8 mM). Morphological modification in the cellular shape was observed. The orange arrows point the formation of fusiform cells and the stratified layers which appeared at D9 and became thicker at D13. ND*: undifferentiated HaCaT, h: hours, D: days.

1- Raman spectroscopy: Spectral features of HaCaT cells

The spectral signatures of HaCaT cells at each differentiation day were achieved using Raman microspectroscopy. In order to focus on the evolution of lipids along differentiation, spectra from cytoplasm were collected. Figure 2 illustrates mean cytoplasm spectra at each differentiation day. The presence of a band at 960 (953-978) cm^{-1} can visually be observed on the spectra. The 960 cm^{-1} band appeared after 6 days of differentiation and clearly increased after 13 days. Spectra collected at D13 on stratified layers showed the highest integrated intensities of this band compared to other regions. The 960 cm^{-1} band could be assigned to an increase in saturated ceramides amounts, as it was mentioned in a comparative study

1
2
3 between unsaturated and saturated ceramides²². Another possible assignment could be an increase in the
4
5 keratin level based on the band attribution in²³ table 1.
6

7
8 In order to highlight significant spectral modifications, Kruskal-Wallis was applied between spectra issued
9
10 from cells at different differentiation days. P-value was fixed to 0.05 and the spectral regions showing
11
12 significant variations were identified and listed in table 1.
13

14
15 A decrease in the bands related to phospholipids at D3 (598 cm⁻¹, 831 cm⁻¹ (v_{asym} O-P-O)), followed by a
16
17 slight increase at D6. While no changes in phospholipids were observed between D6 and D9, a marked
18
19 decrease was shown at D13 (598 cm⁻¹, 831 cm⁻¹ (v_{asym} O-P-O)) (table 1). The decrease in the phospholipids
20
21 relative content between D0 and D3 can also be observed by following alkyl chain related vibrations: vCC
22
23 (1086 cm⁻¹), δ CH₂ twisting (1305 cm⁻¹) and δ CH scissoring (1448 cm⁻¹).
24

25
26 The observation of cholesterol and cholesterol esters related bands showed no modification at D3 while
27
28 higher intensity appeared at D6 (700 cm⁻¹) and the increase continued for D9 (607 cm⁻¹). Cholesterol and
29
30 cholesterol esters are known to markedly increase in the SG.
31

32
33 The most marked evolution was observed in the spectral features related to lipid chains between D9 and
34
35 D13 by mainly a high marked increase in the CH₂ stretching bands (2851 and 2882 cm⁻¹). This increase was
36
37 also observed for CC stretching (1086 cm⁻¹) and δ CH₂ twisting (1305 cm⁻¹). Lipid increase may be related
38
39 to an advanced stage in the differentiation *i.e.* the transition between SG and SC. The decrease in the δ CH
40
41 scissoring (1448 cm⁻¹) was more related to conformational modifications in other molecular components
42
43 (proteins) than lipids²⁴.
44

45
46 The 3059 cm⁻¹ band (=CH stretching) showed a continuous decrease in unsaturated moieties from D6 till
47
48 D13.
49

50
51 In addition to the modifications observed in the lipid related spectral features, changes were observed for
52
53 other molecular features. For instance, between D9 and D13, a marked decrease was automatically identified
54
55

1
2
3 by Kruskal-Wallis in the spectral region (705-736 cm^{-1}) assigned to nucleic acids (table 1). DNA variation
4
5 accords with the observations obtained from lipid related bands suggesting the end of SG phase (figure 3).
6

7
8 An interesting observation was also shown for amino-acids and protein related features. From D0 to D3,
9
10 significant decreases were observed for amino-acids related bands, *i.e.* cytosine (ν_{CS} at 669 cm^{-1}), tryptophan
11
12 (ring breathing at 760 cm^{-1}), tyrosine (855 cm^{-1} (ring breathing), 1044 cm^{-1}) and proline (855 cm^{-1}). The
13
14 decrease in the proteins spectral signature contributes in the decrease in the CH scissoring band at 1448
15
16 cm^{-1} .
17

18
19 From D3 to D6, the significant changes in proteins are mostly structure related. For instance, the increase
20
21 in the 960 cm^{-1} band was explained by an increase in the β -sheets in keratin (table 1). In parallel, a shift to
22
23 higher wavenumbers of the symmetric CH_3 stretching band can be associated with unfolding of proteins.
24
25 The modifications in the secondary structure of proteins was also observed from D6 to D9 with an increase
26
27 in the α helix related sub-band in the Amide I band (table1). An increase in the CH scissoring occurred also.
28
29

30
31 Finally, at D13, a decrease in the relative intensities of CH_3 stretching bands mainly associated to proteins
32
33 is observed. This was described previously^{25, 26} and is associated to skin barrier function.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: At each differentiation day, the mean cytoplasmic spectrum of several cells was calculated ($n=12$). The mean spectra of HaCaT obtained after high calcium concentration exposure at (2H, 3, 6, 9, and 13 days) were compared. Several modifications were observed between the differentiation days. A clear increase in the peak intensity at 960 cm^{-1} was detected at D13 (yellow spectrum) and particularly on the stratification layers (light blue spectrum). * mean spectrum of HaCaT at D13, taken from the stratified layers.

Table 1 : Assignment of spectral bands in a Raman spectrum of HaCaT cells during the differentiation process. ν elongation or stretching, δ deformation.

Wavenumber (cm^{-1})	2H vs D3	D3 vs D6	D6 vs D9	D9 vs D13	Assignment
598 (592-605)	↘	↗	-	↘	Phospholipids ²⁷
607 (605-614)	-	-	↗	-	Cholesterol ²⁸ /Glycerol ²⁹
669 (661-676)	↘	-	-	-	ν (C-S) of cytosine ³⁰
700 (693-705)	-	↗	-	-	Cholesterol, cholesterol ester ²³
725 (705-736)	-	-	-	↘	Ring breathing mode of DNA /RNA bases ^{27, 30}
760 (746-768)	↘	↗	-	-	Ring breathing tryptophan ^{30, 31}
831 (823- 844)	↘	-	-	↘	ν_{asym} (O-P-O) ³⁰ /Out-of-plane ring breathing, tyrosine ³¹
855 (846-871)	↘	-	-	-	Ring breathing mode of tyrosine & ν (C-C) of proline ring ³¹
960 (953-978)	-	↗	-	↗↗	ν (C-C) β -sheet keratin conformation ²³ . ν C-N ²²
1044 (1040-1053)	↘	-	-	-	Tyrosine ³²
1086 (1073-1100)	-	-	-	↗	ν C-C, lipid chains ^{30, 31}
1128 (1116-1140)	↘	-	-	-	ν C-C, lipid chains ^{28, 33}
1305 (1284-1320)	↘	-	-	↗	δCH_2 twisting (lipid) ^{30, 34}
1448 (1406-1494)	↘	-	↗	↘	δ C-H scissoring (proteins and lipid) ^{30, 34}
1558 (1538-1564)	-	-	-	↗	Tryptophan ³⁵

1658 (1630-1744)	-	-	↗↗	-	v (C=O) amide I, α helix of keratin ²³ . Little contribution for v(C=C) lipid ^{34, 36}
2723 (2697-2755)	-	-	-	↘	v (CH) aliphatic ³³
2851 (2833-2856)	-	-	-	↗↗	Lipid, v_{sym} (CH ₂) ³⁶⁻³⁸
2882 (2856-2895)	-	-	-	↗↗	Lipid, v_{asym} (CH ₂) ^{27, 37}
2931 (2902-2953)	-	2934	-	↘	v_{sym} (CH ₃) ³³
2962 (2953-3000)	-	-	-	↘↘	v_{asym} (CH ₃) ^{30, 33}
3059 (3047-3080)	-	↘	↘	↘	v(=CH) of lipid ^{36, 39}

2- Raman microspectroscopy: Spectral images of HaCaT cells

In parallel with spectral analyses, spectral imaging was performed in order to monitor possible sub-cellular changes along with the differentiation days and false-colored images were obtained (figure 3).

In the first column of figure 3, the evolution of DNA to lipid content can be monitored by following the (773-790 cm⁻¹) / (2820-2890 cm⁻¹) ratio.

At D0, after 2 hours of exposure to high calcium concentration medium, nucleus can be clearly identified in green (light blue). Based on the color bar, the pixels presented in green (light blue) are indicative to higher ratio values than dark blue ones. A decrease in the ratio values was observed starting at D6 where the nucleus remained clearly identified. At D9, the decrease of DNA is clearly observed, and the identification of nucleus is much difficult. Finally, at D13, nucleic acids disappeared (figure 3, first column).

The second column of figure 3 illustrates the lipid to protein ratio (2820-2890 cm⁻¹) / (2900-2970 cm⁻¹).

After 2H of high calcium exposure, lipid / protein ratio seems to record the highest intensity (red color) at the cell membrane area, indicating the localization of the lipid at a high level at the cell membrane zone.

After 3 days of differentiation, the red color begins to disperse throughout the cell except the nucleus zone.

In addition, at this differentiation day a very high intensity zone was detected which may reflect the localization of a very high amount of lipid content. Egawa *et al.*, also observed strong signal intensities of lipid surrounding the nucleus (2850 cm⁻¹) and related it to endoplasmic reticulum synthesis of lipid⁴⁰. At D6 and D9, we can clearly see a decrease in the intensity of this ratio in the cytoplasm. In contrast, higher values started to be observed in the nucleus at D6 and continued to increase at D9. At D13, the lipid/protein distribution is more homogenous all over the images with high values. This is in concordance with results

obtained from spectral analyses (table 1) and with Egawa *et al.* observation at the final stage of Keratinocytes differentiation. At this stage, Keratinocytes with digested nuclei form layered structures and present a flattened shape with strong signal intensity of lipids (2850 cm^{-1})⁴⁰.




Figure 3: False-colored Raman images of HaCaT during different period after high calcium exposure. The evolution of DNA/lipid ratio ($773\text{-}799/2820\text{-}2890\text{ cm}^{-1}$) and lipid /protein ratio ($2820\text{-}2890/2900\text{-}2970\text{ cm}^{-1}$) was followed during the differentiation. The green arrow indicates the localization of a high lipid content. *Stratified layers.

3- Lipid biomarkers evolution by HPLC/Corona & HPLC/HR-MS

Ceramides, glucosylceramides and sphingomyelins classes are considered as lipid biomarkers of differentiation stages. The analytical method used in this study was previously developed by our group²¹ and enables to separate epidermis lipid classes in one single run (data not shown). Using this method, we have followed epidermal lipid biomarkers evolution during keratinocytes differentiation. As shown in figure 4, the ratios of chromatographic peak area of glucosylceramides/ceramides and sphingomyelins/ceramides increased throughout the differentiation until reaching a maximal value at D9. A marked decrease of these ratios can be observed at D13.




Figure 4: HaCaT lipid were extracted and analyzed using HPLC/HR-MS and HPLC/Corona. The evolution of lipid biomarkers during keratinocytes differentiation was studied. Ratios of chromatographic peak area of glucosylceramides/ceramides and of sphingomyelins /ceramides calculated at different cells collection step.

Discussion and conclusion

In this study, we induced keratinocytes cell differentiation and followed the process over two weeks using HaCaT cells grown *in vitro* as a 2D cell culture model. Both Raman microspectroscopy and HPLC/HR-MS enabled the detection of several modifications in cell morphology. We also observed spectral signal and biomolecular changes during differentiation and underlined some lipid biomarkers (glucosylceramides and sphingomyelins). Morphological, spectral and biomolecular changes contributed altogether to determine of the differentiation stage. Results demonstrated the ability of HaCaT cells to differentiate *in vitro* (2D) till observing stratified layers, but we did not observe the formation of fully stratified layers, the latter can be obtained when using a 3D culture model.

Our data demonstrated clear biomolecular changes between D9 and D13, which may indicate the passage from the SG to the SC layer. In fact, literature indicates that after 3 weeks of incubation in a low-calcium concentration medium (0.03 mM), HaCaT cells proliferate, take a spindle shape but fail to stratify and to develop intracellular contact. This observation confirm the ability of HaCaT cells to revert back to a basal-like state – under a low-calcium concentration – as previously reported^{12, 18}. In contrast, to induce the differentiation process, HaCaT cells should be subjected to a high-concentration of calcium (2.8 mM). Two hours later, they start to take a more cuboidal shape with cell-cell contact¹⁵. These modifications occur to achieve the morphological characteristics of suprabasal layers^{15, 18}. In addition, the stratification layers and the flatter shape of several cells, which were observed at D9 and D13, may indicate the achievement of suprabasal layers but do not enable to determine morphologically the exact stage of keratinocyte differentiation. Wilson *et al.*, reported that the morphological modification should be confirmed biochemically by immunoblotting and immunofluorescence of protein and genes which are known as differentiation markers such as keratin-1 and involucrin⁸.

1
2
3
4
5 In this work, we are interested in demonstrating the morphological changes by following biomolecular
6 modifications involving lipid, protein, and DNA using Raman microspectroscopy. Raman is a very suitable
7 free label technique to follow cell differentiation at single cell level. The comparison of the mean Raman
8 spectra of each differentiation day demonstrated several modifications in the cells' spectra as shown in
9 figure 2 and table 1. A high increase of the 960 cm^{-1} peak was detected after 13 days and particularly at the
10 stratified layers. This peak has previously been assigned to the C-N vibration of saturated ceramides (such
11 as ceramide III)²². In fact, in the outermost epidermal layer, ceramides are the dominant lipid class by weight
12 (over 50%) of total lipids species besides free fatty acids (FFA) and cholesterol⁹. High ceramide content
13 is considered as a marker of the SC layer. Ceramide class consists of fatty acid chains linked to a sphingoid
14 base. Since in the human SC, the FFA are predominantly saturated⁴¹⁻⁴³, this may explain the high increase
15 in saturated ceramides at the late stage of differentiation. This observation was also confirmed as shown in
16 table (1), by the 3059 cm^{-1} peak at corresponding to the =C-H stretching, this peak decreased during
17 differentiation and may indicate a decrease in the unsaturation chain, related to unsaturated ceramides. In
18 addition, the high increase of the Raman peak at (960 cm^{-1}) is also related to an increase in the $\nu(\text{C-C})$ β -
19 sheet keratin conformation²³. The SC layers (corneocytes) are rich in keratin filament, and a strong increase
20 of the keratin component is a marker of SC layers²³.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

When the keratinocytes reach the SC layer, they lose their nucleus and acquire a flatter shape to become
corneocytes^{7, 44, 45}. Raman spectral images of HaCaT during differentiation permitted to follow the
DNA/lipid ratio ($773\text{-}790\text{ cm}^{-1}/2820\text{-}2890\text{ cm}^{-1}$) as shown in figure 3. Our results confirm the decrease of
this ratio during the differentiation until it reaches its lowest value at D13. This observation is also confirmed
by the characteristic band of nucleic acids at 725 cm^{-1} which decreased significantly at the last day (table 1).
The decrease in the nucleic acids levels may confirm the achievement of a later stage of differentiation (SC).
Lipids are mainly found in the cell membrane zone. During keratinocytes differentiation, lipid biosynthesis
evolves at early stages and forms at advanced stages the inter-corneocytes matrix of the *stratum corneum*.

1
2
3
4
5 In this study, the evolution of the lipid/protein (2820-2890) / (2900-2970) cm^{-1} ratio is an important marker
6
7 to determine the keratinocytes differentiation stage. As shown in figure 3, after 3 days of high-calcium
8
9 exposure, this ratio recorded a high increase throughout the cell (except the nucleus zone) indicating a high
10
11 lipid content compared to that of the proteins. In contrast, the decrease in the intensity at D6 and D9 could
12
13 be due to an increase in the protein content such as keratin at this stage of differentiation. As shown in table
14
15 (1), the Raman peak at 1658 cm^{-1} , which is related to the amide I of protein (α helix of keratin), significantly
16
17 increased at D9. The highest and most homogeneous lipid / protein ratio was observed at D13 (figure 3).
18
19 This increase reveals the formation of the inter-corneocytes lipid matrix (at the SC) which consists in
20
21 particular of ceramides, cholesterol and free fatty acids. Literature links this increase to the decrease in the
22
23 protein content subjected to the proteases of the SC – such as caspase 14 which contributes to the
24
25 metabolization of filaggrin (involved in the terminal differentiation of keratinocytes and the formation of
26
27 the cornified envelope in the *stratum corneum*) – leading to less (intact) protein in the outermost layers of
28
29 the SC^{25, 46}.
30

31
32 We also followed the evolution of lipid biomarkers during differentiation as shown in figure 4 by
33
34 HPLC/corona & HPLC/HR-MS. Sphingomyelins and glucosylceramides are recognized as markers of the
35
36 SG layer¹¹. An increase in the glucosylceramides/ceramides and sphingomyelins/ceramides ratio was
37
38 observed during differentiation. These ratios reached their maximum value at D9, and then started to
39
40 decrease. This observation indicates the passage of keratinocytes from the SG (high glucosylceramides and
41
42 sphingomyelins content at D9) to the SC layer (D 13).
43
44
45

46 To conclude, HaCaT cells subjected to a high-calcium concentration medium differentiated *in vitro* but did
47
48 not show fully stratified layers. We detected, according to our 2D cell culture model, several modifications
49
50 occurring at single cell level during differentiation. Our results demonstrate that Raman microspectroscopy
51
52 is an efficient methodology that allows a free staining observation of epidermal cells. It is a powerful
53
54
55
56
57
58
59
60

1
2
3
4
5 technique to follow the biomolecular changes including lipid, protein, and DNA evolution with a high ability
6
7 to determine their location throughout the cells. The complementary information obtained from both Raman
8
9 microspectroscopy and (HPLC/corona & HPLC/HR-MS) clearly demonstrated molecular modifications
10
11 between D9 and D13. Indeed, HaCaT cells presented a high increase in lipid and keratin contents, a decrease
12
13 in DNA and proteins, and a decrease in the ratio of glucosylceramides/ceramides and
14
15 sphingomyelins/ceramides between D9 and D13. All these observations characterize the differentiation of
16
17 HaCaT cells and their passage from SG to SC between these two days. In perspective, conducting further
18
19 data processing by combing Raman and chromatographic data might be interesting⁴⁷.
20

21 22 23 24 **Acknowledgements**

25
26 The authors would like to thank the SILAB - Jean PAUFIQUE Corporate Foundation of the financial support
27
28 for this work. We also thank Dr Franck Bonnier for his valuable help.
29

30 31 32 **Compliance with ethical standards Conflict of interest**

33
34 The authors declare that they have no conflict of interest.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. B. Breiden, H. Gallala, T. Doering and K. Sandhoff, *Eur J Cell Biol*, 2007, **86**, 657-673.
2. D. D. Bikle, Z. Xie and C. L. Tu, *Expert Rev Endocrinol Metab*, 2012, **7**, 461-472.
3. L. Eckhart, S. Lippens, E. Tschachler and W. Declercq, *Biochim Biophys Acta*, 2013, **1833**, 3471-3480.
4. A. Seo, N. Kitagawa, T. Matsuura, H. Sato and T. Inai, *Histochem Cell Biol*, 2016, **146**, 585-597.
5. M. Kypriotou, M. Huber and D. Hohl, *Exp Dermatol*, 2012, **21**, 643-649.
6. J. A. Bouwstra, P. L. Honeywell-Nguyen, G. S. Gooris and M. Ponec, *Prog Lipid Res*, 2003, **42**, 1-36.
7. M. Weil, M. C. Raff and V. M. Braga, *Curr Biol*, 1999, **9**, 361-364.
8. V. G. Wilson, *Methods Mol Biol*, 2013, **1195**, 33-41.
9. Y. Mizutani, S. Mitsutake, K. Tsuji, A. Kihara and Y. Igarashi, *Biochimie*, 2009, **91**, 784-790.
10. A. Kihara, *The Journal of Biochemistry*, 2012, **152**, 387-395.
11. M. Rabionet, K. Gorgas and R. Sandhoff, *Biochimica Et Biophysica Acta-Molecular and Cell Biology of Lipids*, 2014, **1841**, 422-434.
12. A. F. Deyrieux and V. G. Wilson, *Cytotechnology*, 2007, **54**, 77-83.
13. L. Micallef, 'UNIVERSITE DE LIMOGES 2009.
14. P. J. Houghton, P. J. Hylands, A. Y. Mensah, A. Hensel and A. M. Deters, *J Ethnopharmacol*, 2005, **100**, 100-107.
15. D. D. Bikle, *J Cell Biochem*, 2004, **92**, 436-444.
16. S. E. Lee and S. H. Lee, *Ann Dermatol*, 2018, **30**, 265-275.
17. L. Verzeaux, R. Vyumvuhore, D. Boudier, M. Le Guillou, S. Bordes, M. Essendoubi, M. Manfait and B. Closs, *Exp Dermatol*, 2017, DOI: 10.1111/exd.13388.
18. S. Pillai, D. D. Bikle, M. L. Mancianti, P. Cline and M. Hincenbergs, *J Cell Physiol*, 1990, **143**, 294-302.
19. E. G. Bligh and W. J. Dyer, *Can J Biochem Physiol*, 1959, **37**, 911-917.
20. A. Mieremet, M. Rietveld, S. Absalah, J. van Smeden, J. A. Bouwstra and A. El Ghalbzouri, *PLoS One*, 2017, **12**, e0174478.
21. A. Assi, J. Bakar, D. Libong, E. Sarkees, A. Solgadi, A. Baillet-Guffroy, R. Michael-Jubeli and A. Tfayli, *Anal Bioanal Chem*, 2019, DOI: 10.1007/s00216-019-02301-3.
22. A. Tfayli, E. Guillard, M. Manfait and A. Baillet-Guffroy, *Anal Bioanal Chem*, 2010, **397**, 1281-1296.
23. C. Choe, J. Schleusener, J. Lademann and M. E. Darvin, *Sci Rep*, 2017, **7**, 15900.
24. M. Gniadecka, O. Faurskov Nielsen, D. H. Christensen and H. C. Wulf, *J Invest Dermatol*, 1998, **110**, 393-398.
25. M. Janssens, J. van Smeden, G. J. Puppels, A. P. Lavrijsen, P. J. Caspers and J. A. Bouwstra, *Br J Dermatol*, 2014, **170**, 1248-1255.
26. L. Verzeaux, R. Vyumvuhore, D. Boudier, M. Le Guillou, S. Bordes, M. Essendoubi, M. Manfait and B. Closs, *Exp Dermatol*, 2017, **27**, 403-408.
27. S. Tfayli, C. Gobinet, G. Josse, J. F. Angiboust, M. Manfait and O. Piot, *Analyst*, 2012, **137**, 3673-3682.
28. G. Zhang, D. J. Moore, C. R. Flach and R. Mendelsohn, *Anal Bioanal Chem*, 2007, **387**, 1591-1599.
29. N. Garcia, O. Doucet, M. Bayer, D. Fouchard, L. Zastrow and J. P. Marty, *Int J Cosmet Sci*, 2002, **24**, 25-34.
30. Z. Movasaghi, S. Rehman and I. U. Rehman, *Applied Spectroscopy Reviews*, 2007, **42**, 493-541.

