



**HAL**  
open science

## Atypical cortical hierarchy in $A\beta$ -positive older adults and its reflection in spontaneous speech

Rui He, Jalal Al-Tamimi, Gonzalo Sánchez-Benavides, Guillermo Montaña-Valverde, Juan Domingo Gispert, Oriol Grau-Rivera, Marc Suárez-Calvet, Carolina Minguillon, Karine Fauria, Arcadi Navarro, et al.

### ► To cite this version:

Rui He, Jalal Al-Tamimi, Gonzalo Sánchez-Benavides, Guillermo Montaña-Valverde, Juan Domingo Gispert, et al.. Atypical cortical hierarchy in  $A\beta$ -positive older adults and its reflection in spontaneous speech. *Brain Research*, 2024, 1830, pp.148806. 10.1016/j.brainres.2024.148806 . hal-04529221


**HAL Id: hal-04529221**

**<https://hal.science/hal-04529221>**

Submitted on 2 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Atypical cortical hierarchy in A $\beta$ -positive older adults and its reflection in spontaneous speech

Rui He<sup>a,\*</sup>, Jalal Al-Tamimi<sup>b</sup>, Gonzalo Sánchez-Benavides<sup>c,d,e</sup>, Guillermo Montaña-Valverde<sup>a</sup>, Juan Domingo Gispert<sup>c,d,f,g</sup>, Oriol Grau-Rivera<sup>c,d,e,h</sup>, Marc Suárez-Calvet<sup>c,d,e,h</sup>, Carolina Minguillon<sup>c,d,e</sup>, Karine Fauria<sup>c,e</sup>, Arcadi Navarro<sup>c,i,j,k</sup>, Wolfram Hinzen<sup>a,i</sup>

<sup>a</sup> Department of Translation & Language Sciences, Universitat Pompeu Fabra, 08018 Barcelona, Spain

<sup>b</sup> Université Paris Cité, Laboratoire de Linguistique Formelle (LLF), CNRS, 75013 Paris, France

<sup>c</sup> BarcelonaBeta Brain Research Center (BBRC), Pasqual Maragall Foundation, 08005 Barcelona, Spain

<sup>d</sup> Neurosciences Department, IMIM (Hospital del Mar Medical Research Institute), 08003 Barcelona, Spain

<sup>e</sup> Centro de Investigación Biomédica en Red de Fragilidad y Envejecimiento Saludable (CIBERFES), Instituto de Salud Carlos III, 28029 Madrid, Spain

<sup>f</sup> Department of Medicine and Life Sciences, Universitat Pompeu Fabra, 08003 Barcelona, Spain

<sup>g</sup> Centro de Investigación Biomédica en Red de Bioingeniería, Biomateriales y Nanomedicina (CIBERBBN), Instituto de Salud Carlos III, Madrid, Spain

<sup>h</sup> Servei de Neurologia, Hospital del Mar, 08003 Barcelona, Spain

<sup>i</sup> Catalan Institution of Research and Advanced Studies (ICREA), 08010 Barcelona, Spain

<sup>j</sup> Department of Medicine and Life Sciences, Institute of Evolutionary Biology (UPF-CSIC), Universitat Pompeu Fabra, 08003 Barcelona, Spain

<sup>k</sup> CRG, Centre for Genomic Regulation, Barcelona Institute of Science and Technology (BIST), 08003 Barcelona, Spain

### ARTICLE INFO

#### Keywords:

Cortical gradient  
Functional connectivity  
A $\beta$  amyloid  
Cognitive decline  
Dementia

### ABSTRACT

Abnormal deposition of A $\beta$  amyloid is an early neuropathological marker of Alzheimer's disease (AD), arising long ahead of clinical symptoms. Non-invasive measures of associated early neurofunctional changes, together with easily accessible behavioral readouts of these changes, could be of great clinical benefit. We pursued this aim by investigating large-scale cortical gradients of functional connectivity with functional MRI, which capture the hierarchical integration of cortical functions, together with acoustic-prosodic features from spontaneous speech, in cognitively unimpaired older adults with and without A $\beta$  positivity (total N = 188). We predicted distortions of the cortical hierarchy associated with prosodic changes in the A $\beta$  + group. Results confirmed substantially altered cortical hierarchies and less variability in these in the A $\beta$  + group, together with an increase in quantitative prosodic measures, which correlated with gradient variability as well as digit span test scores. Overall, these findings confirm that long before the clinical stage and objective cognitive impairment, increased risk of cognitive decline as indexed by A $\beta$  accumulation is marked by neurofunctional changes in the cortical hierarchy, which are related to automatically extractable speech patterns and alterations in working memory functions.

### 1. Introduction

Neuropathological processes underlying Alzheimer's disease (AD), including extracellular A $\beta$  peptide deposition and intracellular neurofibrillary tangles (Mahaman et al., 2022), long predate the onset of cognitive decline at a clinical level (Parnetti et al., 2019). Subtle cognitive changes may accompany this process, which are neither subjectively apparent to individuals themselves, nor objectively detectable through standardized neuropsychological evaluation. While amyloid and tau deposition, as well as neurodegeneration, can be reliably

detectable *in vivo* by CSF or neuroimaging, it is not clear yet which non-invasive neurofunctional metrics could pick up these changes and how they might relate to behavioral measures.

A crucial candidate for such behavioral measures is the changes in spontaneous speech, which are objectively detectable non-invasively and ecologically with natural language processing techniques. There is evidence that changes in spontaneous speech or writing may predate the clinical onset of AD several years, if not decades (Iacono et al., 2009; Le et al., 2011), including at the stages of mild cognitive impairment (MCI) (Chapin et al., 2022; Roark et al., 2011; Sherman et al., 2021) and even

\* Corresponding author.

E-mail address: [rui.he@upf.edu](mailto:rui.he@upf.edu) (R. He).

<https://doi.org/10.1016/j.brainres.2024.148806>

Received 1 February 2024; Accepted 7 February 2024

Available online 15 February 2024

0006-8993/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

of healthy ageing with subjective cognitive decline (Verfaillie et al., 2019) or with A $\beta$  positivity (Mueller et al., 2021). Automatically extractable speech parameters at the acoustic-prosodic level, which require no transcription, have been shown to have high potential to accurately detect AD (Fraser et al., 2015), MCI (Fraser et al., 2019), and even earlier stages such as people at risk of AD with subjective cognitive decline (SCD) (He et al., 2023). Speech prosody concerns how speakers organize sounds into comprehensible sequences, using language-specific patterns of stress, rhythm, intonation, and tone of voice (Edelson and Diehl, 2013). Changes in speech prosody have shown sensitivity to cognitive decline in the AD continuum (Lofgren and Hinzen, 2022; López-de-Ipiña et al., 2013; Pistono et al., 2016), and specificity for discriminating between different dementia pathologies, such as primary progressive aphasia (Zimmerer et al., 2020), dementia with Lewy bodies (Yamada et al., 2022), and frontotemporal dementia (Pressman et al., 2015).

While evidence of the potential of a speech biomarker for dissecting early disease progression is accumulating, a crucial lacuna across pathologies is direct evidence of a relation between changes in speech and associated neurofunctional changes. Functional connectivity (FC) within and between brain regions or networks can capture such neurofunctional changes (Tosun et al., 2014) and opens a potential path for a neural validation of automatically extractable behavioral markers of disease progression. Using resting-state functional MRI (rs-fMRI), macroscale brain-functional networks can be shown to exhibit a hierarchical organization, ranging from primary, unimodal and segregated sensorimotor systems, specialized to deal with external stimuli in particular sensory modalities, to transmodal cortices, where information is increasingly *integrated* so as to enable abstract and internally-directed mental functions (memory, thought) (Mesulam, 1998; Taylor et al., 2015; Margulies et al., 2016; Huntenburg et al., 2018). These latter functions are specifically associated with the default mode network (DMN), which lies at the apex of this cortical hierarchy, integrating and orchestrating all other functional networks (Deco et al., 2021). Recent studies have specifically measured the cortical hierarchy using functional connectome gradients, defined as large-scale components of FC obtained with dimension reduction techniques (Huntenburg et al., 2018; Margulies et al., 2016). It turns out that the DMN lies at a maximal cortical 'distance' from sensorimotor cortex regions (at the bottom), while other networks such as the frontoparietal control network (FPN) are interspersed in between. A sufficient such distance between the two poles of the cortical hierarchy is considered to enable or facilitate abstract, higher-order cognitive functions, which depend less on immediate environmental input (Smallwood et al., 2021). This triggers the general expectation that disorders of thought would relate to distortions of the cortical hierarchy, which incipient evidence (specifically in the form of a compression) supports, for multiple mental disorders including AD (Hu et al., 2022), psychosis (Dong et al., 2021), and autism spectrum disorders (Hong et al., 2019).

The purpose of the present study was to identify changes in the cortical hierarchy, as measured with FC gradients, and to link them to automatically extractable speech features. This was motivated by specific reasons for expecting that language changes at the level of spontaneous connected speech should pick up neurofunctional changes in the large-scale hierarchical organization of cognition. In particular, language (and speech, which it incorporates) necessarily *integrates* multiple cognitive functions in its normal use, from memory (e.g. retrieval of semantic concepts under temporal constraints) to social cognition and to complex executive and motor control (Hagoort, 2019; Roger et al., 2022). A distortion in the cross-network organization of these large-scale cognitive systems would therefore be naturally expected to be reflected in speech distortions. In line with this idea, there already is evidence from neurotypical individuals that individual differences in the cortical hierarchy relate to language function (Shao et al., 2022; Wang et al., 2022). In particular, Wang et al. (2022) show that greater cortical distance to primary sensorimotor landmarks predicts higher activations

in a story comprehension task during fMRI. However, in any mental disorders where speech biomarkers have been widely discussed, such as the dementias or psychosis, these potential links between the cortical hierarchy and forms of language dysfunction have not been explored to our knowledge. This specifically applies in the context of AD, where Hu et al. (2022) found that the values of the principal gradient spanning between sensorimotor cortices and the posterior DMN decreased notably in patients with AD relative to both MCI and controls, reflecting a diminished segregation between this subnetwork of the DMN and other large-scale networks. Whether and how these changes are related to language changes remains unknown. The present study aims to fill this gap.


Specifically, our aims were to use naturalistic speech samples obtained from telephone interviews, together with cortical gradients in FC as a proxy of the cortical hierarchy, to (i) test for an effect of early amyloid pathology on both speech prosody and cortical gradient changes; (ii) relate these two metrics (speech and cortical gradient changes) to each other; and (iii) to background these relations against those with standard neuropsychological measures. We pursued these aims in a sample of cognitively unimpaired (CU) individuals, half of which were at increased risk of AD in virtue of A $\beta$ -positivity.

## 2. Materials and methods

### 2.1. Dataset

188 subjects were sub-selected from the Alfa + study based on data availability in all domains relevant to the present investigation (Fig. 1a). Alfa + is a research cohort of middle-aged CU individuals, most of them with a family history of AD, who undergo clinical interviews, lifestyle and risk factors questionnaires, cognitive testing, CSF biomarkers, and neuroimaging procedures every 3 years with the aim to identify the earliest pathophysiological changes in the preclinical AD continuum (Molinuevo et al., 2016). All subjects came from North-East region of Spain with Spanish and/or Catalan as their native language(s). They were all diagnosed as CU, while approximately half had a higher risk of dementia as indicated by A $\beta$  biomarker positivity. Subjects with CSF A $\beta$ 42/40 ratio lower than 0.071 were defined as A $\beta$  pathology positive (A +) in conformity with Milà-Alomà et al. (2020), and else negative (A-).

There were 93 A + subjects and 95 A- subjects. Age and gender were significantly different between groups ( $p < .05$ ), while years in education and language (Spanish vs. Catalan) were not. Table 1 shows the demographic characteristics of all subjects included. Basic sociodemographic data, neuropsychological battery assessment results, magnetic resonance images (MRI), and CSF biomarkers were registered as had been done in the Alfa + study (Molinuevo et al., 2016). In addition, during the COVID pandemic confinement, a team of the BarcelonaBeta Brain Research Center (BBRC) contacted Alfa + participants and asked whether they wanted to participate in a voice characteristics study, to which they were asked orally to give their consent (recorded). If they agreed, they were asked two questions in a recorded telephone interview, one past-directed ('How have you lived these last months of confinement? What activities have you carried out, and how have you felt?'), and one future-directed ('Let's imagine that a vaccine is not found in the next few years. How do you think social relationships would change in the world due to this new post-COVID-19 situation?'). The interviews were recorded over the regular phone line using (mono channel, sample rate at 8000 Hz, 32 bits per sample, bit rate at 16 Kb/s). During the telephone interview, 140 (74.47 %) subjects used Catalan, while 48 (25.53 %) used Spanish. Presence of subjective cognitive decline (SCD) was acknowledged when the subject gave a positive answer to the question: *Do you perceive memory or cognitive difficulties?* (Sánchez-Benavides et al., 2021).



**Fig. 1.** Experiment schematic for this study.

## 2.2. Neuropsychological evaluation

Neuropsychological tests included scores from the Mini-Mental State Examination (MMSE), Semantic Fluency (SF), Trail Making Test (TMT), Free and Cued Selective Reminding Test (FCSRT), Memory Binding Test (MBT), Wechsler Adult Intelligence Scale (WAIS)-IV and Wechsler Memory Scale (WMS)-IV tests, and RBANS Line Orientation assessment. Details of the scores used can be found in [Supplementary Table 1](#). Differences in neuropsychological test scores between groups were

compared with a linear regression model including age, gender, years of education, and group as exogenous variables and each neuropsychological test result as the endogenous variable. *P*-values from multiple comparisons were corrected using false discovery rate (FDR) and reported as *q* values after correction ([Benjamini and Hochberg, 1995](#)). The threshold of statistical significance was set at 0.05 after correction. As the dataset is comprised of two very similar groups, we also set a higher threshold of 0.1 after correction as a marked 'tendency'. The same correction approach and statistical thresholds are applied to other

**Table 1**  
Demographic characteristics of subjects included in the dataset.

Variables	A+	A-	test	p	effect size	
	Mean ± SD	Mean ± SD			Indicator	Value
Age <sup>a</sup>	62.16 ± 4.68	60.44 ± 4.78	Mann-Whitney	0.023	RBC <sup>c</sup>	-0.084
Sex <sup>b</sup>	64.52 %	49.47 %	χ <sup>2</sup> test	0.037	Cramer's V	0.152
Educ. years <sup>a</sup>	13.08 ± 3.74	13.59 ± 3.39	Mann-Whitney	0.314	RBC <sup>c</sup>	0.192
Language <sup>c</sup>	75.27 %	73.68 %	χ <sup>2</sup> test	0.803	Cramer's V	0.018
SCD <sup>d</sup>	34.41 %	24.21 %	χ <sup>2</sup> test	0.124	Cramer's V	0.112

<sup>a</sup> Age and education (Educ.) years are represented by mean ± standard deviation.

<sup>b</sup> Sex is represented by the percentage of female subjects.

<sup>c</sup> The language used in the telephone interview is represented by the percentage of Catalan-speaking subjects.

<sup>d</sup> SCD is represented by the percentage of subjects with presence of Subjective Cognitive Decline.

<sup>e</sup> RBC stands for rank-biserial correlation.

statistical comparisons used in this paper.

### 2.3. Neuroimaging data acquisition and pre-processing

All participants underwent an MRI protocol in a 3 T Philips Ingenia CX scanner which included a 3D T1-weighted sequence (TE/TR = 4.6 / 9.9 ms, flip angle = 8°; voxel size = 0.75 × 0.75 × 0.75 mm<sup>3</sup>) and a 9-minute task-free functional MRI sequence (TE/TR = 35 / 1600 ms, flip angle = 70°; voxel size = 3.0 × 3.0 × 3.1 mm<sup>3</sup>). Results included in this manuscript come from preprocessing performed using fMRIPrep 21.0.1 (Esteban et al., 2018b, 2018a) based on Nipype 1.6.1 (Gorgolewski et al., 2011, 2018). The functional images were spatially standardized with MNI152Nlin2009cAsym space and all confounds were regressed out based on the output of fMRIPrep. Details on the fMRIPrep pre-processing pipeline can be found in the [supplementary materials](#).

### 2.4. Whole-brain functional connectome gradient analysis

For each subject, after pre-processing, the Schaefer 400-parcels (Schaefer et al., 2018) were applied to extract functional time-series from the 400 regions of interest (ROIs). Then, we computed a 400 × 400 connectivity matrix using correlation matrix from the Ledoit-Wolf covariance estimator (Fig. 1b, i.e. the ConnectivityMeasure function from Nilearn package). All connectivity matrices were Fisher-z transformed. For each group, we averaged the connectivity matrices from the subjects within that group to compute a group-level connectivity template. In the group-level connectivity template, for each row, all the negative elements and 90 % smallest elements were zeroed out so that only positive, strong, and less noisy correlations were kept. Next, we obtained an affinity matrix by computing the cosine similarities between the FC of each two ROIs from the sparse FC matrix. These cosine similarity scores were normalized into scores bounded between 0 and 1. Ten gradients were extracted from the affinity matrix for each group using diffusion mapping, to obtain two group-level gradient templates, one for the A + group and another for the A- group. These steps were carried out with the BrainSpace toolbox (Vos de Wael et al., 2020), and the parameters were: number of components = 10, dimension reduction approach = diffusion embedding, kernel = normalized angle, sparsity = 0.9, and random state = 7. The two group-level gradient templates were plotted on the cortical surface. Applying the same toolbox with identical parameters, we built gradient maps at the subject level and aligned these gradient maps to the group-level gradient solution with Procrustes rotation, in order to increase the robustness of subject gradient maps.

While, in line with previous work, the number of gradients was set as 10, only the first three gradients, G1, G2, and G3, were analyzed, due to their more established relationship with cognitive functions (Mckeown et al., 2020), with G1 reflecting the extent of separation between unimodal and transmodal cortex, G2 differentiating the somatosensory and visual cortex, and G3 reflecting the separation between the DMN and executive networks (Shao et al., 2022). The effect of Aβ positivity on these three gradients was investigated with the surface-based linear models (SLM) from the BrainStat toolbox (Larivière et al., 2023). Models included fixed effects of age, gender, years in education, and group (Fig. 1e). P-values from SLM were corrected for multiple comparisons with random field theory and FDR (Benjamini and Hochberg, 1995). In addition to this surface-based comparison, all ROIs were grouped into Yeo's 7 networks: DMN, FPN, limbic network (LN), ventral attention network (VAN), dorsal attention network (DAN), somatomotor network (SMN), and visual network (VN) (Yeo et al., 2011). Gradient scores of ROIs in each network and the whole brain were averaged to generate network-level and brain-level gradient scores. Approximate permutation tests with 10,000 rounds were applied to compare the group differences at the network-level, and brain-level gradient scores. P-values were corrected with FDR.

In addition to the gradient values for the first three gradients, we also measured how similar the subjects' gradient templates were to their corresponding group template (Fig. 1c). Spearman's rank correlation was used to compute the correlation coefficient between the gradient solutions of every subject and the group-level templates of their corresponding groups. These correlation coefficients were transformed with Fisher's z-transformation and are referred to as 'gradient fit' hereafter. Gradient fit indicates the within-group similarity to the gradient templates (Mckeown et al., 2020) and has been previously shown to correlate with behavioral measures (Shao et al., 2022), which motivated the inclusion of such a measure in our analysis pipeline. Similar to neuropsychological comparisons, a linear regression model was used to compare whether the gradient fits significantly differed between the groups, using identical exogenous variables.

### 2.5. Speech prosodic profiling

As the telephone recordings contained voices from both the interviewees and subjects, they were first processed by the pyannote model for speaker diarization (Fig. 1d, 2 speakers, 300 msec for minimum silence length, 200 msec for minimum voice length, -1.00 for speaker clustering preference) (Bredin et al., 2020; Bredin and Laurent, 2021). Although not perfect, this tool satisfactorily partitioned voices from the two speakers into several segments (diarization error rate of 25.50 % on a Spanish news dataset and 32.37 % on an English telephone recording dataset). After randomly quality-checking 10 processed samples, the partitioned segments were deemed as sound for later analysis. For each subject, we concatenated all voice segments only from themselves into a new recording. Interviewee-only recordings were forwarded to Prosogram, which automatically measures prosodic features for individual syllables and prosodic properties of longer stretches of speech (Mertens, 2004). Prosogram extracted 42 prosodic features, where 33 were selected after removing duplicated variables. Definitions and exclusion criteria for the prosodic features can be found in the [Supplementary Table 2](#). Similar to gradient fit comparison, an identical generalized linear model was applied to test whether prosodic measures significantly differed between groups.

### 2.6. Relate connectome gradients to neuropsychological and prosodic measures

Third-order partial Pearson's correlation was applied to investigate the correlations between gradient fits with neuropsychological test scores, using age, gender, and years in education as covariates (Fig. 1e). For prosodic features, only those prosodic features that showed a

tendency of differing in the A + group before correction (uncorrected  $p$  less than 0.1) entered the partial Pearson's correlation to investigate the correlation between gradient fits with the prosodic features. The correlation coefficients were plotted into heatmaps while the  $p$ -values were corrected with FDR.

## 2.7. Data and code availability

The data that support the findings of this study are selected from the Alfa + study and not publicly available due to their containing information that could compromise the privacy of research participants. The pre-processing application fMRIPrep is publicly available. All analyses were carried out with scripts in Python. Codes for gradient analysis and SLM can be found in the tutorials of BrainSpace and BrainStat toolboxes. Pyannote model is freely available on BAS Web Services for speaker diarization (<https://clarin.phonetik.uni-muenchen.de/BASWebServices/interface/SpeakDiar>). Prosogram (v301b, <https://sites.google.com/site/prosogram/home>) is publicly available. The linear model for group comparison and the FDR correction were carried out with the statsmodels package (0.13.5). We applied the pingouin package (0.5.3) for third-order partial correlation analyses. Other statistical tests were carried out with scipy 1.7.3. Codes can be requested from the corresponding author.

## 3. Results

### 3.1. Qualitative comparison of group-level gradient solutions

Fig. 2a displays the first three gradient scores on the cortical surface (fsaverage5 template); the remaining gradients can be found in the supplementary materials (Supplementary Fig. 1 and Supplementary Fig. 2). These first three gradients explained 52 % of the variance as shown by the scree plots (see Supplementary Fig. 3 and Supplementary Fig. 4). Gradients explained similar variance in the A + and A- groups (two-tailed Student's  $t$ -test:  $t = 0.0$ ,  $p = 1.0$ ). Globally, as shown in the histograms with kernel density estimation (Fig. 2b), the gradient distribution was compressed in the A + group, with a thinner tail but higher peak. This global compression in the A + group was significant as indicated by Kolmogorov–Smirnov test ( $KS = 0.03$ ,  $p = .02$ ). The first and the third gradients together are plotted in scatter plots in Fig. 2c, with the color of each dot (=ROI) referring to the specific Yeo's network where it is situated. Changes in distribution in the A + group can be visually observed.

### 3.2. Significant changes take place in gradients and gradient fit

Vertex-wise comparisons from SLM suggested that there were significant increases in G1 values in the A + group, in the anterior frontal lobe, anterior temporal lobe, and posterior parietal lobe, which strongly overlapped with the DMN (Fig. 3a); and significant decreases in the visual cortex. Network-wise comparisons suggested significant decreases in the A + group in the whole brain, DMN, FPN, and LN, and significant decreases in VN and DAN. For G2 (Fig. 3b), there were significant increases in the somatosensory and motor regions and significant decreases in the frontal, middle and inferior temporal, and parietal lobes, and of the visual cortex. Network-wise comparisons suggested significant decreases in the A + group in the whole brain, DMN, FPN, LN, and VN, and significant increases in VAN and SMN. For G3, fewer significant differences were found, which were weaker and scattered sporadically across the frontal and the parietal lobes (Fig. 3c). In the A + group, network-wise analysis suggested significant decreases in FPN, and significant increases in LN and VN. Despite of these differences in gradient component scores, no significant differences were observed in G1 fit (Fig. 3d) and G2 fit (Fig. 3e). By contrast, G3 fit increased in the A + group (Fig. 3f). This difference was significant before correction ( $p = .028$ ) and remained a tendency after correction ( $q = 0.083$ ).

### 3.3. Differences in neuropsychological scores and prosody

Supplementary Table 3 shows results from the linear regression model for neuropsychological scores, prosodic measures, and gradient fit. For the neuropsychological test scores, the A + group had higher performance in semantic fluency ( $p = .046$ ) and in total recall of FCSRT ( $p = .028$ ), but these differences did not survive correction ( $q = 0.511$ ). As for prosody, before correction, the A + group showed longer total syllable nucleus duration ( $p = .034$ ), and tendencies for more syllable nuclei ( $p = .084$ ), more safe syllable nuclei ( $p = .084$ ), longer speech ( $p = .081$ ), and longer internucleus duration ( $p = .097$ ). None of these differences, which are all related to speech timing, survived FDR correction ( $q > 0.05$ ).

### 3.4. Relating gradient fit to neuropsychology and prosody


The results of a partial Pearson's correlation analysis are shown in a heatmap in Fig. 4 with coefficients and significant values available in Supplementary Table 4. For neuropsychology, no significant correlations were found with either G1 or G2 fits. For G3 fit, before correction, there was a significant correlation with WAIS-IV digit span subtest scores ( $r = 0.202$ ,  $p = .006$ ), which did not survive correction. For prosody, the five measures showing pre-correction significance in the group comparisons showed no significant correlation with G1 or G2 fits either. Four of them, however, significantly positively correlated with G3 fit, both before and after correction: number of nuclei ( $r = 0.161$ ,  $q = 0.035$ ), number of safe nuclei ( $r = 0.161$ ,  $q = 0.035$ ), speech time ( $r = 0.172$ ,  $q = 0.035$ ), and total internucleus duration ( $r = 0.162$ ,  $q = 0.035$ ). Only total nucleus duration did not significantly correlate with G3 fit ( $r = 0.116$ ,  $q = 0.117$ ).

## 4. Discussion


In this study, we applied a data-driven approach to investigate changes in speech prosody and functional connectome gradients in a group of CU older adults with A $\beta$  positivity, with a view to explore links behavioral changes in speech prosody to neurofunctional changes. Results showed that gradient changes (including gradient fit) could be observed in numerous brain regions in the A $\beta$  + group. These changes in the large-scale organization of cognitive functions across a cortical hierarchy are not reflected by standard neuropsychological test batteries. The individual gradient fit values, on the other hand, are reflected in the prosodic profile of connected speech samples, which we obtained from a short conversational section of a telephone interview. These profile changes are specific, in that they uniquely concern prosodic features characterizing temporal aspects of speech, such as the number of syllabic nuclei produced, the speech time, and total internucleus duration. By contrast, pitch-related features, which do not involve specific aspects of the hierarchical organization of language such as syllables, words, and sentences, were not affected. Our findings thus not only indicate that a large-scale functional reorganization of the brain accompanies A $\beta$  positivity, but that spontaneous speech indexes these changes and forms an objective read-out of them.

Our findings of functional changes in a population at higher risk of AD are partially in line with previous studies of MCI and AD (He et al., 2022; Hu et al., 2022). Consistent with these studies of populations more progressed on the AD continuum, we qualitatively observed a compressed gradient distribution with a higher peak value already in an A + CU group (Fig. 2b). As speech production is an executive demanding task, and the third gradient did indeed relate to working memory as an aspect of executive functioning, we plotted the principle (first) and third gradients together in a scatter plot (Fig. 2c). Qualitative visual impression of the distribution of the values of these two gradients suggests a 'boomerang' shape in the A- group (left subplot of Fig. 2c). In the positive group (right subplot of Fig. 2c), this shape appears distorted, with the DMN and FPN collapsing onto each other, while VN extricates itself


(a) Cortical surface with G1, G3, and G3 (column 1, 2, 3) of the A- (row 1 & 2) and the A+ (row 3 & 4) groups




(b) Correlations between group gradients and with HCP gradients




(c) Global histogram of gradient



(d) G1-G3 scatter plot of A- and A+ groups



**Fig. 2.** Qualitatively observing the group-level gradient solutions. (a) Plotting the first three gradients on the cortical surface. (b) Distribution of all 10 gradients. The grey histogram refers to the A- group while the red one refers to the A+. (c) Scatter plots of the first (y-axis) and the third (x-axis) gradient. The color refers to Yeo's 7 networks: blue: visual; orange: somatomotor; green: dorsal attention; red: salience/ventral attention; purple: limbic; brown: frontoparietal control; pink: default mode. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.) (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



**Fig. 3.** Investigating the effect of Aβ group. (a-c) Effect of Aβ group on gradient 1, 2, and 3, with t and p values from SLM plotted on the brain surface. For each gradient, t values were plotted on the first row (in blue and yellow, blue for positive values while yellow for negative values). Significant p values (less than 0.05 after FDR correction) were plotted on the second row (in green). The denser the color is, the smaller the p values are. (d-f) Comparing gradient 1, 2, and 3 between groups within the 7 Yeo's networks and the whole brain. (g-i) Effect of Aβ group on G1 fit, G2 fit and G3 shown with strip plots (x axis: group, grey for A- and red for A+; y axis: gradient fit). The  $R^2$ , t value, uncorrected p value, and FDR-corrected p value were annotated with the mean line. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.) (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

out and disperses into a separate small triangle. Upon quantitative scrutiny, however, it turns out that the directions of changes in specific functional networks differ from changes in MCI and AD observed in the previous work. In particular, the principal gradient increased in the DMN and FPN (transmodal regions), while it decreased in VN and SMN (unimodal regions). On the other hand, He et al. (2022) observed decreases in the transmodal regions and increases in the unimodal regions in their MCI and AD groups. This suggests that neurofunctional changes at preclinical stages are unlikely to be continuous with changes seen at clinical changes. Early neuropathology may imply changes in distinct directions than those seen after a period where the brain cannot maintain neurotypical levels of cognitive functioning as assessed by clinical cognitive tests.

Although changes in prosody features, including both quantitative reductions of production and changes in pitch-related features, have previously been found at the clinical end of AD (Qiao et al., 2020; Vincze et al., 2021), this remains controversial for earlier stages. For temporal

aspects of speech production, in particular, the above two studies found *increases* in speech duration in the MCI group (Qiao et al., 2020; Vincze et al., 2021), different from AD. However, decreases in speech duration in MCI have also been observed (Ambrosini et al., 2019; Imre et al., 2022). In the preclinical stage marked by subjective cognitive decline (SCD), a recent study used identical prosodic features as in the present study for a distinct Spanish/Catalan dataset and found speech duration-related features to significantly increase in the preclinical SCD group, but to decrease in MCI and AD groups. Increase of total speech duration, though insignificant, was also found in another study of an Aβ-positive group (Mueller et al., 2021). Together, these previous studies, as well as the tendencies of increases in speech and nuclei durations observed in the present data, suggest that adults with signals for elevated risk of AD may show compensatory effects resulting in longer speech times. Crucially, subjects in our dataset were unaware of their biomarker status, and the SCD proportions in each group were not significantly different. The compensation in question is therefore not the reflection of





**Fig. 4.** Heatmap for correlation between gradient fit and behavior measures. The heatmap displays coefficients of the partial Pearson correlation between G1, G2, and G3 fits and the neuropsychological test scores (the first three rows) and the prosodic measures (the last row). White boxes indicate significant correlation after correction, while white underlines indicate significant correlation only before correction.

a conscious effort, but may index a compensatory brain process triggered by an onset of neural degradation. A potential pattern of slight increases in quantitative speech production features in preclinical stages, switching to a mix of increases and decreases in MCI, followed by a consistent decrease, awaits confirmation in future studies. Compensatory effects in the preclinical high-risk group could also explain why changes in the gradient values ran in different directions than previous studies of functional gradients in MCI and AD.

As changes in gradient values are difficult to interpret, the gradient fits were introduced in the present study as a more interpretable variable illustrating a clustering tendency within both groups in relation to their respective templates. In our data, differences in gradient fit were specifically significant for G3, the topography of which previous studies have related to the hierarchical separation between the executive control and DMN networks. The A + individuals showed a better fit suggesting a more cohesive group. This may indicate that they are less variable by virtue of sharing a cognitive characteristic, which predisposes them to cognitive decline. Moreover, the significant correlations between G3 fit and temporal aspects of speech, as well as that between G3 fit and a working memory measure (before correction), together support the possibility that changes in speech duration may specifically relate to executive functioning, which depends on working memory.

An important desideratum for future work is to extend the present study scheme to aspects of linguistic function depending on text-based analysis. While acoustic-prosodic features have a specific virtue for not depending on transcriptions, and have shown high promise in recent machine-learning studies especially at preclinical stages (He et al., 2023), a neurocognitive investigation needs to ask which specific mechanism in the multidimensional organization of language are affected, at which pathological stage. Semantic processing is a case in point, especially because the DMN strongly overlaps with the semantic network and plays an important role in functional gradients (Andrews-Hanna et al., 2014; Binder et al., 2009; Gordon et al., 2020). Additionally, transferring the gradient analysis to subcortical regions like the hippocampus is a promising direction. Incorporating the neuroimaging findings and language analysis may present more extensive and comprehensive understanding on the underlying mechanism of early cognitive decline.

#### CRedit authorship contribution statement

**Rui He:** Conceptualization, Formal analysis, Methodology, Software, Validation, Visualization, Writing – original draft, Writing – review & editing. **Jalal Al-Tamimi:** Formal analysis, Methodology, Validation, Writing – review & editing. **Gonzalo Sánchez-Benavides:** Data curation, Validation, Writing – review & editing. **Guillermo Montaña-Valverde:** Software, Validation, Writing – review & editing. **Juan Domingo Gispert:** Data curation, Writing – review & editing. **Oriol Grau-Rivera:** Data curation, Writing – review & editing. **Marc Suárez-Calvet:** Data curation, Writing – review & editing. **Carolina Minguiñon:** Data curation, Writing – review & editing. **Karine Fauria:** Data curation, Writing – review & editing. **Arcadi Navarro:** Data curation, Writing – review & editing. **Wolfram Hinzen:** Conceptualization, Funding acquisition, Methodology, Supervision, Validation, Writing – original draft, Writing – review & editing.

#### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

Data will be made available on request.

#### Acknowledgments

We acknowledge all individuals participating in this study. We also want to thank our sponsors for making this project possible. We would like to give special thanks to Mary Lofgren for providing valuable thoughts in interpreting the results.

#### Funding

This research was supported by the China Scholarship Council (CSC202108390062 to RH), the Ministerio de Ciencia, Innovación y Universidades (MCIU) and the Agencia Estatal de Investigación (AEI) (grant PID2019-105241 GB-I00/AEI/10.13039/501100011033 to WH), and partially by French Investissements d’Avenir - Labex EFL program (ANR-10-LABX-0083 to the IdEx Université Paris Cité - ANR-18-IDEX-0001, to JAT).

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brainres.2024.148806>.

#### References

- Ambrosini, E., Cid, M., de Isla, C.G., Salamanca, P., Borghese, N.A., Ferrante, S., Caielli, M., Milis, M., Loizou, C., Azzolino, D., Damanti, S., Bertagnoli, L., Cesari, M., Moccia, S., 2019. Automatic speech analysis to early detect functional cognitive decline in elderly population. *Annu. Int. Conf. IEEE Eng. Med. Biol. Soc. IEEE Eng. Med. Biol. Soc. Annu. Int. Conf.* 2019, 212–216. <https://doi.org/10.1109/EMBC.2019.8856768>.
- Andrews-Hanna, J.R., Smallwood, J., Spreng, R.N., 2014. The default network and self-generated thought: component processes, dynamic control, and clinical relevance. *Ann. N. Y. Acad. Sci.* 1316, 29–52. <https://doi.org/10.1111/nyas.12360>.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* 57, 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>.
- Binder, J.R., Desai, R.H., Graves, W.W., Conant, L.L., 2009. Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cereb. Cortex N. Y. N 1991* (19), 2767–2796. <https://doi.org/10.1093/cercor/bhp055>.
- Bredin, H., Laurent, A., 2021. End-To-End Speaker Segmentation for Overlap-Aware Resegmentation, in: *Interspeech 2021*. Presented at the Interspeech 2021, ISCA, pp. 3111–3115. <https://doi.org/10.21437/Interspeech.2021-560>.
- Bredin, H., Yin, R., Coria, J.M., Gelly, G., Korshunov, P., Lavechin, M., Fustes, D., Titeux, H., Bouaziz, W., Gill, M.-P., 2020. Pyannote.Audio: Neural Building Blocks for Speaker Diarization, in: *ICASSP 2020 - 2020 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*. Presented at the ICASSP 2020 - 2020 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP), pp. 7124–7128. <https://doi.org/10.1109/ICASSP40776.2020.9052974>.
- Chapin, K., Clarke, N., Garrard, P., Hinzen, W., 2022. A finer-grained linguistic profile of Alzheimer’s disease and Mild Cognitive Impairment. *J. Neurolinguistics* 63, 101069. <https://doi.org/10.1016/j.jneuroling.2022.101069>.
- Deco, G., Vidaurre, D., Kringelbach, M.L., 2021. Revisiting the global workspace orchestrating the hierarchical organization of the human brain. *Nat. Hum. Behav.* 5, 497–511. <https://doi.org/10.1038/s41562-020-01003-6>.
- Dong, D., Yao, D., Wang, Y., Hong, S.-J., Genon, S., Xin, F., Jung, K., He, H., Chang, X., Duan, M., Bernhardt, B.C., Margulies, D.S., Sepulcre, J., Eickhoff, S.B., Luo, C., 2021. Compressed sensorimotor-to-transmodal hierarchical organization in schizophrenia. *Psychol. Med.* 1–14. <https://doi.org/10.1017/S0033291721002129>.
- Edelson, L., Diehl, J., 2013. *Prosody*. In: Volkmar, F.R. (Ed.), *Encyclopedia of Autism Spectrum Disorders*. Springer, New York, NY, pp. 2413–2417. [https://doi.org/10.1007/978-1-4419-1698-3\\_366](https://doi.org/10.1007/978-1-4419-1698-3_366).
- Esteban, O., Blair, R., Markiewicz, C.J., Berleant, S.L., Moodie, C., Ma, F., Isik, A.I., Erramuzpe, A., Kent, M., James D. andGoncalves, DuPre, E., Sitek, K.R., Gomez, D.E. P., Lurie, D.J., Ye, Z., Poldrack, R.A., Gorgolewski, K.J., 2018a. fMRIPrep. Software. <https://doi.org/10.5281/zenodo.852659>.
- Esteban, O., Markiewicz, C., Blair, R.W., Moodie, C., Isik, A.I., Erramuzpe Aliaga, A., Kent, J., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S., Wright, J., Durnez, J., Poldrack, R., Gorgolewski, K.J., 2018b. fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat. Methods*. <https://doi.org/10.1038/s41592-018-0235-4>.
- Fraser, K.C., Meltzer, J.A., Rudzicz, F., 2015. Linguistic Features Identify Alzheimer’s Disease in Narrative Speech. *J. Alzheimers Dis.* 49, 407–422. <https://doi.org/10.3233/JAD-150520>.
- Fraser, K.C., Lundholm Fors, K., Eckerström, M., Öhman, F., Kokkinakis, D., 2019. Predicting MCI Status From Multimodal Language Data Using Cascaded Classifiers. *Front. Aging Neurosci.* 11, 205. <https://doi.org/10.3389/fnagi.2019.00205>.
- Gordon, E.M., Laumann, T.O., Marek, S., Raut, R.V., Gratton, C., Newbold, D.J., Greene, D.J., Coalson, R.S., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E.,

- Dosenbach, N.U.F., Nelson, S.M., 2020. Default-mode network streams for coupling to language and control systems. *Proc. Natl. Acad. Sci. U. S. A.* 117, 17308–17319. <https://doi.org/10.1073/pnas.2005238117>.
- Gorgolewski, K., Burns, C.D., Madison, C., Clark, D., Halchenko, Y.O., Waskom, M.L., Ghosh, S., 2011. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in Python. *Front. Neuroinformatics* 5, 13. <https://doi.org/10.3389/fninf.2011.00013>.
- Gorgolewski, K.J., Esteban, O., Markiewicz, C.J., Ziegler, E., Ellis, D.G., Notter, M.P., Jarecka, D., Johnson, H., Burns, C., Manhães-Savio, A., Hamalainen, C., Yvernault, B., Salo, T., Jordan, K., Goncalves, M., Waskom, M., Clark, D., Wong, J., Loney, F., Modat, M., Dewey, B.E., Madison, C., di Oleggio, V., Castello, M., Clark, M. G., Dayan, M., Clark, D., Keshavan, A., Pinsard, B., Gramfort, A., Berleant, S., Nielson, D.M., Bougacha, S., Varoquaux, G., Cipollini, B., Markello, R., Rokem, A., Moloney, B., Halchenko, Y.O., Wassermann, D., Hanke, M., Horea, C., Kuczmarzyk, J., de Hollander, G., DuPre, E., Gillman, A., Mordom, D., Buchanan, C., Tungaraza, R., Pauli, W.M., Iqbal, S., Sikka, S., Mancini, M., Schwartz, Y., Malone, I. B., Dubois, M., Frohlich, C., Welch, D., Forbes, J., Kent, J., Watanabe, A., Cumba, C., Huntenburg, J.M., Kastman, E., Nichols, B.N., Eshaghi, A., Ginsburg, D., Schaefer, A., Acland, B., Giavasis, S., Kleesiek, J., Erickson, D., Küttner, R., Haselgrove, C., Correa, C., Ghayoor, A., Liem, F., Millman, J., Haehn, D., Lai, J., Zhou, D., Blair, R., Glatard, T., Renfro, M., Liu, S., Kahn, A.E., Pérez-García, F., Triplett, W., Lampe, L., Stadler, J., Kong, X.-Z., Hallquist, M., Chetverikov, A., Salvatore, S., Sharp, P., Molina-Romero, M., Weinstein, A., Broderick, W., Saase, V., Andberg, S.K., Harms, R., Schlamp, K., Arias, J., Papadopoulos Orfanos, D., Tarbert, C., Tambini, A., De La Vega, A., Nickson, T., Brett, M., Falkiewicz, M., Podranski, K., Linkersdorfer, J., Flandin, G., Ort, E., Shachnev, D., McNamee, D., Davison, A., Varada, J., Schwabacher, I., Pellman, J., Perez-Guevara, M., Khanuja, R., Pannetier, N., McDermtroee, C., Ghosh, S., 2018. Nipype. Software. <https://doi.org/10.5281/zenodo.596855>.
- Hagoort, P., 2019. The neurobiology of language beyond single-word processing. *Science* 366, 55–58. <https://doi.org/10.1126/science.aax0289>.
- He, R., Chapin, K., Al-Tamimi, J., Bel, N., Marquié, M., Rosende-Roca, M., Pytel, V., Tartari, J.P., Alegret, M., Sanabria, Á., Ruiz, A., Boada, M., Valero, S., Hinzen, W., 2023. Automated classification of cognitive decline and probable Alzheimer's dementia across multiple speech and language domains. Accepted and in press, *Am. J. Speech Lang. Pathol.*
- He, Y., Li, Q., Fu, Z., Zeng, D., Han, Y., Li, S., 2022. Functional Gradients Reveal Altered Functional Segregation in Patients with Amnesic Mild Cognitive Impairment and Alzheimer's Disease. <https://doi.org/10.1101/2022.11.16.516698>.
- Hong, S.-J., Vos de Wael, R., Bethlehem, R.A.I., Larivière, S., Paquola, C., Valk, S.L., Milham, M.P., Di Martino, A., Margulies, D.S., Smallwood, J., Bernhardt, B.C., 2019. Atypical functional connectome hierarchy in autism. *Nat. Commun.* 10, 1022. <https://doi.org/10.1038/s41467-019-08944-1>.
- Hu, Q., Li, Y., Wu, Y., Lin, X., Zhao, X., 2022. Brain network hierarchy reorganization in Alzheimer's disease: A resting-state functional magnetic resonance imaging study. *Hum. Brain Mapp.* 43, 3498–3507. <https://doi.org/10.1002/hbm.25863>.
- Huntenburg, J.M., Bazin, P.-L., Margulies, D.S., 2018. Large-Scale Gradients in Human Cortical Organization. *Trends Cogn. Sci.* 22, 21–31. <https://doi.org/10.1016/j.tics.2017.11.002>.
- Iacono, D., Markesbery, W.R., Gross, M., Pletnikova, O., Rudow, G., Zandi, P., Troncoso, J.C., 2009. The Nun study: clinically silent AD, neuronal hypertrophy, and linguistic skills in early life. *Neurology* 73, 665–673. <https://doi.org/10.1212/WNL.0b013e3181b01077>.
- Imre, N., Balogh, R., Gosztolya, G., Tóth, L., Hoffmann, I., Várkonyi, T., Lengyel, C., Pákási, M., Kálmán, J., 2022. Temporal Speech Parameters Indicate Early Cognitive Decline in Elderly Patients With Type 2 Diabetes Mellitus. *Alzheimer Dis. Assoc. Disord.* 36, 148–155. <https://doi.org/10.1097/WAD.0000000000000492>.
- Larivière, S., Bayrak, Ş., Vos de Wael, R., Benkarim, O., Herholz, P., Rodriguez-Cruces, R., Paquola, C., Hong, S.-J., Mistic, B., Evans, A.C., Valk, S.L., Bernhardt, B.C., 2023. BrainStat: A toolbox for brain-wide statistics and multimodal feature associations. *NeuroImage* 266, 119807. <https://doi.org/10.1016/j.neuroimage.2022.119807>.
- Le, X., Lancashire, I., Hirst, G., Jokel, R., 2011. Longitudinal detection of dementia through lexical and syntactic changes in writing: a case study of three British novelists. *Lit. Linguist. Comput.* 26, 435–461. <https://doi.org/10.1093/lc/fqr013>.
- Lofgren, M., Hinzen, W., 2022. Breaking the flow of thought: Increase of empty pauses in the connected speech of people with mild and moderate Alzheimer's disease. *J. Commun. Disord.* 97, 106214. <https://doi.org/10.1016/j.jcomdis.2022.106214>.
- López-de-Ipiña, K., Alonso, J.-B., Travieso, C., Solé-Casals, J., Egrara, H., Faundez-Zanuy, M., Ezeiza, A., Barroso, N., Ecay-Torres, M., Martínez-Lage, P., Lizardui, U., 2013. On the Selection of Non-Invasive Methods Based on Speech Analysis Oriented to Automatic Alzheimer Disease Diagnosis. *Sensors* 13, 6730–6745. <https://doi.org/10.3390/s130506730>.
- Maham, Y.A.R., Embaye, K.S., Huang, F., Li, L., Zhu, F., Wang, J.-Z., Liu, R., Feng, J., Wang, X., 2022. Biomarkers used in Alzheimer's disease diagnosis, treatment, and prevention. *Ageing Res. Rev.* 74, 101544. <https://doi.org/10.1016/j.arr.2021.101544>.
- Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langa, S., Bezgin, G., Eickhoff, S.B., Castellanos, F.X., Petrides, M., Jefferies, E., Smallwood, J., 2016. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc. Natl. Acad. Sci.* 113, 12574–12579. <https://doi.org/10.1073/pnas.1608282113>.
- Mckeown, B., Strawson, W.H., Wang, H.-T., Karapanagiotidis, T., Vos de Wael, R., Benkarim, O., Turnbull, A., Margulies, D., Jefferies, E., McCall, C., Bernhardt, B., Smallwood, J., 2020. The relationship between individual variation in macroscale functional gradients and distinct aspects of ongoing thought. *NeuroImage* 220, 117072. <https://doi.org/10.1016/j.neuroimage.2020.117072>.
- Mertens, P., 2004. The prosgogram: semi-automatic transcription of prosody based on a tonal percepting model, in: *Proceedings of Speech Prosody 2004*. Nara (Japan).
- Mesulam, M.M., 1998. From sensation to cognition. *Brain J. Neurol.* 121 (Pt 6), 1013–1052. <https://doi.org/10.1093/brain/121.6.1013>.
- Milà-Alomà, M., Salvadó, G., Gispert, J.D., Vilor-Tejedor, N., Grau-Rivera, O., Sala-Vila, A., Sánchez-Benavides, G., Arenaza-Urquijo, E.M., Crous-Bou, M., González-de-Echávarri, J.M., Minguillon, C., Fauria, K., Simon, M., Kollmorgen, G., Zetterberg, H., Blennow, K., Suárez-Calvet, M., Molinuevo, J.L., 2020. Amyloid beta, tau, synaptic, neurodegeneration, and glial biomarkers in the preclinical stage of the Alzheimer's continuum. *Alzheimers Dement.* 16, 1358–1371. <https://doi.org/10.1002/alz.12131>.
- Molinuevo, J.L., Gramunt, N., Gispert, J.D., Fauria, K., Esteller, M., Minguillon, C., Sánchez-Benavides, G., Huesa, G., Morán, S., Dal-Ré, R., Camí, J., 2016. The ALFA project: A research platform to identify early pathophysiological features of Alzheimer's disease. *Alzheimers Dement. Transl. Res. Clin. Interv.* 2, 82–92. <https://doi.org/10.1016/j.trci.2016.02.003>.
- Mueller, K.D., Van Hulle, C.A., Kosciak, R.L., Jonaitis, E., Peters, C.C., Betthausen, T.J., Christian, B., Chin, N., Hermann, B.P., Johnson, S., 2021. Amyloid beta associations with connected speech in cognitively unimpaired adults. *Alzheimers Dement. Amst. Neth.* 13, e12203.
- Parnetti, L., Chipi, E., Salvadori, N., D'Andrea, K., Eusebi, P., 2019. Prevalence and risk of progression of preclinical Alzheimer's disease stages: a systematic review and meta-analysis. *Alzheimers Res. Ther.* 11, 7. <https://doi.org/10.1186/s13195-018-0459-7>.
- Pistono, A., Jucla, M., Barbeau, E.J., Saint-Aubert, L., Lemesle, B., Calvet, B., Köpke, B., Puel, M., Pariante, J., 2016. Pauses During Autobiographical Discourse Reflect Episodic Memory Processes in Early Alzheimer's Disease. *J. Alzheimers Dis.* 50, 687. <https://doi.org/10.3233/JAD-150408>.
- Pressman, P., Levenson, R., Miller, B., Tempini, M.G., 2015. Prosodic Expression in Frontotemporal Dementia (P1.229). *Neurology* 84.
- Qiao, Y., Xie, X.-Y., Lin, G.-Z., Zou, Y., Chen, S.-D., Ren, R.-J., Wang, G., 2020. Computer-Assisted Speech Analysis in Mild Cognitive Impairment and Alzheimer's Disease: A Pilot Study from Shanghai, China. *J. Alzheimers Dis.* 75, 211–221. <https://doi.org/10.3233/JAD-191056>.
- Roark, B., Mitchell, M., Hosom, J.-P., Hollingshead, K., Kaye, J., 2011. Spoken Language Derived Measures for Detecting Mild Cognitive Impairment. *IEEE Trans. Audio Speech Lang. Process.* 19, 2081–2090. <https://doi.org/10.1109/TASL.2011.2112351>.
- Roger, E., Banjac, S., Thiebaut de Schotten, M., Baciu, M., 2022. Missing links: The functional unification of language and memory (LUM). *Neurosci. Biobehav. Rev.* 133, 104489. <https://doi.org/10.1016/j.neubiorev.2021.12.012>.
- Sánchez-Benavides, G., Suárez-Calvet, M., Milà-Alomà, M., Arenaza-Urquijo, E.M., Grau-Rivera, O., Operto, G., Gispert, J.D., Vilor-Tejedor, N., Sala-Vila, A., Crous-Bou, M., González-de-Echávarri, J.M., Minguillon, C., Fauria, K., Simon, M., Kollmorgen, G., Zetterberg, H., Blennow, K., Molinuevo, J.L., 2021. Amyloid-β positive individuals with subjective cognitive decline present increased CSF neuroinflammation light levels that relate to lower hippocampal volume. *Neurobiol. Aging* 104, 24–31. <https://doi.org/10.1016/j.neurobiolaging.2021.02.026>.
- Schaefer, A., Kong, R., Gordon, E.M., Laumann, T.O., Zuo, X.-N., Holmes, A.J., Eickhoff, S.B., Yeo, B.T.T., 2018. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb. Cortex N. Y. N* 1991 (28), 3095–3114. <https://doi.org/10.1093/cercor/bhx179>.
- Shao, X., Mckeown, B., Karapanagiotidis, T., Vos de Wael, R., Margulies, D.S., Bernhardt, B., Smallwood, J., Krieger-Redwood, K., Jefferies, E., 2022. Individual differences in gradients of intrinsic connectivity within the semantic network relate to distinct aspects of semantic cognition. *Cortex* 150, 48–60. <https://doi.org/10.1016/j.cortex.2022.01.019>.
- Sherman, J.C., Henderson, C.R., Flynn, S., Gair, J.W., Lust, B., 2021. Language Decline Characterizes Amnesic Mild Cognitive Impairment Independent of Cognitive Decline. *J. Speech Lang. Hear. Res. JSLHR* 64, 4287–4307. <https://doi.org/10.1044/2021.JSLHR-20-00503>.
- Smallwood, J., Bernhardt, B.C., Leech, R., Bzdok, D., Jefferies, E., Margulies, D.S., 2021. The default mode network in cognition: a topographical perspective. *Nat. Rev. Neurosci.* 22, 503–513. <https://doi.org/10.1038/s41583-021-00474-4>.
- Taylor, P., Hobbs, J.N., Burroni, J., Siegelmann, H.T., 2015. The global landscape of cognition: hierarchical aggregation as an organizational principle of human cortical networks and functions. *Sci. Rep.* 5, 18112. <https://doi.org/10.1038/srep18112>.
- Tosun, D., Joshi, S., Weiner, M.W., the Alzheimer's Disease Neuroimaging Initiative, 2014. Multimodal MRI-based Imputation of the Aβ+ in Early Mild Cognitive Impairment. *Ann. Clin. Transl. Neurol.* 1, 160–170. <https://doi.org/10.1002/acn3.40>.
- Verfaillie, S.C.J., Witteman, J., Slot, R.E.R., Pruis, I.J., Vermaat, L.E.W., Prins, N.D., Schiller, N.O., van de Wiel, M., Scheltens, P., van Berckel, B.N.M., van der Flier, W. M., Sikkes, S.A.M., 2019. High amyloid burden is associated with fewer specific words during spontaneous speech in individuals with subjective cognitive decline. *Neuropsychologia* 131, 184–192. <https://doi.org/10.1016/j.neuropsychologia.2019.05.006>.
- Vince, V., Szałóczyki, G., Tóth, L., Gosztolya, G., Pákási, M., Hoffmann, I., Kálmán, J., 2021. Telltale silence: temporal speech parameters discriminate between prodromal dementia and mild Alzheimer's disease. *Clin. Linguist. Phon.* 35, 727–742. <https://doi.org/10.1080/02699206.2020.1827043>.

- Vos de Wael, R., Benkarim, O., Paquola, C., Larivière, S., Royer, J., Tavakol, S., Xu, T., Hong, S.-J., Lams, G., Valk, S., Misić, B., Milham, M., Margulies, D., Smallwood, J., Bernhardt, B.C., 2020. BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Commun. Biol.* 3, 1–10. <https://doi.org/10.1038/s42003-020-0794-7>.
- Wang, X., Krieger-Redwood, K., Zhang, M., Cui, Z., Wang, X., Karapanagiotidis, T., Du, Y., Leech, R., Bernhardt, B.C., Margulies, D.S., Smallwood, J., Jefferies, E., 2022. Physical distance to sensory-motor landmarks predicts language function. *Cereb. Cortex N. Y. N 1991 bhac344* <https://doi.org/10.1093/cercor/bhac344>.
- Yamada, Y., Shinkawa, K., Nemoto, M., Ota, M., Nemoto, K., Arai, T., 2022. Speech and language characteristics differentiate Alzheimer's disease and dementia with Lewy bodies. *Alzheimers Dement. Amst. Neth.* 14, e12364.
- Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zöllei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R. L., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106, 1125–1165. <https://doi.org/10.1152/jn.00338.2011>.
- Zimmerer, V.C., Hardy, C.J.D., Eastman, J., Dutta, S., Varnet, L., Bond, R.L., Russell, L., Rohrer, J.D., Warren, J.D., Varley, R.A., 2020. Automated profiling of spontaneous speech in primary progressive aphasia and behavioral-variant frontotemporal dementia: An approach based on usage-frequency. *Cortex* 133, 103–119. <https://doi.org/10.1016/j.cortex.2020.08.027>.