

Single vs. multiple evaluations of satisfaction paired with Just-About-Right scales on full portions of beer at home

Takahiro Wakihira, Michel Visalli, Pascal Schlich

▶ To cite this version:

Takahiro Wakihira, Michel Visalli, Pascal Schlich. Single vs. multiple evaluations of satisfaction paired with Just-About-Right scales on full portions of beer at home. 15TH PANGBORN SENSORY SCIENCE SYMPOSIUM - MEETING NEW CHALLENGES IN A CHANGING WORLD, Aug 2023, Nantes, France. hal-04533767

HAL Id: hal-04533767 https://hal.science/hal-04533767

Submitted on 5 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Single vs. multiple evaluations of satisfaction paired with Just-About-Right scales on full portions of beer at home

Takahiro Wakihira¹, Michel Visalli^{2,3}, Pascal Schlich^{2,3}

¹ Brewing Science Laboratories, ASAHI QUALITY & INNOVATIONS, LTD. ² Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE, Université Bourgogne ³ INRAE, PROBE Research Infrastructure, ChemoSens Facility

Introduction and Objective

✓ Consumers' responses are usually collected retrospectively after consumption of a product in home use tests, which requires the consumers to recall their perceptions on aromas or tastes.

✓ Eating and drinking experiences are rather variable during consumption of the full portion, which implies the importance of collecting data over different periods of consumption.

Y The objective of this study was to determine if a 3-step multiple evaluation with full consumption of a product can provide additional information compared to a single evaluation after full consumption in home use tests.

Materials and Methods

Test design


Asahi

- A total of 447 Japanese beer consumers were recruited to taste four beers, whose alcohol by volume was 5% (P1), 0.5% (P2), 0.5% (P3), and 0% (P4), each one on a different day at home.
- Y P1 and P2 were tested twice (P1_rep, P2_rep) by each consumer without letting them know that they were the same (i.e., different sample codes). Therefore, the consumers had to taste six samples on six different days that they could choose as they wanted. Tasting order of the six samples was designed to be balanced according to Williams Latin squares.
- Y The consumers were separated into two panels. Both panels evaluated the beers based on satisfaction paired with Just-About-Right (JAR) scales, with their own smartphones at home. One of the panels (SING panel) evaluated the beer only once, after finishing the full portion. The other panel (MULT panel) evaluated the same questions three times repeatedly, after the first sip,



after half of the portion, and after the full portion (i.e., 3-step evaluations).

- ✓ The beer consumers included both users and non-users (20-39 years old) of beer alternative products with less than 1% alcohol content.
- ✓ Following the instructions displayed on their own smartphone screens, each consumer tasted a full portion of a beer at their own drinking pace.
- \checkmark Data were collected using the TimeSens[©] V2 web app (INRAE, Dijon, France).



Data analysis

- ANOVA model for Satisfaction: Satisfaction = Product + Method + User + Product*User + Product*Use Note: "Method" refers to SING or MULT panel. "User" refers to users of beer alternative products. Data collected after the full can were used for both panels in this model.
- ✓ ANOVA models for JAR scales: Attribute = Product + Method + User + Product*Method + Product*User + Method*User + error

Panelist: a random effect. All the others: fixed effects.

Note: "Attribute" refers to percentages of Just-About-Right, Not Enough, Too Much, respectively for each attribute. Data collected after the full can were used for both panels in this model.


ANOVA model for evolution of Wanting: Wanting = Product + Step + User + Product*Step + Product*User + Step*User + Product*Step*User + Panelist(User) + error

Note: "Step" refers to "after the first sip," "after half of the portion," and "after the full portion." Data collected from the MULT panel were used in this model. (SING panel data not used.)

Results

ANOVA of Satisfaction

- Y Product rankings and statistical differences of satisfaction were consistent among SING and MULT panels. No interaction was observed between product and method, indicating no methodological bias caused by the repeated questions in the 3-step approach.
- Inter MULT panel showed higher satisfaction scores than the SING panel. Satisfaction scores after the 1st sip among the MULT panel were higher than those of the SING panel after the full can, and the MULT panel had unchanged levels of satisfaction for the tested beers across the three evaluations. (Data tables not shown in this poster.)



| Method panel p | banel |
|---------------------------------|--|
| 70 0.973 0.579a 0 | .588a |
| 02 0.601 0.612a 0 | .657a |
| 2 <mark>5</mark> 0.127 0.529a 0 | .566b |
| 0.302 0.529a 0 | .616b |
| 0.183 0.644a 0 | .659a |
| 16 0.311 0.458a 0 | .529b |
| 04 0.065 0.497a 0 | .574b |
| 0.592a 0.592a 0 | .642b |
| 04 0.774 0.605a 0 | .671b |
| 18 0.273 0.551a 0 | .601a |
| | Methodpanelp700.9730.579a0020.6010.612a0250.1270.529a0020.3020.529a0120.1830.644a0460.3110.458a0040.0650.497a0040.7740.605a0 |

| p-values of tests of fixed effects | | | | | | User LS-Means | | |
|------------------------------------|---------|---------|------------------|------------------|---------------|---------------------------|---------------|--------|
| Product | Step | User | Product *Step | Product* User | Step* User | Product *Step* User | Non- users | Users |
| < 0.001 | < 0.001 | < 0.001 | < 0.001 | 0.013 | 0.005 | 0.820 | 6.141a | 6.878b |

| Before the 1 st sip | | After a half can | + | After the full can | | | | |
|--------------------------------------|--|------------------------|---|--------------------------|--|--|--|--|
| Evaluation Steps | | | | | | | | |

Conclusions

- ✓ Comparing results of the SING and the MULT panels proved the consistency in both hedonic and sensory evaluations toward the beers.
- ✓ The MULT panel, who had unchanged levels of satisfaction across the three evaluations, showed higher satisfaction scores than the SING panel.
- ✓ Non-users of beer alternative products with less than 1% alcohol content were less satisfied with the tested beers than users.
- ✓ The results from 3-step multiple evaluations revealed the changes in *wanting* for drinking a beer during a whole drinking experience.
- ✓ The research design applied in this study was useful at home, and would add greater value to product development and marketing.

Presenter: Takahiro Wakihira. E-mail: takahiro.wakihira@asahi-qi.co.jp