

# Branching Brownian motion versus Random Energy Model in the supercritical phase: overlap distribution and temperature susceptibility

Benjamin Bonnefont, Michel Pain, Olivier Zindy

# ▶ To cite this version:

Benjamin Bonnefont, Michel Pain, Olivier Zindy. Branching Brownian motion versus Random Energy Model in the supercritical phase: overlap distribution and temperature susceptibility. 2024. hal-04537499

# HAL Id: hal-04537499 https://hal.science/hal-04537499

Preprint submitted on 8 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Branching Brownian motion versus Random Energy Model in the supercritical phase: overlap distribution and temperature susceptibility

Benjamin BONNEFONT<sup>\*</sup>, Michel PAIN<sup>†</sup> and Olivier ZINDY<sup>‡</sup>

#### Abstract

In comparison with Derrida's REM, we investigate the influence of the so-called decoration processes arising in the limiting extremal processes of numerous log-correlated Gaussian fields. In particular, we focus on the branching Brownian motion and two specific quantities from statistical physics in the vicinity of the critical temperature. The first one is the two-temperature overlap, whose behavior at criticality is smoothened by the decoration process—unlike the one-temperature overlap which is identical—and the second one is the temperature susceptibility, as introduced by Sales and Bouchaud, which is strictly larger in the presence of decorations and diverges, close to the critical temperature, at the same speed as for the REM but with a different multiplicative constant. We also study some general decorated cases in order to highlight the fact that the BBM has a critical behavior in some sense to be made precise.

# Contents

1	Introduction and results							
	1.1	Motivations	2					
	1.2	Definitions and some results	3					
	1.3	Main results	6					
	1.4	Organization of the paper	8					
<b>2</b>	Preliminaries							
	2.1	Properties of $Z(\beta)$	8					
	2.2	Change of measure in the general decorated case	10					
	2.3	The decoration of the BBM	12					
3	Ove	erlap distribution at two temperatures	15					
	3.1	Results in the general decorated case	15					
	3.2	Some tools for the proofs	19					
	3.3	Proof of the general theorems	21					
	3.4	The REM case	23					
	3.5	The BBM case	24					

\*Université de Genève, Section de Mathématiques, Rue du Conseil-Général 7-9, 1205 Genève, Suisse. Email: benjamin.bonnefont@unige.ch. Partially supported by GdR Branchement.

<sup>†</sup>Institut Mathématiques de Toulouse (UMR 5219), Université de Toulouse, France, CNRS. Email: michel.pain@math.univ-toulouse.fr. Partially supported by GdR Branchement and by PEPS JCJC grant n°246644.

<sup>‡</sup>Sorbonne Université, Sorbonne Paris Cité, CNRS, Laboratoire de Probabilités Statistique et Modélisation, LPSM, F-75005 Paris, France. Email: olivier.zindy@sorbonne-universite.fr. Partially supported by GdR Branchement.

<b>4</b>	Temperature susceptibility				
	4.1	A first formula for $\kappa_d$	26		
	4.2	The REM case	27		
	4.3	Comparison wih the REM in the general decorated case	29		
	4.4	The BBM case	31		
Aı	ppen	dix	38		
$\mathbf{A}$	Pro	of of technical results on the decoration of the BBM	38		
	A.1	Uniform bounds for moments of level sets	38		
	A.2	First moment of level sets on a particular event	39		
	A.3	Cross-moments of level sets	42		
	A.4	Small moments of $S_{\beta}$	44		

# 1 Introduction and results

## 1.1 Motivations

In order to shed some light on the mysteries of the Parisi theory for mean field spin glasses, Derrida introduced in the 80's the random energy models (REMs) [24], where the Gaussian energy levels are assumed to be independent, and its generalizations, the generalized random energy models (GREMs) [25], whose correlations are given by a tree structure of finite depth. One question of central interest in spin glass theory is to understand the structure of pure states, which translates into the analysis of the extremal process in the language of extreme value theory of stochastic processes.

These two tractable models have been extensively studied and allowed, in particular, to investigate the phenomenon of replica symmetry breaking. Indeed the REM seems to be the foremost representative of a universality class: in spin glass terminology one may call this the 1-step replica symmetry breaking (1-RSB) class, or REM-class. More precisely, a spin glass model displays a 1-RSB if there exists some critical  $\beta_c > 0$  such that, asymptotically, the overlap between two points chosen independently according to the Gibbs measure at inverse temperature  $\beta > 0$  is concentrated at 0 for  $\beta \leq \beta_c$ , but is supported by 0 and 1, when  $\beta > \beta_c$ . This phenomenon is a consequence of the fact that the REM-class undergo what physicists refer to as the *REM-freezing transition*: there is a phase transition for the free energy at some  $\beta_c > 0$ meaning that, for  $\beta \leq \beta_c$ , there is an exponentially large number of configurations, with energy level strictly less than the extremes, contributing to free energy and the Gibbs measure is roughly uniformly distributed among such configurations while, for  $\beta > \beta_c$ , the relevant configurations are the ones with the largest energies and the free energy becomes dominated by a relatively small set of configurations. Another striking fact characterizing the REM-class is that, for  $\beta > \beta_c$ , the ordered weights of the pure states under the Gibbs measure at inverse temperature  $\beta > 0$  follow asymptotically a *Poisson-Dirichlet distribution* of parameter  $\beta_c/\beta$ . We refer to Bolthausen [11, 8] and Kistler [32] for surveys on the REMs, GREMs and connections to spin glass theory. Finally, let us mention that, despite the simplicity of the freezing transition, rather sophisticated models are known, or conjectured, to belong to the REM-class, such is the case for the extremes of the Riemann zeta-function along the critical line, see Arguin [2] for a survey.

Natural hierarchical models with an infinite number of levels are the branching Brownian motion (BBM) and the branching random walk (BRW), see e.g. the seminal paper by Derrida and Spohn [27], who introduced directed polymers on trees (a BRW with i.i.d. displacements) as an infinite hierarchical extension of the GREMs for spin glasses. Physicists suggested that Gaussian BRW and BBM should belong to a universality class called *log-correlated Gaussian* 

fields, or log-*REMs*, which is in some sense a "subclass" of the REM-class. These models are not necessarily hierarchical but admit correlations that decay approximately like the logarithm of the inverse of the distance between index points. We refer to the works by Carpentier and Le Doussal [18], Fyodorov and Bouchaud [29, 30] and Fyodorov & al [31] for connection between log-REMs and spin glass theory. Furthermore, these processes play an essential role in Liouville quantum gravity as well as models of three-dimensional turbulence or finance, see the review on Gaussian Multiplicative Chaos by Rhodes and Vargas [44] for discussions.

Another line of research heavily relies on relations between log-REMs and traveling wave equations of Fisher-Kolmogorov-Petrovsky-Piscounov (FKPP) type. Such a relation is exact for a particular instance of log-REM: BBM mentioned above. Therefore BBM was studied over the last 50 years as a subject of interest in its own right, with contributions by McKean [39], Bramson [17, 14], Lalley and Sellke [34], Chauvin and Rouault [19, 20]. From the probabilistic point of view, the full picture was recently obtained by Aïdékon & al [1] and Arguin & al [3] separately, while Cortines & al [21, 22] obtained a third description of the decorations' distribution. Indeed it is now known that, in the thermodynamic limit, the extremal process tends to a randomly shifted decorated Poisson point process (SDPPP), see [46] for a precise definition. Compared with the Poisson point process which describes the extremes of the (uncorrelated) REM, the decorations appearing here describe the internal structure of blocks of extreme values which share a near *ancestor*, and thus are highly correlated. For instance, this confirms the observation made by Bovier and Kurkova [13] that BBM is a particularly interesting example, lying right at the borderline where correlations begin to influence the behavior of the extremes of the process. Finally one can say that BBM is the prototype of hierarchical log-REMs and therefore will naturally be the model of interest in this paper.

Let us also mention that another important and famous example of log-correlated Gaussian fields is the two-dimensional discrete Gaussian Free Field (GFF), which possesses a complicated (non-hierachical) structure of extreme values, but it turns out to be possible to compare it with that of the branching random walk. By comparison to analoguous results for branching random walks, many deep results have been recently established by Bolthausen, Deuschel and Giacomin [9], Daviaud [23], Arguin and Zindy [4], Bolthausen, Deuschel and Zeitouni [10], Bramson and Zeitouni [16], Bramson, Ding and Zeitouni [15], Biskup and Louidor [6, 7]. We refer to the excellent notes by Biskup [5] for more details.

While from a probabilistic point of view, the difference between BBM and REM is perfectly known, it is not the same when quantities from statistical physics are considered. Indeed it is not clear when the decorations of the log-correlated models are felt at the level of the Gibbs measure. For example, Bonnefont [12] recently proved for the BBM that the distribution of the overlap between two points sampled independently according to Gibbs measures at different temperatures is different than the REM's one, while it is the same for both models if the two points are sampled at the same temperature. This raises questions about the influence of these decorations. Motivated by the recent work of Derrida and Mottishaw [26], one seeks to compare both models by studying carefuly the overlap distribution in the neighborhood of the critical temperature. Another quantity of interest is the notion of temperature susceptibility introduced and studied for the REM (with exponentially distributed energies) by Sales and Bouchaud [45].

### **1.2** Definitions and some results

A probabilistic point of view. On one side, the (binary) branching Brownian motion (BBM) is a branching Markov process defined on some general probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  as follows. Initially, there is one single particle at the origin which moves according to a standard Brownian

motion during an exponentially distributed time of parameter  $1/2^1$  and then splits into two new particles. These new particles start the same process from their place of birth, behaving independently of the others and the system goes on indefinitely. Let  $L_t$  denote the set of alive particles at time  $t \ge 0$  and  $h_t(x)$  the position of the particle x at time t. The position of the highest particle has been studied by Bramson [17, 14] and Lalley and Sellke [34]. A new step has recently been taken with the proof of the convergence of the extremal process in the space of Radon measures on  $\mathbb{R}$  endowed with the vague topology, to a randomly shifted decorated Poisson point process. More precisely, Aïdékon et al. [1] and Arguin et al. [3] proved simultaneously that

$$\sum_{x \in L_t} \delta_{h_t(x) - t + \frac{3}{2}\log t - \log(c_d D_\infty)} \xrightarrow[t \to \infty]{(d)} \sum_{i,j} \delta_{\xi_i + d_{ik}}, \qquad t \to +\infty,$$
(1.1)

for some positive constant  $c_d > 0$  and where  $D_{\infty} := \lim_{t \to +\infty} \sum_{x \in L_t} (t - h_t(x)) e^{h_t(x) - t} > 0$ (almost-surely) is the limiting *derivative martingale*,  $\sum_{i \ge 1} \delta_{\xi_i}$  is a Poisson point process on  $\mathbb{R}$  with intensity measure  $e^{-x} dx$  independent of  $(\sum_{k \ge 0} \delta_{d_{ik}})_{i \ge 1}$ , which are are i.i.d. copies of a point process  $\mathcal{D}$  on  $(-\infty, 0]$  which has a.s. an atom at 0. The point process  $\mathcal{D}$  is called *decoration process*<sup>2</sup>.

On the other side, the REM of interest is defined as follows in order to be comparable to the BBM introduced above: it consists of  $n_t := \lfloor e^{t/2} \rfloor$  i.i.d. centered Gaussian random variables of variance t, denoted by  $(g_t(k); 1 \le k \le n_t)$ . It is well known that the extremal process for the REM satisfies the following convergence in the space of Radon measures on  $\mathbb{R}$  endowed with the vague topology:

$$\sum_{1 \le k \le n_t} \delta_{g_t(k) - t + \frac{1}{2}\log t - \log c_0} \xrightarrow[t \to \infty]{(d)} \sum_i \delta_{\xi_i}, \qquad t \to +\infty,$$
(1.2)

for some numerical positive constant  $c_0 > 0$  and where again the  $(\xi_i)_{i\geq 1}$  are the atoms of a Poisson point process on  $\mathbb{R}$  with intensity measure  $e^{-x} dx$ . We refer to Kistler [32] for a recent survey on the REM.

Looking first at the convergences in Equations (1.1) and (1.2) allows to compare the BBM and the REM from a probabilistic point of view: both model's maxima share the same first order t but the correlations for the BBM start to affect the second order, namely  $-\frac{3}{2} \log t$  for the BBM is smaller than  $-\frac{1}{2} \log t$  for the REM. And finally Equations (1.1) and (1.2) also complete the picture by telling that the limiting extremal process for the REM is a standard Poisson point process while the BBM's one is a randomly shifted decorated Poisson point process. The additional ingredient for the BBM is mainly the decoration process, which describes the internal structure of blocks of extremal particles sharing a near ancestor and thus highly correlated.

A statistical physics approach. In statistical physics, it is common to consider first the partition function  $Z_{t,d}(\beta)$  of the model, here the BBM ( $\beta$  stands for the inverse-temperature):

$$Z_{t,d}(\beta) \coloneqq \sum_{x \in L_t} e^{\beta h_t(x)}, \qquad \forall \beta > 0,$$

and the *free energy* 

$$f_{t,d}(\beta) \coloneqq \frac{1}{t} \log Z_{t,d}(\beta), \qquad \forall \beta > 0.$$

<sup>&</sup>lt;sup>1</sup>This choice is made such that the critical inverse temperature introduced later in this article and denoted by  $\beta_c$  equals 1.

<sup>&</sup>lt;sup>2</sup>We will also consider the case of general decoration processes, see Section 2.2 and use the same notation  $(\sum_{k>0} \delta_{d_{ik}})_{i\geq 1}$ . If the case of general decoration processes is studied, this will be specified, otherwise it is only the BBM's decoration process.

It is well known, see [27], that the BBM exhibits a phase transition at the level of the free energy and that this latter is the same as for the REM, namely

$$f_d(\beta) \coloneqq \lim_{t \to \infty} f_{t,d}(\beta) = \begin{cases} 1 + \beta^2, & \text{if } \beta < \beta_c \coloneqq 1, \\ 2\beta, & \text{if } \beta \ge \beta_c, \end{cases} \quad \text{a.s. and in } L^1.$$

In particular, the model undergoes *freezing* above  $\beta_c$  in the sense that the quantity  $f(\beta)/\beta$  is constant. More importantly, one considers the *Gibbs measure*; this is the random probability measure  $\mathcal{G}_{\beta,t,d}$  on  $L_t$  defined, at inverse temperature  $\beta > 0$ , by

$$\mathcal{G}_{\beta,t,d}(x) \coloneqq \frac{\mathrm{e}^{\beta h_t(x)}}{Z_{t,d}(\beta)}, \qquad \forall x \in L_t.$$

By design, the Gibbs measure concentrates at low temperature, i.e. when  $\beta > \beta_c$ , on the high points of the Gaussian field. With spin glasses in mind, one also considers the normalized covariance or *overlap* 

$$q_{t,d}(x,y) \coloneqq \frac{1}{t} \mathbb{E}[h_t(x)h_t(y)] = \frac{1}{t} \sup\{s \ge 0 : x, y \text{ share a common ancestor in } L_s\}, \quad \forall x, y \in L_t.$$

A fundamental object, that records the correlations of high points, is the distribution function of the overlap sampled from the Gibbs measure, i.e.  $\mathcal{G}_{\beta,t,d} \otimes \mathcal{G}_{\beta',t,d}(q_{t,d}(u,v) \geq a)$  for any  $a \in (0,1)$ , where u (respectively v) is sampled according to  $\mathcal{G}_{\beta,t,d}$  (respectively  $\mathcal{G}_{\beta',t,d}$ ). Bonnefont [12] recently proved that, if  $\beta \leq \beta_c$  or  $\beta' \leq \beta_c$ , then  $\mathcal{G}_{\beta,t,d} \otimes \mathcal{G}_{\beta',t,d}(q_{t,d}(u,v) \geq a)$  tends to 0 in  $L^1$  for all  $a \in (0,1)$ , while if  $\beta > \beta_c$  and  $\beta' > \beta_c$ , then, for all  $a \in (0,1)$ ,

$$\mathcal{G}_{\beta,t,d} \otimes \mathcal{G}_{\beta',t,d}(q_{t,d}(u,v) \ge a) \xrightarrow[t \to \infty]{(d)} \frac{\sum_{i} \left( e^{\beta \xi_{i}} \sum_{k} e^{\beta d_{ik}} \right) \left( e^{\beta' \xi_{i}} \sum_{k} e^{\beta' d_{ik}} \right)}{\left( \sum_{i,k} e^{\beta \xi_{i}} e^{\beta d_{ik}} \right) \left( \sum_{i,k} e^{\beta' \xi_{i}} e^{\beta' d_{ik}} \right)} \Longrightarrow Q_{d}(\beta,\beta'), \quad (1.3)$$

where the  $(\xi_i)_i$  and the  $(d_{ik})_{i,k}$  were introduced for the description of the limiting extremal process of the BBM, see Equation (1.1). In other words, this result proves the convergence of the pushforward of the measure  $\mathcal{G}_{\beta,t,d} \otimes \mathcal{G}_{\beta',t,d}$  on  $L_t^2$  by the function  $q_{t,d}$ , which is a random measure on [0, 1]. The limit is either  $\delta_0$  if  $\beta \wedge \beta' \leq \beta_c$ , or  $(1 - Q_d(\beta, \beta'))\delta_0 + Q_d(\beta, \beta')\delta_1$  otherwise. Note that, when  $\beta > \beta_c$  and  $\beta' > \beta_c$ , the random variables  $(e^{\beta\xi_i} \sum_k e^{\beta d_{ik}} / \sum_j e^{\beta\xi_j} \sum_k e^{\beta d_{jk}})_{i\geq 1}$  are the asymptotic weights of the clusters under  $\mathcal{G}_{\beta,t,d}$ , such that  $Q_d(\beta, \beta')$  is simply the probability of choosing two points in the same cluster (when they are chosen proportionally to their Gibbs weights with inverse temperature  $\beta$  and  $\beta'$  respectively).

For the REM, the overlap is simply given by

$$q_t(k,\ell) \coloneqq \frac{1}{t} \mathbb{E}[g_t(k)g_t(\ell)] = \mathbb{1}_{\{k=\ell\}}, \qquad \forall 1 \le k, \ell \le n_t,$$

and the Gibbs measure at inverse temperature  $\beta > 0$  is defined by

$$\mathcal{G}_{\beta,t}(k) \coloneqq \frac{\mathrm{e}^{\beta g_t(k)}}{Z_t(\beta)}, \qquad \forall 1 \le k \le n_t,$$

where  $Z_t(\beta) := \sum_{1 \le k \le n_t} e^{\beta g_t(k)}$ . Kurkova [33] proved that, if  $\beta \le \beta_c$  or  $\beta' \le \beta_c$ , then  $\mathcal{G}_{\beta,t} \otimes \mathcal{G}_{\beta',t}(q_t(u,v) \ge a)$  tends to 0 in  $L^1$  for all  $a \in (0,1)$ , while if  $\beta > \beta_c$  and  $\beta' > \beta_c$ , then, for all  $a \in (0,1)$ ,

$$\mathcal{G}_{\beta,t} \otimes \mathcal{G}_{\beta',t}(q_t(u,v) \ge a) \xrightarrow[t \to \infty]{(\mathrm{d})} \frac{\sum_i \mathrm{e}^{\beta\xi_i} \mathrm{e}^{\beta'\xi_i}}{(\sum_i \mathrm{e}^{\beta\xi_i})(\sum_i \mathrm{e}^{\beta'\xi_i})} \eqqcolon Q(\beta,\beta').$$

It is well known that in the case  $\beta' = \beta$ , the random variables  $Q_d(\beta, \beta)$  and  $Q(\beta, \beta)$  have the same distribution (see [42, Equation (1.2)] for more details). Therefore a natural question one

may ask is whether  $Q_d(\beta, \beta')$  and  $Q(\beta, \beta')$  still have the same distribution when  $\beta \neq \beta' > \beta_c$ . Following the work by Pain and Zindy [42] for the two-dimensional discrete Gaussian free field, Bonnefont [12] proved, for the BBM, that

$$F_d(\beta, \beta') \coloneqq \mathbb{E}[Q_d(\beta, \beta')] < \mathbb{E}[Q(\beta, \beta')] \rightleftharpoons F(\beta, \beta'), \tag{1.4}$$

meaning that the answer is negative. In this paper, our aim is to study and compare the functions  $\beta \mapsto F_d(\beta, \beta')$  and  $\beta \mapsto F(\beta, \beta')$ , when  $\beta' > \beta_c$  is fixed. We will specially focus on their behavior when  $\beta$  tends to  $\beta_c^+$ .

Before introducing the second quantity of interest let us define, for any  $\beta > \beta_c$ , the partition functions associated with both limiting extremal processes

$$Z(\beta) \coloneqq \sum_{i \ge 1} e^{\beta \xi_i}, \qquad Z_d(\beta) \coloneqq \sum_{i \ge 1} \sum_{k \ge 0} e^{\beta(\xi_i + d_{ik})}.$$
(1.5)

Then, let us consider the correlation coefficient between the free energies at two temperatures, introduced by Fisher and Huse [28] and given for both models by

$$\mathbf{C}(\beta,\beta') \coloneqq \frac{\operatorname{Cov}(\log Z(\beta), \log Z(\beta'))}{\sigma(\log Z(\beta)) \,\sigma(\log Z(\beta'))}, \qquad \mathbf{C}_d(\beta,\beta') \coloneqq \frac{\operatorname{Cov}(\log Z_d(\beta), \log Z_d(\beta'))}{\sigma(\log Z_d(\beta)) \,\sigma(\log Z_d(\beta'))},$$

where  $\sigma(X) \coloneqq \sqrt{\operatorname{Var}(X)}$ . The susceptibility to temperature we are interested in is defined by Sales and Bouchaud [45] as the coefficient  $\kappa_d(\beta)$  for the BBM and  $\kappa(\beta)$  for the REM such that

$$\begin{aligned} \mathbf{C}(\beta,\beta+h) &= 1 - \kappa(\beta) \, h^2 + o(h^2), \qquad h \to 0, \\ \mathbf{C}_d(\beta,\beta+h) &= 1 - \kappa_d(\beta) \, h^2 + o(h^2), \qquad h \to 0 \end{aligned}$$

See Corollary 4.2 for a justification that this coefficient exists. Let us emphasize that Sales and Bouchaud [45] work with a slightly different convention, which results in their temperature susceptibility to equal  $\kappa(\beta)\beta^2$  with our notation.

**Notation.** Throughout the paper, C and c denote a positive constant that does not depend on the parameters and can change from line to line. For  $f: (1, \infty) \to \mathbb{R}$  and  $g: (1, \infty) \to \mathbb{R}^*$ , we say, as  $\beta \downarrow 1$ , that  $f(\beta) = o(g(\beta))$  if  $\lim_{\beta \downarrow 1} f(\beta)/g(\beta) = 0$ , that  $f(\beta) = O(g(\beta))$  if  $\lim_{\beta \downarrow 1} |f(\beta)|/g(\beta) < \infty$ , that  $f(\beta) \sim g(\beta)$  if  $\lim_{\beta \downarrow 1} f(\beta)/g(\beta) = 1$  and that  $f(\beta) \asymp g(\beta)$  if  $0 < \liminf_{\beta \downarrow 1} f(\beta)/g(\beta) \le \limsup_{\beta \downarrow 1} f(\beta)/g(\beta) < \infty$ .

#### 1.3 Main results

Near-critical two-temperature overlap. Our first results concern the behavior of the mean overlap  $F(\beta, \beta')$  or  $F_d(\beta, \beta')$  at two supercritical inverse temperatures  $\beta$  and  $\beta'$ . Recall from Equation (1.4) that it has already been proved that  $F_d(\beta, \beta') < F(\beta, \beta')$ . Our goal here is to quantify this difference in the regime where  $\beta'$  is fixed but  $\beta$  approaches the critical inverse temperature  $\beta_c = 1$ .

We start with the REM case. It is not hard to see that the function  $\beta \mapsto F(\beta, \beta')$  has an infinite right-derivative at 1, see Remark 3.10. We establish a sharper estimate in the following result.

**Theorem 1.1** (REM case). For any  $\beta' > \beta_c = 1$ , as  $\beta \downarrow 1$ ,

$$F(\beta, \beta') = (\beta - 1) \log \frac{1}{\beta - 1} + O(\beta - 1).$$

We compare this behavior to the one arising for the BBM. Surprisingly, the decorations have the effect of smoothing the function  $\beta \mapsto F_d(\beta, \beta')$  at criticality: more precisely,  $F_d(\beta, \beta')$  is of order  $(\beta - 1)$  as  $\beta \downarrow 1$ , as shown in the following result. It would be interesting to obtain an asymptotic equivalent instead, which would prove that  $\beta \mapsto F_d(\beta, \beta')$  is right-differentiable at  $\beta = 1$ , but this seems out of reach with the techniques used here.

**Theorem 1.2** (BBM case). For any  $\beta' > \beta_c = 1$ ,

$$0 < \liminf_{\beta \downarrow 1} \frac{F_d(\beta, \beta')}{\beta - 1} \le \limsup_{\beta \downarrow 1} \frac{F_d(\beta, \beta')}{\beta - 1} < +\infty.$$

In order to shed some light on these different behaviors, we study in Section 3.1 the case of general decoration processes  $\mathcal{D} = \sum_{k\geq 0} \delta_{d_k}$ . We do not state these results precisely here, some of them having very technical assumptions, but rather give an informal description of some of the conclusions:

- If the sum  $\sum_{k\geq 0} e^{\beta d_k}$  has bounded  $(1+\varepsilon)$ -moment as  $\beta \downarrow 1$ , then the behavior of  $F_d(\beta, \beta')$  is the same as for the REM in Theorem 1.1, see Corollary 3.2.
- If the γ-th moment of the sum Σ<sub>k≥0</sub> e<sup>βdk</sup> explodes like (β − 1)<sup>-ψ(γ)</sup> as β ↓ 1, then different behaviors appear depending on the value of ψ'(1) − ψ(1): the larger this value—in other words, the more moments of Σ<sub>k≥0</sub> e<sup>βdk</sup> are governed by rare events—the faster F<sub>d</sub>(β,β') decreases as β ↓ 1. More precisely, when ψ'(1) − ψ(1) < 1, F<sub>d</sub>(β,β') is still of order (β−1) log <sup>1</sup>/<sub>β−1</sub> like in the REM case, but possibly with another multiplicative constant. At the critical value ψ'(1) − ψ(1) = 1, the first order constant vanishes and F<sub>d</sub>(β,β') becomes of order (β − 1). Above the critical value, F<sub>d</sub>(β,β') decreases faster than (β − 1)<sup>1+η</sup> for η > 0 in some explicit interval, see Corollary 3.5 for a precise statement.

It turns out that the BBM corresponds exactly to the critical case described in the second point. This fact is a consequence of some fine properties of the decoration of the BBM, proved by building upon results by Cortines, Hartung and Louidor [21, 22], who showed that the mean of level sets of the decoration process is driven by rare events in which the decoration is much bigger than its typical behavior. We believe that this should be the case of other models falling in the log-correlated fields universality class, such as general branching random walks or the 2D discrete Gaussian free field.

**Temperature susceptibility.** We now turn to results concerning the temperature susceptibility. We first compute it explicitly for the REM in the supercritical phase and study its asymptotic behavior close to the critical temperature, as well as in the low temperature regime.

**Theorem 1.3** (REM case). For any  $\beta > \beta_c = 1$ ,

$$\kappa(\beta) = \frac{1}{2} \left( \frac{1}{\beta^2 - 1} + \frac{6}{\pi^2 \beta^3 (\beta + 1)} \left( \frac{\Gamma''\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)} - \left(\frac{\Gamma'\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)}\right)^2 \right) - \frac{\beta^2}{(\beta^2 - 1)^2} \right)$$

and

$$\kappa(\beta) \underset{\beta \downarrow 1}{\sim} \left(\frac{3}{2\pi^2} - \frac{1}{8}\right) \frac{1}{(\beta - 1)^2}, \qquad \qquad \kappa(\beta) \underset{\beta \to +\infty}{\sim} \left(\frac{6}{\pi^2} \zeta(3) - \frac{1}{2}\right) \frac{1}{\beta^5}.$$

Remark 1.4. This question has already been studied in the physics literature by Sales and Bouchaud [45] in the case of the REM with exponentially distributed energies. The formula they obtain for  $\kappa(\beta)$  (see their appendix) does not seem to match the one above, but we are not able to find the source of this discrepancy. In particular, the exponents appearing as  $\beta \downarrow 1$  and  $\beta \to \infty$  are different.

Then, we study the temperature susceptibility of the BBM. The following results shows that it is strictly larger than the one of the REM and gives the behavior of the temperature susceptibility close to the critical temperature, showing that it diverges at the same speed as for the REM but with a different multiplicative constant. We do not investigate the behavior at low temperature.

**Theorem 1.5** (BBM case). For any  $\beta > \beta_c = 1$ , one has  $\kappa_d(\beta) > \kappa(\beta)$ . Moreover, we have

$$\frac{3}{2\pi^2} - \frac{1}{8} < \liminf_{\beta \downarrow 1} \, (\beta - 1)^2 \, \kappa_d(\beta) \le \limsup_{\beta \downarrow 1} \, (\beta - 1)^2 \, \kappa_d(\beta) \le \frac{3}{\pi^2} - \frac{1}{8}.$$

As for the two-temperature overlap, we also investigate the general decorated case. Proposition 4.5 shows that  $\kappa_d(\beta)$  can be written as the sum of  $\kappa(\beta)$  and a nonnegative term, which is positive for most non-deterministic decoration processes. Showing that this additional term is growing like  $(\beta - 1)^{-2}$  as  $\beta \downarrow 1$ , in particular the lower bound, is the main task in the proof of Theorem 1.5. In general, this term can be negligible w.r.t.  $(\beta - 1)^{-2}$  for "small enough" decorations, and it can be much larger than  $(\beta - 1)^{-2}$  for "big enough" decorations. So again, the BBM seems to belong to a critical window, even if here this critical window seems to include more models than for the two-temperature overlap, see Example 4.6.

#### 1.4 Organization of the paper

Section 2 includes preliminary results on the partition function of the REM, on a change of measures used to study the decorated case and on the decoration process of the BBM. Section 3 is dedicated to the two-temperature overlap, including proofs of Theorems 1.1 and 1.2, as well as results in more general decorated cases in Section 3.1. In Section 4, we investigate the temperature susceptibility, proving Theorems 1.3 and 1.5. Appendix A contains the proof of some results needed about the decoration process of the BBM throughout the paper.

# 2 Preliminaries

## **2.1** Properties of $Z(\beta)$

Recall that the partition function of the REM is given by  $Z(\beta) = \sum e^{\beta \xi_k}$  where  $(\xi_k)_{k\geq 1}$  are the atoms ranked in non-increasing order of a PPP( $e^{-x}dx$ ). It will be convenient to use the notation  $\eta_k := e^{-\xi_k}$  because these are the atoms of a homogeneous Poisson point process on  $\mathbb{R}_+$ and allows us to use the law of large numbers. Hence  $Z(\beta)$ , can be rewritten

$$Z(\beta) = \sum_{k \ge 1} \eta_k^{-\beta}, \qquad \forall \beta > 1.$$
(2.1)

We first show that  $Z(\beta)$  has a stable distribution. We use the following convention: the stable distribution  $S_{\alpha}(c, s, \mu)^3$  has a characteristic function given by

$$t \mapsto \exp\{it\mu - |ct|^{\alpha}(1 - is\operatorname{sgn}(t)\Phi)\}, \quad \text{where} \quad \Phi = \begin{cases} \tan(\frac{\pi\alpha}{2}), & \text{if } \alpha \neq 1, \\ -\frac{2}{\pi}\log|t|, & \text{if } \alpha = 1. \end{cases}$$

<sup>&</sup>lt;sup>3</sup>Here we do not use  $\beta$  for the skewness parameter because the letter is used for the inverse temperature.

**Lemma 2.1.** The Laplace transform of  $Z(\beta)$  is given, for any  $t \ge 0$ , by

$$\mathbb{E}\left[\mathrm{e}^{-tZ(\beta)}\right] = \exp\left\{-\Gamma\left(\frac{\beta-1}{\beta}\right)t^{\frac{1}{\beta}}\right\}.$$

In other words, the distribution of  $Z(\beta)$  is  $S_{1/\beta}\Big(\Big(\Gamma(1-\frac{1}{\beta})\cos\frac{\pi}{2\beta}\Big)^{\beta}, 1, 0\Big).$ 

*Proof.* The formula for Laplace transforms of Poisson point processes applied to Equation (2.1) gives

$$\mathbb{E}\left[\mathrm{e}^{-tZ(\beta)}\right] = \exp\left\{-\int_0^\infty (1 - \mathrm{e}^{-tx^{-\beta}})\,\mathrm{d}x\right\} = \exp\left\{-\Gamma\left(\frac{\beta - 1}{\beta}\right)t^{\frac{1}{\beta}}\right\},\,$$

which is the Laplace transform of the desired stable law.

**Lemma 2.2.** For any  $\beta > 1$  and any  $\alpha > -\frac{1}{\beta}$ ,

$$\mathbb{E}[Z(\beta)^{-\alpha}] = \frac{\Gamma(\alpha\beta + 1)}{\Gamma(\alpha + 1)\Gamma(\frac{\beta - 1}{\beta})^{\alpha\beta}}.$$

*Proof.* For  $\alpha > 0$ , this is proved by taking the expectation of

$$Z(\beta)^{-\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} \mathrm{e}^{-tZ(\beta)} \,\mathrm{d}t,$$

and then applying Lemma 2.1. For  $-1/\beta < \alpha < 0$ , we use instead the following representation

$$Z(\beta)^{-\alpha} = \frac{1}{\Gamma(1+\alpha)} \int_0^\infty t^\alpha Z(\beta) \,\mathrm{e}^{-tZ(\beta)} \,\mathrm{d}t,$$

and note that  $\mathbb{E}[Z(\beta)e^{-tZ(\beta)}]$  is obtained by differentiating  $\mathbb{E}[e^{-tZ(\beta)}]$  with respect to t.

The formula established in the previous lemma yields to the following moment estimates in the regime  $\beta \downarrow 1$ , which are used throughout the paper.

**Corollary 2.3.** Let  $\alpha > -1$ . Then, as  $\beta \downarrow 1$ ,

$$\mathbb{E}[Z(\beta)^{-\alpha}] = (\beta - 1)^{\alpha} \left( 1 + O\left((\beta - 1)\log\frac{1}{\beta - 1}\right) \right), \qquad (2.2)$$

$$\mathbb{E}\left[Z(\beta)^{-\alpha/\beta}\right] = (\beta - 1)^{\alpha}(1 + O(\beta - 1)), \qquad (2.3)$$

$$\mathbb{E}\left[Z(\beta)^{-1/\beta}\log\left(Z(\beta)^{1/\beta}\right)\right] = (\beta - 1)\log\frac{1}{\beta - 1} + O(\beta - 1).$$
(2.4)

*Proof.* The first two expansions are obtained by expanding the Gamma function in the formula given by Lemma 2.2. For the third one, note that, for any  $\beta > 1$  and  $\alpha > -1/\beta$ ,  $\frac{d}{d\alpha} \mathbb{E}[Z(\beta)^{-\alpha}] = -\alpha \mathbb{E}[Z(\beta)^{-\alpha} \log Z(\beta)]$ .

The expansions in the previous corollary are governed by the fact that  $Z(\beta)$  concentrates around  $1/(\beta - 1)$  as  $\beta \downarrow 1$ . More precisely, we have surprisingly the following almost sure expansion.

**Lemma 2.4.** When  $\beta \downarrow 1$ , we have

$$Z(\beta) = \frac{1}{\beta - 1} + S + o(1), \quad almost \ surrely,$$

where  $S \sim S_1(\frac{\pi}{2}, 1, 1 - \gamma)$  and  $\gamma$  is the Euler constant.

*Proof.* With the Riemann  $\zeta$  function defined by  $\zeta(\beta) = \sum_{k \ge 1} \frac{1}{k^{\beta}}$ , one has

$$Z(\beta) - \zeta(\beta) = \sum_{k \ge 1} \frac{k^{\beta} - \eta_k^{\beta}}{k^{\beta} \eta_k^{\beta}}.$$

If  $\beta \leq 2$ , one has

$$\frac{k^{\beta} - \eta_k^{\beta}}{k^{\beta} \eta_k^{\beta}} \le \frac{\beta \left( \eta_k^{\beta-1} + k^{\beta-1} \right) |\eta_k - k|}{k^{\beta} \eta_k^{\beta}} \le 5 \frac{|\eta_k - k|}{k^2},$$

for k large enough. This last term is summable since  $\eta_k = k + O(k^{\frac{3}{4}})$  a.s. Hence, the dominated convergence theorem (for series) implies that  $Z(\beta) - \zeta(\beta)$  has an a.s. limit when  $\beta$  tends to 1. The almost sure expansion follows from the fact that  $\zeta(\beta) = \frac{1}{\beta-1} + \gamma + o(1)$  when  $\beta \downarrow 1$ . Then, computing the characteristic function enables to identify the limiting law: if t > 0, one has

$$\begin{split} \mathbb{E}\Big[\mathrm{e}^{itZ(\beta)}\Big] &= \exp\Big\{-\Gamma\Big(\frac{\beta-1}{\beta}\Big)\cos\frac{\pi}{2\beta}t^{\frac{1}{\beta}}\Big(1-i\tan\frac{\pi}{2\beta}\Big)\Big\}\\ &= \exp\Big\{-\Gamma\Big(\frac{\beta-1}{\beta}\Big)t^{\frac{1}{\beta}}\mathrm{e}^{-i\frac{\pi}{2\beta}}\Big\}\\ &= \exp\Big\{it\Big(\frac{1}{\beta-1}-\log(t)+1-\gamma+i\frac{\pi}{2}\Big)+O(\beta-1)\Big\}, \end{split}$$

as  $\beta \downarrow 1$ , therefore

$$\mathbb{E}\left[\mathrm{e}^{it\left(Z(\beta)-\frac{1}{\beta-1}\right)}\right] \xrightarrow[\beta\downarrow 1]{} \exp\left\{it(1-\gamma)-\frac{\pi}{2}t\left(1+i\frac{2}{\pi}\log t\right)\right\},\$$

which is the expected characteristic function. The case t < 0 is similar.

*Remark* 2.5. These 1-stable fluctuations are reminiscent of some recent results on the BBM: such fluctuations appear for the critical Gibbs measure [37, 38], as well as for the limiting extremal process, when re-centered around a low position [41]. The result above shows that 1-stable fluctuations already appear in the much simpler context of the extremal process of the REM.

## 2.2 Change of measure in the general decorated case

In the paper, we sometimes consider the *general decorated case*. This means that we work with a limiting extremal process which is of the form

$$\sum_{i\geq 1}\sum_{k\geq 0}\delta_{\xi_i+d_{ik}}$$

where the  $(\xi_i)_{i\geq 1}$  are the atoms of a PPP $(e^{-x} dx)$  independent of  $(\sum_{k\geq 0} \delta_{d_{ik}})_{i\geq 1}$ , which are i.i.d. copies of a point process  $\mathcal{D} = \sum_{k\geq 0} \delta_{d_k}$  on  $(-\infty, 0]$  which has a.s. an atom at 0, but we do not assume that  $\mathcal{D}$  is the decoration process of the BBM. In particular, for  $\beta, \beta' > \beta_c = 1$  the

limiting partition function  $Z_d(\beta)$  is defined as in Equation (1.5) and the limiting mass at 1 of the two-temperature overlap  $Q_d(\beta, \beta')$  is defined as in Equation (1.3).

Moreover, we set

$$S_{\beta} \coloneqq \sum_{k \ge 0} \mathrm{e}^{\beta d_k} \qquad \mathrm{and} \qquad S_{\beta,i} \coloneqq \sum_{k \ge 0} \mathrm{e}^{\beta d_{ik}}, \qquad \forall \, i \ge 1.$$

For  $\beta > 1$ , assuming  $\mathbb{E}\left[S_{\beta}^{1/\beta}\right] < \infty$ , we introduce a new probability measure  $\mathbb{P}_{\beta}$  such that

• the distribution of  $\mathcal{D}$  under  $\mathbb{P}_{\beta}$  is characterized by

$$\mathbb{E}_{\beta}[F(\mathcal{D})] = \frac{\mathbb{E}\left[S_{\beta}^{1/\beta}F(\mathcal{D})\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]},$$
(2.5)

for any measurable bounded function F from the space of Radon measures on  $\mathbb{R}$  to  $\mathbb{R}$ ;

• under  $\mathbb{P}_{\beta}$ , the  $(\xi_i)_{i\geq 1}$  are still the atoms of a Poisson point process on  $\mathbb{R}$  with intensity measure  $e^{-x} dx$  independent of  $(\sum_{k\geq 0} \delta_{d_{ik}})_{i\geq 1}$ , which are i.i.d. copies of  $\mathcal{D}$  under  $\mathbb{P}_{\beta}$ .

With this definition in hand, the following fact holds.

**Lemma 2.6.** Let 
$$\beta > 1$$
. Assume  $\mathbb{E}\left[S_{\beta}^{1/\beta}\right] < \infty$ . Then, with  $c_{\beta} \coloneqq \log \mathbb{E}\left[S_{\beta}^{1/\beta}\right]$ ,  
 $\left(\sum_{i\geq 0} \delta_{\xi_{i}+\frac{1}{\beta}\log S_{\beta,i}}, \left(\sum_{k\geq 0} \delta_{d_{ik}}\right)_{i\geq 1}\right)$  under  $\mathbb{P} \stackrel{(\mathrm{d})}{=} \left(\sum_{i\geq 0} \delta_{\xi_{i}+c_{\beta}}, \left(\sum_{k\geq 0} \delta_{d_{ik}}\right)_{i\geq 1}\right)$  under  $\mathbb{P}_{\beta}$ .

*Proof.* This can be obtained via a direct Laplace transform calculation, or as a consequence of [43, Lemma 2.1] applied with  $u_i = e^{\beta \xi_i}$ ,  $m = 1/\beta$ ,  $X_i = S_{\beta,i}$  and  $Y_i = \sum_{k \ge 0} \delta_{d_{ik}}$ .

As a consequence, we get the following simple distribution for the decorated partition function

$$Z_d(\beta) \coloneqq \sum_{i \ge 1} \sum_{k \ge 0} e^{\beta(\xi_i + d_{ik})} = \sum_{i \ge 1} e^{\beta\xi_i} S_{\beta,i}.$$

However, note that one cannot relate the joint distribution of  $(Z_d(\beta), Z_d(\beta'))$  with the one of  $(Z(\beta), Z(\beta'))$  in such a way (otherwise the two-temperature overlap distribution would not depend on the decoration).

**Corollary 2.7.** Let  $\beta > 1$ . Assume  $\mathbb{E}\left[S_{\beta}^{1/\beta}\right] < \infty$ . Then, under  $\mathbb{P}$ ,  $Z_d(\beta) \stackrel{\text{(d)}}{=} \mathbb{E}\left[S_{\beta}^{1/\beta}\right]^{\beta} Z(\beta).$ 

*Proof.* By Lemma 2.6,  $Z_d(\beta)$  under  $\mathbb{P}$  has the same distribution as  $e^{\beta c_\beta} Z(\beta)$  under  $\mathbb{P}_\beta$ . But  $Z(\beta)$  has the same law under  $\mathbb{P}$  and under  $\mathbb{P}_\beta$ , so the result follows.

Remark 2.8. When  $\mathbb{E}[S_{\beta}^{1/\beta}] = \infty$ , one can show that  $Z_d(\beta) = \infty$  a.s. This follows for example from the previous result applied to  $S_{\beta} \wedge M$  and then letting  $M \to \infty$ . Therefore, the assumption that  $\mathbb{E}[S_{\beta}^{1/\beta}] < \infty$  for any  $\beta > 1$  is a very minimalist one when working with the decorated partition function.

#### 2.3 The decoration of the BBM

A description of the decoration. In this section, we recall some results on the law of the decoration point process

$$\mathcal{D} = \sum_{k \ge 0} \delta_{d_k} \,,$$

appearing in the limit of the extremal process of the BBM. As mentioned in the introduction (see Equation (1.1)), convergence of this extremal process has been established in [1] and [3], which give two different descriptions of the law of  $\mathcal{D}$ . However, we use here a third description obtained in [21, Lemma 5.1], as well as several other results shown in this paper and its sequel [22]. Note that the authors work with a BBM with branching rate 1 (instead of 1/2 for us here), but that both processes have the same distribution up to a time-space scaling. In particular, one can check that the decoration process  $\mathcal{C}$  appearing in this case (branching rate 1) can be related to  $\mathcal{D}$  as follows:

$$\mathcal{C} = \sum_{k \ge 0} \delta_{d_k/\sqrt{2}} \,.$$

For the sake of clarity, we give below the description of the law of C (instead of D) so that we can work exactly in the same setting as [21] when we adapt some of their proofs.

We consider a BBM with a spine defined under a new probability measure  $\mathbb{P}$  as follows. It starts with one particle at 0 at time 0 which is part of the spine. Particles along the spine branch at rate 2 and move according to a standard Brownian motion. When they branch into two particles, one of them, chosen uniformly at random, is part of the spine and the other one starts a standard BBM with branching rate 1. We denote by  $L_t$  the set of particles alive at time t, by  $X_t$  the particle at time t which is part of the spine, and by  $h_t(x)$  for  $x \in L_t$  the position of particle x at time t.

Moreover, we set

$$m_t \coloneqq \sqrt{2}t - \frac{3}{2\sqrt{2}}\log t$$

which is the position of the maximum of the BBM (with branching rate 1) at time t up to O(1) fluctuations, and, for  $0 < r \le t$  and  $x \in L_t$ ,

$$\mathcal{C}_{t,r}(x) \coloneqq \sum_{y \in L_t : d(x,y) < r} \delta_{h_t(y) - h_t(x)}$$

where  $d(x, y) := \inf\{s \ge 0 : x \text{ and } y \text{ share a common ancestor in } L_{t-s}\}$ . For brevity, we introduce the new probability measure

$$\widetilde{\mathbb{P}}_t(\cdot) \coloneqq \widetilde{\mathbb{P}}\bigg( \cdot \ \left| \ h_t(X_t) = \max_{x \in L_t} h_t(x) = m_t \bigg),$$
(2.6)

as well as

$$\mathcal{C}_{t,r_t}^* \coloneqq \mathcal{C}_{t,r_t}(X_t).$$

Then, for any positive function  $t \mapsto r_t$  such that both  $r_t$  and  $t - r_t$  tend to  $\infty$  as  $t \to \infty$ , [21, Lemma 5.1] with u = 0 establishes that, for the vague convergence on the set of Radon measures on  $\mathbb{R}$ ,

$$\mathcal{C}_{t,r_t}^* \text{ under } \widetilde{\mathbb{P}}_t \xrightarrow[t \to \infty]{(d)} \mathcal{C} \text{ under } \mathbb{P}.$$
(2.7)

In their papers [21, 22], Cortines, Hartung and Louidor develop tools to study  $C_{t,r_t}^*$  under  $\mathbb{P}_t$  and therefore obtain results on the distribution of C.

Level sets of the decoration. Some of the main results in [21] concern the level sets of the decoration point process itself. Recall that C is supported on  $(-\infty, 0]$  and has a.s. an atom at 0. In [21, Proposition 1.5], they prove the existence of constants  $C_{\star}, C > 0$  such that

$$\mathbb{E}\Big[\mathcal{D}([-\sqrt{2}x,0])\Big] = \mathbb{E}[\mathcal{C}([-x,0])] \sim C_{\star} e^{\sqrt{2}x}, \quad \text{as } x \to \infty, \quad (2.8)$$

and, for any  $x \ge 0$ ,

$$\mathbb{E}\Big[\mathcal{D}([-\sqrt{2}x,0])^2\Big] = \mathbb{E}\Big[\mathcal{C}([-x,0])^2\Big] \le C(x+1)\,\mathrm{e}^{2\sqrt{2}x}.$$
(2.9)

Note also that, as a consequence of Equation (2.8), there exists C > 0 such that, for any  $x \ge 0$ ,

$$\mathbb{E}\Big[\mathcal{D}([-\sqrt{2}x,0])\Big] = \mathbb{E}[\mathcal{C}([-x,0])] \le C e^{\sqrt{2}x}.$$
(2.10)

As a consequence, Cortines, Hartung and Louidor deduce a law of large numbers for large level sets of the whole limiting extremal process. This law of large numbers, as well as the 1-stable fluctuations appearing at the next order, have been subsequently obtained via PDE techniques by Mytnik, Roquejoffre and Ryzhik [41].

Remark 2.9. The second moment bound, see Equation (2.9), is actually of the right order, as proved by Cortines, Hartung and Louidor in a second paper [22, Proposition 1.1], which means that the first and second moments of  $\mathcal{C}([-x, 0])$  are dominated by an unlikely event, as  $x \to \infty$ . This event already appeared in the proof of Equation (2.8) in [21], when working with  $\mathcal{C}^*_{t,r_t}([-x, 0])$  under  $\widetilde{\mathbb{P}}_t$ , before taking the limit  $t \to \infty$ . It consists in the fact that the spine is going sufficiently high at a time t - s with s of order  $x^2$ , more precisely, it can be defined, for some small  $\eta > 0$  and large M > 0, as

$$\left\{\max_{s\in[\eta x^2,\eta^{-1}x^2]}(h_{t-s}(X_{t-s})-m_t+m_s)\in[-M,M]\right\},$$
(2.11)

or in the same way but with  $[-M, \infty)$  instead of  $[-M, M]^4$ . This event has a probability of order 1/x and, on this event,  $\mathcal{C}^*_{t,r_t}([-x, 0])$  is typically of order  $xe^{\sqrt{2}x}$ . Moreover, it is the dominating event in the first and second moments of  $\mathcal{C}^*_{t,r_t}([-x, 0])$  and this implies, up to multiplicative constants, that

$$\mathbb{E}\Big[\mathcal{C}^*_{t,r_t}([-x,0])\Big] \simeq \mathrm{e}^{\sqrt{2}x} \quad \text{and} \quad \mathbb{E}\Big[\mathcal{C}^*_{t,r_t}([-x,0])^2\Big] \simeq x \mathrm{e}^{2\sqrt{2}x}.$$

in accordance with Equations (2.8) and (2.9). See [22, Section 1.3] for a more detailed version of this heuristic picture, which plays a crucial role in the arguments used in this paper for the proofs of Theorems 1.2 and 1.5. We also mention here that the question of the typical size of  $\mathcal{C}([-x, 0])$  has been investigated in the physics literature [40, 36], where it is conjectured that it differs from  $e^{\sqrt{2}x}$  by a stretched-exponentially small factor  $e^{-cx^{2/3}}$ .

From these estimates on level sets of the decoration, we can deduce a first moment estimate and a second moment bound for  $S_{\beta} = \sum_{k\geq 0} e^{\beta d_k}$  stated in the following proposition.

**Proposition 2.10.** Let  $C_{\star} > 0$  be the constant appearing in Equation (2.8). As  $\beta \downarrow 1$ , we have

$$\mathbb{E}[S_{\beta}] \sim \frac{C_{\star}}{\beta - 1},\tag{2.12}$$

and there exists a constant C > 0 such that, for any  $\beta \in (1, 2]$ ,

$$\mathbb{E}\left[S_{\beta}^{2}\right] \leq \frac{C}{(\beta-1)^{3}}.$$
(2.13)

<sup>&</sup>lt;sup>4</sup>To see that both choices are roughly equivalent, note that, if  $h_{t-s}(X_{t-s}) \simeq m_t - m_s + y$ , then the particles branching from the spine around time s typically (under  $\widetilde{\mathbb{P}}$ ) have descendants at time t close to  $m_t + y$ . Hence, the conditioning in the definition of  $\widetilde{\mathbb{P}}_t$  implies that  $h_{t-s}(X_{t-s})$  cannot be much larger than  $m_t - m_s$ .

*Proof.* We write  $S_{\beta} = \int_0^{\infty} \mathcal{D}([-x,0])\beta e^{-\beta x} dx$ . Therefore, by Fubini's theorem and Equation (2.8),

$$\mathbb{E}[S_{\beta}] = \int_0^\infty (C_{\star} \mathrm{e}^x + o(\mathrm{e}^x))\beta \mathrm{e}^{-\beta x} \mathrm{d}x \underset{\beta \downarrow 1}{\sim} \frac{C_{\star}}{\beta - 1}.$$

Similarly, applying Fubini's theorem and then Cauchy-Schwarz inequality together with Equation (2.9) yields

$$\begin{split} \mathbb{E}\Big[S_{\beta}^{2}\Big] &= \int_{0}^{\infty} \int_{0}^{\infty} \mathbb{E}[\mathcal{D}([-x,0])\mathcal{D}([-y,0])] \,\beta^{2} \mathrm{e}^{-\beta(x+y)} \mathrm{d}x \mathrm{d}y \\ &\leq C \int_{0}^{\infty} \int_{0}^{\infty} \sqrt{(x+1)(y+1)} \,\mathrm{e}^{x+y} \,\beta^{2} \mathrm{e}^{-\beta(x+y)} \mathrm{d}x \mathrm{d}y \\ &= C \Big(\int_{0}^{\infty} \sqrt{x+1} \,\mathrm{e}^{x} \,\beta \mathrm{e}^{-\beta x} \mathrm{d}x\Big)^{2} \leq \frac{C}{(\beta-1)^{3}}, \end{split}$$

which concludes the proof.

Remark 2.11. As for the level sets of the decoration C (see Remark 2.9), the moments of  $S_{\beta}$  are governed by a rare event when  $\beta \downarrow 1$ . Heuristically, everything behaves as if  $S_{\beta}$  was of order 1 except on an event of small probability  $(\beta - 1)$  on which it is of order  $1/(\beta - 1)^2$ . This event can be described when working on  $C_{t,r_t}^*$  (before taking the limit  $t \to \infty$  in the description of the decoration (2.7)) and is given by the event in Equation (2.11) with  $x = 1/(\beta - 1)$ . This heuristic picture is consistent with Proposition 2.10 and also with the further results in Lemma 3.12 and Corollary 3.13.

**Corollary 2.12.** Let  $C_* > 0$  be the constant appearing in Equation (2.8). As  $\beta \downarrow 1$ , we have

$$\mathbb{E}\left[S_{\beta}^{1/\beta}\right] \sim \mathbb{E}\left[S_{\beta}^{2-1/\beta}\right] \sim \frac{C_{\star}}{\beta-1}.$$

*Proof.* This is a consequence of the following inequalities: for any real random variables  $X \ge 1$  and  $Y \ge 0$  and any  $\varepsilon \in (0, 1)$ ,

$$\mathbb{E}[Y]^{1+\varepsilon}\mathbb{E}[XY]^{-\varepsilon} \le \mathbb{E}[X^{-\varepsilon}Y] \le \mathbb{E}[Y] \quad \text{and} \quad \mathbb{E}[Y] \le \mathbb{E}[X^{\varepsilon}Y] \le \mathbb{E}[Y]^{1-\varepsilon}\mathbb{E}[XY]^{\varepsilon}, \quad (2.14)$$

which follows on one side simply from the fact that  $X \ge 1$  and on the other side from Hölder's inequality. These inequalities with  $X = Y = S_{\beta}$  and  $\varepsilon = 1 - 1/\beta$  together with Proposition 2.10 yield the desired result.

To conclude this section we state two new results on level sets of the decoration. The first one is a bound on cross-moments of level sets, which can be of independent interest. Its proof is postponed to Section A.3.

**Proposition 2.13.** There exists C > 0 such that, for any  $v \ge v' \ge 0$ ,

$$\mathbb{E}[\mathcal{C}([-v,0])\mathcal{C}([-v',0])] \le C(v'+1)e^{\sqrt{2}(v+v')}.$$

The last result of this section is a uniform bound on the first and second moment of the level sets at finite t, which follows from the proofs of Equations (2.8) and (2.9) in [21], as explained in Section A.1.

**Lemma 2.14.** There exists C > 0 such that for any  $v \ge 0$  and  $t \ge 1$ ,

$$\widetilde{\mathbb{E}}_t \left[ \mathcal{C}^*_{t,r_t}([-v,0]) \right] \le C \operatorname{e}^{\sqrt{2}v},$$
$$\widetilde{\mathbb{E}}_t \left[ \mathcal{C}^*_{t,r_t}([-v,0])^2 \right] \le C(v+1) \operatorname{e}^{2\sqrt{2}v}.$$

# **3** Overlap distribution at two temperatures

### 3.1 Results in the general decorated case

We study here the influence of the decorations on the behavior of  $F_d(\beta, \beta') = \mathbb{E}[Q_d(\beta, \beta')]$ , when  $\beta' > \beta_c = 1$  is fixed and  $\beta \downarrow 1$  and show that they can change the leading order drastically. We work in the general decorated case (see Section 2.2) in order to highlight the fact that the BBM has a critical behavior in some sense to be made precise below.

We assume that

$$\mathbb{E}\left[S_{\beta}^{1/\beta}\right] < \infty, \qquad \forall \beta > 1.$$
(3.1)

Recall from Corollary 2.7 and Remark 2.8 that it is a necessary and sufficient condition to have  $Z_d(\beta) < \infty$  almost surely. In particular, this ensures that  $Q_d(\beta, \beta')$ , introduced in Equation (1.3) and which can be written as

$$Q_d(\beta,\beta') = \frac{\sum_i e^{(\beta+\beta')\xi_i} S_{\beta,i} S_{\beta',i}}{(\sum_i e^{\beta\xi_i} S_{\beta,i}) (\sum_i e^{\beta'\xi_i} S_{\beta',i})},$$
(3.2)

is well-defined, and henceforth  $F_d(\beta, \beta')$  as well.

In the following results, assumptions are stated in terms of the following random variables

$$T_{\beta} \coloneqq \frac{S_{\beta}^{1/\beta}}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]}, \qquad \forall \beta > 1.$$

Note that  $\mathbb{E}[T_{\beta}] = 1$ . The fact that the assumptions of the following results are stated in terms of  $T_{\beta}$  shows that the behavior of  $F_d(\beta, \beta')$  as  $\beta \downarrow 1$  is not governed by the growth rate of  $\mathbb{E}[S_{\beta}^{1/\beta}]$ as  $\beta \downarrow 1$ , but rather by how much  $S_{\beta}^{1/\beta}$  is fluctuating around its mean. In the forthcoming assumptions, these fluctuations are controlled in terms of how fast  $\mathbb{E}[T_{\beta}^{1+\varepsilon}]$  explodes or  $\mathbb{E}[T_{\beta}^{1-\varepsilon}]$ vanishes.

The first result concerns the case of weakly fluctuating decorations in which  $\mathbb{E}[T_{\beta}^{1+\varepsilon}]$  does not explode too fast.

**Theorem 3.1.** Assume (3.1) and the following

- (i) There exists  $\alpha \in [0,1]$  such that  $\mathbb{E}[T_{\beta} \log T_{\beta}] = (\alpha + o(1)) \log \frac{1}{\beta 1}$ , as  $\beta \downarrow 1$ .
- (ii) There exists  $\varepsilon > 0$  such that  $\mathbb{E}[T_{\beta}^{1+\varepsilon}] = O((\beta 1)^{-\varepsilon})$ , as  $\beta \downarrow 1$ .
- (iii) For any  $\beta' > 1$ ,  $\mathbb{E}[T_{\beta} \log T_{\beta'}] = O(1)$ , as  $\beta \downarrow 1$ .

Then, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,

$$F_d(\beta, \beta') = (1 - \alpha + o(1))(\beta - 1)\log \frac{1}{\beta - 1}.$$

If moreover, we assume

(i') There exists  $\alpha \in [0,1]$  such that  $\mathbb{E}[T_{\beta} \log T_{\beta}] = \alpha \log \frac{1}{\beta-1} + O(1)$ , as  $\beta \downarrow 1$ ,

then, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,

$$F_d(\beta, \beta') = (1 - \alpha)(\beta - 1)\log \frac{1}{\beta - 1} + O(\beta - 1).$$

**Corollary 3.2.** Assume there exists  $\varepsilon > 0$  such that  $\mathbb{E}[S_{\beta}^{1+\varepsilon}] = O(1)$ , as  $\beta \downarrow 1$ . Then, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,

$$F_d(\beta, \beta') = (\beta - 1) \log \frac{1}{\beta - 1} + O(\beta - 1).$$

*Proof.* Noting that  $T_{\beta} \leq S_{\beta}^{1/\beta} \leq S_{\beta}$ ,  $\log T_{\beta} \leq C T_{\beta}^{\varepsilon}$  and  $S_{\beta'} \leq S_{\beta}$  as soon as  $\beta \leq \beta'$ , Assumptions (i')-(ii)-(iii) of Theorem 3.1 are satisfied with  $\alpha = 0$  and the result follows.

In the case  $\alpha = 1$ , which we call the critical case, the previous theorem yields an upper bound for  $F_d(\beta, \beta')$ , but does not identify the main order. The following result proves a lower bound.

**Theorem 3.3.** Assume (3.1) and that there exists  $\varepsilon > 0$  such that  $\mathbb{E}[T_{\beta}^{1+\varepsilon}] = O((\beta - 1)^{-\varepsilon})$ , as  $\beta \downarrow 1$ . Then, for any  $\beta' > 1$ ,

$$\liminf_{\beta \downarrow 1} \frac{F_d(\beta, \beta')}{\beta - 1} > 0.$$

Finally, we consider strongly fluctuating cases where  $F_d(\beta, \beta')$  can vanish faster than  $(\beta - 1)$ , as  $\beta \downarrow 1$ . In that case, the function  $\beta \mapsto F_d(\beta, \beta')$  has a zero derivative at  $\beta = 1$ .

**Theorem 3.4.** Assume (3.1) and that there exists  $\theta \in (0,1)$  and  $\eta \ge 0$  such that

$$\mathbb{E}\Big[T^{1-\theta}_{\beta}T^{\theta}_{\beta'}\Big] = O\Big((\beta-1)^{\theta+\eta}\Big)$$

Then, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,

$$F_d(\beta, \beta') = O\left((\beta - 1)^{1+\eta}\right).$$

In the three theorems above, assumptions were stated exactly as we need them in the proof. In the following corollary, we work instead under a more readable set of assumptions without seeking any optimality. Recall that, for  $f: (1, \infty) \to \mathbb{R}$  and  $g: (1, \infty) \to \mathbb{R}^*_+$ , we write  $f(\beta) \asymp g(\beta)$  if  $0 < \liminf_{\beta \downarrow 1} f(\beta)/g(\beta) \le \limsup_{\beta \downarrow 1} f(\beta)/g(\beta) < \infty$ .

**Corollary 3.5.** Assume there exist  $\gamma_{-} < 1 < \gamma_{+}$  and a function  $\psi: (\gamma_{-}, \gamma_{+}) \rightarrow [0, \infty)$  such that, for any  $\gamma \in (\gamma_{-}, \gamma_{+})$ ,

$$\mathbb{E}\left[S_{\beta}^{\gamma}\right] \asymp (\beta - 1)^{-\psi(\gamma)}, \qquad as \ \beta \downarrow 1.$$
(3.3)

Assume moreover that  $\psi$  is differentiable at 1. Then, the following holds, with all asymptotic notation meant to hold as  $\beta \downarrow 1$ .

(i) If  $\psi'(1) < \psi(1) + 1$  and, for any  $\beta' > 1$ ,  $\mathbb{E}[S_{\beta} \log S_{\beta'}] = O(\mathbb{E}[S_{\beta}])$ , then, setting  $\alpha := \psi'(1) - \psi(1) \in [0, 1)$ , we have, for any  $\beta' > 1$ ,

$$F_d(\beta, \beta') \sim (1 - \alpha)(\beta - 1)\log \frac{1}{\beta - 1}.$$

(ii) If  $\psi'(1) = \psi(1) + 1$ ,  $\psi$  is linear on a neighborhood of 1 and, for any  $\beta' > 1$ ,  $\mathbb{E}[S_{\beta} \log S_{\beta'}] = O(\mathbb{E}[S_{\beta}])$ , then, for any  $\beta' > 1$ ,

$$F_d(\beta, \beta') \asymp (\beta - 1).$$

(iii) If 
$$\psi'(1) > \psi(1) + 1$$
 and, for any  $\gamma \in (\gamma_{-}, 1)$  and  $\beta' > 1$ ,  $\mathbb{E}[S^{\gamma}_{\beta}S^{1-\gamma}_{\beta'}] = O(\mathbb{E}[S^{\gamma}_{\beta}])$ , then  

$$\eta_{0} \coloneqq \sup_{\gamma \in (\gamma_{-}, 1)} [\psi(1)\gamma - \psi(\gamma) - 1 + \gamma] > 0,$$

and, for any  $\beta' > 1$ ,

$$F_d(\beta, \beta') \le (\beta - 1)^{1 + \eta_0 + o(1)}.$$

*Proof.* Before distinguishing cases, we first make some general remarks. Firstly, for any  $\gamma \in (\gamma_{-}, \gamma_{+})$ , we have

$$\mathbb{E}\left[S_{\beta}^{\gamma/\beta}\right] \sim \mathbb{E}\left[S_{\beta}^{\gamma}\right] \asymp (\beta - 1)^{-\psi(\gamma)},\tag{3.4}$$

as a consequence of the first part of Equation (2.14) with  $X = S_{\beta}^{\gamma_0 - \gamma}$ ,  $Y = S_{\beta}^{\gamma}$  and  $\varepsilon = \gamma(\beta - 1)/(\beta(\gamma_0 - \gamma))$  for some  $\gamma_0 \in (\gamma, \gamma_+)$ . Secondly, recalling the definition of  $\mathbb{P}_{\beta}$  in Equation (2.5), we have, for any h > 0,

$$\mathbb{E}[T_{\beta}\log S_{\beta}^{1/\beta}] = \frac{1}{h} \frac{\mathbb{E}[S_{\beta}^{1/\beta}\log S_{\beta}^{h/\beta}]}{\mathbb{E}[S_{\beta}^{1/\beta}]} = \frac{1}{h} \mathbb{E}_{\beta}[\log S_{\beta}^{h/\beta}] \le \frac{1}{h}\log \mathbb{E}_{\beta}[S_{\beta}^{h/\beta}] = \frac{1}{h}\log \frac{\mathbb{E}[S_{\beta}^{(1+h)/\beta}]}{\mathbb{E}[S_{\beta}^{1/\beta}]}.$$

Similarly, we have

$$\mathbb{E}[T_{\beta}\log S_{\beta}^{1/\beta}] = -\frac{1}{h}\mathbb{E}_{\beta}[\log S_{\beta}^{-h/\beta}] \ge -\frac{1}{h}\log \mathbb{E}_{\beta}[S_{\beta}^{-h/\beta}] = -\frac{1}{h}\log \frac{\mathbb{E}[S_{\beta}^{(1-h)/\beta}]}{\mathbb{E}[S_{\beta}^{1/\beta}]}.$$

By Assumption (3.3), we deduce the following inequalities, for any fixed h > 0 such that  $\gamma_{-} < 1 - h$  and  $1 + h < \gamma_{+}$ ,

$$\frac{\psi(1) - \psi(1-h)}{h} \log \frac{1}{\beta - 1} + O(1) \le \mathbb{E}[T_{\beta} \log S_{\beta}^{1/\beta}] \le \frac{\psi(1+h) - \psi(1)}{h} \log \frac{1}{\beta - 1} + O(1).$$
(3.5)

We now treat the different cases separately.

<u>Part (i)</u>. First note that  $\psi'(1) \geq \psi(1)$ , as a consequence of the fact that  $\mathbb{E}[S_{\beta}^{\gamma}] \geq \mathbb{E}[S_{\beta}]^{\gamma}$  for any  $\gamma > 1$ . This implies  $\alpha \geq 0$ . We now check that the assumptions of Theorem 3.1 are satisfied. First note that it follows from Equation (3.5) by letting  $h \to 0$  that  $\mathbb{E}[T_{\beta} \log S_{\beta}^{1/\beta}] = (\psi'(1) + o(1)) \log \frac{1}{\beta - 1}$ . Together with Equation (3.4), this shows Assumption (i). Then, Equation (3.4) implies, for any  $\varepsilon \in (0, \gamma_{+} - 1)$ ,

$$\mathbb{E}[T_{\beta}^{1+\varepsilon}] \asymp (\beta-1)^{-\psi(1+\varepsilon)+(1+\varepsilon)\psi(1)}, \tag{3.6}$$

so Assumption (ii) follows from  $\psi'(1) < \psi(1) + 1$ . Finally, Assumption (iii) follows from  $\mathbb{E}[S_{\beta} \log S_{\beta'}] = O(\mathbb{E}[S_{\beta}])$ , together with Equation (3.4) again. Therefore, the result follows from Theorem 3.1.

<u>Part (ii)</u>. We check again that the assumptions of Theorem 3.1 are satisfied. Since  $\psi$  is linear in a neighborhood of 1, we have  $\psi(1 \pm h) = \psi(1) \pm h\psi'(1)$  for h small enough and Equation (3.5) implies  $\mathbb{E}[T_{\beta} \log S_{\beta}^{1/\beta}] = \psi'(1) \log \frac{1}{\beta-1} + O(1)$ . Combining this with Equation (3.4), we get Assumption (i') with  $\alpha = \psi'(1) - \psi(1) = 0$ . Assumption (ii) follows from Equation (3.6) and the fact that  $\psi(1+\varepsilon) - (1+\varepsilon)\psi(1) = \varepsilon(\psi'(1)-\psi(1)) = \varepsilon$ , and Assumption (iii) is obtained as before. Hence, Theorem 3.1 implies  $F_d(\beta, \beta') = O(\beta - 1)$ . But under these assumptions Theorem 3.3 can also be applied and yield the desired lower bound. <u>Part (iii)</u>. The fact that  $\eta_0 > 0$  is a consequence of  $\psi'(1) > \psi(1) + 1$ . Now, for some fixed  $\eta \in \overline{(0,\eta_0)}$ , there exists  $\gamma \in (\gamma_-, 1)$  such that  $\psi(1)\gamma - \psi(\gamma) - 1 + \gamma \ge \eta$ . Then, with  $\theta = 1 - \gamma$ , we have

$$\mathbb{E}\Big[T_{\beta}^{1-\theta}T_{\beta'}^{\theta}\Big] = \frac{\mathbb{E}\Big[S_{\beta}^{\gamma/\beta}S_{\beta'}^{(1-\gamma)/\beta}\Big]}{\mathbb{E}\Big[S_{\beta}^{1/\beta}\Big]^{\gamma}\mathbb{E}\Big[S_{\beta'}^{1/\beta}\Big]^{1-\gamma}} \le \frac{\mathbb{E}\Big[S_{\beta}^{\gamma}S_{\beta'}^{1-\gamma}\Big]}{\mathbb{E}\Big[S_{\beta}^{1/\beta}\Big]^{\gamma}} = O\left(\frac{\mathbb{E}\Big[S_{\beta}^{\gamma}\Big]}{\mathbb{E}\Big[S_{\beta}^{1/\beta}\Big]^{\gamma}}\right),$$

using  $S_{\beta'} \ge 1$  in the inequality and then the assumption of Part (iii). Applying Equations (3.3) and (3.4), we get

$$\mathbb{E}\Big[T_{\beta}^{1-\theta}T_{\beta'}^{\theta}\Big] = O\Big((\beta-1)^{-\psi(\gamma)+\gamma\psi(1)}\Big) = O\Big((\beta-1)^{\theta+\eta}\Big),$$

by our choice of  $\gamma$ . Hence, we can apply Theorem 3.4 and get  $F_d(\beta, \beta') = O((\beta - 1)^{1+\eta})$ , which proves the result.

Example 3.6. We introduce a family of decoration processes  $(\mathcal{D}^{a,b})_{a,b>0}$  and study the behavior of  $F_d(\beta,\beta')$  for these cases. For a,b>0, let  $X_a \ge 1$  be a random variable with law defined by

$$\mathbb{P}(X_a \ge x) = x^{-a}, \text{ for } x \ge 1$$

and let  $\mathcal{D}^{a,b}$  be a point process such that, conditionally on  $X_a$ ,  $\mathcal{D}^{a,b}$  is the sum of a Dirac mass at 0 and a Poisson point process with intensity  $|x|^{b-1}e^{-x}\mathbb{1}_{x\in[-X_a,0]} dx$ . This decoration is a toy model of the decoration of the BBM when a = 1 and b = 2, see Remarks 2.9 and 2.11. We are going to check that, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,

$$F_{d}(\beta,\beta') \begin{cases} \sim (\beta-1)\log\frac{1}{\beta-1}, & \text{if } b < a, \\ \sim (1-a)(\beta-1)\log\frac{1}{\beta-1}, & \text{if } b > a \text{ and } a < 1, \\ \approx (\beta-1), & \text{if } b > a = 1, \\ = O((\beta-1)^{1+(a-1)(1-\frac{a}{b})+o(1)}), & \text{if } b > a > 1. \end{cases}$$
(3.7)

Note that one does not treat the case a = b, which do not fits in the theorems of this section. In the last case, by taking for example b = 2a large enough, one can have  $F_d(\beta, \beta')$  vanishing faster than any power of  $(\beta - 1)$ .

Let  $(d_k)_{k\geq 0}$  be the atoms of  $\mathcal{D}^{a,b}$  ranked in decreasing order (with the convention  $d_k = -\infty$  when there are no atoms left). One has

$$S_{\beta} = \sum_{k \ge 0} e^{\beta d_k} = 1 + \underbrace{\int_0^{X_a} x^{b-1} e^{-(\beta-1)x} dx}_{=:R_{\beta}} + \underbrace{\sum_{k \ge 1} e^{\beta d_k} - \int_0^{X_a} x^{b-1} e^{-(\beta-1)x} dx}_{=:V_{\beta}}.$$

One can check that  $\mathbb{E}[V_{\beta}^2]$  remains bounded when  $\beta \downarrow 1$ , so we focus on the behavior of  $R_{\beta}$ . Let us write

$$R_{\beta} = \frac{1}{(\beta - 1)^{b}} \int_{0}^{(\beta - 1)X_{a}} u^{b - 1} e^{-u} du = \frac{1}{h^{b}} G(hX_{a}),$$

where  $h \coloneqq \beta - 1$  and  $G(x) \coloneqq \int_0^x u^{b-1} e^{-u} du$ . Then, if  $\gamma > 0$ , one has

$$\mathbb{E}[R^{\gamma}_{\beta}] = \frac{1}{h^{b\gamma}} \int_{1}^{\infty} G(hx)^{\gamma} \frac{a \mathrm{d}x}{x^{a+1}} = \frac{a}{h^{b\gamma-a}} \int_{h}^{\infty} \frac{G(t)^{\gamma}}{t^{a+1}} \mathrm{d}t.$$

Using the fact that G is bounded and  $G(t) \sim t^b/b$  when  $t \to 0$ , one gets, up to constants  $C = C(a, b, \gamma)$ ,

$$\mathbb{E}[R^{\gamma}_{\beta}] \underset{\beta \downarrow 1}{\sim} \begin{cases} Ch^{a-b\gamma}, & \text{if } b\gamma > a, \\ C\log \frac{1}{h}, & \text{if } b\gamma = a, \\ C, & \text{if } b\gamma < a. \end{cases}$$

Let us write  $||Y||_{\gamma} := \mathbb{E}[|Y|^{\gamma}]^{1/\gamma}$ , for any  $\gamma > 0$ . One has

$$\left\|R_{\beta}\right\|_{\gamma} - \left\|1 + V_{\beta}\right\|_{\gamma} \le \left\|S_{\beta}\right\|_{\gamma} \le \left\|R_{\beta}\right\|_{\gamma} + \left\|1 + V_{\beta}\right\|_{\gamma},$$

which extends the previous results to  $S_{\beta}$  when  $\gamma \leq 2$  using that  $\mathbb{E}[V_{\beta}^2] = O(1)$ :

$$\mathbb{E}[S_{\beta}^{\gamma}] \underset{\beta \downarrow 1}{\simeq} \begin{cases} h^{a-b\gamma}, & \text{if } b\gamma > a, \\ \log \frac{1}{h}, & \text{if } b\gamma = a, \\ 1, & \text{if } b\gamma < a. \end{cases}$$

Note that, if b < a, then there exists  $\gamma > 1$  such that  $\mathbb{E}[S_{\beta}^{\gamma}] = O(1)$ , so the first part of (3.7) follows from Corollary 3.2.

So we now focus on the case b > a. Then, we have  $\mathbb{E}[S_{\beta}^{\gamma}] \simeq (\beta - 1)^{-\psi(\gamma)}$  with  $\psi(\gamma) = b\gamma - a$  for  $\gamma \in (a/b, 2]$ . We want to apply Corollary 3.5 noting that  $\psi'(1) - \psi(1) = a$ . For this we first prove that, for any  $\beta' > 1$  and  $\gamma \in (0, 1]$ , as  $\beta \downarrow 1$ ,

$$\mathbb{E}\Big[S^{\gamma}_{\beta}S_{\beta'}\Big] = O\Big(\mathbb{E}\Big[S^{\gamma}_{\beta}\Big]\Big). \tag{3.8}$$

To see this, we first use subadditivity of  $x \mapsto x^{\gamma}$  to get

$$\mathbb{E}\Big[S_{\beta}^{\gamma}S_{\beta'}\Big] \leq \mathbb{E}\Big[\Big(1+R_{\beta}^{\gamma}+|V_{\beta}|^{\gamma}\Big)\Big(1+R_{\beta'}+|V_{\beta'}|\Big)\Big] \leq C\Big(1+\mathbb{E}\Big[R_{\beta}^{\gamma}\Big]+\mathbb{E}\Big[R_{\beta}^{\gamma}\cdot|V_{\beta'}|\Big]\Big),$$

where  $C = C(a, b, \beta')$  and we used  $\mathbb{E}[V_{\beta}^2] \leq C$  and  $R_{\beta'} \leq C$ . Then, we bound

$$\mathbb{E}\Big[R_{\beta}^{\gamma} \cdot |V_{\beta'}|\Big] = \mathbb{E}\Big[R_{\beta}^{\gamma} \mathbb{E}\big[|V_{\beta'}||X_a\big]\Big] \le \mathbb{E}\Big[R_{\beta}^{\gamma} \mathbb{E}\big[V_{\beta'}^2\Big|X_a\big]^{1/2}\Big] \le C\mathbb{E}\Big[R_{\beta}^{\gamma}\Big].$$

Since  $\mathbb{E}[R_{\beta}^{\gamma}] \simeq \mathbb{E}[S_{\beta}^{\gamma}]$ , this implies Equation (3.8). In particular, this proves  $\mathbb{E}[S_{\beta} \log S_{\beta'}] \le \mathbb{E}[S_{\beta}S_{\beta'}] = O(\mathbb{E}[S_{\beta}])$ , so we can apply Corollary 3.5.(i) if a < 1 and Corollary 3.5.(ii) if a = 1 to get the second and third parts of Equation (3.7). On the other hand, Equation (3.8) implies  $\mathbb{E}[S_{\beta}^{\gamma}S_{\beta'}^{1-\gamma}] = O(\mathbb{E}[S_{\beta}^{\gamma}])$  for  $\gamma \in (0, 1]$ , so we can apply Corollary 3.5.(i) if a > 1, with

$$\eta_0 = \sup_{\gamma \in (a/b,1)} [(b-a)\gamma - (b\gamma - a) - 1 + \gamma] = (a-1)\left(1 - \frac{a}{b}\right),$$

which yields the fourth part of Equation (3.7).

## 3.2 Some tools for the proofs

The starting point for the proof of the results stated in the previous section is the following expression for  $F_d(\beta, \beta')$ .

**Lemma 3.7.** Let  $\beta, \beta' > 1$ . Define

$$R = R(\beta, \beta') \coloneqq \left(\frac{Z_d(\beta)}{S_\beta}\right)^{1/\beta} \left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'}.$$

Then, we have

$$F_d(\beta,\beta') = \mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'} \int_0^\infty \frac{\mathrm{d}x}{(1+(Rx)^\beta)(1+x^{\beta'})}\right].$$

*Proof.* Starting from Equation (3.2) and recalling that  $\eta_k = e^{-\xi_k}$ , we get

$$\begin{split} F(\beta,\beta') &= \mathbb{E}\left[\sum_{k\geq 1} \eta_k^{-(\beta+\beta')} S_{\beta,k} S_{\beta',k} \frac{1}{\left(\eta_k^{-\beta} S_{\beta,k} + \sum_{j\neq k} \eta_j^{-\beta} S_{\beta,j}\right) \left(\eta_k^{-\beta'} S_{\beta',k} + \sum_{j\neq k} \eta_j^{-\beta'} S_{\beta',j}\right)}\right] \\ &= \int_{(\mathbb{R}_+)^3} \mathbb{E}\left[x^{-(\beta+\beta')} ss' \frac{1}{(x^{-\beta}s + Z_d(\beta))(x^{-\beta'}s' + Z_d(\beta'))}\right] \mathrm{d}x \,\mathrm{d}\nu(s,s'), \end{split}$$

where  $\nu$  denotes the law of  $(S_{\beta}, S_{\beta'})$  and one applied Palm formula (see Proposition 3.8 stated below) to the Poisson point process  $\sum_{i} \delta_{(\eta_i, S_{\beta,i}, S_{\beta',i})}$  on  $(\mathbb{R}_+)^3$ , which has intensity  $dx \otimes d\nu(s, s')$ . Using Fubini's theorem together with the fact that  $(S_{\beta}, S_{\beta'})$  is independent of  $(Z_d(\beta), Z_d(\beta'))$ yields

$$F_d(\beta,\beta') = \mathbb{E}\left[\int_0^\infty \frac{\mathrm{d}x}{(1+x^\beta Z_d(\beta)/S_\beta)(1+x^{\beta'}Z_d(\beta')/S_{\beta'})}\right].$$

Then the result follows from an obvious change of variable.

**Proposition 3.8** (Palm formula, see Theorem 4.1 in [35]). Let  $\Pi$  be a  $PPP(\mu)$  where  $\mu$  is a non-zero  $\sigma$ -finite positive measure on  $\mathbb{R}$ . Let  $\mathcal{M}$  denote the set of Radon measures on  $\mathbb{R}$  and  $f: \mathbb{R} \times \mathcal{M} \to \mathbb{R}$  be a positive mesurable function. Then, we have

$$\mathbb{E}\left[\sum_{X\in\Pi}f(X,\Pi\setminus\{X\})\right] = \int_{\mathbb{R}}\mathbb{E}[f(x,\Pi)]\mu(\mathrm{d}x)\,.$$

We now study the integral appearing in Lemma 3.7 in a deterministic fashion.

**Lemma 3.9.** Let  $\beta' \ge \beta > 1$ . For  $r \ge 0$ , we set

$$I(r) \coloneqq \int_0^\infty \frac{\mathrm{d}x}{(1 + (rx)^\beta)(1 + x^{\beta'})}$$

- (i) For any  $r \ge 0$ ,  $0 \le I(r) \le C(\beta')$ , where  $C(\beta')$  denotes a constant depending only on  $\beta'$ .
- (ii) For any  $r \geq 1$ ,

$$\left|I(r) - \frac{\log r}{r}\right| \le \frac{4}{r} + (\beta - 1)\frac{\log^2 r}{2r}.$$

(iii) For any  $\delta \in (0,1)$  there exists  $C(\beta', \delta)$ , depending only on  $\beta'$  and  $\delta$  such that, for any r > 0,

$$\left|I(r) - \frac{\log r}{r}\right| \le C(\beta', \delta) \left(\frac{1}{r} + \frac{(\beta - 1)}{r^{1 - \delta}} + \frac{1}{r^{1 + \delta}}\right)$$

(iv) For any  $r \ge 0$ ,  $I(r) \ge \frac{1}{4}(1 \land \frac{1}{r})$ .

(v) For any  $\delta \in (0,1)$ , there exists  $C(\beta', \delta)$  depending only on  $\beta'$  and  $\delta$  such that, for any r > 0,  $I(r) \leq C(\beta', \delta)r^{\delta-1}$ .

*Proof.* Part (i). The fact that I(r) is nonnegative is trivial and the upper bound follows from the inequality  $I(r) \leq \int_0^\infty \frac{\mathrm{d}x}{1+x^{\beta'}}$ .

Part (ii). We first split I(r) into three pieces, keeping the central part as the main one:

$$\left| I(r) - \int_{1/r}^{1} \frac{\mathrm{d}x}{(1 + (rx)^{\beta})(1 + x^{\beta'})} \right| \le \int_{0}^{1/r} \mathrm{d}x + \int_{1}^{\infty} \frac{\mathrm{d}x}{r^{\beta} x^{\beta + \beta'}} \le \frac{2}{r},$$

using  $1/r^{\beta} \leq 1/r$  (recall that r is assumed to be larger than 1). We now focus on the integral from 1/r to 1. Using that for any u, u' > 0,

$$\left|\frac{1}{(1+u)(1+u')} - \frac{1}{u}\right| = \frac{1+u'+uu'}{u(1+u)(1+u')} \le \frac{1}{u^2} + \frac{u'}{u},$$

we get

$$\left| \int_{1/r}^{1} \frac{\mathrm{d}x}{(1+(rx)^{\beta})(1+x^{\beta'})} - \int_{1/r}^{1} \frac{\mathrm{d}x}{(rx)^{\beta}} \right| \le \int_{1/r}^{\infty} \frac{\mathrm{d}x}{(rx)^{2\beta}} + \int_{0}^{1} \frac{x^{\beta'} \,\mathrm{d}x}{(rx)^{\beta}} \le \frac{2}{r}$$

Finally, note that

$$\int_{1/r}^{1} \frac{\mathrm{d}x}{(rx)^{\beta}} = \frac{r^{\beta-1}-1}{r^{\beta}(\beta-1)} = \frac{1 - \mathrm{e}^{-(\beta-1)\log r}}{r(\beta-1)},$$

which yields Part (ii) using  $|1 - e^{-t} - t| \le t^2/2$ .

Part (iii). For  $r \ge 1$ , we apply Part (ii) and use that  $\log^2 r \le C(\delta)r^{\delta}$ . For  $r \in (0,1)$ , we write

$$\left|I(r) - \frac{\log r}{r}\right| \le I(r) + \frac{\left|\log r\right|}{r} \le C(\beta') + \frac{C(\delta)}{r^{1+\delta}}$$

using Part (i), and then, using  $1 \le 1/r$  for the first term on the right-hand side proves Part (iii).

Part (iv). For  $x \in [0, 1 \land \frac{1}{r}]$ , we have  $(1 + (rx)^{\beta})(1 + x^{\beta'}) \le 4$ . So keeping only this part of the integral yields Part (iv).

Part (v). This follows from Part (i) if  $r \in (0, 1]$ , and from Part (ii) if  $r \ge 1$ .

## **3.3** Proof of the general theorems

In this section, we prove Theorems 3.1, 3.3 and 3.4.

Proof of Theorem 3.1. Applying Lemma 3.7 and Lemma 3.9.(iii) we get, for some fixed  $\delta \in (0, 1)$  to be chosen small enough later,

$$F_d(\beta,\beta') = \mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'} \frac{\log R}{R}\right] + O\left(\mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'} \left(\frac{1}{R} + \frac{(\beta-1)}{R^{1-\delta}} + \frac{1}{R^{1+\delta}}\right)\right]\right)$$
  
=:  $E_0 + O(E_1 + E_2 + E_3).$  (3.9)

We estimate these four terms successively, in increasing order of difficulty.

<u>Term E<sub>1</sub></u>. Using the definition of R and then Corollary 2.7 together with the independence of  $S_{\beta}$  and  $Z_d(\beta)$ , we get

$$E_1 = \mathbb{E}\left[\left(\frac{S_\beta}{Z_d(\beta)}\right)^{1/\beta}\right] = \mathbb{E}\left[Z(\beta)^{-1/\beta}\right] \sim \beta - 1, \qquad (3.10)$$

by Equation (2.3).

<u>Term E\_3</u>. We use first the definition of R and the independence of  $(S_{\beta}, S_{\beta'})$  and  $(Z_d(\beta), Z_d(\beta'))$ , and then  $S_{\beta'} \geq 1$  and Cauchy–Schwarz inequality to get

$$E_{3} = \mathbb{E}\left[\frac{S_{\beta}^{(1+\delta)/\beta}}{S_{\beta'}^{\delta/\beta'}}\right] \mathbb{E}\left[\frac{Z_{d}(\beta')^{\delta/\beta'}}{Z_{d}(\beta)^{(1+\delta)/\beta}}\right] \leq \mathbb{E}\left[S_{\beta}^{(1+\delta)/\beta}\right] \mathbb{E}\left[Z_{d}(\beta')^{2\delta/\beta'}\right]^{1/2} \mathbb{E}\left[\frac{1}{Z_{d}(\beta)^{2(1+\delta)/\beta}}\right]^{1/2}$$

Using Corollary 2.7, Lemma 2.2 and  $\mathbb{E}[S_{\beta'}^{1/\beta'}] < \infty$ , we get  $\mathbb{E}[Z_d(\beta')^{2\delta/\beta'}] = O(1)$  as soon as  $\delta < 1/2$ . Therefore, by Corollary 2.7 and Equation (2.3),

$$E_3 = O\left(\mathbb{E}\left[S_{\beta}^{(1+\delta)/\beta}\right] \cdot \frac{(\beta-1)^{1+\delta}}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]^{1+\delta}}\right) = O(\beta-1), \tag{3.11}$$

by Assumption (ii) as soon as  $\delta \leq \varepsilon$ . Indeed, if Assumption (ii) holds for some  $\varepsilon > 0$  then, for any  $\delta \in (0, \varepsilon]$  it holds with  $\delta$  replacing  $\varepsilon$  by Hölder's inequality<sup>5</sup>.

<u>Term  $E_2$ </u>. We proceed as for  $E_3$ , using Cauchy–Schwarz inequality to bound the second expectation below:

$$E_2 = (\beta - 1)\mathbb{E}\Big[S_{\beta}^{(1-\delta)/\beta}S_{\beta'}^{\delta/\beta'}\Big]\mathbb{E}\Big[\frac{Z_d(\beta')^{-\delta/\beta'}}{Z_d(\beta)^{(1-\delta)/\beta}}\Big] = O\left(\mathbb{E}\Big[S_{\beta}^{(1-\delta)/\beta}S_{\beta'}^{\delta/\beta'}\Big] \cdot \frac{(\beta - 1)^{2-\delta}}{\mathbb{E}[S_{\beta}^{1/\beta}]^{1-\delta}}\right)$$

Using  $\mathbb{E}[X^{1-\delta}Y^{\delta}] \leq \mathbb{E}[X]^{1-\delta}\mathbb{E}[Y]^{\delta}$  with  $X = S_{\beta}^{1/\beta}$  and  $Y = S_{\beta'}^{1/\beta'}$  yields  $E_2 = O(\beta - 1)$ .

<u>Term  $E_0$ </u>. We split this term into three parts:

$$E_{0} = \mathbb{E}\left[\left(\frac{S_{\beta}}{Z_{d}(\beta)}\right)^{1/\beta} \left(\log\frac{Z_{d}(\beta)^{1/\beta}}{S_{\beta}^{1/\beta}} + \frac{1}{\beta'}\log S_{\beta'} - \frac{1}{\beta'}\log Z_{d}(\beta')\right)\right] =: E_{00} + E_{01} - E_{02},$$

where  $E_{00}$  is the main term. By Corollary 2.7, we get

$$E_{00} = \mathbb{E}\left[\frac{\log Z(\beta)^{1/\beta}}{Z(\beta)^{1/\beta}}\right] - \mathbb{E}\left[Z(\beta)^{-1/\beta}\right] \mathbb{E}\left[\frac{S_{\beta}^{1/\beta}}{\mathbb{E}[S_{\beta}^{1/\beta}]}\log\frac{S_{\beta}^{1/\beta}}{\mathbb{E}[S_{\beta}^{1/\beta}]}\right].$$

Then, using Equations (2.4) for the first term and Equation (2.3) together with Assumption (i) or (i') for the second one, we get

$$E_{00} = \begin{cases} (1 - \alpha + o(1))(\beta - 1)\log\frac{1}{\beta - 1} & \text{under Assumption (i),} \\ (1 - \alpha)(\beta - 1)\log\frac{1}{\beta - 1} + O(\beta - 1) & \text{under Assumption (i').} \end{cases}$$

On the other hand, we have  $E_{01} = O(\beta - 1)$  by Corollary 2.7 together with Equation (2.3) and Assumption (iii) (note that it implies  $\mathbb{E}[T_{\beta} \log S_{\beta'}] = O(1)$ ). Finally, we have

$$E_{02} = \frac{1}{\beta'} \mathbb{E} \Big[ S_{\beta}^{1/\beta} \Big] \mathbb{E} \Big[ \frac{\log Z_d(\beta')}{Z_d(\beta)^{1/\beta}} \Big] = O \Big( \mathbb{E} \Big[ \log^2 \Big( Z(\beta') \mathbb{E} \Big[ S_{\beta'}^{1/\beta'} \Big] \Big) \Big]^{1/2} \mathbb{E} \Big[ Z(\beta)^{-2/\beta} \Big]^{1/2} \Big),$$

using Cauchy–Schwarz inequality and Corollary 2.7. The first expectation on the right-hand side is a O(1) so Equation (2.3) implies  $E_{02} = O(\beta - 1)$ . This concludes the proof.

<sup>&</sup>lt;sup>5</sup>More precisely this follows from the inequality  $\mathbb{E}[T_{\beta}^{1+\delta}] \leq \mathbb{E}[T_{\beta}^{1+\varepsilon}]^{\delta/\varepsilon} \mathbb{E}[T_{\beta}]^{(\varepsilon-\delta)/\varepsilon} = \mathbb{E}[T_{\beta}^{1+\varepsilon}]^{\delta/\varepsilon}.$ 

Proof of Theorem 3.3. Applying Lemma 3.7 and Lemma 3.9.(iv), we get

$$F_d(\beta,\beta') \ge \frac{1}{4} \mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'} \left(1 \wedge \frac{1}{R}\right)\right] = \frac{1}{4} \mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'} \wedge \left(\frac{S_{\beta}}{Z_d(\beta)}\right)^{1/\beta}\right].$$
(3.12)

Moreover, for any  $a, b, \delta > 0$  and M > 1, we have

$$a \wedge b \ge \frac{(Ma) \wedge b}{M} \ge \frac{b}{M} \mathbb{1}_{b \le Ma} = \frac{1}{M} (b - b \mathbb{1}_{b > Ma}) \ge \frac{1}{M} \left( b - \frac{b^{1+\delta}}{(Ma)^{\delta}} \right).$$

Applying this to (3.12) yields, for any  $\delta > 0$  and M > 1,

$$F_d(\beta,\beta') \ge \frac{1}{4M} \left( \mathbb{E}\left[ \left( \frac{S_\beta}{Z_d(\beta)} \right)^{1/\beta} \right] - \frac{1}{M^{\delta}} \mathbb{E}\left[ \left( \frac{Z_d(\beta')}{S_{\beta'}} \right)^{\delta/\beta'} \left( \frac{S_\beta}{Z_d(\beta)} \right)^{(1+\delta)/\beta} \right] \right).$$
(3.13)

The first expectation in Equation (3.13) equals the term  $E_1$  appearing in the proof of Theorem 3.1, and hence is asymptotically equivalent to  $\beta - 1$ , see Equation (3.10). The second expectation equals the term  $E_3$  and so is a  $O(\beta - 1)$  if  $\delta \leq \frac{1}{2} \wedge \varepsilon$  with  $\varepsilon$  given by the assumption of the theorem. Choosing M large enough, the first expectation dominates and the result follows.  $\Box$ 

Proof of Theorem 3.4. Let  $\theta \in (0,1)$  be given by the assumption of the theorem. Applying Lemma 3.7 and Lemma 3.9.(iii) with  $\delta = \theta$ , we get

$$F_d(\beta,\beta') = O\left(\mathbb{E}\left[\left(\frac{S_{\beta'}}{Z_d(\beta')}\right)^{1/\beta'}\frac{1}{R^{1-\theta}}\right]\right) = O\left(\mathbb{E}\left[S_{\beta}^{(1-\theta)/\beta}S_{\beta'}^{\theta/\beta'}\right]\mathbb{E}\left[\frac{Z_d(\beta')^{-\theta/\beta'}}{Z_d(\beta)^{(1-\theta)/\beta}}\right]\right), \quad (3.14)$$

using the definition of R and the independence of  $(S_{\beta}, S_{\beta'})$  and  $(Z_d(\beta), Z_d(\beta'))$ . By Cauchy–Schwarz inequality and Corollary 2.7,

$$\mathbb{E}\left[\frac{Z_d(\beta')^{-\theta/\beta'}}{Z_d(\beta)^{(1-\theta)/\beta}}\right] \le \frac{\mathbb{E}\left[Z(\beta')^{-2\theta/\beta'}\right]^{1/2}}{\mathbb{E}\left[S_{\beta'}^{1/\beta'}\right]^{\theta}} \frac{\mathbb{E}\left[Z(\beta)^{-2(1-\theta)/\beta}\right]^{1/2}}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]^{1-\theta}} = O\left(\frac{(\beta-1)^{1-\theta}}{\mathbb{E}\left[S_{\beta'}^{1/\beta'}\right]^{\theta}\mathbb{E}\left[S_{\beta}^{1/\beta}\right]^{1-\theta}}\right),$$

by Equation (2.3). Coming back to Equation (3.14) and applying the assumption of the theorem concludes the proof.  $\hfill \Box$ 

#### 3.4 The REM case

First note that Theorem 1.1 is simply a particular case of Corollary 3.2.

*Proof of Theorem 1.1.* For the REM,  $S_{\beta} = 1$  so the result follows from Corollary 3.2.

In this subsection, we add some comments and further results concerning the REM case. Remark 3.10. It follows from Lemma 2.4 that a.s., as  $\beta \downarrow 1$ ,

$$Q(\beta, \beta') = \frac{Z(\beta + \beta')}{Z(\beta)Z(\beta')} \sim \frac{Z(1 + \beta')}{Z(\beta')}(\beta - 1),$$

proving that the function  $Q(\cdot, \beta')$  is almost surely right differentiable at 1. On the other hand, Theorem 1.1 implies that  $F(\cdot, \beta') = \mathbb{E}[Q(\cdot, \beta')]$  has an infinite right-derivative at 1. There is however a simple way to show this, without aiming for the first order of  $F(\beta, \beta')$  as  $\beta \downarrow 1$ : indeed, using Fatou's lemma,

$$\liminf_{\beta \downarrow 1} \frac{F(\beta,\beta')}{\beta-1} \geq \mathbb{E} \left[ \liminf_{\beta \downarrow 1} \frac{Q(\beta,\beta')}{\beta-1} \right] = \mathbb{E} \left[ \frac{Z(1+\beta')}{Z(\beta')} \right]$$

and it is not hard to see that this last expectation is infinite using Palm formula.

We have seen in the previous remark that, as  $\beta \downarrow 1$ ,  $Q(\beta, \beta')$  is of order  $\beta-1$ , but Theorem 1.1 shows that  $\mathbb{E}[Q(\beta, \beta')]$  is much larger. Hence, the expectation is dominated by an unlikely event, in which the highest point in the extremal process is exceptionally high, more precisely at a position of order  $\log \frac{1}{\beta-1}$ , but not higher than  $\log \frac{1}{\beta-1}$ , as proved in the following proposition.

**Proposition 3.11.** Let  $\varepsilon > 0$  and

$$E_{\beta} \coloneqq \left\{ \xi_1 \in \left[ \varepsilon \log \frac{1}{\beta - 1}, \log \frac{1}{\beta - 1} \right] \right\}.$$

Then, as  $\beta \downarrow 1$ ,

$$\mathbb{E}\Big[Q(\beta,\beta')\mathbb{1}_{E_{\beta}^{c}}\Big] \leq \varepsilon(\beta-1)\log\frac{1}{\beta-1} + O(\beta-1).$$

*Proof.* On the one hand, noting that  $Q(\beta, \beta') \leq 1$ , we have

$$\mathbb{E}\left[Q(\beta,\beta')\mathbb{1}_{\{\xi_1>\log\frac{1}{\beta-1}\}}\right] \le \mathbb{P}\left(\xi_1>\log\frac{1}{\beta-1}\right) = 1 - \exp\left\{-\int_{\log\frac{1}{\beta-1}}^{\infty} e^{-x} dx\right\} = O(\beta-1).$$

On the other hand, recalling that  $\eta_k = e^{-\xi_k}$  for all  $k \ge 1$ ,

$$\begin{split} \mathbb{E}\Big[Q(\beta,\beta')\mathbbm{1}_{\{\xi_1<\varepsilon\log\frac{1}{\beta-1}\}}\Big] &\leq \mathbb{E}\Bigg[\sum_{k\geq 1}\eta_k^{-(\beta+\beta')}\mathbbm{1}_{\{\eta_k>(1-\beta)^\varepsilon\}}\frac{1}{\left(\eta_k^{-\beta}+\sum_{j\neq k}\eta_j^{-\beta}\right)\left(\eta_k^{-\beta'}+\sum_{j\neq k}\eta_j^{-\beta'}\right)}\Bigg]\\ &= \mathbb{E}\Bigg[\int_{(1-\beta)^\varepsilon}^\infty\frac{\mathrm{d}x}{(1+x^\beta Z(\beta))(1+x^{\beta'}Z(\beta'))}\Bigg],\end{split}$$

by Palm's formula as in the proof of Lemma 3.7. This is at most

$$\mathbb{E}\left[\int_{(1-\beta)^{\varepsilon}}^{1} \frac{\mathrm{d}x}{x^{\beta}Z(\beta)} + \int_{1}^{\infty} \frac{\mathrm{d}x}{x^{\beta+\beta'}Z(\beta)Z(\beta')}\right] \leq \mathbb{E}\left[Z(\beta)^{-1}\right]\varepsilon\log\frac{1}{\beta-1} + \mathbb{E}\left[Z(\beta)^{-1}Z(\beta')^{-1}\right]$$
$$= \varepsilon(\beta-1)\log\frac{1}{\beta-1} + O(\beta-1),$$

using Cauchy–Schwarz inequality for the second term and then Lemma 2.2 and Equation (2.2) to estimate the two expectations.

#### 3.5 The BBM case

In this section, we work in the BBM case and prove Theorem 1.2 as an application of Theorems 3.1 and 3.3. In order to check the assumptions of these theorems, we first prove the following upper bounds for small moments of  $S_{\beta}$ . Only the case  $\gamma \in (1/2, 1)$  is useful for our purposes, but the other bounds follow from the same proof, so we include them as well. In particular, by the monotone convergence theorem, this shows that the random variable  $S_1 = \sum_{k\geq 0} e^{d_k}$ is in  $L^{\gamma}$  for  $\gamma \in (0, 1/2)$  and therefore is finite a.s. for the BBM.

**Lemma 3.12** (BBM case). For any  $\gamma \in (0,1)$ , there exists  $C = C(\gamma) > 0$  such that, for any  $\beta \in (1,2]$ ,

$$\mathbb{E}\Big[S_{\beta}^{\gamma}\Big] \leq \begin{cases} C(\beta-1)^{1-2\gamma}, & \text{if } \gamma \in (1/2,1), \\ C\log\frac{1}{\beta-1}, & \text{if } \gamma = 1/2, \\ C, & \text{if } \gamma \in (0,1/2). \end{cases}$$

*Proof.* We postpone most of the work to Lemma A.7. With this result in hand, setting  $f_{\beta}(x) = e^{\beta\sqrt{2}x}$ , we have  $S_{\beta} = \mathcal{C}(f_{\beta})$  and therefore it is enough to show

$$\mathbb{E}[\mathcal{C}(f_{\beta})^{\gamma}] \leq \limsup_{t \to \infty} \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}_{t,r_t}^*(f_{\beta})^{\gamma} \Big].$$
(3.15)

To see this, we consider, for K > 0, a continuous function  $\chi_K$  such that  $\mathbb{1}_{[-K,K]} \leq \chi_K \leq \mathbb{1}_{[-(K+1),K+1]}$ . On the one hand, it follows from the vague convergence stated in Equation (2.7) that  $\mathcal{C}^*_{t,r_t}(\chi_K f_\beta)$  under  $\widetilde{\mathbb{P}}_t$  converges in distribution to  $\mathcal{C}(\chi_K f_\beta)$  under  $\mathbb{P}$ . On the other hand, we have  $\mathcal{C}^*_{t,r_t}(\chi_K f_\beta)^{\gamma} \leq e^{\gamma\beta\sqrt{2}(K+1)}\mathcal{C}^*_{t,r_t}([-(K+1),0])^{\gamma}$ , which is bounded in  $L^2$  by Lemma 2.14. So  $\mathcal{C}^*_{t,r_t}(\chi_K f_\beta)^{\gamma}$  is uniformly integrable in t and we get

$$\mathbb{E}[\mathcal{C}(\chi_K f_\beta)^{\gamma}] = \lim_{t \to \infty} \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}_{t,r_t}^* (\chi_K f_\beta)^{\gamma} \Big] \le \limsup_{t \to \infty} \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}_{t,r_t}^* (f_\beta)^{\gamma} \Big].$$

Applying the monotone convergence theorem to let  $K \to \infty$  on the left-hand side yields Equation (3.15) and concludes the proof.

We note in the following corollary that combining the estimates from Proposition 2.10 and Lemma 3.12, we get the order of  $\mathbb{E}[S_{\beta}^{\gamma}]$  for any  $\gamma \in (1/2, 2]$ .

**Corollary 3.13.** For any 
$$\gamma \in (1/2, 2]$$
, there exists  $0 < c < C$  such that, for any  $\beta \in (1, 2]$ ,  
 $c(\beta - 1)^{1-2\gamma} \leq \mathbb{E}\left[S_{\beta}^{\gamma}\right] \leq C(\beta - 1)^{1-2\gamma}.$ 

*Proof.* The result holds for  $\gamma = 1$  by Proposition 2.10. The upper bound holds for  $\gamma \in (1/2, 1)$  by Lemma 3.12 and for  $\gamma = 2$  by Proposition 2.10. For  $\gamma \in (1, 2)$ , the upper bound then follows from Hölder's inequality:

$$\mathbb{E}\Big[S_{\beta}^{\gamma}\Big] \leq \mathbb{E}[S_{\beta}]^{2-\gamma} \mathbb{E}[S_{\beta}^{2}]^{\gamma-1}.$$

It remains to prove the lower bound for  $\gamma \in (1/2, 2] \setminus \{1\}$ . If  $\gamma \in (1/2, 1)$ , it follows from this consequence of Hölder's inequality

$$\mathbb{E}\left[S_{\beta}^{\gamma}\right] \geq \frac{\mathbb{E}[S_{\beta}]^{2-\gamma}}{\mathbb{E}[S_{\beta}^{2}]^{1-\gamma}},$$

together with the lower bound on  $\mathbb{E}[S_{\beta}]$  and the upper bound on  $\mathbb{E}[S_{\beta}^2]$ . If  $\gamma \in (1, 2]$ , it follows from from this consequence of Hölder's inequality, for some  $\gamma_0 \in (1/2, 1)$  ( $\gamma_0 = 3/4$  for example),

$$\mathbb{E}\left[S_{\beta}^{\gamma}\right] \geq \frac{\mathbb{E}[S_{\beta}]^{(\gamma-\gamma_{0})/(1-\gamma_{0})}}{\mathbb{E}[S_{\beta}^{\gamma_{0}}]^{(\gamma-1)/(1-\gamma_{0})}}.$$

together with the lower bound on  $\mathbb{E}[S_{\beta}]$  and the upper bound on  $\mathbb{E}[S_{\beta}^{\gamma_0}]$ .

Proof of Theorem 1.2. We prove that the decoration of the BBM satisfies the assumptions of Corollary 3.5.(ii). Assumption (3.3) has been checked in Corollary 3.13, with  $\psi(\gamma) = 2\gamma - 1$  for  $\gamma \in (1/2, 2)$ . In particular, we have  $\psi'(1) = 2 = \psi(1) + 1$  and  $\psi$  is linear on a right-neighborhood of 1. It remains to check that, for any  $\beta' > 1$ , as  $\beta \downarrow 1$ ,  $\mathbb{E}[S_{\beta} \log S_{\beta'}] = O(\mathbb{E}[S_{\beta}])$ . To prove this, note that

$$\mathbb{E}[S_{\beta}\log S_{\beta'}] \leq \mathbb{E}[S_{\beta}S_{\beta'}] = \beta\beta' \int_{0}^{\infty} \int_{0}^{\infty} \mathbb{E}[\mathcal{D}([-x,0])\mathcal{D}([-y,0])] e^{-\beta x} e^{-\beta' y} dx dy$$
$$\leq C\beta\beta' \int_{0}^{\infty} \int_{0}^{\infty} ((x \wedge y) + 1) e^{x+y} e^{-\beta x} e^{-\beta' y} dx dy,$$

where we used Proposition 2.13. Integrating first w.r.t. y and then w.r.t. x, it follows that  $\mathbb{E}[S_{\beta} \log S_{\beta'}] = O(\frac{1}{\beta-1})$ , which is enough since  $\mathbb{E}[S_{\beta}] \sim C_{\star}/(\beta-1)$  by Proposition 2.10.

# 4 Temperature susceptibility

# 4.1 A first formula for $\kappa_d$

We work here in the general decorated case (see Section 2.2) to prove a first formula for the susceptibility, which is useful for both the REM and the BBM. Recall that we consider

$$\mathbf{C}_d(\beta, \beta+h) = \frac{\operatorname{Cov}(\log Z_d(\beta), \log Z_d(\beta+h))}{\sigma(\log Z_d(\beta)) \,\sigma(\log Z_d(\beta+h))} \tag{4.1}$$

and define the susceptibility in temperature, if it exists, as the coefficient  $\kappa_d(\beta)$  such that

$$\mathbf{C}_d(\beta,\beta+h) = 1 - \kappa_d(\beta) h^2 + o(h^2), \qquad h \to 0.$$
(4.2)

,

The main technical tool to prove existence of  $\kappa_d(\beta)$  is the following lemma.

**Lemma 4.1.** Assume that  $\mathbb{E}[S_{\beta}] < \infty$ , for every  $\beta > 1$ . Then, for any  $\beta' > 1$ , the functions

$$\beta \in (1,\infty) \mapsto f_1(\beta) \coloneqq \mathbb{E}\left[\log Z_d(\beta') \log Z_d(\beta)\right]$$
$$\beta \in (1,\infty) \mapsto f_2(\beta) \coloneqq \mathbb{E}\left[\log^2 Z_d(\beta)\right],$$
$$\beta \in (1,\infty) \mapsto f_3(\beta) \coloneqq \mathbb{E}[\log Z_d(\beta)],$$

are of class  $C^2$  and their first and second derivatives are given by taking the derivative under  $\mathbb{E}$ .

*Proof.* Let us write

$$Z_d(\beta) = \mathrm{e}^{\beta \xi_1} \sum_{k \ge 1} \mathrm{e}^{\beta E_k},$$

where  $(E_k)_{k\geq 1}$  is the decreasing reordering of  $\{\xi_i + d_{i,j} - \xi_1; i, j\}$ . Note that  $E_1 = 0$  and rewrite  $f_1$  as

$$f_1(\beta) = \mathbb{E}\Big[\log Z_d(\beta')\log\Big(e^{\beta\xi_1}\sum e^{\beta E_k}\Big)\Big]$$
$$= \beta \mathbb{E}[\xi_1 \log Z_d(\beta')] + \mathbb{E}\Big[\log Z_d(\beta')\log \sum e^{\beta E_k}\Big].$$

Thus  $f_1$  is  $C^1$  with

$$f_1'(\beta) = \mathbb{E}[\xi_1 \log Z_d(\beta')] + \mathbb{E}\left[\log Z_d(\beta') \frac{\sum E_k e^{\beta E_k}}{\sum e^{\beta E_k}}\right],$$

since, for any given  $\beta_1 > 1$ , one has, for all  $\beta > \beta_1$ ,

$$\left|\log Z_d(\beta') \frac{\sum E_k e^{\beta E_k}}{\sum e^{\beta E_k}}\right| \le \left|\log Z_d(\beta')\right| \frac{\sum |E_k| e^{\beta_1 E_k}}{\sum e^{\beta_1 E_k}},$$

and those two last terms have finite second moments. The first one because  $\log Z_d(\beta')$  has the same law as  $\log Z(\beta)$  up to translation and the second one since

$$\left(\frac{\sum |E_k| e^{\beta_1 E_k}}{\sum e^{\beta_1 E_k}}\right)^2 \le \frac{\sum E_k^2 e^{\beta_1 E_k}}{\sum e^{\beta_1 E_k}} \le \sum E_k^2 e^{\beta_1 E_k} \le C \sum e^{\beta_2 E_k},$$

for some  $\beta_2 \in (1, \beta_1)$  and C > 0. For the second derivative, one has

$$f_1''(\beta) = \mathbb{E}\left[\log Z_d(\beta') \left(\frac{\sum E_k^2 e^{\beta E_k}}{\sum e^{\beta E_k}} - \left(\frac{\sum E_k e^{\beta E_k}}{\sum e^{\beta E_k}}\right)^2\right)\right],$$

since, for any given  $\beta_1 > 1$ , one has, for all  $\beta > \beta_1$ ,

$$\left|\log Z_d(\beta')\left(\frac{\sum E_k^2 e^{\beta E_k}}{\sum e^{\beta E_k}} - \left(\frac{\sum E_k e^{\beta E_k}}{\sum e^{\beta E_k}}\right)^2\right)\right| \le \left|\log Z_d(\beta')\right| \frac{\sum E_k^2 e^{\beta_1 E_k}}{\sum e^{\beta_1 E_k}},$$

which has again finite expectation with the same reasoning as above. One can then check that the first and second derivatives are the same than the ones obtained by derivating under  $\mathbb{E}$ . The computations are very similar for  $f_2$  and  $f_3$ , the only extra argument one needs for  $f_2$  is

$$\Big|\log \sum e^{\beta E_k} \Big| \frac{\sum E_k^2 e^{\beta E_k}}{\sum e^{\beta E_k}} \le \Big|\log \sum e^{\beta_1 E_k} \Big| \frac{\sum E_k^2 e^{\beta_1 E_k}}{\sum e^{\beta_1 E_k}}$$

for  $\beta > \beta_1 > 1$  using the fact  $\beta \mapsto \log \sum e^{\beta E_k}$  is decreasing and positive.

We can now deduce the main result of this section.

**Corollary 4.2.** Assume that  $\mathbb{E}[S_{\beta}] < \infty$ , for every  $\beta > 1$ . Then, for any  $\beta > 1$ ,  $\kappa_d(\beta)$  is well-defined and given by

$$\kappa_d(\beta) = \frac{1}{2} \left( \frac{\operatorname{Var}\left(\frac{Z'_d(\beta)}{Z_d(\beta)}\right)}{\operatorname{Var}\left(\log Z_d(\beta)\right)} - \left(\frac{\operatorname{Cov}\left(\log Z_d(\beta), \frac{Z'_d(\beta)}{Z_d(\beta)}\right)}{\operatorname{Var}\left(\log Z_d(\beta)\right)}\right)^2 \right).$$

*Proof.* We expand  $\mathbf{C}_d(\beta, \beta + h)$  as  $h \to 0$ , starting from Equation (4.1) and using Lemma 4.1, writing  $Z_d$  instead of  $Z_d(\beta)$  for brevity,

$$\begin{split} \mathbf{C}_{d}(\beta,\beta+h) &= \frac{\operatorname{Cov}\left(\log Z_{d},\log Z_{d} + \frac{Z'_{d}}{Z_{d}}h + \left(\frac{Z''_{d}}{Z_{d}} - \left(\frac{Z'_{d}}{Z_{d}}\right)^{2}\right)h^{2}\right)}{\sigma(\log Z_{d})\,\sigma\left(\log Z_{d} + \frac{Z'_{d}}{Z_{d}}h + \left(\frac{Z''_{d}}{Z_{d}} - \left(\frac{Z'_{d}}{Z_{d}}\right)^{2}\right)h^{2}\right)} + o\left(h^{2}\right)\\ &= 1 - \frac{1}{2}\left(\frac{\operatorname{Var}\frac{Z'_{d}}{Z_{d}}}{\operatorname{Var}\log Z_{d}} - \left(\frac{\operatorname{Cov}\left(\log Z_{d}, \frac{Z'_{d}}{Z_{d}}\right)}{\operatorname{Var}\log Z_{d}}\right)^{2}\right)h^{2} + o\left(h^{2}\right), \end{split}$$

which yields the result.

## 4.2 The REM case

In the case of the REM, the quantities appearing in the expression for the susceptibility given in Corollary 4.2 are explicit. To see this, we start by establishing the following formulae. Note that the formula for  $\mathbb{E}[\log Z(\beta)]$  already appears in the initial paper by Derrida [24]. **Lemma 4.3.** For any  $\beta > 1$ ,

$$\mathbb{E}[\log Z(\beta)] = \gamma(\beta - 1) + \beta \log \Gamma\left(1 - \frac{1}{\beta}\right), \tag{4.3}$$

$$\operatorname{Var}(\log Z(\beta)) = \frac{\pi^2}{6} (\beta^2 - 1).$$
(4.4)

*Proof.* Let  $\beta > 1$  be fixed. The Laplace transform of  $\log Z(\beta)$  has been computed in Lemma 2.2 and is finite in a neighborhood of 0. Hence, differentiating it at 0 yields the formula for  $\mathbb{E}[\log Z(\beta)]$ , and differentiating once more, we get

$$\mathbb{E}\Big[\log^2 Z(\beta)\Big] = \frac{\pi^2}{6} \Big(\beta^2 - 1\Big) + \gamma^2 (\beta - 1)^2 + \beta \log \Gamma \Big(1 - \frac{1}{\beta}\Big) \Big(2\gamma(\beta - 1) + \beta \log \Gamma \Big(1 - \frac{1}{\beta}\Big)\Big),$$
  
hich yields the formula for the variance.

which yields the formula for the variance.

The following formula is a bit more tricky to obtain.

**Lemma 4.4.** For  $\beta > 1$ , we have

$$\operatorname{Var}\left(\frac{Z'(\beta)}{Z(\beta)}\right) = \frac{\pi^2}{6} + \frac{\beta - 1}{\beta^3} \left(\frac{\Gamma''\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)} - \left(\frac{\Gamma'\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)}\right)^2\right).$$

*Proof.* To lighten notations, let us denote  $\Gamma = \Gamma(\frac{\beta-1}{\beta})$  and similarly  $\Gamma' = \Gamma'(\frac{\beta-1}{\beta}), \Gamma'' = \Gamma''(\frac{\beta-1}{\beta})$ . Differentiating once Equation (4.3) gives

$$\mathbb{E}\left[\frac{Z'(\beta)}{Z(\beta)}\right] = \log \Gamma + \frac{1}{\beta} \frac{\Gamma'}{\Gamma} + \gamma, \qquad (4.5)$$

and differentiating once more yields

$$\mathbb{E}\left[\frac{Z''(\beta)}{Z(\beta)} - \left(\frac{Z'(\beta)}{Z(\beta)}\right)^2\right] = \frac{1}{\beta^3} \left(\frac{\Gamma''}{\Gamma} - \left(\frac{\Gamma'}{\Gamma}\right)^2\right).$$
(4.6)

Therefore, we now aim at computing  $\mathbb{E}[Z''(\beta)/Z(\beta)]$ . Applying Palm formula (see Proposition 3.8), we get

$$\begin{split} \mathbb{E}\Big[\frac{Z''(\beta)}{Z(\beta)}\Big] &= \int_0^\infty \log^2(x) \,\mathbb{E}\Big[\frac{1}{1+x^\beta Z(\beta)}\Big] \,\mathrm{d}x \\ &= \int_0^\infty \mathrm{d}x \,\log^2(x) \int_0^\infty \mathrm{e}^{-t} \,\mathbb{E}\Big[\mathrm{e}^{-tx^\beta Z(\beta)}\Big] \,\mathrm{d}t \\ &= \int_0^\infty \mathrm{d}t \mathrm{e}^{-t} \int_0^\infty \log^2(x) \mathrm{e}^{-\Gamma \,t^{1/\beta}x} \,\mathrm{d}x \\ &= \frac{1}{\Gamma} \int_0^\infty \mathrm{d}t \,t^{-1/\beta} \,\mathrm{e}^{-t} \int_0^\infty \Big(\log u - \log t^{1/\beta} \,\Gamma\Big)^2 \,\mathrm{e}^{-u} \,\mathrm{d}u \\ &= \frac{1}{\Gamma} \int_0^\infty \mathrm{d}t \,t^{-1/\beta} \,\mathrm{e}^{-t} \Big(\Gamma''(1) - 2 \,t^{1/\beta} \,\Gamma'(1) \log \Gamma + t^{1/\beta} \,\log^2 \Gamma\Big) \\ &= \gamma^2 + \frac{\pi^2}{6} + 2\gamma \log \Gamma + \log^2 \Gamma + \frac{2\gamma}{\beta} \frac{\Gamma'}{\Gamma} + \frac{2}{\beta} \frac{\Gamma'}{\Gamma} \log \Gamma + \frac{1}{\beta^2} \frac{\Gamma''}{\Gamma} \,. \end{split}$$

Combining this with Equation (4.6) gives

$$\mathbb{E}\left[\left(\frac{Z'(\beta)}{Z(\beta)}\right)^2\right] = \gamma^2 + \frac{\pi^2}{6} + 2\gamma \log \Gamma + \log^2 \Gamma + \frac{2\gamma}{\beta} \frac{\Gamma'}{\Gamma} + \frac{2}{\beta} \frac{\Gamma'}{\Gamma} \log \Gamma + \frac{\beta - 1}{\beta^3} \frac{\Gamma''}{\Gamma} + \frac{1}{\beta^3} \left(\frac{\Gamma'}{\Gamma}\right)^2,$$
  
hich together with Equation (4.5) yields the result.

which together with Equation (4.5) yields the result.

We can now conclude the section by proving Theorem 1.3 concerning the susceptibility for the REM.

*Proof of Theorem 1.3.* We start from the formula given by Corollary 4.2. Differentiating Equation (4.4) shows that

$$\operatorname{Cov}\left(\log Z(\beta), \frac{Z'(\beta)}{Z(\beta)}\right) = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}\beta} \operatorname{Var}(\log Z_d(\beta)) = \frac{\pi^2}{6}\beta.$$

Together, with Equation (4.4) and Lemma 4.4, this yields

$$\kappa(\beta) = \frac{1}{2} \left( \frac{1}{\beta^2 - 1} + \frac{6}{\pi^2 \beta^3 (\beta + 1)} \left( \frac{\Gamma''\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)} - \left(\frac{\Gamma'\left(\frac{\beta - 1}{\beta}\right)}{\Gamma\left(\frac{\beta - 1}{\beta}\right)}\right)^2 \right) - \frac{\beta^2}{(\beta^2 - 1)^2} \right)$$

and using the asymptotic expansion of  $\Gamma$  at 0 and at 1 together with extra computations yield the desired asymptotic equivalents of  $\kappa(\beta)$  as  $\beta \downarrow 1$  and  $\beta \to \infty$ .

## 4.3 Comparison wih the REM in the general decorated case

In this section, we come back to the general decorated case. Our aim is to prove the following proposition which compares the temperature susceptibility  $\kappa_d(\beta)$  with the one of the REM.

**Proposition 4.5.** Assume that  $\mathbb{E}[S_{\beta}] < \infty$ , for every  $\beta > 1$ . Then, for any  $\beta > 1$ ,

$$\kappa_d(\beta) = \kappa(\beta) + \frac{3}{\pi^2 \beta(\beta+1)} \operatorname{Var}_\beta \left(\frac{1}{\beta} \log S_\beta - \frac{S'_\beta}{S_\beta}\right)$$

where  $\operatorname{Var}_{\beta}$  denotes the variance under  $\mathbb{P}_{\beta}$ , which was introduced in Subsection 2.2, and  $S'_{\beta} = \sum_{k\geq 0} d_k e^{\beta d_k}$ .

Proof. We start from the expression given by Corollary 4.2. By Corollary 2.7, we have

$$\operatorname{Var}\left(\log Z_d(\beta)\right) = \operatorname{Var}\left(\log\left(\mathbb{E}\left[S_{\beta}^{1/\beta}\right]^{\beta}Z(\beta)\right)\right) = \operatorname{Var}\left(\log Z(\beta)\right) = \frac{\pi^2}{6}(\beta^2 - 1)$$

and, by differentiating with respect to  $\beta$ ,

$$\operatorname{Cov}\left(\log Z_d(\beta), \frac{Z'_d(\beta)}{Z_d(\beta)}\right) = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}\beta} \operatorname{Var}\left(\log Z_d(\beta)\right) = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}\beta} \operatorname{Var}\left(\log Z(\beta)\right) = \operatorname{Cov}\left(\log Z(\beta), \frac{Z'(\beta)}{Z(\beta)}\right).$$

Therefore, we get

$$\kappa_d(\beta) = \kappa(\beta) + \frac{3}{\pi^2(\beta^2 - 1)} \left( \operatorname{Var}\left(\frac{Z'_d(\beta)}{Z_d(\beta)}\right) - \operatorname{Var}\left(\frac{Z'(\beta)}{Z(\beta)}\right) \right).$$
(4.7)

We shall now focus on  $\operatorname{Var}\left(\frac{Z'_d(\beta)}{Z_d(\beta)}\right)$ . Notice first that

$$Z'_d(\beta) = \sum_{i,k} (\xi_i + d_{ik}) \mathrm{e}^{\beta(\xi_i + d_{ik})}$$

Therefore, it follows from Lemma 2.6 that

$$\frac{Z'_{d}(\beta)}{Z_{d}(\beta)} \text{ under } \mathbb{P} \stackrel{\text{(d)}}{=} \frac{\sum_{i,k} \left(\xi_{i} + c_{\beta} - \frac{1}{\beta} \log S_{\beta,i} + d_{ik}\right) \mathrm{e}^{\beta(\xi_{i} - \frac{1}{\beta} \log S_{\beta,i} + d_{ik})}}{\sum_{i} \mathrm{e}^{\beta\xi_{i}}} \text{ under } \mathbb{P}_{\beta}.$$

Moreover, the quantity on the right-hand side of the last equation equals

$$c_{\beta} + \frac{\sum_{i} \xi_{i} \mathrm{e}^{\beta\xi_{i}}}{\sum_{i} \mathrm{e}^{\beta\xi_{i}}} + \frac{\sum_{i} \mathrm{e}^{\beta\xi_{i}} (S'_{\beta,i} / S_{\beta,i} - \frac{1}{\beta} \log S_{\beta,i})}{\sum_{i} \mathrm{e}^{\beta\xi_{i}}}.$$

Hence, using the law of total variance, we get

$$\operatorname{Var}\left(\frac{Z_{d}'(\beta)}{Z_{d}(\beta)}\right) = \operatorname{Var}_{\beta}\left(\mathbb{E}_{\beta}\left[\frac{\sum_{i}\xi_{i}e^{\beta\xi_{i}}}{\sum_{i}e^{\beta\xi_{i}}} + \frac{\sum_{i}e^{\beta\xi_{i}}(S_{\beta,i}'/S_{\beta,i} - \frac{1}{\beta}\log S_{\beta,i})}{\sum_{i}e^{\beta\xi_{i}}} \middle| \xi\right]\right) \\ + \mathbb{E}_{\beta}\left[\operatorname{Var}_{\beta}\left(\frac{\sum_{i}\xi_{i}e^{\beta\xi_{i}}}{\sum_{i}e^{\beta\xi_{i}}} + \frac{\sum_{i}e^{\beta\xi_{i}}(S_{\beta,i}'/S_{\beta,i} - \frac{1}{\beta}\log S_{\beta,i})}{\sum_{i}e^{\beta\xi_{i}}} \middle| \xi\right)\right].$$

Then, using the fact that the  $(S_{\beta,i}, S'_{\beta,i})_{i\geq 0}$  are i.i.d. and independent of  $\xi$ , we get

$$\operatorname{Var}\left(\frac{Z_{d}'(\beta)}{Z_{d}(\beta)}\right) = \operatorname{Var}_{\beta}\left(\frac{\sum_{i} \xi_{i} e^{\beta\xi_{i}}}{\sum_{i} e^{\beta\xi_{i}}}\right) + \mathbb{E}_{\beta}\left[\frac{\sum_{i} e^{2\beta\xi_{i}}}{(\sum_{i} e^{\beta\xi_{i}})^{2}}\right] \operatorname{Var}_{\beta}\left(\frac{S_{\beta}'}{S_{\beta}} - \frac{1}{\beta}\log S_{\beta}\right)$$
$$= \operatorname{Var}\left(\frac{Z'(\beta)}{Z(\beta)}\right) + \left(1 - \frac{1}{\beta}\right) \operatorname{Var}_{\beta}\left(\frac{S_{\beta}'}{S_{\beta}} - \frac{1}{\beta}\log S_{\beta}\right).$$

Coming back to Equation (4.7), this concludes the proof.

Example 4.6. We investigate here the behavior of the temperature susceptibility for the family of decoration processes  $(\mathcal{D}^{a,b})_{a>0,b>0}$  introduced in Example 3.6. By Proposition 4.5, one has

$$\kappa_d(\beta) = \kappa(\beta) + \frac{3}{\pi^2 \beta(\beta+1)} \operatorname{Var}_\beta \left(\frac{1}{\beta} \log S_\beta - \frac{S'_\beta}{S_\beta}\right).$$

The main contribution as  $\beta \downarrow 1$  will comes from  $\operatorname{Var}_{\beta}(S'_{\beta}/S_{\beta})$ . In order to keep the example relatively concise, we make the estimates for  $\operatorname{Var}_{\beta}(R'_{\beta}/R_{\beta})$ , where

$$R_{\beta} \coloneqq \int_0^X x^{b-1} \mathrm{e}^{-(\beta-1)x} \mathrm{d}x$$

approximates well  $S_{\beta}$  (see Example 3.6). We expect the same results to hold for  $S_{\beta}$ .

As in Example 3.6, using  $G_b(x) \coloneqq \int_0^x u^{b-1} e^{-u} du$ , one has

$$R_{\beta} = \frac{1}{h^b} G_b(hX), \qquad R'_{\beta} = -\frac{1}{h^{b+1}} G_{b+1}(hX),$$

Therefore,

$$\mathbb{E}\left[\frac{(R_{\beta}')^2}{R_{\beta}}\right] = \frac{1}{h^{b+2}} \mathbb{E}\left[\frac{G_{b+1}(hX)^2}{G_b(hX)}\right] = \frac{1}{h^{b-a+2}} \int_h^\infty \frac{G_{b+1}(t)^2}{G_b(t)} \frac{a}{t^{a+1}} \,\mathrm{d}t.$$

Then,

$$\mathbb{E}\left[\frac{(R_{\beta}')^2}{R_{\beta}}\right] \underset{\beta\downarrow1}{\sim} \begin{cases} \frac{c_{a,b}}{h^{b-a+2}}, & \text{if } b-a > -2, \\ c_{a,b}\log\frac{1}{h}, & \text{if } b-a = -2, \\ c_{a,b}, & \text{if } b-a < -2, \end{cases}$$

where  $c_{a,b} \coloneqq \int_0^{+\infty} \frac{G_{b+1}(t)^2}{G_b(t)} \frac{a}{t^{a+1}} dt$  if b-a > -2. Similarly, one finds

$$\mathbb{E}_{\beta}\left[\left(\frac{R_{\beta}'}{R_{\beta}}\right)^{2}\right] = \frac{1}{\mathbb{E}\left[R_{\beta}^{1/\beta}\right]} \cdot \mathbb{E}\left[\frac{(R_{\beta}')^{2}}{R_{\beta}^{2-1/\beta}}\right] \underset{\beta\downarrow1}{\sim} \begin{cases} \frac{c_{a,b}'}{h^{2}}, & \text{if } b-a > 0, \\ \frac{c_{a,b}'}{h^{b-a+2}}, & \text{if } b-a \in (-2,0) \\ c_{a,b}', & \text{if } b-a < -2, \end{cases}$$

where  $c'_{a,b} \coloneqq \int_0^{+\infty} \frac{G_{b+1}(t)^2}{G_b(t)} \frac{a}{t^{a+1}} \mathrm{d}t \left( \int_0^{+\infty} G_b(t) \frac{a}{t^{a+1}} \mathrm{d}t \right)^{-1}$  if b-a > 0, and

$$\mathbb{E}_{\beta} \left[ \frac{R_{\beta}'}{R_{\beta}} \right]^2 = \frac{1}{\mathbb{E} \left[ R_{\beta}^{1/\beta} \right]^2} \cdot \mathbb{E} \left[ \frac{R_{\beta}'}{R_{\beta}^{1-1/\beta}} \right]^2 \underset{\beta \downarrow 1}{\sim} \left\{ \begin{array}{ll} \frac{c_{a,b}'}{h^2}, & \text{if } b-a > 0, \\ \frac{c_{a,b}'}{h^{2(b-a)+2}}, & \text{if } b-a \in (-1,0), \\ c_{a,b}'', & \text{if } b-a < -1, \end{array} \right\}$$

with  $c_{a,b}' \coloneqq \left(\int_0^{+\infty} G_{b+1}(t) \frac{a}{t^{a+1}} dt\right)^2 \left(\int_0^{+\infty} G_b(t) \frac{a}{t^{a+1}} dt\right)^{-2}$  when b-a > -1. Now notice that by Cauchy-Schwarz inequality, one has

$$\left(\int_0^{+\infty} G_{b+1}(t) \frac{a \mathrm{d}t}{t^{a+1}}\right)^2 < \int_0^{+\infty} \frac{G_{b+1}(t)^2}{G_b(t)} \frac{a \mathrm{d}t}{t^{a+1}} \int_0^{+\infty} G_b(t) \frac{a \mathrm{d}t}{t^{a+1}},$$

thus  $c'_{a,b} > c''_{a,b}$  and one gets, up to constants,

$$\operatorname{Var}_{\beta}\left(\frac{1}{\beta}\log R_{\beta} - \frac{R_{\beta}'}{R_{\beta}}\right) \underset{\beta\downarrow1}{\sim} \begin{cases} \frac{1}{h^2}, & \text{if } b - a > 0, \\ \frac{1}{h^{b-a+2}}, & \text{if } b - a \in (-2,0), \\ 1, & \text{if } b - a < -2. \end{cases}$$

Therefore, we see mainly two behaviors for the temperature susceptibility  $\kappa_d(\beta)$  as  $\beta \downarrow 1$ . If b < a, then the contribution of the variance above is negligible and we have  $\kappa_d(\beta) \sim \kappa(\beta)$  by Proposition 4.5. On the other hand, if b > a, then the variance grows at the same speed as  $\kappa(\beta)$  as  $\beta \downarrow 1$ , so  $\kappa_d(\beta)$  is of the same order as  $\kappa(\beta)$ , but with a different multiplicative constant.

## 4.4 The BBM case

In this section, our goal is to prove Theorem 1.5 which concerns the BBM.

**Lemma 4.7.** For the decoration arising in the BBM and the constant  $C_{\star}$  appearing in Equation (2.8), we have, as  $\beta \downarrow 1$ ,

$$\mathbb{E}\Big[S'_{\beta}\Big] \sim -\frac{C_{\star}}{(\beta-1)^2} \qquad and \qquad \mathbb{E}\Big[S''_{\beta}\Big] \sim \frac{2C_{\star}}{(\beta-1)^3} \,.$$

*Proof.* This follows from Equation (2.8) in a similar way as the proof of Proposition 2.10.

Proof of Theorem 1.5. We first prove that  $\kappa_d(\beta) > \kappa(\beta)$ , when  $\beta > 1$  is fixed. By Proposition 4.5, it is sufficient to show that

$$\operatorname{Var}_{\beta}\left(\frac{1}{\beta}\log S_{\beta} - \frac{S_{\beta}'}{S_{\beta}}\right) > 0.$$

$$(4.8)$$

We proceed by contradiction and assume that this variance equals zero. Then, there exists  $c \in \mathbb{R}$  such that,  $\mathbb{P}_{\beta}$ -a.s.,

$$\frac{1}{\beta}\log S_{\beta} = \frac{S_{\beta}'}{S_{\beta}} + c, \qquad (4.9)$$

But  $\mathbb{P}$  is absolutely continuous w.r.t.  $\mathbb{P}_{\beta}$  such that Equation (4.9) holds also  $\mathbb{P}$ -a.s. On the one hand, using  $S_{\beta} = \sum_{k>0} e^{\beta d_k}$ , we have

$$S_{\beta} \log S_{\beta} > \sum_{k \ge 0} \mathrm{e}^{\beta d_k} \beta d_k = \beta S'_{\beta}$$

Thus, we necessarily have c > 0. On the other hand, fix some  $\beta' \in (1, \beta)$ . Then, there exists C > 0 such that

$$\forall x \le 0, \qquad (1 \lor |x|) e^{\beta x} \le C e^{\beta' x}. \tag{4.10}$$

Now, for some  $\varepsilon > 0$ , consider the event

$$\left\{\sum_{k\geq 1} \mathrm{e}^{\beta' d_k} \leq \varepsilon\right\}.$$

This event has positive  $\mathbb{P}$ -probability by [12, Proposition 3.4]. So, on this event intersected with the one where Equation (4.9) holds (this intersection being non-empty), we have

$$c = \frac{1}{\beta} \log S_{\beta} - \frac{S_{\beta}'}{S_{\beta}} \le \frac{1}{\beta} \log \left( 1 + \sum_{k \ge 1} e^{\beta d_k} \right) + \sum_{k \ge 1} d_k e^{\beta d_k} \le \frac{1}{\beta} \log(1 + C\varepsilon) + C\varepsilon,$$

using Equation (4.10). Letting  $\varepsilon \to 0$  shows  $c \leq 0$ , which contradicts c > 0 and concludes our proof of Equation (4.8).

We study now the regime  $\beta \downarrow 1$ . Since the asymptotics of  $\kappa(\beta)$  are given by Theorem 1.3, it remains to study the behavior of

$$\operatorname{Var}_{\beta}\left(\frac{1}{\beta}\log S_{\beta} - \frac{S_{\beta}'}{S_{\beta}}\right) = \frac{1}{\beta^{2}}\operatorname{Var}_{\beta}\left(\log S_{\beta}\right) - \frac{2}{\beta}\operatorname{Cov}_{\beta}\left(\log S_{\beta}, \frac{S_{\beta}'}{S_{\beta}}\right) + \operatorname{Var}_{\beta}\left(\frac{S_{\beta}'}{S_{\beta}}\right).$$
(4.11)

As we will see, the main term is the third one, which is of order  $(\beta - 1)^{-2}$ , while the other terms are negligible.

For the first term on the right-hand side of Equation (4.11), observe that

$$\operatorname{Var}_{\beta}\left(\log S_{\beta}\right) \leq \mathbb{E}_{\beta}\left[\log^{2} S_{\beta}\right] \leq \mathbb{E}_{\beta}\left[\log^{2}(e+S_{\beta})\right] \leq \log^{2}(e+\mathbb{E}_{\beta}[S_{\beta}]),$$

by Jensen's inequality. Then, by definition of  $\mathbb{E}_{\beta}$ , we get

$$\mathbb{E}_{\beta}[S_{\beta}] = \frac{\mathbb{E}\left[S_{\beta}^{1+1/\beta}\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \le \frac{\mathbb{E}\left[S_{\beta}^{2}\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} = O\left(\frac{1}{(\beta-1)^{2}}\right),$$

as  $\beta \downarrow 1$  by Equation (2.13) and Corollary 2.12. This proves

$$\operatorname{Var}_{\beta}\left(\log S_{\beta}\right) = O\left(\log^{2} \frac{1}{\beta - 1}\right).$$

$$(4.12)$$

We now consider the third term on the right-hand side of Equation (4.11). Using the definition of  $\mathbb{E}_{\beta}$  and then that  $S_{\beta} \geq 1$ , we get

$$\mathbb{E}_{\beta}\left[\frac{S_{\beta}'}{S_{\beta}}\right] = \frac{\mathbb{E}\left[S_{\beta}^{1/\beta-1}S_{\beta}'\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \sim -\frac{1}{\beta-1},\tag{4.13}$$

as  $\beta \downarrow 1$  by Corollary 2.12 and the fact that  $\mathbb{E}\left[S_{\beta}^{1/\beta-1}(-S_{\beta}')\right] \sim C_{\star}(\beta-1)^{-2}$  as a consequence of the first part of Equation (2.14) (with  $X = S_{\beta}$  and  $Y = -S_{\beta}'$ ) together with Lemma 4.7. On the other hand, it follows from Cauchy–Schwarz inequality that

$$\frac{S_{\beta}^{\prime 2}}{S_{\beta}} = S_{\beta} \cdot \left(\sum_{k \ge 0} d_k \frac{\mathrm{e}^{\beta d_k}}{S_{\beta}}\right)^2 \le S_{\beta} \cdot \sum_{k \ge 0} d_k^2 \frac{\mathrm{e}^{\beta d_k}}{S_{\beta}} = S_{\beta}^{\prime\prime},$$

and therefore

$$\mathbb{E}_{\beta}\left[\left(\frac{S_{\beta}'}{S_{\beta}}\right)^{2}\right] = \frac{\mathbb{E}\left[S_{\beta}^{1/\beta-2}S_{\beta}'^{2}\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \le \frac{\mathbb{E}\left[S_{\beta}'^{2}/S_{\beta}\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \le \frac{\mathbb{E}\left[S_{\beta}''\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \sim \frac{2}{(\beta-1)^{2}},\tag{4.14}$$

using Lemma 4.7 and Corollary 2.12. For the lower bound, using the first part of Equation (2.14) with  $X = S_{\beta}$  and  $Y = (S'_{\beta})^2/S_{\beta}$ , we get

$$\mathbb{E}_{\beta}\left[\left(\frac{S_{\beta}'}{S_{\beta}}\right)^{2}\right] = \frac{\mathbb{E}\left[S_{\beta}^{1/\beta-2}S_{\beta}'^{2}\right]}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]} \ge \frac{\mathbb{E}\left[S_{\beta}'^{2}/S_{\beta}\right]^{2-1/\beta}}{\mathbb{E}\left[S_{\beta}^{1/\beta}\right]\mathbb{E}\left[S_{\beta}\right]^{1-1/\beta}}.$$

Hence, applying Lemma 4.8, Proposition 2.10 and Corollary 2.12, we get

$$\liminf_{\beta \downarrow 1} (\beta - 1)^2 \mathbb{E}_{\beta} \left[ \left( \frac{S_{\beta}'}{S_{\beta}} \right)^2 \right] > 1.$$
(4.15)

Combining Equations (4.13), (4.14) and (4.15) yields

$$0 < \liminf_{\beta \downarrow 1} (\beta - 1)^2 \operatorname{Var}_{\beta} \left( \frac{S_{\beta}'}{S_{\beta}} \right) \leq \limsup_{\beta \downarrow 1} (\beta - 1)^2 \operatorname{Var}_{\beta} \left( \frac{S_{\beta}'}{S_{\beta}} \right) \leq 1.$$
(4.16)

Finally, for the second term on the right-hand side of Equation (4.11), by Cauchy–Schwarz inequality and then Equations (4.12) and (4.16), we get

$$\left|\operatorname{Cov}_{\beta}\left(\log S_{\beta}, \frac{S_{\beta}'}{S_{\beta}}\right)\right| \leq \left(\operatorname{Var}_{\beta}(\log S_{\beta}) \operatorname{Var}_{\beta}\left(\frac{S_{\beta}'}{S_{\beta}}\right)\right)^{1/2} = O\left(\frac{\log \frac{1}{\beta-1}}{\beta-1}\right),$$

which proves that this term is negligible in Equation (4.11) and concludes the proof.

Lemma 4.8. We have

$$\liminf_{\beta \downarrow 1} (\beta - 1)^3 \mathbb{E}\left[\frac{(S_{\beta}')^2}{S_{\beta}}\right] > C_{\star}.$$

*Remark* 4.9. A lower bound with a weak inequality could be easily obtained via Cauchy–Schwarz inequality:

$$\mathbb{E}\left[\frac{(S_{\beta}')^2}{S_{\beta}}\right] \geq \frac{\mathbb{E}[S_{\beta}']^2}{\mathbb{E}[S_{\beta}]} \sim \frac{C_{\star}}{(\beta-1)^3},$$

using Lemma 4.7 and Proposition 2.10. Equality at the first order in this inequality would suggest that  $S'_{\beta}$  and  $S_{\beta}$  are collinear at first order (on the events that are significant for the first moment). Therefore, the idea of the proof below is to find an event such that the first moments of  $S'_{\beta}$  and  $S_{\beta}$  given this event have a different ratio than the one for the non-conditional first moments.

Proof. Step 1: Working at finite t. Fix some  $\beta > 1$ . Setting  $f_{\beta}(x) := e^{\beta\sqrt{2}x}$  and  $\partial_{\beta}f_{\beta}(x) := \sqrt{2}xe^{\beta\sqrt{2}x}$ , note that  $S_{\beta} = \mathcal{C}(f_{\beta})$  and  $S'_{\beta} = \mathcal{C}(\partial_{\beta}f_{\beta})$ . Our first aim in this step is to show

$$\mathbb{E}\left[\frac{(S_{\beta}')^2}{S_{\beta}}\right] = \lim_{t \to \infty} \widetilde{\mathbb{E}}_t \left[\frac{(\mathcal{C}_{t,r_t}^*(\partial_{\beta}f_{\beta}))^2}{\mathcal{C}_{t,r_t}^*(f_{\beta})}\right].$$
(4.17)

For K > 0, let  $\chi_K$  denote a continuous function such that  $\mathbb{1}_{[-K,K]} \leq \chi_K \leq \mathbb{1}_{[-(K+1),K+1]}$ . It follows from the vague convergence stated in Equation (2.7) that

$$\left(\mathcal{C}_{t,r_t}^*(\chi_K f_\beta), \mathcal{C}_{t,r_t}^*(\chi_K \partial_\beta f_\beta)\right)$$
 under  $\widetilde{\mathbb{P}}_t \xrightarrow{(\mathrm{d})} (\mathcal{C}(\chi_K f_\beta), \mathcal{C}(\chi_K \partial_\beta f_\beta))$  under  $\mathbb{P}$ .

Moreover, recalling  $\mathcal{C}^*_{t,r_t}$  is supported on  $(-\infty, 0]$  under  $\widetilde{\mathbb{P}}_t$  we get

$$\left|\frac{\mathcal{C}_{t,r_t}^*(\chi_K\partial_\beta f_\beta)^2}{\mathcal{C}_{t,r_t}^*(\chi_K f_\beta)}\right| \le 2(K+1)^2 \mathcal{C}_{t,r_t}^*(\chi_K f_\beta) \le 2(K+1)^2 \mathcal{C}_{t,r_t}^*([-(K+1),0]),$$

which is bounded in  $L^2$  by Lemma 2.14. Hence,  $(\mathcal{C}^*_{t,r_t}(\chi_K \partial_\beta f_\beta)^2 / \mathcal{C}^*_{t,r_t}(\chi_K f_\beta))_t$  is uniformly integrable and we get

$$\mathbb{E}\left[\frac{\mathcal{C}(\chi_K\partial_\beta f_\beta)^2}{\mathcal{C}(\chi_K f_\beta)}\right] = \lim_{t \to \infty} \widetilde{\mathbb{E}}_t \left[\frac{\mathcal{C}_{t,r_t}^*(\chi_K\partial_\beta f_\beta)^2}{\mathcal{C}_{t,r_t}^*(\chi_K f_\beta)}\right].$$
(4.18)

We now control the differences between the expectations in Equation (4.17) and Equation (4.18) and show they are small when K is large. We have

$$\left|\frac{\mathcal{C}(\partial_{\beta}f_{\beta})^{2}}{\mathcal{C}(f_{\beta})} - \frac{\mathcal{C}(\chi_{K}\partial_{\beta}f_{\beta})^{2}}{\mathcal{C}(\chi_{K}f_{\beta})}\right| \leq \frac{|\mathcal{C}(\partial_{\beta}f_{\beta})^{2} - \mathcal{C}(\chi_{K}\partial_{\beta}f_{\beta})^{2}|}{\mathcal{C}(f_{\beta})} + \mathcal{C}(\chi_{K}\partial_{\beta}f_{\beta})^{2} \left|\frac{1}{\mathcal{C}(f_{\beta})} - \frac{1}{\mathcal{C}(\chi_{K}f_{\beta})}\right| \\ \leq 2\mathcal{C}(\partial_{\beta}f_{\beta})\mathcal{C}\left(\mathbb{1}_{(-\infty, -K]}\partial_{\beta}f_{\beta}\right) + (K+1)^{2}\mathcal{C}\left(\mathbb{1}_{(-\infty, -K]}\partial_{\beta}f_{\beta}\right),$$

$$(4.19)$$

where, for the first term, we used  $C(f_{\beta}) \geq 1$  and the fact that  $\partial_{\beta}f_{\beta}$  is of constant sign on  $(-\infty, 0]$  which is the support of C and, for the second term, we used  $|C(\chi_K \partial_{\beta}f_{\beta})|/C(f_{\beta}) \leq |C(\chi_K \partial_{\beta}f_{\beta})|/C(\chi_K f_{\beta}) \leq K+1$ . Then, writing  $\partial_{\beta}f_{\beta}(x) = \int_{-\infty}^{x} \sqrt{2}(\beta\sqrt{2}y+1)e^{\beta\sqrt{2}y} dy$  and using Fubini's theorem, we have

$$\begin{split} & \mathbb{E}\Big[\mathcal{C}(\partial_{\beta}f_{\beta})\mathcal{C}(\mathbb{1}_{(-\infty,-K]}\partial_{\beta}f_{\beta})\Big] \\ &= \int_{-\infty}^{0}\int_{-\infty}^{-K}\mathbb{E}[\mathcal{C}([y,0])\,\mathcal{C}([z,-K])]\,2(\beta\sqrt{2}z+1)\,\mathrm{e}^{\beta\sqrt{2}z}(\beta\sqrt{2}y+1)\mathrm{e}^{\beta\sqrt{2}y}\,\mathrm{d}z\,\mathrm{d}y \\ &\leq C(\beta)(K+1)^{3/2}\mathrm{e}^{-(\beta-1)\sqrt{2}K}, \end{split}$$

where we bounded the last expectation by  $(|y| + 1)^{1/2}(|z| + 1)^{1/2}e^{-\sqrt{2}(y+z)}$  using Cauchy–Schwarz inequality and Equation (2.9), and where  $C(\beta)$  denotes a constant depending only on  $\beta$  and which can change from line to line. Similarly, we have  $\mathbb{E}[\mathcal{C}(\mathbb{1}_{(-\infty,-K]}\partial_{\beta}f_{\beta})] \leq C(\beta)(K+1)e^{-(\beta-1)\sqrt{2}K}$  by using Equation (2.10). Coming back to Equation (4.19), we get

$$\left| \mathbb{E} \left[ \frac{\mathcal{C}(\partial_{\beta} f_{\beta})^{2}}{\mathcal{C}(f_{\beta})} \right] - \mathbb{E} \left[ \frac{\mathcal{C}(\chi_{K} \partial_{\beta} f_{\beta})^{2}}{\mathcal{C}(\chi_{K} f_{\beta})} \right] \right| \leq C(\beta) (K+1)^{3} \mathrm{e}^{-(\beta-1)\sqrt{2}K},$$

and the same holds true for  $C_{t,r_t}^*$  instead of C, uniformly in  $t \ge 1$ , by replacing Equations (2.9) and (2.10) by Lemma 2.14 in the proof. Therefore, letting  $K \to \infty$  in Equation (4.18) yields Equation (4.17).

<u>Step 2:</u> Using Cauchy–Schwarz inequality conditionally on a well-chosen event. We fix some parameters a > 0 and 0 < b < B. For any  $\beta > 1$ , letting  $r = r(\beta) \coloneqq a(\beta - 1)^{-2}$ , and for any t > r, we introduce the event

$$B_{r,t} = \{h_{t-r}(X_{t-r}) - m_t + m_r \in [-B\sqrt{r}, -b\sqrt{r}]\}.$$
(4.20)

Then, we use Cauchy–Schwarz inequality given  $B_{r,t}$  or given  $B_{r,t}^c$  to get

$$\widetilde{\mathbb{E}}_{t}\left[\frac{(\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta}))^{2}}{\mathcal{C}_{t,r_{t}}^{*}(f_{\beta})}\right] = \widetilde{\mathbb{E}}_{t}\left[\frac{(\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta}))^{2}}{\mathcal{C}_{t,r_{t}}^{*}(f_{\beta})} \middle| B_{r,t}\right] \widetilde{\mathbb{P}}_{t}(B_{r,t}) + \widetilde{\mathbb{E}}_{t}\left[\frac{(\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta}))^{2}}{\mathcal{C}_{t,r_{t}}^{*}(f_{\beta})} \middle| B_{r,t}\right]^{2}}\right] \widetilde{\mathbb{P}}_{t}(B_{r,t}) + \frac{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta}) \middle| B_{r,t}\right]^{2}}{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(f_{\beta}) \middle| B_{r,t}\right]^{2}} \widetilde{\mathbb{P}}_{t}(B_{r,t}) + \frac{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta}) \middle| B_{r,t}\right]^{2}}{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta})\mathbb{1}_{B_{r,t}}\right]^{2}} + \frac{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(\partial_{\beta}f_{\beta})\mathbb{1}_{B_{r,t}}\right]^{2}}{\widetilde{\mathbb{E}}_{t}\left[\mathcal{C}_{t,r_{t}}^{*}(f_{\beta})\mathbb{1}_{B_{r,t}}\right]^{2}}, \qquad (4.21)$$

which yields a lower bound for the right-hand side of Equation (4.17), that we now have to estimate.

Step 3: Estimating the expectations. Let  $\varepsilon > 0$ . Our aim in this step consists in proving that, for  $\beta > 1$  close enough to 1, there exists  $t_0 > r$  such that, for any  $t \ge t_0$ ,

$$\left| (\beta - 1) \widetilde{\mathbb{E}}_t \left[ \mathcal{C}^*_{t,r_t}(f_\beta) \mathbb{1}_{B_{r,t}} \right] - C_\star \kappa \right| \le \varepsilon, \left| (\beta - 1)^2 \widetilde{\mathbb{E}}_t \left[ \mathcal{C}^*_{t,r_t}(\partial_\beta f_\beta) \mathbb{1}_{B_{r,t}} \right] + C_\star \kappa' \right| \le \varepsilon,$$

$$(4.22)$$

where  $\kappa, \kappa'$  are constants depending only on the parameters a, b, B defined as follows

$$\kappa \coloneqq \frac{1}{2\sqrt{\pi}} \int_0^\infty \varphi_{b,B} \left( \frac{a \vee w}{|a - w|} \right) \left( \int_0^\infty u \mathrm{e}^{-u - u^2/(4w)} \,\mathrm{d}u \right) \frac{\mathrm{d}w}{w^{3/2}},$$
  

$$\kappa' \coloneqq \frac{1}{2\sqrt{\pi}} \int_0^\infty \varphi_{b,B} \left( \frac{a \vee w}{|a - w|} \right) \left( \int_0^\infty u^2 \mathrm{e}^{-u - u^2/(4w)} \,\mathrm{d}u \right) \frac{\mathrm{d}w}{w^{3/2}},$$
(4.23)

with  $\varphi_{b,B}$  defined by

$$\varphi_{b,B}(v) \coloneqq \int_{b\sqrt{v}}^{B\sqrt{v}} \sqrt{\frac{2}{\pi}} y^2 \mathrm{e}^{-y^2/2} \,\mathrm{d}y \,, \qquad v \ge 0.$$

$$(4.24)$$

Note that  $\kappa, \kappa' \in (0, 1)$  and they tend to 1 as  $b \to 0$  and  $B \to \infty$  (which means  $\mathbb{1}_{B_{r,t}} \to 1$ ). To prove the first inequality in Equation (4.22), we write

$$\widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}(f_\beta) \mathbb{1}_{B_{r,t}} \Big] = \int_0^\infty \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}([-x,0]) \mathbb{1}_{B_{r,t}} \Big] \beta \sqrt{2} \, \mathrm{e}^{-\beta \sqrt{2}x} \, \mathrm{d}x.$$

Then, we apply Lemma 4.10 for some  $\theta \in (0, 1)$  and with  $\varepsilon > 0$  introduced earlier to estimate the part  $x \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]$  of the integral and use Lemma 2.14 to bound the remaining part. Note also that  $\varphi_{b,B} \leq 1$  and  $\int_0^\infty \frac{x e^{-x^2/(2s)}}{\sqrt{2\pi s^{3/2}}} ds = 1$ . Therefore, for  $\beta > 1$  close enough to 1 (equivalently r large enough), there exists  $t_0 > r$  such that, for any  $t \geq t_0$ ,

$$\begin{split} & \left| \widetilde{\mathbb{E}}_{t} \Big[ \mathcal{C}_{t,r_{t}}^{*}(f_{\beta}) \mathbb{1}_{B_{r,t}} \Big] - C_{\star} \int_{0}^{\infty} \left( \int_{0}^{\infty} \varphi_{b,B} \Big( \frac{r \lor s}{|r-s|} \Big) \frac{x \mathrm{e}^{-x^{2}/(2s)}}{\sqrt{2\pi} s^{3/2}} \, \mathrm{d}s \Big) \beta \sqrt{2} \, \mathrm{e}^{-(\beta-1)\sqrt{2}x} \, \mathrm{d}x \right| \\ & \leq \varepsilon \int_{0}^{\infty} \beta \sqrt{2} \mathrm{e}^{-(\beta-1)\sqrt{2}x} \, \mathrm{d}x + C \int_{[\theta \sqrt{r}, \theta^{-1}\sqrt{r}]^{c}} \beta \sqrt{2} \, \mathrm{e}^{-(\beta-1)\sqrt{2}x} \, \mathrm{d}x \\ & = \frac{\beta}{\beta-1} \Big( \varepsilon + C \Big( 1 - \mathrm{e}^{-\theta\sqrt{a}} + \mathrm{e}^{-\theta^{-1}\sqrt{a}} \Big) \Big), \end{split}$$
(4.25)

recalling that  $r = a(\beta - 1)^{-2}$ . Choosing  $\theta$  small enough and considering  $\beta < 2$ , the right-hand side of Equation (4.25) is smaller that  $3\varepsilon/(\beta - 1)$ . Then, we rewrite the double integral on the left-hand side of Equation (4.25) by using Fubini's theorem and changing variables with  $u = (\beta - 1)\sqrt{2}x$  and  $w = s(\beta - 1)^2$ , which shows that this double integral equals  $\beta \kappa/(\beta - 1)$ . This proves the first inequality in Equation (4.22) (with  $4\varepsilon$  instead of  $\varepsilon$ ). The second inequality is proved by writing

$$\widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}(f_\beta) \mathbb{1}_{B_{r,t}} \Big] = -\int_0^\infty \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}([-x,0]) \mathbb{1}_{B_{r,t}} \Big] \sqrt{2} (\beta \sqrt{2}x - 1) \mathrm{e}^{-\beta \sqrt{2}x} \,\mathrm{d}x$$

and then proceeding similarly (note that  $\beta\sqrt{2}x - 1$  can be replaced by  $\beta\sqrt{2}x$  up to a negligible error as  $\beta \downarrow 1$ ).

Step 4: Conclusion. We first claim that for  $\theta \in (0,1)$  and  $\epsilon > 0$ , there exists  $r_0 > 0$  such that, for any  $r \ge r_0$ , there exists  $t_0 > 0$  such that, for any  $t \ge t_0$  and any  $v \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]$ ,

$$\left|\widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}([-v,0]) \Big] - C_\star \mathrm{e}^{\sqrt{2}v} \right| \le \varepsilon \, \mathrm{e}^{\sqrt{2}v}.$$

This is a slightly stronger version of [21, Lemma 5.2] where no uniformity for  $v \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]$ is stated: however, the aforementioned claim follows from their proof (more precisely, it follows from [21, Lemma 5.6] in the same way as Lemma 4.10 below follows from Lemma A.1). Then, we deduce from this claim that, for  $\beta > 1$  close enough to 1, up to a modification of  $t_0$ , we also have, for any  $t \ge t_0$ ,

$$\begin{split} \left| (\beta - 1) \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t, r_t}(f_\beta) \Big] - C_\star \right| &\leq \varepsilon, \\ \left| (\beta - 1)^2 \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t, r_t}(\partial_\beta f_\beta) \Big] + C_\star \right| &\leq \varepsilon, \end{split}$$

where these inequalities are obtained in the same way as Equations (4.22) have been obtained from Lemma 4.10 in Step 3. Combining this with Equations (4.17), (4.21) and (4.22), we get

$$\mathbb{E}\left[\frac{(S_{\beta}')^2}{S_{\beta}}\right] \geq \frac{1}{(\beta-1)^3} \left(\frac{(C_{\star}\kappa'-\varepsilon)^2}{C_{\star}\kappa+\varepsilon} + \frac{(C_{\star}(1-\kappa')-\varepsilon)^2}{C_{\star}(1-\kappa)+\varepsilon}\right).$$

Letting  $\varepsilon \to 0$ , this proves

$$\liminf_{\beta \downarrow 1} (\beta - 1)^3 \mathbb{E}\left[\frac{(S_{\beta}')^2}{S_{\beta}}\right] \ge C_{\star}\left(\frac{(\kappa')^2}{\kappa} + \frac{(1 - \kappa')^2}{1 - \kappa}\right).$$
(4.26)

By Lemma 4.11, we can choose a, b, B such that  $\kappa \neq \kappa'$  and, together with the fact that  $\kappa, \kappa' \in (0, 1)$ , this implies that the right-hand side of Equation (4.26) is larger than  $C_{\star}$ .

The proof of the following lemma is postponed to Subsection A.2.

**Lemma 4.10.** Let  $\theta \in (0,1)$ ,  $\varepsilon, a > 0$  and 0 < b < B. There exists  $r_0 > 0$  such that, for any  $r \ge r_0$ , there exists  $t_0 > 0$  such that, for any  $t \ge t_0$  and any  $v \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]$ ,

$$\left| \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t} ([-v,0]) \mathbbm{1}_{B_{r,t}} \Big] - C_\star \mathrm{e}^{\sqrt{2}v} \int_0^\infty \varphi_{b,B} \Big( \frac{r \lor s}{|r-s|} \Big) \frac{v \mathrm{e}^{-v^2/(2s)}}{\sqrt{2\pi} s^{3/2}} \, \mathrm{d}s \right| \le \varepsilon \mathrm{e}^{\sqrt{2}v}.$$

**Lemma 4.11.** Recall the definition of  $\kappa$  and  $\kappa'$  in Equation (4.23). There exist a > 0 and 0 < b < B such that  $\kappa \neq \kappa'$ .

*Proof.* We proceed by contradiction: assume that, for any a > 0 and 0 < b < B,  $\kappa = \kappa'$ . Then, differentiating the relation  $\kappa = \kappa'$  w.r.t. B, we get, for any a, B > 0, the following quantity is the same for k = 1 and k = 2:

$$\int_0^\infty \frac{a \vee w}{|a-w|} \exp\left(-\frac{B^2}{2} \frac{a \vee w}{|a-w|}\right) \left(\int_0^\infty u^k \mathrm{e}^{-u-u^2/(4w)} \,\mathrm{d}u\right) \frac{\mathrm{d}w}{w^{3/2}}.$$

On the other hand, a direct calculation shows that, for  $k \in \{1, 2\}$ ,

$$\int_0^\infty \left( \int_0^\infty u^k e^{-u - u^2/(4w)} \, \mathrm{d}u \right) \frac{\mathrm{d}w}{w^{3/2}} = 2\sqrt{\pi}$$

and therefore is the same for k = 1 and k = 2. Therefore, we deduce that the following quantity is the same for k = 1 and k = 2:

$$\int_0^\infty \left( \exp\left( -\frac{B^2}{2} \frac{a \vee w}{|a - w|} \right) - e^{-B^2/2} \right) \left( \int_0^\infty u^k e^{-u - u^2/(4w)} \, \mathrm{d}u \right) \frac{\mathrm{d}w}{w^{3/2}}.$$
 (4.27)

Our goal is now to study the behavior as  $a \to 0$  of this quantity, to find a contradiction. We decompose the main integral in Equation (4.27) into a part  $w \in [0, a)$  in which we change w to x = a/(a - w) and u to  $v = u(x/(a(x - 1)))^{1/2}$ , and a part  $w \in (a, \infty)$  in which we change w to x = w/(w - a) and u to  $v = u((x - 1)/(ax))^{1/2}$ . This yields that the following quantity is the same for k = 1 and k = 2:

$$a^{k/2} \int_{1}^{\infty} \left( x \mathrm{e}^{-xB^{2}/2} - \mathrm{e}^{-B^{2}/2} \right) \left( \frac{(x-1)^{k/2-1}}{x^{k/2+1}} \int_{0}^{\infty} v^{k} \mathrm{e}^{-v^{2}/4 - v(\frac{a(x-1)}{x})^{1/2}} \,\mathrm{d}v + \frac{x^{k/2-1}}{(x-1)^{k/2+1}} \int_{0}^{\infty} v^{k} \mathrm{e}^{-v^{2}/4 - v(\frac{ax}{x-1})^{1/2}} \,\mathrm{d}v \right) \,\mathrm{d}x.$$

$$(4.28)$$

For k = 1, both integrals w.r.t. v in (4.28) converge as  $a \to 0$  towards  $\int_0^\infty v e^{-v^2/4} dv = 2$  by dominated convergence. Hence, by dominated convergence again but in the integral w.r.t. x, (4.28) equals

$$2a^{1/2} \int_{1}^{\infty} \left( x e^{-xB^2/2} - e^{-B^2/2} \right) \left( \frac{(x-1)^{-1/2}}{x^{3/2}} + \frac{x^{-1/2}}{(x-1)^{3/2}} \right) dx + o(a^{1/2}),$$

as  $a \to 0$ . For k = 2, we cannot use the same argument: the second domination cannot be justified. Instead, we bound the first integral w.r.t. v by a constant and, for the second one, for some  $\varepsilon > 0$ , we write

$$\int_0^\infty v^2 \mathrm{e}^{-v^2/4 - v(\frac{ax}{x-1})^{1/2}} \,\mathrm{d}v = \left(\frac{x-1}{ax}\right)^{3\varepsilon} \int_0^\infty u^2 \mathrm{e}^{-u^2(\frac{x-1}{ax})^{2\varepsilon}/4 - u(\frac{ax}{x-1})^{1/2-\varepsilon}} \,\mathrm{d}u \le C \left(\frac{x-1}{ax}\right)^{3\varepsilon},$$

by bounding the last integral by  $\int_0^\infty u^2 e^{-u^2/4} du$  if  $(x-1)/ax \ge 1$  and by  $\int_0^\infty u^2 e^{-u} du$  otherwise. With  $\varepsilon < 1/6$ , this proves that (4.28) is a  $o(a^{1/2})$  for k = 2. Since (4.28) is the same for k = 1 and k = 2, this implies

$$\int_{1}^{\infty} \left( x e^{-xB^2/2} - e^{-B^2/2} \right) \left( \frac{(x-1)^{-1/2}}{x^{3/2}} + \frac{x^{-1/2}}{(x-1)^{3/2}} \right) dx = 0,$$
(4.29)

for any B > 0. But the left-hand side of Equation (4.29) tends to infinity as  $B \to \infty$ , so this is a contradiction and concludes the proof.

## A Proof of technical results on the decoration of the BBM

This section is dedicated to the proof of several technical results concerning the decoration of the branching Brownian motion, mostly based on ideas introduced in [21]. Therefore, we explain how to adapt their argument and use their notation without introducing it.

#### A.1 Uniform bounds for moments of level sets

*Proof of Lemma 2.14.* The bound on the second moment is a direct consequence of Lemma A.6, so we focus here on the first moment. However, it could also be deduced from the proof of [21, Lemma 5.3], in a similar way as what is done below for the first moment.

To bound uniformly  $\widetilde{\mathbb{E}}_t[\mathcal{C}^*_{t,r_t}([-v,0])]$ , we follow the proof of [21, Lemma 5.2], which establishes an asymptotic equivalent for the first moment of  $\mathcal{C}^*_{t,r_t}([-v,0])$  as  $t \to \infty$  and then  $v \to \infty$ . The proof begins by writing

$$\widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}([-v,0]) \Big] = \frac{\widetilde{\mathbb{E}} \Big[ \mathcal{C}^*_{t,r_t}([-v,0]) \mathbb{1}_{\{\max_{x \in L_t} h_t(x) \le m_t\}} \Big| h_t(X_t) = m_t \Big]}{\widetilde{\mathbb{P}}(\max_{x \in L_t} h_t(x) \le m_t \mid h_t(X_t) = m_t)},$$
(A.1)

where the denominator satisfies, for some constant  $C_1 > 0$ , as  $t \to \infty$ ,

$$\widetilde{\mathbb{P}}\left(\max_{x\in L_t} h_t(x) \le m_t \ \middle| \ h_t(X_t) = m_t\right) \sim \frac{C_1}{t},\tag{A.2}$$

by [21, Lemmata 3.1 and 3.4] (the constant  $C_1$  equals  $2f^{(0)}(0)g(0)$  with notation of [21, Lemma 3.4]). In particular, there exists a constant c > 0 such that the probability in Equation (A.2) is at least c/t for any  $t \ge 1$ . Therefore, it remains to prove that there exist C > 0 and  $t_0 \ge 0$  such that, for any  $t \ge t_0$  and  $v \ge 0$ ,

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*([-v,0])\mathbb{1}_{\{\max_{x\in L_t}h_t(x)\leq m_t\}} \mid h_t(X_t) = m_t\Big] \leq \frac{C}{t} e^{\sqrt{2}v}.$$
(A.3)

Indeed, for  $t \leq t_0$ , bounding the indicator function by 1 and  $C^*_{t,r_t}([-v,0]) \leq \#L_{t_0}$ , the left-hand side of Equation (A.3) is at most

$$\widetilde{\mathbb{E}}[\#L_{t_0} \mid h_t(X_t) = m_t] = \widetilde{\mathbb{E}}[\#L_{t_0}] = \int_0^{t_0} e^{t_0 - s} \cdot 2 \, \mathrm{d}s = C(t_0), \tag{A.4}$$

where the first equality follows from the fact that displacement of the spine and branching of the BBM are independent and the second equality uses the facts that the spine branches at rate 2, giving birth to standard BBMs and that a standard BBM has in mean  $e^r$  particles at time r.

We now focus on proving Equation (A.3). Applying successively [21, Lemmata 5.4 and 5.5] (note that  $j_{t,v}(s) = j_{t,v}^{\geq 0}(s)$ ), we get for any t such that  $r_t \leq t/2$  and any  $v \geq 0$ ,

$$\begin{split} \widetilde{\mathbb{E}}\Big[\mathcal{C}^*_{t,r_t}([-v,0])\mathbbm{1}_{\{\max_{x\in L_t}h_t(x)\leq m_t\}} \ \Big| \ h_t(X_t) = m_t\Big] &= 2\int_0^{r_t} j_{t,v}(s) \,\mathrm{d}s \\ &\leq \frac{C}{t}(v+1) \,\mathrm{e}^{\sqrt{2}v} \int_0^\infty \frac{\mathrm{e}^{-v^2/(16s)} + \mathrm{e}^{-v/2}}{(s+1)\sqrt{s}} \,\mathrm{d}s. \end{split}$$

This last integral is smaller than  $C((v+1)^{-1} + e^{-v/2})$  for any  $v \ge 0$  (see [21, Equations (5.44)-(5.45)] for details) and so we get Equation (A.3).

#### A.2 First moment of level sets on a particular event

We prove in this subsection Lemma 4.10. This is a (non-trivial) refinement of the proof of [21, Lemma 5.2].

Proof of Lemma 4.10. Recalling the definition of  $\widetilde{\mathbb{P}}_t$  in Equation (2.6), we first have

$$\widetilde{\mathbb{E}}_{t}\Big[\mathcal{C}_{t,r_{t}}^{*}([-v,0])\mathbb{1}_{B_{r,t}}\Big] = \frac{\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_{t}}^{*}([-v,0])\mathbb{1}_{B_{r,t}}\mathbb{1}_{\{\max_{x\in L_{t}}h_{t}(x)\leq m_{t}\}} \mid h_{t}(X_{t})=m_{t}\Big]}{\widetilde{\mathbb{P}}(\max_{x\in L_{t}}h_{t}(x)\leq m_{t}\mid h_{t}(X_{t})=m_{t})}.$$
(A.5)

Recall from Equation (A.2) that the denominator in Equation (A.5) is asymptotically equivalent to  $C_1/t$ , so we now focus on the numerator.

We introduce the event

$$\mathcal{B}_{r,t} \coloneqq \Big\{ \widehat{W}_{t,r} \in [-B\sqrt{r}, -b\sqrt{r}] \Big\},\,$$

which is analog to  $B_{r,t}$  (see Equation (4.20)) but for the process  $(\widehat{W}_{t,s})_{s\in[0,t]}$  defined in [21, Equation (3.1)] as

$$\widehat{W}_{t,s} \coloneqq W_s - \gamma_{t,s}, \quad \text{with } \gamma_{t,s} \coloneqq \frac{3}{2\sqrt{2}} \left( \log_+ s - \frac{s}{t} \log_+ t \right), \tag{A.6}$$

for any  $0 \le s \le t$ , with W a standard Brownian motion under  $\mathbb{P}$ . Defining, for  $v \ge 0$  and  $0 \le s, r \le t$ ,

$$j_{t,v,r}(s) \coloneqq \mathbb{E}\Big[J_{t,v}(s)\mathbb{1}_{\mathcal{B}_{r,t}}\Big|\widehat{W}_{t,0} = \widehat{W}_{t,t} = 0\Big]$$

where  $J_{t,v}(s)$  is introduced in [21, Equation (5.9)], it follows from the proof of [21, Lemma 5.4], that

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*([-v,0])\mathbb{1}_{B_{r,t}}\mathbb{1}_{\{\max_{x\in L_t}h_t(x)\leq m_t\}} \mid h_t(X_t) = m_t\Big] = 2\int_0^{r_t} j_{t,v,r}(s)\,\mathrm{d}s.$$
(A.7)

As in [21, Equation (5.17)], for any  $M \ge 0$ , we split  $j_{t,v,r}(s)$  into

$$\begin{aligned} j_{t,v,r}^{$$

We postpone the estimate of  $j_{t,v,r}^{\leq M}(s)$  to Lemma A.1 (which replace Lemma 5.6 in [21]). With this lemma in hand, we can conclude the proof. Let  $\delta \in (0, \theta \wedge \frac{1}{2})$  and M > 0. Considering t large enough such that  $\delta^{-1}r \leq r_t$ , we decompose the right-hand side of Equation (A.7) as

$$2\int_{[\delta r,\delta^{-1}r]} j_{t,v,r}^{\leq M}(s) \,\mathrm{d}s + 2\int_{[\delta r,\delta^{-1}r]} j_{t,v,r}^{\geq M}(s) \,\mathrm{d}s + 2\int_{[\delta r,\delta^{-1}r]^c} j_{t,v,r}(s) \,\mathrm{d}s,$$

where the two last terms are negligible. Indeed, it is proved in [21, Equation (5.41)] (and the paragraph around) that

$$\limsup_{\eta \to 0} \limsup_{M \to \infty} \limsup_{v \to \infty} \limsup_{t \to \infty} \frac{t}{\mathrm{e}^{\sqrt{2}v}} \cdot \left( 2 \int_{[\eta v^2, \eta^{-1}v^2]} j_{t,v}^{\geq M}(s) \,\mathrm{d}s + 2 \int_{[\eta v^2, \eta^{-1}v^2]^c} j_{t,v}(s) \,\mathrm{d}s \right) = 0,$$

and therefore

 $\limsup_{\delta \to 0} \limsup_{M \to \infty} \limsup_{r \to \infty} \limsup_{t \to \infty} \sup_{v \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]} \left( 2 \int_{[\delta r, \delta^{-1} r]} j_{t,v,r}^{\geq M}(s) \, \mathrm{d}s + 2 \int_{[\delta r, \delta^{-1} r]^c} j_{t,v,r}(s) \, \mathrm{d}s \right) = 0.$ (A.8)

Now, setting  $S_{\delta} := [\delta r, (1 - \delta)r] \cup [(1 + \delta)r, \delta^{-1}r]$ , we have

$$2\int_{[\delta r,\delta^{-1}r]} j_{t,v,r}^{< M}(s) \,\mathrm{d}s = 2\int_{S_{\delta}} j_{t,v,r}^{< M}(s) \,\mathrm{d}s + 2\int_{(1-\delta)r}^{(1+\delta)r} j_{t,v,r}^{< M}(s) \,\mathrm{d}s.$$

This last integral is also negligible for  $\delta$  small enough (in the same sense as in Equation (A.8)), because

$$j_{t,v,r}^{\leq M}(s) \leq j_{t,v}(s) = j_{t,v}^{\geq 0}(s) \leq C \, \frac{(v+1)\mathrm{e}^{\sqrt{2}v}}{s^{3/2}t} \leq C(\theta) \, \frac{\mathrm{e}^{\sqrt{2}v}}{rt} \,,$$

for any M > 0,  $0 \le r \le t/4$ ,  $s \in [(1 - \delta)r, (1 + \delta)r]$  and  $v \in [\theta\sqrt{r}, \theta^{-1}\sqrt{r}]$  by [21, Lemma 5.5]. Finally, using Lemma A.1, we have, as  $t \to \infty$ , then  $r \to \infty$  and then  $M \to \infty$ , uniformly in  $v \in [\theta\sqrt{r}, \theta^{-1}\sqrt{r}]$ ,

$$2\int_{S_{\delta}} j_{t,v,r}^{\leq M}(s) \,\mathrm{d}s \sim \frac{2C_2}{t} v \mathrm{e}^{\sqrt{2}v} \int_{S_{\delta}} \frac{\mathrm{e}^{-v^2/(2s)}}{s^{3/2}} \varphi_{b,B}\left(\frac{r \vee s}{|r-s|}\right) \mathrm{d}s.$$

Coming back to Equation (A.5) and letting  $\delta \to 0$ , this proves that there exists  $r_0 > 0$  such that, for any  $r \ge r_0$ , there exists  $t_0 > 0$  such that, for any  $t \ge t_0$  and any  $v \in [\theta \sqrt{r}, \theta^{-1} \sqrt{r}]$ ,

$$\left| \widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t} ([-v,0]) \mathbb{1}_{B_{r,t}} \Big] - \frac{2C_2}{C_1} v \mathrm{e}^{\sqrt{2}v} \int_0^\infty \varphi_{b,B} \left( \frac{r \vee s}{|r-s|} \right) \frac{\mathrm{e}^{-v^2/(2s)}}{s^{3/2}} \,\mathrm{d}s \right| \le \varepsilon \,\mathrm{e}^{\sqrt{2}v}. \tag{A.9}$$

Using the relation<sup>6</sup>  $C_{\star} = 2\sqrt{2\pi} C_2/C_1$  gives the desired result.

**Lemma A.1.** Let a > 0, 0 < b < B and  $\delta \in (0, 1/2)$ . As  $t \to \infty$ , then  $r \to \infty$  and then  $M \to \infty$ , we have

$$j_{t,v,r}^{< M}(s) \sim \frac{C_2}{ts^{3/2}} v e^{\sqrt{2}v - v^2/(2s)} \varphi_{b,B}\left(\frac{r \lor s}{|r-s|}\right),$$

uniformly  $s \in [\delta r, (1-\delta)r] \cup [(1+\delta)r, \delta^{-1}r]$  and  $v \in [\delta\sqrt{r}, \delta^{-1}\sqrt{r}]$ , with

$$C_2 := \frac{2\sqrt{2}}{\pi} f^{(0)}(0)g(0) \int_{\mathbb{R}} f(z)g(z)^2 e^{\sqrt{2}z} \, \mathrm{d}z \in (0,\infty),$$

where  $f^{(0)}$ , f, g are positive functions introduced in [21, Lemma 3.4].

<sup>&</sup>lt;sup>6</sup>The authors of [21] do not keep precisely track of constants but this relation can be deduced from a careful reading of their paper. Alternatively letting  $b \to 0$  and  $B \to \infty$ , we have  $\mathbb{1}_{B_{r,t}} \to 1$  and  $\varphi_{b,B} \to 1$  and Equation (A.9) gives an alternative proof of their [21, Lemma 5.2], showing that the constant C appearing there equals  $2\sqrt{2\pi}C_2/C_1$ . Then, a quick look at the proof of [21, Proposition 1.5] ensures that this constant C is actually  $C_*$ .

*Proof.* We follow ideas from the proof of [21, Lemma 5.6], but instead of only distinguishing according to the value of  $\widehat{W}_{t,s}$ , we also distinguish according to the value of  $\widehat{W}_{t,r}$ . For comparison, the constant C appearing in the statement of [21, Lemma 5.6] equals  $C_2$  introduced here.

We start with the case  $r \leq s$ , that is  $s \in [(1 + \delta)r, \delta^{-1}r]$ . Then, we have, with the notations from [21],

$$j_{t,v,r}^{$$

The function  $(y, z) \mapsto p_t((r, y); (s, z))$  is the density of  $(\widehat{W}_{t,r}, \widehat{W}_{t,s})$  given  $\widehat{W}_{t,0} = \widehat{W}_{t,t} = 0$ . It is explicitly given by (recall the definition of  $\gamma_{t,s}$  in Equation (A.6))

$$p_t((r,y);(s,z)) = \frac{1}{2\pi} \sqrt{\frac{t}{r(s-r)(t-s)}} \exp\left(-\frac{(s(y+\gamma_{t,r}) - r(z+\gamma_{t,s}))^2}{2rs(s-r)} - \frac{t(z+\gamma_{t,s})^2}{2s(t-s)}\right) \\ \sim \frac{1}{2\pi\sqrt{r(s-r)}} \exp\left(-\frac{sy^2}{2r(s-r)}\right),$$

as  $t \to \infty$  and then  $r \to \infty$ , uniformly in  $s \in [(1 + \delta)r, \delta^{-1}r]$ ,  $y \in [-B\sqrt{r}, -b\sqrt{r}]$  and  $z \in [-M, M]$ . Furthermore, it follows from [21, Lemma 3.4] that

$$q_t((0,0);(r,y)) = q_r((0,0);(r,y)) \sim \frac{2(-y)f^{(0)}(0)}{r}$$
$$q_t((r,y);(s,z)) = q_s((r,y);(s,z)) \sim \frac{2(-y)g(z)}{s-r},$$
$$q_t((s,z);(t,0)) \sim \frac{2f^{(s)}(z)g(0)}{t-s} \sim \frac{2f(z)g(0)}{t},$$

as  $t \to \infty$  and then  $r \to \infty$ , uniformly in s, y and z as before. Finally, it follows from [21, Lemma 4.2] (see also [21, Equation (5.37)]) that

$$e_{s,v}(z) \sim v \mathrm{e}^{\sqrt{2}v - v^2/(2s)} \frac{g(z)}{\sqrt{\pi}} \mathrm{e}^{\sqrt{2}z},$$

as  $r \to \infty$ , uniformly in  $v \in [\delta \sqrt{r}, \delta^{-1} \sqrt{r}]$  and s, z as before. Altogether, this proves

$$j_{t,v,r}^{$$

as  $t \to \infty$  and then  $r \to \infty$ , uniformly in s and v as before. A change of variable in the first integral and letting  $M \to \infty$  in the second integral (the fact that this integral converges to a finite limit is justified at the end of the proof of [21, Lemma 5.6]) yields the result.

The case  $s \leq r$ , that is  $s \in [\delta r, (1 - \delta)r]$  is similar: we write

$$j_{t,v,r}^{$$

and use the same asymptotics as before, the main difference being

$$p_t((s,z);(r,y)) = \frac{1}{2\pi} \sqrt{\frac{t}{s(r-s)(t-r)}} \exp\left(-\frac{(r(z+\gamma_{t,s})-s(y+\gamma_{t,r}))^2}{2rs(r-s)} - \frac{t(y+\gamma_{t,r})^2}{2r(t-r)}\right) \\ \sim \frac{1}{2\pi\sqrt{s(r-s)}} \exp\left(-\frac{y^2}{2(s-r)}\right).$$

This yields the result in that case.

#### A.3 Cross-moments of level sets

Our aim in this section is to prove Proposition 2.13. For this, we follow the proof of [21, Proposition 1.5], which bounds  $\mathbb{E}[\mathcal{C}([-v,0])^2]$ . This proof is based on a series of five lemmas, that we re-state here in a new version tuned for our purpose of dealing with two level sets of different levels v and v'.

In the following lemma, as in [21, Lemma 4.3], we work with a 2-spine BBM defined under the probability measure  $\tilde{\mathbb{P}}^{(2)}$  as follows. It starts with one particle at 0 at time 0 which is part of the spines 1 and 2. Particles belonging to m spines branch at rate  $2^m$  and move according to a standard Brownian motion. At a branching point, for each  $k \in \{1, 2\}$ , if the parent was part of spine k, then one of both children is chosen uniformly at random to be part of spine k. We denote  $X_t(k)$  the particle at time t which is part of the spine k. y For  $x, y \in L_t$ , we write d(x, y) = r if the most recent common ancestor of x and y died at time t - r.

**Lemma A.2.** There exists C > 0 such that, for any  $0 \le r \le t$  and  $v \le v' \le u$ ,

$$\widetilde{\mathbb{P}}^{(2)}\left(\widehat{h}_t(X_t(1)) \ge v, \, \widehat{h}_t(X_t(2)) \ge v', \, \widehat{h}_t^* \le u \mid d(X_t(1), X_t(2)) = r\right)$$

$$\le \frac{C \mathrm{e}^{-t-r}}{1 + (r \wedge (t-r))^{3/2}} (u_+ + 1) \mathrm{e}^{\sqrt{2}u} (u - v + 1) (u - v' + 1) \mathrm{e}^{-\sqrt{2}(v+v')} \left(\mathrm{e}^{-(u-v)^2/(4t)} + \mathrm{e}^{-(u-v)/2}\right).$$

*Proof.* This is a new version of [21, Lemma 4.3] and we explain how to adapt its proof. Similarly as [21, Equation (4.15)], the probability in the statement equals

$$\int_{\mathbb{R}} \widetilde{\mathbb{P}} \Big( \widehat{h}_r(X_r) \ge v - z, \widehat{h}_r^* \le u - z \Big) \widetilde{\mathbb{P}} \Big( \widehat{h}_r(X_r) \ge v - z, \widehat{h}_r^* \le u - z \Big) \\ \times \widetilde{\mathbb{P}} \Big( \widehat{h}_{t-r}(X_{t-r}) - m_{t,r} \in \mathrm{d}z, \widehat{h}_t^*(\mathrm{B}(X_t)^c) \le u \Big).$$
(A.10)

Following [21], we split the integral according to  $z \leq u$  and z > u.

For z > u, using [21, Equation (4.2)] for the first one and [21, Equation (4.3)] for the second one, we bound the product of the two first probabilities in Equation (A.10) by

$$Ce^{-2r}(u-v+1)(u-v'+1)e^{-\sqrt{2}(v+v')}e^{2\sqrt{2}z-\frac{3}{2}(z-u)}\left(e^{-(v-z)^2/(4t)}+e^{(v-z)/2}\right).$$

This is exactly the same as [21, Equation (4.20)] up to the factor  $(u - v' + 1)e^{-\sqrt{2}v'}$  where primes have been added. Therefore, this part of the integral is dealt with exactly the same way as in [21].

For  $z \leq u$ , we use [21, Equation (4.2)] for both first probabilities in Equation (A.10) and note that, because  $v \leq v'$ ,

$$\left( e^{-(v-z)^2/(4t)} + e^{(v-z)/2} \right) \left( e^{-(v'-z)^2/(4t)} + e^{(v'-z)/2} \right) \le \left( e^{-(v-z)^2/(4t)} + e^{(v-z)/2} \right) \left( 1 + e^{(v-z)/2} \right)$$
$$\le \left( e^{-(v-z)^2/(4t)} + e^{v-z} + 2e^{(v-z)/2} \right).$$

This shows the product of the two first probabilities in Equation (A.10) is at most

$$Ce^{-2r}(u-v+1)(u-v'+1)e^{-\sqrt{2}(v+v')}e^{2\sqrt{2}z}(u-z+1)^2\left(e^{-(v-z)^2/(4t)}+e^{(v-z)/2}+e^{(v-z)/2}\right).$$

There are two differences with [21, Equation (4.19)]: the factor  $(u - v' + 1)e^{-\sqrt{2}v'}$  where primes have been added (but this adds no new difficulty), and the additional term  $e^{(v-z)/2}$  in the last

parentheses. This latter gives rise to the following new term, which should be added to the integral in [21, Equation (4.21)],

$$\int_{-\infty}^{u} e^{\frac{v}{2} + (\sqrt{2} - \frac{1}{2})z)} (u - z + 1)^3 dz \le C e^{\sqrt{2}u} e^{-(u - v)/2},$$

which can be included in the upper bound of the statement of the lemma after taking care of the other factors (see [21, Equation (4.22)]: this is exactly how another part of the integral in [21, Equation (4.21)] is bounded).  $\Box$ 

**Lemma A.3.** There exists C > 0 such that, for any  $t \ge 0$  and  $v \le v' \le u$ ,

$$\mathbb{E}\Big[\mathcal{E}_t([v,\infty))\mathcal{E}_t([v',\infty)); \hat{h}_t^* \le u\Big] \\ \le C(u_++1)\mathrm{e}^{\sqrt{2}u}(u-v+1)(u-v'+1)\mathrm{e}^{-\sqrt{2}(v+v')}\Big(\mathrm{e}^{-(u-v)^2/(4t)}+\mathrm{e}^{-(u-v)/2}\Big).$$

*Proof.* This follows from Lemma A.2 in the same way as [21, Lemma 4.4] follows from [21, Lemma 4.3].  $\Box$ 

Lemma A.4. For any  $v, v' \ge 0$ ,

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}^*_{t,r_t}([-v,0]) \,\mathcal{C}^*_{t,r_t}([-v',0])\,;\, \widehat{h}^*_t \le 0 \mid \widehat{h}_t(X_t) = 0\Big] \\ = 4 \int_0^{r_t} \int_0^{r_t} j_{t,v,v'}(s,s') \,\mathrm{d}s' \,\mathrm{d}s + 2 \int_0^{r_t} j_{t,v,v'}(s,s) \,\mathrm{d}s,$$

where  $j_{t,v,v'}(s,s') \coloneqq \widehat{\mathbb{E}}_t[J_{t,v}(s)J_{t,v'}(s')].$ 

*Proof.* This follows directly from the proof of [21, Lemma 5.4].

**Lemma A.5.** There exists C > 0 such that, for any  $t \ge 0$ ,  $s, s' \in [0, t/2]$  and  $v \ge v' \ge 0$ ,

$$j_{t,v,v'}(s,s') \le \frac{C(v+1)(v'+1)\mathrm{e}^{\sqrt{2}(v+v')}}{t(s\wedge s'+1)\sqrt{s\wedge s'}(|s'-s|+1)\sqrt{|s'-s|} + \mathbb{1}_{s=s'}} \Big(\mathrm{e}^{-v^2/(16s)} + \mathrm{e}^{-v/4}\Big).$$

*Proof.* This is a new version of [21, Lemma 5.5].

For the case  $s \neq s'$ , we first assume that s < s' but do not assume  $v \geq v'$ . Then it follows directly from the proof in [21] that

$$j_{t,v,v'}(s,s') \le \frac{C(v+1)(v'+1)\mathrm{e}^{\sqrt{2}(v+v')}}{t(s+1)\sqrt{s}(s'-s+1)\sqrt{s'-s+1}\mathrm{I}_{s=s'}} \Big(\mathrm{e}^{-v^2/(16s)} + \mathrm{e}^{-v/4}\Big) \Big(\mathrm{e}^{-(v')^2/(16s')} + \mathrm{e}^{-v'/4}\Big).$$

Then, we use  $j_{t,v,v'}(s,s') = j_{t,v',v}(s',s)$  to cover the case s > s' (this is fine because we removed the assumption  $v \ge v'$ ). Finally, for  $v \ge v'$ , we bound  $e^{-(v')^2/(16s')} + e^{-v'/4} \le 2$ . This yields the desired result.

The case s = s' is also identical to the proof of [21, Lemma 5.5], applying here Lemma A.3 instead of [21, Lemma 4.4].

-	-	-	-	

**Lemma A.6.** There exists C > 0 such that, for any  $t \ge 1$  and  $v \ge v' \ge 0$ ,

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*([-v,0])\mathcal{C}_{t,r_t}^*([-v',0]) \ \Big| \ \widehat{h}_t^* = \widehat{h}_t(X_t) = 0\Big] \le C(v'+1)\mathrm{e}^{-\sqrt{2}(v+v')}.$$

*Proof.* Proceeding as in Equations (A.1), (A.2), (A.3) and (A.4)<sup>7</sup>, it is enough to prove that there exist C > 0 and  $t_0 \ge 0$  such that, for any  $t \ge t_0$  and  $v \ge 0$ ,

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*([-v,0])\mathcal{C}_{t,r_t}^*([-v',0])\mathbb{1}_{\widehat{h}_t^* \le 0} \ \Big| \ \widehat{h}_t(X_t) = 0\Big] \le \frac{C}{t}(v'+1)\mathrm{e}^{\sqrt{2}v}.$$
(A.11)

We choose  $t_0$  such that, for any  $t \ge t_0$ ,  $r_t \le t/2$ . Using Lemmas A.4 and A.5, the left-hand side of Equation (A.11) is at most

$$\begin{split} \frac{C}{t}(v+1)(v'+1)\mathrm{e}^{\sqrt{2}(v+v')} & \left(\int_{0}^{r_{t}}\int_{0}^{r_{t}}\frac{(\mathrm{e}^{-v^{2}/(16s)}+\mathrm{e}^{-v/4})}{(s\wedge s'+1)\sqrt{s\wedge s'}(|s'-s|+1)\sqrt{|s'-s|}}\,\mathrm{d}s'\,\mathrm{d}s \right. \\ & \left.+\int_{0}^{r_{t}}\frac{(\mathrm{e}^{-v^{2}/(16s)}+\mathrm{e}^{-v/4})}{(s+1)\sqrt{s}}\,\mathrm{d}s\right) \\ & \leq \frac{C}{t}(v+1)(v'+1)\mathrm{e}^{\sqrt{2}(v+v')}\int_{0}^{r_{t}}\frac{(\mathrm{e}^{-v^{2}/(16s)}+\mathrm{e}^{-v/4})}{(s+1)\sqrt{s}}\,\mathrm{d}s, \end{split}$$

by integrating w.r.t. s' first. Then, proceeding as in the proof of [21, Lemma 5.3], this last integral is at most C/(v+1) and this yields the result.

*Proof of Proposition 2.13.* This follows from Lemma A.6 in the same way as [21, Proposition 1.5] follows from [21, Lemma 5.3].  $\Box$ 

#### A.4 Small moments of $S_{\beta}$

**Lemma A.7.** For any  $\gamma \in (0,1)$ , there exists  $C = C(\gamma) > 0$  and  $t_0 \ge 0$ , such that, for any  $t \ge t_0$  and any  $\beta \in (1,2]$ ,

$$\widetilde{\mathbb{E}}_t \Big[ \mathcal{C}^*_{t,r_t}(f_\beta)^\gamma \Big] \le \begin{cases} C(\beta-1)^{1-2\gamma} & \text{if } \gamma \in (1/2,1), \\ C\log\frac{1}{\beta-1} & \text{if } \gamma = 1/2, \\ C & \text{if } \gamma \in (0,1/2), \end{cases}$$

where  $f_{\beta} \colon x \in \mathbb{R} \mapsto e^{\beta \sqrt{2}x}$ .

*Proof.* Proceeding as in Equations (A.1) and (A.2), it is enough to prove that there exist C > 0 such that, for any  $t \ge t_0$  and  $\beta \in (1, 2]$ ,

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*(f_{\beta})^{\gamma}\mathbb{1}_{\{\widehat{h}_t^* \le 0\}} \ \Big| \ \widehat{h}_t(X_t) = 0\Big] \le \frac{C}{t} \times \begin{cases} (\beta - 1)^{1-2\gamma} & \text{if } \gamma \in (1/2, 1), \\ \log \frac{1}{\beta - 1} & \text{if } \gamma = 1/2, \\ 1 & \text{if } \gamma \in (0, 1/2), \end{cases}$$
(A.12)

where  $t_0$  is chosen such that, for any  $t \ge t_0$ ,  $r_t \le t/2$ .

<sup>&</sup>lt;sup>7</sup>For this step note that we end up here with  $\mathbb{E}[(\#L_{t_0})^2]$ , which is also a finite constant depending on  $t_0$  using similar arguments, including the fact that the second moment of the number of particles at time r in a standard BBM is finite (more precisely it equals  $2e^{2r} - e^r$ ).

We first decompose  $C_{t,r_t}^*(f_\beta)$  along the spine as it is done for level sets in [21, Lemma 5.4]. Setting (this replaces  $J_{t,v}(s)$  defined in [21, Equation (5.9)])

$$K_{t,\beta}(s) \coloneqq \mathcal{E}_s^s(f_{\beta}(\cdot + \widehat{W}_{t,s})) \times \mathbb{1}_{\{\widehat{h}_s^{s*} \le -\widehat{W}_{t,s}\}} \times \mathbb{1}_{\mathcal{A}_t}$$

and  $\widehat{\mathbb{P}}_t \coloneqq \mathbb{P}(\,\cdot\,|\widehat{W}_{t,0} = \widehat{W}_{t,t} = 0)$ , we have

$$\mathcal{C}_{t,r_t}^*(f_{\beta})\mathbb{1}_{\{\widehat{h}_t^* \le 0\}} \text{ under } \widetilde{\mathbb{P}}(\cdot | \widehat{h}_t(X_t) = 0) \stackrel{\text{(d)}}{=} \int_0^{r_t} K_{t,\beta}(s) \mathcal{N}(\mathrm{d}s) \text{ under } \widehat{\mathbb{P}}_t,$$

where  $\mathcal{N}$  is a Poisson point process on  $\mathbb{R}_+$  with intensity 2 ds. Using subadditivity of  $x \mapsto x^{\gamma}$  (note that the integral above is actually a finite sum) and then proceeding as in the proof of [21, Lemma 5.4], we get

$$\widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_t}^*(f_\beta)^{\gamma}\mathbb{1}_{\{\widehat{h}_t^* \le 0\}} \ \Big| \ \widehat{h}_t(X_t) = 0\Big] \le \widehat{\mathbb{E}}_t\Big[\int_0^{r_t} K_{t,\beta}(s)^{\gamma} \mathcal{N}(\mathrm{d}s)\Big] = 2\int_0^{r_t} k_{t,\beta,\gamma}(s) \,\mathrm{d}s, \qquad (A.13)$$

with  $k_{t,\beta,\gamma}(s) \coloneqq \widehat{\mathbb{E}}_t[K_{t,v}(s)^{\gamma}].$ 

We now aim at proving that there exists C > 0 such that, for any  $t \ge 0$ ,  $s \in [0, t/2]$  and  $\beta \in (1, 2]$ ,

$$k_{t,\beta,\gamma}(s) \le \frac{C}{t(s+1)\sqrt{s}} \Big( (s+1) \wedge (\beta-1)^{-2} \Big)^{\gamma}.$$
 (A.14)

For this, we follow the ideas from the proof of [21, Lemma 5.5] (in the case M = 0). Conditioning on  $\widehat{W}_{t,s}$ , we get, similarly as [21, Equation (5.20)],

$$k_{t,\beta,\gamma}(s) = \int_{\mathbb{R}} q_t((0,0);(s,z)) \mathbb{E} \Big[ \mathcal{E}_s(f_\beta(\cdot+z))^{\gamma} \mathbb{1}_{\{\widehat{h}_s^* \le -z\}} \Big] q_t((s,z);(t,0)) p_t(s,z) \, \mathrm{d}z.$$
(A.15)

The single difference with [21] is inside the expectation. Using Jensen's inequality and then writing  $f_{\beta}(x+z) = \int_0^\infty \beta \sqrt{2} e^{-\beta \sqrt{2}v} \mathbb{1}_{\{x \ge -v-z\}} dv$  for  $x \le -z$ , we get

$$\mathbb{E}\Big[\mathcal{E}_{s}(f_{\beta}(\cdot+z))^{\gamma}\mathbb{1}_{\{\widehat{h}_{s}^{*}\leq-z\}}\Big]^{1/\gamma} \leq \mathbb{E}\Big[\mathcal{E}_{s}(f_{\beta}(\cdot+z))\mathbb{1}_{\{\widehat{h}_{s}^{*}\leq-z\}}\Big]$$
$$=\int_{0}^{\infty}\beta\sqrt{2}\mathrm{e}^{-\beta\sqrt{2}v}\mathbb{E}\Big[\mathcal{E}_{s}([-v,0]-z)\mathbb{1}_{\{\widehat{h}_{s}^{*}\leq-z\}}\Big]\,\mathrm{d}v.$$
(A.16)

Using [21, Lemma 4.2], the right-hand side of Equation (A.16) is at most

$$C(z_{-}+1)\mathrm{e}^{\sqrt{2}z} \int_{0}^{\infty} (v+1)\mathrm{e}^{-(\beta-1)\sqrt{2}v} \left(\mathrm{e}^{-(v+z)^{2}/4s} + \mathrm{e}^{-(v+z)/2}\right) \mathrm{d}v$$

The part of the last integral due to the term  $e^{-(v+z)/2}$  is bounded by  $Ce^{-z/2}$ . For the other part, it can be bounded by  $\int_0^\infty (v+1)e^{-(\beta-1)\sqrt{2}v} dv \le C(\beta-1)^{-2}$  or by  $\int_{\mathbb{R}} (v+1)e^{-(v+z)^2/4s} dv \le C(|z|+1)(s+1)$ . Therefore, we proved

$$\mathbb{E}\Big[\mathcal{E}_s(f_{\beta}(\cdot+z))^{\gamma}\mathbb{1}_{\{\widehat{h}_s^*\leq -z\}}\Big] \leq C(|z|+1)^{2\gamma} \mathrm{e}^{\gamma\sqrt{2}z} \Big(\mathrm{e}^{-z/2} + \Big((s+1)\wedge(\beta-1)^{-2}\Big)\Big)^{\gamma}.$$

Coming back to Equation (A.15) and bounding the other factors in the integrand as in [21, Equations (5.23) and (5.24)], we get

$$k_{t,\beta,\gamma}(s) \le \frac{C}{t(s+1)\sqrt{s}} \int_{\mathbb{R}} \left( z_{-} + e^{-\frac{3}{2}z_{+}} \right) (|z|+1)^{2\gamma+1} e^{\gamma\sqrt{2}z} \left( e^{-\frac{\gamma}{2}z} + \left( (s+1) \wedge (\beta-1)^{-2} \right)^{\gamma} \right) \mathrm{d}z,$$

and Equation (A.14) follows.

Finally, we come back to Equation (A.13) and apply Equation (A.14) (we use here  $r_t \leq t/2$ ) to get

$$\begin{split} \widetilde{\mathbb{E}}\Big[\mathcal{C}_{t,r_{t}}^{*}(f_{\beta})^{\gamma}\mathbb{1}_{\widehat{h}_{t}^{*}\leq0} \ \Big| \ \widehat{h}_{t}(X_{t}) &= 0\Big] \leq \frac{C}{t} \int_{0}^{\infty} \frac{1}{(s+1)\sqrt{s}} \Big((s+1) \wedge (\beta-1)^{-2}\Big)^{\gamma} \,\mathrm{d}s \\ &\leq \frac{C}{t} \left(\int_{0}^{(\beta-1)^{-2}} \frac{\mathrm{d}s}{(s+1)^{1-\gamma}\sqrt{s}} + (\beta-1)^{-2\gamma} \int_{(\beta-1)^{-2}}^{\infty} \frac{\mathrm{d}s}{(s+1)\sqrt{s}}\right) \end{split}$$

and Equation (A.12) follows.

# Acknowledgements

The authors warmly thank Bernard Derrida for very stimulating discussions and the GdR Branchement for financial support. Michel Pain also acknowledges support from the PEPS JCJC grant n°246644.

## References

- E. Aïdékon, J. Berestycki, É. Brunet, and Z. Shi. Branching Brownian motion seen from its tip. *Probab. Theory Related Fields*, 157(1-2):405–451, 2013.
- [2] L.-P. Arguin. Extrema of Log-correlated Random Variables: Principles and Examples, pages 166–204. Cambridge University Press, 2016.
- [3] L.-P. Arguin, A. Bovier, and N. Kistler. The extremal process of branching Brownian motion. *Probab. Theory Related Fields*, 157(3-4):535–574, 2013.
- [4] L.-P. Arguin and O. Zindy. Poisson-Dirichlet statistics for the extremes of the twodimensional discrete Gaussian free field. *Electron. J. Probab.*, 20:no. 59, 19, 2015.
- [5] M. Biskup. Extrema of the two-dimensional discrete Gaussian free field. In Random graphs, phase transitions, and the Gaussian free field, volume 304 of Springer Proc. Math. Stat., pages 163–407. Springer, Cham, [2020] ©2020.
- [6] M. Biskup and O. Louidor. Extreme local extrema of two-dimensional discrete Gaussian free field. Comm. Math. Phys., 345(1):271–304, 2016.
- [7] M. Biskup and O. Louidor. Full extremal process, cluster law and freezing for the twodimensional discrete Gaussian Free Field. Adv. Math., 330:589–687, 2018.
- [8] E. Bolthausen. Random media and spin glasses: an introduction into some mathematical results and problems. In *Spin glasses*, volume 1900 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 2007.
- [9] E. Bolthausen, J.-D. Deuschel, and G. Giacomin. Entropic repulsion and the maximum of the two-dimensional harmonic crystal. Ann. Probab., 29(4):1670–1692, 2001.
- [10] E. Bolthausen, J. D. Deuschel, and O. Zeitouni. Recursions and tightness for the maximum of the discrete two dimensional Gaussian free field. *Electron. Commun. Probab.*, 16:114–119, 2011.
- [11] E. Bolthausen and A.-S. Sznitman. Ten lectures on random media, volume 32 of DMV Seminar. Birkhäuser Verlag, Basel, 2002.
- [12] B. Bonnefont. The overlap distribution at two temperatures for the branching Brownian motion. *Electron. J. Probab.*, 27:Paper No. 116, 21, 2022.
- [13] A. Bovier and I. Kurkova. Derrida's generalized random energy models. II. Models with continuous hierarchies. Ann. Inst. H. Poincaré Probab. Statist., 40(4):481–495, 2004.

- [14] M. Bramson. Convergence of solutions of the Kolmogorov equation to travelling waves. Mem. Amer. Math. Soc., 44(285):iv+190, 1983.
- [15] M. Bramson, J. Ding, and O. Zeitouni. Convergence in law of the maximum of the twodimensional discrete Gaussian free field. Comm. Pure Appl. Math., 69(1):62–123, 2016.
- [16] M. Bramson and O. Zeitouni. Tightness of the recentered maximum of the two-dimensional discrete Gaussian free field. Comm. Pure Appl. Math., 65(1):1–20, 2012.
- [17] M. D. Bramson. Maximal displacement of branching Brownian motion. Comm. Pure Appl. Math., 31(5):531–581, 1978.
- [18] D. Carpentier and P. Le Doussal. Glass transition of a particle in a random potential, front selection in nonlinear renormalization group, and entropic phenomena in Liouville and sinh-Gordon models. *Phys. Rev. E*, 63:026110, Jan 2001.
- [19] B. Chauvin and A. Rouault. KPP equation and supercritical branching Brownian motion in the subcritical speed area. Application to spatial trees. *Probab. Theory Related Fields*, 80(2):299–314, 1988.
- [20] B. Chauvin and A. Rouault. Supercritical branching Brownian motion and K-P-P equation in the critical speed-area. *Math. Nachr.*, 149:41–59, 1990.
- [21] A. Cortines, L. Hartung, and O. Louidor. The structure of extreme level sets in branching Brownian motion. Ann. Probab., 47(4):2257 – 2302, 2019.
- [22] A. Cortines, L. Hartung, and O. Louidor. More on the structure of extreme level sets in branching Brownian motion. *Electron. Commun. Probab.*, 26:Paper No. 2, 14, 2021.
- [23] O. Daviaud. Extremes of the discrete two-dimensional Gaussian free field. Ann. Probab., 34(3):962–986, 2006.
- [24] B. Derrida. Random-energy model: an exactly solvable model of disordered systems. Phys. Rev. B (3), 24(5):2613–2626, 1981.
- [25] B. Derrida. A generalization of the random energy model which includes correlations between energies. J. Physique Lett., 46(9):401–407, 1985.
- [26] B. Derrida and P. Mottishaw. One step replica symmetry breaking and overlaps between two temperatures. Journal of Physics A: Mathematical and Theoretical, 54(4):045002, jan 2021.
- [27] B. Derrida and H. Spohn. Polymers on disordered trees, spin glasses, and traveling waves. J. Statist. Phys., 51(5-6):817–840, 1988. New directions in statistical mechanics (Santa Barbara, CA, 1987).
- [28] D. S. Fisher and D. A. Huse. Directed paths in a random potential. Phys. Rev., B 43:10728, 1991.
- [29] Y. V. Fyodorov and J.-P. Bouchaud. Freezing and extreme-value statistics in a random energy model with logarithmically correlated potential. J. Phys. A, 41(37):372001, 12, 2008.
- [30] Y. V. Fyodorov and J.-P. Bouchaud. Statistical mechanics of a single particle in a multiscale random potential: Parisi landscapes in finite-dimensional Euclidean spaces. J. Phys. A, 41(32):324009, 25, 2008.
- [31] Y. V. Fyodorov, P. Le Doussal, and A. Rosso. Statistical mechanics of logarithmic REM: duality, freezing and extreme value statistics of 1/f noises generated by Gaussian free fields. J. Stat. Mech. Theory Exp., (10):P10005, 32, 2009.
- [32] N. Kistler. Derrida's random energy models. From spin glasses to the extremes of correlated random fields. In *Correlated random systems: five different methods*, volume 2143 of *Lecture Notes in Math.*, pages 71–120. Springer, Cham, 2015.
- [33] I. Kurkova. Temperature dependence of the Gibbs state in the random energy model. J. Statist. Phys., 111(1-2):35–56, 2003.

- [34] S. P. Lalley and T. Sellke. A conditional limit theorem for the frontier of a branching Brownian motion. Ann. Probab., 15(3):1052–1061, 1987.
- [35] G. Last and M. Penrose. Lectures on the Poisson Process. Institute of Mathematical Statistics Textbooks. Cambridge University Press, 2017.
- [36] A. D. Le, A. H. Mueller, and S. Munier. Probabilistic picture for particle number densities in stretched tips of the branching Brownian motion. *Europhysics Letters*, 140(5):51003, 2022.
- [37] P. Maillard and M. Pain. 1-stable fluctuations in branching Brownian motion at critical temperature I: The derivative martingale. Ann. Probab., 47(5):2953–3002, 2019.
- [38] P. Maillard and M. Pain. 1-stable fluctuations in branching Brownian motion at critical temperature II: general functionals. 2021. arXiv:2103.10412.
- [39] H. P. McKean. Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov. Comm. Pure Appl. Math., 28(3):323–331, 1975.
- [40] A. H. Mueller and S. Munier. Particle-number distribution in large fluctuations at the tip of branching random walks. *Phys. Rev. E*, 102(2):Paper No. 022104, 15, 2020.
- [41] L. Mytnik, J.-M. Roquejoffre, and L. Ryzhik. Fisher-KPP equation with small data and the extremal process of branching Brownian motion. Adv. Math., 396:Paper No. 108106, 58, 2022.
- [42] M. Pain and O. Zindy. Two-temperatures overlap distribution for the 2D discrete Gaussian free field. Ann. Inst. H. Poincaré Probab. Statist., 57(2):685 – 699, 2021.
- [43] D. Panchenko and M. Talagrand. On one property of Derrida-Ruelle cascades. C. R. Math. Acad. Sci. Paris, 345(11):653–656, 2007.
- [44] R. Rhodes and V. Vargas. Gaussian multiplicative chaos and applications: a review. Probab. Surv., 11:315–392, 2014.
- [45] M. Sales and J.-P. Bouchaud. Rejuvenation in the random energy model. Europhys. Lett., 56(2):181–186, 2001.
- [46] E. Subag and O. Zeitouni. Freezing and decorated Poisson point processes. Comm. Math. Phys., 337(1):55–92, 2015.